

Central Lancashire Online Knowledge (CLoK)

Title	Treatment of urinary incontinence after stroke in adults
Type	Article
URL	https://clock.uclan.ac.uk/1109/
DOI	##doi##
Date	2009
Citation	Thomas, Lois H orcid iconORCID: 0000-0001-5218-6546, Cross, Stephen, Barrett, James, French, Beverley, Leathley, Michael, Sutton, Chris J orcid iconORCID: 0000-0002-6406-1318 and Watkins, Caroline orcid iconORCID: 0000-0002-9403-3772 (2009) Treatment of urinary incontinence after stroke in adults. Cochrane Database of Systematic reviews, 1 (-). pp. 1-43. ISSN 1469-493X
Creators	Thomas, Lois H, Cross, Stephen, Barrett, James, French, Beverley, Leathley, Michael, Sutton, Chris J and Watkins, Caroline

It is advisable to refer to the publisher's version if you intend to cite from the work. ##doi##

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Treatment of urinary incontinence after stroke in adults (Review)

Thomas LH, Cross S, Barrett J, French B, Leathley M, Sutton CJ, Watkins C



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2009, Issue 1

<http://www.thecochranelibrary.com>



TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	2
OBJECTIVES	3
METHODS	3
RESULTS	5
DISCUSSION	11
AUTHORS' CONCLUSIONS	12
ACKNOWLEDGEMENTS	13
REFERENCES	13
CHARACTERISTICS OF STUDIES	15
DATA AND ANALYSES	24
Analysis 1.1. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 1 Number of people with incontinence after treatment.	26
Analysis 1.2. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 2 Severity of incontinence: mean number of incontinence episodes in 24 hours.	27
Analysis 1.3. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 3 Urinary symptoms - overall.	28
Analysis 1.4. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 4 Urinary symptoms - frequency (continuous variables).	29
Analysis 1.5. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 5 Urinary symptoms - frequency (dichotomous variables).	29
Analysis 1.6. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 6 Urinary symptoms - urgency.	30
Analysis 1.7. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 7 Urinary symptoms - nocturia (continuous variables).	31
Analysis 1.8. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 8 Urinary symptoms - nocturia (dichotomous variables).	31
Analysis 1.9. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 9 Urological measures: mean function of the pelvic floor muscle.	32
Analysis 1.10. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 10 Health status: mean total score SF36.	33
Analysis 1.11. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 11 Quality of life.	33
Analysis 1.12. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 12 Function: mean Barthel score (continuous variables).	34
Analysis 1.13. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 13 Function: people independent in transfer from wheelchair to toilet (dichotomous variables).	34
Analysis 1.14. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 14 Patient satisfaction.	35
Analysis 1.15. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 15 Cost/service use.	35
Analysis 3.1. Comparison 3 INTERVENTION VERSUS PLACEBO, Outcome 1 Severity of incontinence: mean number of incontinence episodes per week.	36
Analysis 5.1. Comparison 5 COMBINED INTERVENTION VERSUS SINGLE INTERVENTION, Outcome 1 Number of people with incontinence after treatment.	37
Analysis 5.2. Comparison 5 COMBINED INTERVENTION VERSUS SINGLE INTERVENTION, Outcome 2 Severity of incontinence: mean number of incontinence episodes.	37
APPENDICES	37
WHAT'S NEW	39

HISTORY	39
CONTRIBUTIONS OF AUTHORS	39
DECLARATIONS OF INTEREST	40
SOURCES OF SUPPORT	40
INDEX TERMS	40

[Intervention Review]

Treatment of urinary incontinence after stroke in adults

Lois H Thomas¹, Stephen Cross², James Barrett³, Beverley French¹, Michael Leathley¹, Christopher J Sutton⁴, Caroline Watkins¹

¹Department of Nursing, University of Central Lancashire, Preston, UK. ²Stroke Rehabilitation Unit, The Lakes Care centre, Dunkinfield, UK. ³Clatterbridge Hospital, Wirral, UK. ⁴Lancashire School of Health and Postgraduate Medicine, University of Central Lancashire, Preston, UK

Contact address: Lois H Thomas, Department of Nursing, University of Central Lancashire, Room 434, Brook Building, Preston, Lancashire, PR1 2HE, UK. lhthomas@uclan.ac.uk.

Editorial group: Cochrane Incontinence Group.

Publication status and date: Edited (no change to conclusions), published in Issue 1, 2009.

Review content assessed as up-to-date: 13 November 2007.

Citation: Thomas LH, Cross S, Barrett J, French B, Leathley M, Sutton CJ, Watkins C. Treatment of urinary incontinence after stroke in adults. *Cochrane Database of Systematic Reviews* 2008, Issue 1. Art. No.: CD004462. DOI: 10.1002/14651858.CD004462.pub3.

Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Urinary incontinence can affect 40-60% of people admitted to hospital after a stroke, with 25% still having problems on hospital discharge and 15% remaining incontinent at one year.

Objectives

To determine the optimal methods for treatment of urinary incontinence after stroke in adults.

Search methods

We searched the Cochrane Incontinence and Stroke Groups specialised registers (searched 15 March 2007 and 5 March 2007 respectively), CINAHL (January 1982 to January 2007), national and international trial databases for unpublished data, and the reference lists of relevant articles.

Selection criteria

Randomised or quasi-randomised controlled trials evaluating the effects of interventions designed to promote continence in people after stroke.

Data collection and analysis

Data extraction and quality assessment were undertaken by two reviewers working independently. Disagreements were resolved by a third reviewer.

Main results

Twelve trials with a total of 724 participants were included in the review. Participants were from a mixture of settings, age groups and phases of stroke recovery.

Behavioural interventions

Three trials assessed behavioural interventions, such as timed voiding and pelvic floor muscle training. All had small sample sizes and confidence intervals were wide.

Specialised professional input interventions

Two trials assessed variants of professional input interventions. Results tended to favour the intervention groups: in a small trial in early rehabilitation, fewer people had incontinence at discharge from hospital after structured assessment and management than in a control group (1/21 vs. 10/13; RR 0.06, 95% CI 0.01 to 0.43); in the second trial, assessment and management by Continence Nurse Advisors was associated with fewer participants having urinary symptoms (48/89 vs. 38/54; RR 0.77, 95% CI 0.59 to 0.99) and statistically significantly more being satisfied with care.

Complementary therapy interventions

Three small trials all reported fewer participants with incontinence after acupuncture therapy (overall RR 0.44; 95% 0.23 to 0.86), but there were particular concerns about study quality.

Pharmacotherapy and hormonal interventions

There were three small trials that included groups allocated meclfenoxate, oxybutinin or oestrogen. There were no apparent differences other than in the trial of meclfenoxate where fewer participants had urinary symptoms in the active group than in the control group (9/40 vs. 27/40; RR 0.33, 95% CI 0.18 to 0.62).

Authors' conclusions

Data from the available trials are insufficient to guide continence care of adults after stroke. However, there was suggestive evidence that professional input through structured assessment and management of care and specialist continence nursing may reduce urinary incontinence and related symptoms after stroke. Better quality evidence is required of the range of interventions that have been suggested for continence care after stroke.

PLAIN LANGUAGE SUMMARY

Treatment of urinary incontinence after stroke in adults

Urinary incontinence is a common consequence of stroke and has many causes. In early stroke rehabilitation, structured assessment and management of care shows promise in reducing the number of people with urinary incontinence. In the later phases of stroke recovery the use of specialist advisors may be helpful in reducing symptoms associated with urinary incontinence. Even late after stroke, interventions targeted at specific causes of incontinence may be helpful. Unfortunately, all the conclusions were limited by a lack of robust information.

BACKGROUND

Urinary incontinence is defined as the complaint of any involuntary leakage of urine (Abrams 2002). 40-60% of people admitted to hospital after a stroke can have problems with urinary incontinence, with 25% of stroke survivors still having problems on hospital discharge, and 15% remaining incontinent after one year (Barrett 2001). Not all the studies reviewed by Barrett 2001 excluded people with pre-morbid incontinence, however, so figures presented may include old as well as new cases. Addressing problems with continence whilst the person is still in hospital may prevent long-term problems for the patient and family.

The more severe the stroke, the greater is the likelihood of urinary incontinence (Burney 1996a). Other risk factors for urinary in-

continence include older age, female sex, speech difficulties, motor weakness, visual field defects or cognitive impairment (Barrett 2001). Problems experienced may range from urinary retention to complete incontinence. The most likely pattern of incontinence is urinary frequency, urgency (a sudden compelling desire to pass urine which is difficult to defer) and urge incontinence (involuntary leakage) (Marinkovic 2001). This is generally the result of detrusor overactivity (Arunabh 1993), although this may depend on the site of the stroke lesion (Burney 1996b).

Damage to the frontal lobe, the area believed to be responsible for control of micturition, has been identified in several studies as associated with urinary dysfunction after stroke. However, evidence

suggests that the size of the lesion, rather than its location, is more likely to predict urinary incontinence (Brittain 1999). It is unclear whether incontinence is a direct (i.e. site of brain lesion) or indirect (e.g. functional impairment preventing access) consequence of stroke. Other non-neurological factors contributing to urinary incontinence, including pre-morbid continence state, sphincter incompetence and polyuria (Barrett 2001), are regarded as prevalent in the population of patients with stroke (Brittain 1998b).

Because of their severity, the symptoms of urinary incontinence are reported to have more of an effect on the lives of stroke survivors, when compared with other groups of people with incontinence (Brittain 2000b). Urinary symptoms had more impact on sleep, daily activities, quality of life, physical discomfort, social life and relationships. Incontinence is not just a physical problem, but impacts on what people can do and how they feel. Depression is twice as common in stroke survivors who are incontinent (Brittain 1998a). Urinary incontinence is distressing for both those affected and their carers (Williams 1993).

Continuing incontinence is associated with poor outcome in both stroke survivor and carer (Nakayama 1997). Conversely, stroke outcome is better in those stroke survivors who remain continent or regain continence (Barer 1989). Improvement is common over time (Marinkovic 2001), which suggests that problems with continence may be transient in some stroke survivors, and/or amenable to intervention. Factors predicting early improvements in continence status are less impairment on admission, and the site of the stroke lesion (Ween 1996). Factors associated with poor recovery of continence include stroke type and being aged 75 or over (Patel 2001).

Incontinence is a strong predictor of stroke functional outcome (Meijer 2003). While there are problems with attributing better stroke outcome to improvements in continence, it is possible that recovery from incontinence may improve morale and self esteem and therefore speed overall stroke recovery (Barer 1989; Patel 2001). While differences in incontinence rates between centres will reflect difference in the case mix of individuals and the methods of reporting (Barrett 2001), they might also indicate variations in the processes of continence assessment and management.

It is known that early rehabilitation intervention in stroke results in better outcome overall (Cifu 1999). Some of the studies on which this conclusion is based have included incontinence as a measure of functional outcome. For example, a trial of a multidisciplinary rehabilitation intervention aiming to improve functional independence after stroke showed a positive impact on incontinence rates (Wikander 1998a). Other management techniques include biofeedback, pelvic floor muscle training, electrical stimulation, drug treatments, surgical interventions and mechanical devices.

Current guidelines for the management of urinary incontinence recommend an assessment to guide management (Thuroff 1999).

This begins with physical assessment and history-taking including identification of urological problems before the stroke occurred, such as bladder outlet obstruction in men or stress incontinence in women. The choice of method to promote continence will then depend on the individual's history and type of incontinence. Bladder training and/or anticholinergic drugs may be appropriate for urge incontinence, while problems with retention may require intermittent catheterisation. Alternatively, problems with memory loss or restriction of movement may benefit more from toileting assistance programmes such as prompted or timed voiding or habit retraining (Eustice 2000; Ostaszkiwicz 2004a; Ostaszkiwicz 2004b; Roe 2007).

A systematic review of methodologically robust studies - randomised controlled trials - is needed to identify interventions to promote continence that are effective in the stroke population. Trials that have evaluated different management strategies also need to be reviewed in relation to subgroups of stroke patients with specific patterns of incontinence, because the effects of management are likely to depend on the type of stroke that the person has suffered and the urinary problems they experienced.

OBJECTIVES

The objective of this review was to determine the optimal methods for the treatment of urinary continence after stroke in adults.

The following hypotheses were addressed:

1. intervention is more effective than no intervention;
2. intervention is more effective than placebo;
3. a specific intervention is more effective in comparison with another intervention;
4. combined interventions are more effective than single interventions.

Within each hypothesis, interventions were considered within four categories:

1. behavioural interventions
2. specialised professional input interventions
3. complementary therapy interventions
4. pharmacotherapy interventions.

METHODS

Criteria for considering studies for this review

Types of studies

All randomised and quasi-randomised trials evaluating the effects of interventions designed to promote continence in people who have had a stroke. Quasi-random methods include: allocation by the person's date of birth, by the day of the week or month of the year, by a person's medical record number, or just allocating every alternate person.

Types of participants

Adults (i.e. 18 years of age and over) with a diagnosis of stroke, including people with incontinence who have had a stroke identified as a subgroup within a larger group for whom relevant data are reported.

Types of interventions

One arm of the study must include an intervention designed to promote urinary continence. Trials evaluating any of the following were included in the review:

- pharmacotherapy e.g. anticholinergics, adrenergics, hormonal treatment;
- physical therapy e.g. electrical stimulation, biofeedback;
- physical aids e.g. catheters, pads, pessaries, other appliances;
- behavioural interventions e.g. prompted or scheduled voiding, bladder training, habit retraining (i.e. identification of voiding pattern and development of an individualised toileting schedule), pelvic floor muscle training or other behavioural management programmes;
 - environmental or lifestyle interventions e.g. voiding position, diet and fluid management, alternative communication devices;
 - specialised professional input interventions e.g. provision of information or education, assessment schedules, generic multidisciplinary rehabilitation programmes, Continence Advisors, home-support programmes, nurse practitioners;
 - complementary interventions e.g. homeopathy, acupuncture.

Trials relating solely to surgical or physical interventions for pre-existing continence problems that are not associated with stroke (e.g. transurethral resection of the prostate) were excluded, unless it was a co-intervention in a wider trial testing an included method of continence promotion. Trials relating to urological diagnosis, or to the management of incontinence or retention of urine in the acute phase of stroke were also excluded. The acute phase was defined as up to one month post stroke. Studies were excluded if solely in the acute phase because the aim of intervention is commonly to contain incontinence, monitor urine output and prevent adverse events, rather than promote continence.

Types of outcome measures

The primary outcome of interest was in/continence, measured by:

1. Participant symptoms

- Number of participants regaining continence
- Number of incontinent episodes over 24 hours (indicated by bladder charts, total and mean number of episodes)
- Severity of incontinence e.g. index score
- Perception of improvement or cure in continence (as reported by participant or caregiver)

2. Physical measures

- Pad tests of quantified leakage
- Volume of urine loss
- Total and mean number of pads used

3. Secondary outcomes

- a) Symptom scores or participant /carer report of other urinary symptoms including frequency, urgency, dysuria, polyuria, nocturia, discomfort, pain
- b) Physical measures e.g. post-void retention of urine, time to voiding onset, void volume, urodynamic measures
- c) Health status or measures of psychological health
Impact of incontinence e.g. Incontinence Impact Questionnaire (IIQ), Short Form 36 Health Survey Questionnaire (SF36), functional ability, knowledge, satisfaction, quality of life
- d) Economic outcomes
Impact of continence promotion interventions on cost or service use
- e) Other outcomes
Any other outcomes subsequently deemed appropriate to the review.

Search methods for identification of studies

The review used the search strategies developed for both the Cochrane Incontinence Group and the Cochrane Stroke Group. Relevant trials were initially identified from the Groups' specialised registers of controlled trials described under the groups' details in *The Cochrane Library* ([For more details please see the 'Specialized Register' section of the Incontinence Group's module in The Cochrane Library](#)) ([For details of the Stroke Group's Specialised Register please see the 'Specialized Register' section of their module](#)). The dates of the most recent searches of the registers for this review update were conducted on 15 March 2007 (Stroke Register) and 5 March 2007 (Incontinence Register).

To search the Stroke Group register the following search terms were used: “incontinence (urine/faecal) or (nursing and bladder care) or urinary retention or urinary tract infection”.

The terms used to search the Incontinence Group trials register are given below:

```
(({Topic.urine.incon.Stroke.} OR
{TOPIC.URINE.INCON.UNKNOWN.STROKE.}
OR {TOPIC.URINE.INCON.NEUROGENIC.} (in keywords
field))
OR
({stroke\*} OR {cerebrovascular\*} (in title1 field))
OR
({TOPIC.URINE.NEUROGENIC.} OR
{TOPIC.URINE.NEUROGENIC.stroke.} (in keywords field)))
AND
({DESIGN.CCT*} OR {DESIGN.RCT*} (in keywords field))
```

(All searches were in Reference Manager 9.5 N, ISI ResearchSoft).

The following extra specific searches were performed for this review. For more details, including the search terms used please see [Appendix 1](#).

We did not impose any language or other restrictions on any of these searches.

Electronic searches

Due to the comprehensive nature of the searches already performed by the Cochrane Stroke Group, no additional searches were performed other than on CINAHL, combining stroke terms with terms for urinary incontinence without a research methods filter. The search was combined with the CINAHL search from the Cochrane Stroke Group. This was done because of the potentially poor indexing of nursing research. The CINAHL search covered the years January 1982 to January 2007 (on Dialog Datarstar). More details, including search terms are given in [Appendix 1](#).

Recent unpublished trial data were also searched for on national and international databases i.e. NHS National Research Register, NHS Research Findings Register, US Community of Science NIH Grants, MetaRegister of Controlled Clinical Trials and CRISP, by adapting terms drawn from the existing search strategies of the Incontinence and Stroke Review Groups.

Searching other resources

The reference lists of all relevant reviews and trial reports were searched to identify further relevant studies. Major investigators were contacted to ask for any other possible relevant trials, published or unpublished. In addition, contact was made with the authors of other relevant Cochrane reviews to ascertain whether defined subgroups of stroke survivors were identified in trials testing methods of promoting continence in a general population.

The review was published on the following websites: Joanna Briggs Institute, Royal College of Nursing Research Society, Royal Col-

lege of Nursing Continence Interest Group, Association of Continence Advisors, Sigma Theta Tau.

Data collection and analysis

Trials were considered for inclusion independently by two reviewers (LT, BF). Studies were excluded from the review if they were not randomised or quasi-randomised, or if on more detailed examination, they did not meet the review inclusion criteria. These studies are listed in the 'Characteristics of Excluded Studies Table'. Data extraction forms, based on the protocol, were piloted and checked for the coding of intervention, outcome and quality assessment. Data extraction and review of the methodological quality of the eligible studies was independently conducted by two reviewers for each study (SC, BF, ML, LT) using the quality assessment tool described by the Cochrane Incontinence Review Group. Extracted data and quality assessment were cross checked and any disagreements discussed and if necessary resolved by a third reviewer (LT).

Attempts were made to obtain missing data, as well as data collected but not reported, by contacting trialists.

Outcomes are reported as unfavourable events. Planned subgroup analyses for the effect of urological diagnosis (i.e. detrusor over-activity versus other) and time from stroke onset (1 to 6 months, greater than 7 months) were not possible as the data were not available in the original studies. Included trial data were processed as described in the Cochrane Authors' Handbook ([Deeks 2006](#)), and analysed using the statistical analysis package RevMan Analyses. Effect estimates for continuous outcomes were summarised using weighted mean difference (WMD) and dichotomous outcomes were summarised using relative risk (RR). Where deemed appropriate, effects were summarised across studies using fixed or random effects meta-analysis techniques appropriate to the form of the data. Random effects meta-analysis (DerSimonian and Laird method) was used if the studies showed heterogeneity (defined by the studies' effects having an I-squared statistic of greater than 50%), otherwise a fixed effect analysis (Mantel-Haenszel for dichotomous and inverse variance for continuous data) was used. For continuous outcome data, if change from baseline data were available, these were used; otherwise the raw outcome data were used.

RESULTS

Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

Included/excluded studies

The search identified 1234 papers and 126 studies were retrieved. The search initially identified ten potential trials evaluating the effectiveness of methods to promote continence in adults after stroke (Brittain 2000a; Chu 1997; Cook 1998; Gelber 1997a; Gelber 1997b; Gross 1990; Judge 1969; Lewis 1990; Tekeoglu 1998; Wikander 1998b).

One trial was published in Chinese, one in Korean, and the rest in English. Information from the Chinese trial was translated and extracted by nursing lecturers from the Shanghai Military Medical University. On further examination, three trials (Cook 1998; Gross 1990; Tekeoglu 1998) were excluded for reasons listed in the 'Characteristics of Excluded Studies Table'.

The review update identified a further five studies (Liu 2006; Tibaek 2005; Zhang 2002; Zhou 1999; Zhu 2003). With the exception of Tibaek 2005, all were published in Chinese and were translated by a native speaking Chinese nursing student undertaking postgraduate study in England.

Twelve trials were therefore included in the review update: seven were full reports (Brittain 2000a; Chu 1997; Judge 1969; Liu 2006; Tibaek 2005; Wikander 1998b; Zhu 2003), although one Chinese trial (Chu 1997) was only briefly described in a one page report; the remaining five trials were reported only in conference abstracts (Gelber 1997a; Gelber 1997b; Lewis 1990; Zhang 2002; Zhou 1999). Two of the trials (Gelber 1997a; Gelber 1997b) were in parts of the same report.

Description of interventions

The trials tested the following interventions:

Behavioural interventions

- timed voiding versus void on request (Gelber 1997a);
- oxybutynin versus timed voiding (Gelber 1997b);
- pelvic floor muscle training versus usual care (Tibaek 2005);
- sensory-motor biofeedback device (Uristop) combined with timed voiding against timed voiding alone (Lewis 1990).

Specialised professional input interventions:

- care from a Continence Nurse Advisor (CNP) versus usual care provided by the general practitioner (GP) (Brittain 2000a);
- a special intervention programme based on assessment using the Functional Independence Measure (FIM) versus usual rehabilitation care (Wikander 1998b).

Complementary therapy interventions:

- scalp acupuncture versus no scalp acupuncture (Chu 1997);
- eye and scalp acupuncture versus no acupuncture (Zhou 1999);
- acupuncture versus usual care (Zhang 2002);
- ginger-salt-partitioned moxibustion (involving filling the navel with salt, adding a piece of ginger and a taper and setting the taper alight) plus routine acupuncture versus routine acupuncture (Liu 2006).

Pharmacotherapy interventions:

- oestrogen versus placebo (Judge 1969);
- meclofenoxate (designed to improve glucose utilisation of brain cells) plus salvia miltirrhiza versus salvia miltirrhiza (Zhu 2003);
- oxybutynin versus timed voiding (Gelber 1997b);

Physical therapy interventions:

- sensory-motor biofeedback device (Uristop) combined with timed voiding against timed voiding alone (Lewis 1990).

Intervention versus no intervention / usual care

Nine trials tested an intervention versus no intervention or usual care (Brittain 2000a; Chu 1997; Gelber 1997a; Liu 2006; Tibaek 2005; Wikander 1998b; Zhang 2002; Zhou 1999; Zhu 2003).

Behavioural interventions

One trial (Gelber 1997a) tested a behavioural intervention comprising timed voiding versus void on request for participants with normal urodynamic studies. Normal urodynamic studies was not defined and no further details are given of the intervention. One trial (Tibaek 2005) tested an intensive pelvic floor muscle training programme, comprising individual and group exercises and feedback to participants, compared against normal rehabilitation with no specific treatment of urinary incontinence.

Specialised professional input interventions

Two trials (Brittain 2000a; Wikander 1998b) tested specialised professional input interventions. Brittain 2000a compared care given by a Continence Nurse Advisor against usual care provided by a general practitioner and existing specialised services for the management of continence. Wikander 1998b compared the impact of a special intervention programme based on assessment using the Functional Independence Measure against usual rehabilitation care based on the Bobath method.

Complementary therapy interventions

Four trials tested complementary interventions. Chu 1997 tested scalp acupuncture plus usual care, compared against usual care with no scalp acupuncture; Zhang 2002 tested acupuncture against general treatment and Zhou 1999 tested eye and electriferous scalp acupuncture plus medication therapy against medication therapy only. Liu 2006 tested ginger-salt-partitioned moxibustion plus routine acupuncture against routine acupuncture.

Pharmacotherapy interventions

One trial (Zhu 2003) tested meclofenoxate plus salvia miltirrhiza against salvia miltirrhiza alone.

Intervention versus placebo

One cross-over trial (Judge 1969) tested a pharmacotherapy intervention (oestrogen) against placebo.

Specific intervention versus another intervention

One trial (Gelber 1997b) tested a specific intervention against another intervention, comparing the anticholinergic oxybutynin against timed voiding in participants with bladder hyperreflexia.

Combined intervention versus single intervention

One trial (Lewis 1990) tested a combined intervention against a single intervention. The study tested a sensory-motor biofeedback device combined with timed voiding, against timed voiding alone.

Participants

A total of 724 participants were enrolled in the trials. The numbers in individual trials ranged from 13 to 242. Two trials (Judge 1969; Tibaek 2005) enrolled only female participants. Three trials (Gelber 1997a; Gelber 1997b; Lewis 1990) gave no details of age or gender. The remaining trials (Brittain 2000a; Chu 1997; Liu 2006; Wikander 1998b; Zhang 2002; Zhou 1999; Zhu 2003) ranged from 40% to 61% males. The mean age of the study participants overall cannot be reported, but the highest mean age (82) was in the trial by Judge (Judge 1969), with the widest age range of 40-96 in the Brittain trial (Brittain 2000a).

Diagnosis

Three trials (Liu 2006; Tibaek 2005; Wikander 1998b) only included participants who were continent prior to the stroke. Nine trials (Brittain 2000a; Chu 1997; Gelber 1997a; Gelber 1997b; Judge 1969; Lewis 1990; Zhang 2002; Zhou 1999; Zhu 2003) did not specify whether urinary incontinence was subsequent to the occurrence of stroke, although Lewis 1990 described participants as having "post-stroke urinary urge incontinence".

Diagnostic criteria for incontinence were given in only two trials (Brittain 2000a; Tibaek 2005). Four trials specified a urological diagnosis, given as normal urodynamic studies (Gelber 1997a), bladder hyperreflexia (Gelber 1997b), urge incontinence (Lewis 1990) and urge, stress and mixed stress/urge incontinence (Tibaek 2005). Judge 1969 reported data for two groups of participants defined as mildly or severely incontinent. Liu 2006 reported data for three groups of participants, classified according to the Barthel continence item: completely incontinent, partially incontinent and independent. Chu 1997 included participants who had "urinary frequency or urinary incontinence" but did not define urinary frequency further. Two trials (Brittain 2000a; Wikander 1998b) did not include a urological diagnosis for participants, or group them by type of incontinence.

The trial by Tibaek 2005 stated that participants must be diagnosed according to the World Health Organisation definition of ischaemic stroke verified by CT scan. Three trials (Gelber 1997a; Gelber 1997b; Wikander 1998b) reported that strokes were unilateral, and one trial (Chu 1997) reported the stroke type as multi-

focal infarction. Liu 2006 and Zhou 1999 included participants with cerebral infarction and haemorrhage. Two trials (Zhang 2002; Zhu 2003) included only participants with infarction, and the site of infarction is specified for all participants in the trial by Zhang 2002. Judge 1969 included participants with cerebrovascular accident or "multiple little strokes". Brittain 2000a included participants who self-reported that they had had a stroke, subarachnoid haemorrhage or transient ischaemic attack on a screening questionnaire. One trial (Wikander 1998b) included side of stroke lesion in baseline comparison between groups.

It was not possible in all but one of the trials (Tibaek 2005) to determine whether only participants with a first stroke were included. Participants in two trials (Wikander 1998b; Zhu 2003) were in the acute phase of stroke (i.e. 0 to 1 month) on enrolment to the study, but in one study (Wikander 1998b) the study period extended into early rehabilitation, i.e. up to three months. Participants in the trial by Liu 2006 were 70.74 ± 35.26 days post-stroke, while Zhou 1999 presented findings for participants who were less than or more than three months post-stroke. Participants in three trials (Brittain 2000a; Judge 1969; Tibaek 2005) were also less likely to be in the early rehabilitation phase, as they were either occupying long-stay geriatric hospital beds or living at home. It is difficult to identify the phase of stroke recovery for participants in the other trials (Chu 1997; Gelber 1997a; Gelber 1997b; Lewis 1990; Zhang 2002).

Five trials (Brittain 2000a; Judge 1969; Liu 2006; Tibaek 2005; Zhu 2003) reported exclusion criteria. The remaining seven trials did not detail inclusion or exclusion criteria (Chu 1997; Gelber 1997a; Gelber 1997b; Lewis 1990; Wikander 1998b; Zhou 1999; Zhang 2002). Three trials (Liu 2006; Tibaek 2005; Wikander 1998b) reported detailed baseline comparisons.

Setting

Care was provided in a hospital setting in five of the trials: on a rehabilitation ward in a Department of Geriatrics in Sweden (Wikander 1998b); in two long stay geriatric wards in two hospitals in Scotland (Judge 1969) and in hospitals in China (Liu 2006; Zhu 2003) and Denmark (Tibaek 2005) (although participants were outpatients at the time of this study). Brittain 2000a included participants living in the community in England, excluding those living in residential care. The six remaining trials do not specify the setting of care (Chu 1997; Gelber 1997a; Gelber 1997b; Lewis 1990; Zhang 2002; Zhou 1999).

Description of outcomes

Outcomes addressed by the trials were diverse. Six trials expressed the primary outcome as the number of people with urinary incontinence (Brittain 2000a; Chu 1997; Lewis 1990; Wikander 1998b; Zhang 2002; Zhou 1999), with six trials reporting number of incontinent episodes (Gelber 1997a; Gelber 1997b; Judge

1969; Lewis 1990; Liu 2006; Tibaek 2005). One trial (Brittain 2000a) also reported a measure of the severity of incontinence. The trials also measured primary outcomes at different time points. In terms of secondary outcomes, four trials measured urinary symptoms (Brittain 2000a; Liu 2006; Tibaek 2005; Zhu 2003). One trial (Tibaek 2005) included a physical measure of incontinence in the form of pad tests of quantified leakage. Four trials measured functional and psychological outcomes (Brittain 2000a; Tibaek 2005; Wikander 1998b; Zhu 2003). One trial included an economic measure (Wikander 1998b). All outcomes reported in the original trials are included in the review, except global functional index scores reported by Wikander 1998b.

Risk of bias in included studies

Methodological quality was assessed using the quality assessment tool described by the Cochrane Incontinence Review Group.

Allocation

Three of the trials (Brittain 2000a; Judge 1969; Tibaek 2005) were classed as A (adequate concealment of allocation). Eight of the trials (Chu 1997; Lewis 1990; Liu 2006; Gelber 1997a; Gelber 1997b; Wikander 1998b; Zhang 2002; Zhou 1999) were classed as B (unclear). The trial by Zhu (Zhu 2003) was a quasi-experimental study with subjects allocated in sequence according to the time of admission; allocation was therefore not concealed. The trial by Brittain (Brittain 2000a) had a 2:1 randomisation.

Blinding

In the trials of complementary therapies (Chu 1997; Liu 2006; Zhang 2002; Zhou 1999) and meclofenoxate (Zhu 2003), blinding of staff, trial participants and outcome assessment was unclear. In the trials of professional input interventions (Brittain 2000a; Wikander 1998b) and pelvic floor muscle training (Tibaek 2005), health care providers and trial participants could not be blind to treatment status. Brittain 2000a and Tibaek 2005 used independent outcome assessors. In the trial by Wikander (Wikander 1998b), outcomes were assessed by ward staff. In the cross-over trial of oestrogen versus placebo (Judge 1969), blinding of staff and trial participants is likely to have been effective, but it was unclear who undertook outcome assessment.

Follow-up and exclusions

Loss to follow up was unclear in the trial of scalp acupuncture (Chu 1997). The trial of rehabilitation governed by a functional independence measure (Wikander 1998b) reported no loss to follow up. It is unclear why there were fewer patients in the control group (n=13) than the intervention group (n=21). There was

also no reported loss to follow up in the trials of acupuncture versus usual care (Zhang 2002), eye and scalp acupuncture versus no acupuncture (Zhou 1999) and meclofenoxate plus salvia miltirrhiza versus salvia miltirrhiza (Zhu 2003). The cross-over trial of oestrogen versus placebo (Judge 1969) and the trials of pelvic floor muscle training versus usual care (Tibaek 2005) and ginger-salt-partitioned moxibustion plus routine acupuncture versus routine acupuncture (Liu 2006) state numbers and reasons for withdrawals. In the trial of a Continence Nurse Advisor versus general practitioner care (Brittain 2000a), the numbers of withdrawals at three and six months are stated and reasons given, but are not reported separately for intervention and control group. There was missing data at both three and six months, because some participants were not yet due for their assessment when results were reported. The proportions for which data are reported are 82% at three months, and 63% at six months. Comparisons on age and gender for those excluded from analyses showed no significant differences.

Power calculations and intention-to-treat analyses were not reported in any of the trials, although four trials (Wikander 1998b; Zhang 2002; Zhou 1999; Zhu 2003) reported complete data and are therefore likely to have applied intention to treat.

Effects of interventions

I. INTERVENTION VERSUS NO INTERVENTION/USUAL CARE (Comparison 01)

Nine trials (Brittain 2000a; Chu 1997; Gelber 1997a; Liu 2006; Tibaek 2005; Wikander 1998b; Zhang 2002; Zhou 1999; Zhu 2003) with a total of 670 participants compared an intervention to promote urinary continence against no intervention or usual care. The interventions included:

Behavioural interventions:

- timed voiding versus void on request (Gelber 1997a);
- pelvic floor muscle training versus usual care (Tibaek 2005).

Specialised professional input interventions:

- a special intervention programme based on assessment using the Functional Independence Measure (FIM) versus usual rehabilitation care (Wikander 1998b);
- care from a Continence Nurse Advisor (CNP) versus usual care provided by the general practitioner (GP) (Brittain 2000a).

Complementary therapy interventions:

- scalp acupuncture versus no scalp acupuncture (Chu 1997);
- eye and scalp acupuncture versus no acupuncture (Zhou 1999);
- acupuncture versus usual care (Zhang 2002);
- ginger-salt-partitioned moxibustion plus routine acupuncture versus routine acupuncture (Liu 2006);

Pharmacotherapy interventions:

- meclufenoxate plus salvia miltirrhiza versus salvia miltirrhiza (Zhu 200)

Incontinence

Five trials (Brittain 2000a; Liu 2006; Wikander 1998b; Zhang 2002; Zhou 1999) recruiting 485 participants measured number of people with incontinence after treatment; random effects models were used for the meta-analysis because of heterogeneity. All five trials were of specialised professional input interventions or of complimentary therapy interventions.

Specialist professional input interventions

In both the two trials of structured assessment and management, the rate of incontinence was lower in the intervention group than in the control group (Comparison 01.01.01). There was, however, marked heterogeneity between the trials: Wikander (Wikander 1998b), 1/21 vs. 10/13 (RR 0.06; 95% CI 0.01 to 0.43); Brittain (Brittain 2000a), 40/73 vs. 31/48 (RR 0.85; 95% CI 0.63 to 1.14). Although the overall RR (0.26) was consistent with a large treatment effect, applying a random effects model (because of the heterogeneity) led to a wide confidence interval which was not statistically significant (0.01 to 4.67; $p = 0.36$).

Complementary therapy interventions

In the three trials of complementary therapy interventions for which data were available (Liu 2006; Zhang 2002; Zhou 1999), the rate of incontinence was reported to be lower in the intervention group and this difference was statistically significant for two of the studies. This comparison also showed heterogeneity ($I^2 = 80.8\%$). Applying a random effects model, the overall RR was 0.44 (95% CI 0.23 to 0.86; $p = 0.02$).

Severity of incontinence

Behavioural interventions

One trial (Tibaek 2005) measured the mean number of incontinence episodes in 24 hours. The impact of pelvic floor muscle training indicated no significant difference (WMD -1.00, 95% CI -2.74, 0.74, Comparison 01.02.01).

In the trial by Gelber (Gelber 1997a), timed voiding was compared against void on request (which was interpreted as usual care). The data reported were too few even for tentative conclusions, and no further data could be obtained from the investigators. The study was reported as ongoing, but for the purposes of this review was considered closed.

Urinary symptoms

Behavioural interventions

The impact of pelvic floor muscle training on mean daytime voiding frequency (Tibaek 2005) indicated a significant difference (WMD -2.30, 95% CI -4.14, -0.46, Comparisons 01.04.01).

The impact on mean nighttime voiding frequency indicated no significant difference (WMD -0.10, 95% CI -0.83, 0.63, Comparison 01.07.01).

Specialised professional input interventions

The impact of the Continence Nurse Advisor intervention on the number of people not cured of urinary symptoms at three months (Brittain 2000a) indicated a borderline significant difference (48/89 treatment versus 38/54 control, RR 0.77, 95% CI 0.59, 0.99, Comparison 01.03.01). No significant difference was found in the number of people with urinary frequency at three months (98/120 treatment versus 59/67 control, RR 0.93, 95% CI 0.82, 1.05, Comparison 01.05.01), the number of people with urinary urgency at three months (95/121 treatment versus 50/67 control, RR 1.05, 95% CI 0.89, 1.24, Comparison 01.06.01) or the number of people with nocturia at three months (102/119 treatment versus 60/67 control, RR 0.96, 95% CI 0.86, 1.07, Comparison 01.08.01).

There were no data suitable for analysis in RevMan Analyses in relation to changes in the mean number of symptoms (see 'Characteristics of Included Studies Table'). However, the trial report suggested that the evidence favoured a Continence Nurse Advisor compared with usual care in terms of reducing the total number of symptoms experienced ($p < 0.01$) at three months, with weak evidence in favour of the Continence Nurse Advisor in reducing the total number of symptoms at six months ($p = 0.06$) (Brittain 2000a).

There were no data suitable for analysis in RevMan Analyses in relation to changes in day and night-time leakage scores (see 'Characteristics of Included Studies Table'). However, the trial report suggested that the evidence favoured a Continence Nurse Advisor compared with usual care in terms of day-time severity of leakage at three months ($p = 0.038$).

Complementary therapy interventions

The impact of ginger-salt-partitioned moxibustion on mean daytime voiding frequency (Liu 2006) indicated a significant difference (WMD -5.57, 95% CI -7.00, -4.14, Comparison 01.04.02). There was also a significant difference on mean nighttime voiding frequency (WMD -3.18, 95% CI -3.95, -2.41, Comparison 01.07.02).

There were no data suitable for analysis in RevMan Analyses in the trial of scalp acupuncture versus no scalp acupuncture (Chu 1997). After two weeks and two courses of treatment, the investigators reported a reduction in urinary frequency or incontinence of 90.3% in the intervention group, with two people not regaining "normal urine", 12 people partly regaining "normal urine" and 16 people regaining "normal urine". A significant difference

between the experimental and control groups was reported as “p (0.05–0.001)”. No results are reported for the control group. No further data could be obtained for this study.

Pharmacotherapy interventions

The impact of meclufenoxate indicated a significant difference in the number of people whose incontinence did not improve (Zhu 2003) (9/40 treatment versus 27/40 control, RR 0.33, 95% CI 0.18, 0.62, Comparison 01.03.02).

Urological measures

Behavioural interventions

One trial (Tibaek 2005) measured mean function of the pelvic floor muscle. The impact of pelvic floor muscle training indicated no significant difference (WMD 0.10, 95% CI -0.48, 0.68, Comparison 01.09.01).

Health status and quality of life

Behavioural interventions

Tibaek 2005 measured health status in terms of the mean total score on the SF36. The impact of pelvic floor muscle training indicated no significant difference (WMD -28.00, 95% CI -169.66, 113.66, Comparison 01.10.01). Quality of life was measured using the mean total score on the IIQ; no significant difference was found (WMD -3.00, 95% CI -47.87, 41.87, Comparison 01.11.02).

Specialised professional input interventions

Wikander 1998b measured quality of life in terms of mean change in psychological well-being measured by the Psychological General Well-Being Index. The impact of a functional assessment programme indicated a significant difference (WMD -39.00, 95% CI -51.19, -26.81, Comparison 01.11.01).

Function

Specialised professional input interventions

Wikander 1998b measured function in terms of the number of people independent in transfer from wheelchair to toilet. The impact of a functional assessment programme indicated a significant difference (6/21 treatment versus 11/13 control, RR 0.34, 95% CI 0.17 to 0.69, Comparison 01.13.01).

Pharmacotherapy interventions

One trial measured function in terms of mean Barthel score (Zhu 2003). The impact of meclufenoxate indicated a significant difference (WMD 3.40, 95% CI 2.85 to 3.95, Comparison 01.12.01).

Patient satisfaction

Specialised professional input interventions

One trial (Brittain 2000a) measured satisfaction in terms of people not satisfied with the service at three months. The impact of the Continence Nurse Advisor indicated a significant difference (13/109 versus 17/45 control, RR 0.32, 95% CI 0.17, 0.59, Comparison 01.14.01).

Cost/service use

Specialised professional input interventions

One trial (Wikander 1998b) measured service use in terms of people discharged to a setting other than their home. The impact of a functional assessment programme indicated a significant difference (3/21 treatment versus 8/13 control, RR 0.23, 95% CI 0.07 to 0.72, Comparison 01.15.01).

Six month follow-up

Behavioural interventions

Tibaek 2005 measured outcomes at six months and found a pelvic floor muscle training programme indicated no significant difference in terms of mean SF36 score (WMD -46.00, 95% CI -165.05, 73.05) and mean IIQ score (WMD -4.00, 95% CI -56.17, 48.17).

Specialised professional input interventions

Brittain 2000a measured outcomes at six months and found some evidence of an impact of the Continence Nurse Advisor in terms of people with urinary symptoms at 6 months (48/89 treatment versus 38/54 control, RR 0.77, 95% CI 0.59 to 0.99), but little difference in terms of the number of people with incontinence after treatment (75/91 treatment versus 47/55 control, RR 0.96, 95% CI 0.83, 1.11), urinary frequency (73/89 treatment versus 47/54 control, RR 0.94, 95% CI 0.82, 1.09), urinary urgency (65/91 treatment versus 40/54 control, RR 0.96, 95% CI 0.79, 1.18) and nocturia (77/89 treatment versus 46/53 control, RR 1.00, 95% CI 0.87, 1.14). There were no data suitable for analysis in RevMan Analyses in relation to changes in the total number of symptoms; however, the trial report suggested that the evidence favoured a Continence Nurse Advisor compared with usual care in terms of reducing the total number of symptoms at six months (p=0.06).

2. INTERVENTION VERSUS PLACEBO (Comparison 02)

The only trial assessing this comparison was in the *pharmacotherapy interventions* category.

One small cross-over trial (Judge 1969) compared an intervention designed to promote urinary continence (oestrogen) against placebo in a long term care setting in 13 women with a history of stroke. Results were reported separately for people with mild or severe incontinence and in view of the significant heterogeneity between the two groups; a random effects model was used. There was evidence in favour of oestrogen in both groups, with a stronger result in severe incontinence:

- mild incontinence: paired samples mean difference in number of incontinent episodes per week -1.75; 95% CI -3.31 to -0.19, Comparison 02.01.01 (Judge 1969);
- severe incontinence: paired samples mean difference -6.4; 95% CI -9.47 to -3.33. Comparison 02.01.02 (Judge 1969).

Combined results favoured oestrogen but were not statistically significant (paired samples mean -3.88 95% CI -8.42 to 0.66) (Judge 1969).

3. SPECIFIC INTERVENTION VERSUS ANOTHER INTERVENTION (Comparison 03)

The only trial assessing this comparison compared a behavioural intervention (timed voiding) with a pharmacotherapy intervention (oxybutinin) (Gelber 1997b). The data were too few for useful analysis, and no further data could be obtained from the investigators. The study was reported as ongoing, but for the purposes of this review was considered closed.

4. COMBINED INTERVENTION VERSUS SINGLE INTERVENTION (Comparison 04)

The only trial assessing this comparison compared a behavioural intervention with a physical therapy intervention; no studies assessed combinations involving any of the other categories of intervention. Lewis 1990 compared a combined intervention (sensory motor biofeedback plus timed voiding) designed to promote urinary continence against a single intervention (timed voiding). The number of incontinence episodes were fewer in the control group (WMD 2.20; 95% CI 0.12 to 4.28, Comparison 04.02.01) (Lewis 1990).

DISCUSSION

Summary of main results

The review aimed to consider all interventions designed to treat urinary incontinence after stroke in adults, rather than testing a specific hypothesis. Experimental studies were found testing a wide range of interventions.

Behavioural interventions

The small trial testing pelvic floor muscle training against general rehabilitation (Tibaek 2005) found evidence in favour of the treatment group in terms of mean voiding frequency over 24 hours and mean daytime voiding frequency; however findings were only significant when outcomes were measured over three rather than two days. No significant differences were found in the primary outcome (24 hour pad test) or secondary outcomes including vaginal palpation of the pelvic floor muscle, the SF36 or the IIQ.

Specialised professional input interventions

One of the two trials testing specialised professional input (Wikander 1998b) suggests that structured assessment and management is promising in helping people to regain continence in the early post-stroke period: this type of intervention merits further research. Some features of this trial have to be considered in interpreting the large effects suggested. Because the intervention was delivered on a separate ward, equivalence of treatment other than the intervention could not be assessed, particularly in relation to staff mix and numbers. The sample size was small, it is not clear why the numbers in the two trial groups differed, and outcome measurement was not blinded. Furthermore, if there is an effect, it is not possible to separate out the influences of structured assessment which could include training of the staff, multi-disciplinary coordination or the focus provided to staff by regular feedback of outcome, from the involvement of the patient in the assessment and management of their care (Wikander 1998b).

The other trial of specialised professional input (Brittain 2000a) found that outcomes from Continence Nurse Advisor and usual care groups were similar at all data points, with the exception of the number of people cured of all urinary symptoms at six months and satisfaction with the service at three months, where findings favoured the treatment group (although the confidence intervals were wide). The intervention used by the Continence Nurse Advisor was well defined, with a structured assessment and intervention protocol. This was particularly appropriate where reasons for continuing problems with continence had not been investigated. The generalisability of these results has to take into consideration the wide definition of urinary incontinence used, which included people suffering from a range of other urinary symptoms such as frequency or urgency. While the trial overall was relatively large and well designed, the proportion of participants for whom results were available at six months was quite low (63%) and some of the data could not be analysed.

If there are differences between the two professional input intervention trials, these could be related both to the intensity of the interventions, and to the type of trial participant. Participants in the trial by Wikander 1998b were between 11-19 days post-stroke when the hospital-based intervention began, whereas the trial by Brittain 2000a was conducted in a community setting. This is likely to have included people with a much less recent stroke. Par-

ticipants in the trial by Wikander 1998b were also known to have developed continence problems only subsequent to their stroke, while this is not known for the other trial (Brittain 2000a).

Complementary therapy interventions

Three trials of complementary therapies found significant differences favouring the treatment group: acupuncture was more effective than usual care in terms of people with incontinence after treatment (Zhang 2002), eye and scalp acupuncture was more effective than no acupuncture in terms of people with incontinence after treatment (Zhou 1999) and ginger-salt-partitioned moxibustion plus routine acupuncture was more effective than routine acupuncture in terms of mean daytime voiding frequency and mean nighttime voiding frequency (Liu 2006). However, these trials reported minimal methodological detail and it is likely they were of poor quality. It is also not clear whether, and how, these interventions are transferable to the health services of other countries.

Pharmacotherapy interventions

In the small cross-over trial testing an intervention (oral oestrogen) against placebo (Judge 1969), there were fewer incontinence episodes per week during oestrogen treatment. However, it is only known that the participants had a history of stroke. The high mean age of the participants in the trial, the presence of confusion and lack of mobility, and the setting in which the trial took place (two geriatric hospitals) suggest that continence problems may have been secondary to other conditions as well as stroke. For this reason it is problematic to generalise these results to women after stroke. Furthermore, the dose prescribed is not that currently recommended, hormone replacement therapy is anyway now widely considered to be contraindicated in patients who have had, or are at risk of, stroke, and there is evidence that oestrogens increase incontinence in postmenopausal women (Hendrix 2005).

It is notable that the most promising results for regaining continence were from the Wikander 1998b trial, where the time since stroke onset of the participants was short. Early intervention could have greatest impact on the numbers regaining continence. However, 25% of people discharged from hospital after stroke still have continence problems. There is a lack of evidence on whether the many other forms of intervention that have been shown to have some value in the general population would also be relevant to people after stroke. In particular, there are no usable results from trials testing the use of bladder relaxants such as anticholinergics, or programmes of timed voiding or bladder training.

Overall completeness and applicability of the evidence

The major point of interest in the included studies is the mix of categories of intervention, including physical, behavioural and complementary therapies, drugs, and professional input interventions such as using trained personnel and specific methods of managing care. However, within categories it was generally not possible to combine studies due to use of different outcome measures.

Quality of the evidence

The trials included were generally small (n=724 participants in total; median =49 participants); only three trials (Brittain 2000a; Judge 1969; Tibaek 2005) had adequate allocation concealment and many were limited by poor reporting. For most, it was impossible to judge the extent to which they might have been prone to bias. The small sample sizes meant that confidence intervals were wide and did not rule out clinically important differences when there was no statistically significant difference. Even the largest trial (Brittain 2000a) is not fully reported, and it has not been possible to obtain full data from the investigators. The review is also limited by the lack of full data for four other trials (Chu 1997; Gelber 1997a; Gelber 1997b; Lewis 1990).

Potential biases in the review process

The original protocol for this review specified inclusion criteria for participants as having had a stroke in the previous 12 months, but this definition proved unworkable, because most of the trials did not specify or report time since stroke. There was also a lack of clarity about whether urinary incontinence was subsequent to stroke. Only three trials (Liu 2006; Tibaek 2005; Wikander 1998b) specified that urinary continence problems were subsequent to stroke, and detailed the time since stroke onset. For this reason, the extent to which the results for the remaining trials can be generalised to people with continence problems solely as the result of a stroke is unclear.

AUTHORS' CONCLUSIONS

Implications for practice

There is very little evidence from stroke-specific studies to guide practice. The lack of trials testing the same category of intervention means that recommendations for practice are based on the results of a few, usually small, trials. The Wikander 1998b and Brittain 2000a trials provide some evidence to suggest that specialised professional input using systematic methods to assess and manage continence problems may improve some outcomes. The limited evidence suggests that the greatest impact on urinary incontinence may be in the acute phase of rehabilitation after stroke.

However, the Brittain (Brittain 2000a) trial suggests that specialist input and individualised care management may improve the number of symptoms of urinary incontinence even in the longer term.

Implications for research

Evidence suggesting that beneficial outcomes may be achieved by structuring the management of care for people with urinary continence problems following a stroke points to the need for larger trials. Given the variety of problems that can hinder the maintenance of continence after stroke, the use of individualised assessment and goal setting to tailor interventions to the neurological and functional problems of the individual would seem to be especially worthy of consideration.

Methods of managing continuing urinary incontinence such as intermittent catheterisation or the use of catheter valves are also needed.

Three trials of complementary therapies, namely acupuncture (Zhou 1999; Zhang 2002) and ginger-salt-partitioned moxibustion (Liu 2006), suggest these interventions may be worth investigating further with more rigorous study design.

There is a need for more well-designed studies. Further research should use standardised definitions and classification systems to record details of the type and severity of stroke, and the type and severity of urinary incontinence. Pre-stroke continence status, time since stroke and stroke recurrence should also be recorded, with clear inclusion criteria for continence status. Exclusion cri-

teria should be given for comorbidities and clinical indicators of underlying urogenital or systemic conditions such as infection. Specific details of structured assessment and intervention protocols need to be given, with standardisation of treatment, measures of between groups contamination or differences, and tailoring of intervention to the early or later phases of rehabilitation. Outcome measures of urinary incontinence and of urinary symptoms should be standardised, with attention to their validity and reliability and the blinding of outcome assessment. The measurement of changes in health related quality of life would be valuable. The time periods for review should be standardised for the acute, early and later phases of rehabilitation. Lastly, sample size calculations and secure randomisation at either the cluster or individual patient level should be used appropriately.

ACKNOWLEDGEMENTS

We wish to thank members of the Editorial team in the Incontinence Review Group, in particular Sheila Wallace and Cathryn Glazener, and Hazel Fraser and Brenda Thomas of the Stroke Review Group. Our thanks also to the Faculty of Health of the University of Central Lancashire for supporting this review, to trial authors for kindly responding to our requests for information and to the members of the Stroke Incontinence Interest Group for their valuable comments. Thank you to Gui Li of the Shanghai Medical Military University and Luyan Fang of the University of Central Lancashire for kindly translating some of our included studies.

REFERENCES

References to studies included in this review

Brittain 2000a {published data only}

Brittain KR, Potter JF. The treatment of urinary incontinence in stroke survivors (MS9). Report for NHS R&D Programme on Cardiovascular Disease and Stroke Project, Division of Medicine for the Elderly, Dept of Medicine, University of Leicester, in collaboration with the MRC Incontinence Study 2000.

Chu 1997 {published data only}

Chu M, Feng J. Discussion on treating frequent urine due to multiple cerebral embolism with scalp acupuncture (Translation from Chinese). *Information on Traditional Chinese Medicine* 1997;5:42.

Gelber 1997a {published data only}

Gelber DA, Swords L. Treatment of post-stroke urinary incontinence (Abstract). *Journal of Neurologic Rehabilitation* 1997;11(2):131.

Gelber 1997b {published data only}

Gelber DA, Swords L. Treatment of post-stroke urinary incontinence (Abstract). *Journal of Neurologic Rehabilitation*

1997;11:131.

Judge 1969 {published data only}

Judge TG. The use of quinestradol in elderly incontinence women, a preliminary report. *Gerontologica Clinica* 1969; 11:159–64.

Lewis 1990 {published data only}

Lewis AM, Travis ML, Gordon AL, Weaver HB, Reding MJ. Sensory-motor biofeedback for the treatment of urinary urge incontinence following stroke (Abstract). *Clinical Research* 1990;38(1):A10.

Liu 2006 {published data only}

* Liu H, Wang L. Randomized controlled study on ginger-salt-partitioned moxibustion at Shenque (CV 8) on urination disorders poststroke. *Chinese Acupuncture & Moxibustion* 2006;26(9):621–624.

Tibaek 2005 {published and unpublished data}

Tibaek S, Gard G, Jensen R. Is there a long-lasting effect of pelvic floor muscle training in women with

urinary incontinence after ischemic stroke?. *International Urogynecology Journal* 2007;**18**:281–287.

* Tibaek S, Gard G, Jensen R. Pelvic floor muscle training is effective in women with urinary incontinence after stroke: a randomised, controlled and blinded study. *Neurology and Urodynamics* 2005;**24**:348–357.

Tibaek S, Jensen R, Lindskov G. Can quality of life be improved by pelvic floor muscle training in women with urinary incontinence after ischemic stroke? A randomised, controlled and blinded study. *International Urogynecology Journal* 2004;**15**:117–123.

Wikander 1998b {published data only}

Wikander B, Ekelund P, Milsom I. An evaluation of multidisciplinary intervention governed by functional independence measure (FIMSM) in incontinent stroke patients. *Scandinavian Journal of Rehabilitation Medicine* 1998;**30**(1):15–21.

Zhang 2002 {published data only}

* Zhang Z, Ma F, Ma Y. Observation on the effects of acupuncture in the treatment of urinary retention due to cerebral infarction in 36 patients. *Heilongjiang Medicine and Pharmacy*. 2002; Vol. 25, issue 3:71.

Zhou 1999 {published data only}

* Zhou G, Wu D. 40 examples of using eye acupuncture and electriferous scalp acupuncture to treat urinary incontinence after cerebrovascular accident. *Journal of Clinical Acupuncture*. 1999; Vol. 15, issue 9:33–34.

Zhu 2003 {published data only}

* Zhu Y, Zhu X, Zhu D, Jin Z. Meclofenoxate in treating urinary incontinence after acute cerebral infarction. *Chinese Journal of New Drugs and Clinical Remedies* 2003;**9**: 520–522.

References to studies excluded from this review

Cook 1998 {published data only}

Cook D, Hubokly E, Hasskarl J, Hochrien S, Reding M. Effect of voiding position on urinary retention post stroke. *Journal of Stroke and Cerebrovascular Diseases* 1998;**7**:382.

Gross 1990 {published data only}

Gross J C. Bladder dysfunction after a stroke: it's not always inevitable. *Journal of Gerontological Nursing* 1990;**16**:20–5, 41–2.

Tekeoglu 1998 {published data only}

Tekeoglu Y, Adak B, Goksoy T. Effect of transcutaneous electrical nerve stimulation (TENS) on Barthel Activities of Daily Living (ADL) Index score following stroke. *Clinical Rehabilitation* 1998;**12**:277–80.

References to studies awaiting assessment

Engberg 2002 {published and unpublished data}

Engberg S, Sereika SM, McDowell J, Weber E, Brodak I. Effectiveness of prompted voiding in treating urinary incontinence in cognitively impaired homebound older adults. *Journal of Wound, Ostomy and Continence Nursing* 2002;**29**(5):252–265.

McDowell 1999 {published and unpublished data}

McDowell BJ, Engberg S, Sereika S, Donovan N, Jubeck ME, Weber E, Engberg R. Effectiveness of behavioral therapy to treat incontinence in homebound older adults. *Journal of the American Geriatrics Society* 1999;**47**:309–318.

Additional references

Abrams 2002

Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, et al. The standardisation of terminology of lower urinary tract function: report from the standardisation subcommittee of the International Continence Society. *Neurourology and Urodynamics* 2002;**21**:167–68.

Arunabh 1993

Arunabh MB, Badlani GH. Urologic problems in cerebrovascular accidents. *Probl Urol* 1993;**7**:41.

Barer 1989

Barer DH. Continence after stroke: useful predictor or goal of therapy?. *Age & Ageing* 1989;**18**(3):183–91.

Barrett 2001

Barrett JA. Bladder and bowel problems after a stroke. *Reviews in Clinical Gerontology* 2001;**12**:253–67.

Brittain 1998a

Brittain KR. Urinary symptoms and depression in stroke survivors. *Age & Ageing* 1998;**27**(Suppl 1):116–7.

Brittain 1998b

Brittain KR, Peet SM, Castleden MD. Stroke and incontinence. *Stroke* 1998;**29**:524–8.

Brittain 1999

Brittain KR, Peet SM, Potter JF, Castleden CM. Prevalence and management of urinary incontinence in stroke survivors. *Age & Ageing* 1999;**28**(6):509–11.

Brittain 2000b

Brittain KR, Perry SI, Peet SM, Shaw C, Dallosso H, Assassa RP, et al. Prevalence and impact of urinary symptoms among community dwelling stroke survivors. *Stroke* 2000; **31**:886–91.

Burney 1996a

Burney TL, Senapati M, Desai S, Choudary ST, Badlani GH. Effects of cerebrovascular accident on micturition. *Urologic Clinics of North America* 1996;**23**:483–90.

Burney 1996b

Burney TL, Senapati M, Desai S, Choudhary MS, Badlani GH. Acute cerebrovascular accident and lower urinary tract dysfunction: a prospective correlation of the site of brain injury with urodynamic findings. *Journal of Urology* 1996; **156**:1748.

Cifu 1999

Cifu DX, Stewart DG. Factors affecting functional outcome after stroke: a critical review of rehabilitation interventions. *Archives of Physical Medicine and Rehabilitation* 1999;**80** (Suppl):S35–9.

Deeks 2006

Deeks JJ, Higgins JPT, Altman DG, editors. Analysing and presenting results. In: Higgins JPT, Green S editor

- (s). *Cochrane Reviewers' Handbook 4.2.6* [updated September 2006]; Section 8. <http://www.cochrane.org/resources/handbook/hbook.htm> (accessed 1 August 2007). Chichester, UK: John Wiley & Sons, Ltd., 2006.
- Eustice 2000**
Eustice S, Roe B, Paterson J. Prompted voiding for the management of urinary incontinence in adults. *Cochrane Database of Systematic Reviews* 2000, Issue 2. [DOI: 10.1002/14651858.CD002113]
- Hendrix 2005**
Hendrix SL, Cochrane BB, Nygaard IE, Handa VL, Barnabei VM, Iglesia C, et al. Effects of estrogen with and without progestin on urinary incontinence. *JAMA* 2005; **293**(8):935–48.
- Marinkovic 2001**
Marinkovic S, Badlani G. Voiding and sexual dysfunction after cerebrovascular accidents. *Journal of Urology* 2001; **165**(2):359–70.
- Meijer 2003**
Meijer R, Ihnenfeldt DS, de Groot IJM, van Limbeek J, Vermeulen M, de Haan RJ. Prognostic factors for ambulation and activities of daily living in the subacute phase after stroke. A systematic review of the literature. *Clinical Rehabilitation* 2003; **17**(2):119–29.
- Nakayama 1997**
Nakayama H, Jorgensen HS, Pedersen PM, Raaschou HO, Olsen TS. Prevalence and risk factors of incontinence after stroke. The Copenhagen Stroke Study. *Stroke* 1997; **28**(1): 58–62.
- Ostaszkiwicz 2004a**
Ostaszkiwicz J, Johnston L, Roe B. Habit retraining for the management of urinary incontinence in adults. *Cochrane Database of Systematic Reviews* 2004, Issue 2. [DOI: 10.1002/14651858.CD002801.pub2]
- Ostaszkiwicz 2004b**
Ostaszkiwicz J, Johnston L, Roe B. Timed voiding for the management of urinary incontinence in adults. *Cochrane Database of Systematic Reviews* 2004, Issue 1. [DOI: 10.1002/14651858.CD002802.pub2]
- Patel 2001**
Patel M, Coshall C, Lawrence E, Rudd AG, Wolfe CD. Recovery from poststroke urinary incontinence: associated factors and impact on outcome. *Journal of the American Geriatrics Society* 2001; **49**(9):1229–33.
- Roe 2007**
Roe B, Ostaszkiwicz J, Milne J, Wallace S. Systematic review of bladder training and voiding programmes in adults: a synopsis of findings from data analysis and outcomes using metastudy techniques. *Journal of Advanced Nursing* 2007; **57**(1):15–31.
- Thuroff 1999**
Thuroff JW, Abrams P, Artibani W, Haab F, Khoury S, Madersbacher H. Clinical Guidelines For The Management Of Incontinence. In: Abrams P, Khoury S, Wein A editor(s). *Incontinence: 1st International Consultation on Incontinence, Monaco*. Plymouth, UK: Health Publication Ltd, 1999: 933–43.
- Ween 1996**
Ween JE, Alexander MP, D'Esposito M, Roberts M. Incontinence after stroke in a rehabilitation setting: outcome associations and predictive factors. *Neurology* 1996; **47**(3):659–63.
- Wikander 1998a**
Wikander B, Ekelund P, Milsom I. An evaluation of multidisciplinary intervention governed by functional independence measure (FIMSM) in incontinent stroke patients. *Scandinavian Journal of Rehabilitation Medicine* 1998; **30**(1):15–21.
- Williams 1993**
Williams A. Caregivers of persons with stroke: their physical and emotional well-being. *Quality of Life Research* 1993; **2**: 213–20.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Brittain 2000a

Methods	RCT Setting: at home
Participants	232 male (61%) and female (39%) adult stroke survivors with urinary incontinence. Inclusion: Stroke diagnosed by self-report (questionnaire) as stroke, subarachnoid haemorrhage or transient ischaemic attack. Self-reported clinically significant urinary symptoms, including leakage of urine, stress incontinence, frequency, nocturia or urgency. Exclusion: Pregnancy, physical causes of urinary tract dysfunction (prolapse, urethral stricture, prostatic obstruction, pelvic mass or malignancy in past 5 years), active treatment for incontinence in past 6 months, neurogenic retention, raised post-residual volumes, glycosuria or haematuria Age 40 to 96 years, mean 70
Interventions	A: 152 people allocated to Continence Nurse Practitioner (CNP) assessment and treatment. CNP received 3 months training, and used formatted assessment and guidelines for treatment, which could include habit retraining, pelvic floor awareness, dietary advice, the provision of continence garments, and GP referral for treatment of atrophic vaginitis, candidiasis or constipation. The intervention comprised six contacts and five treatment visits at weeks 1, 2, 4, 6 and 8. Week 1: History + physical exam, Week 2: Diagnostic visit + development of management plan, Weeks 4 and 6: review, adapt + reinforce management plan, Week 8: re-assessment. B: 80 people allocated to usual care provided by general practitioner and referral to existing services for the management of continence
Outcomes	Incontinence after treatment at 3 months: A: 40/73, B: 31/48 Incontinence after treatment at 6 months: A: 75/91, B: 47/55 Mean day-time leakage severity at 3 months: A: -0.27, B: +0.05 Mean day-time leakage severity at 6 months: A: -0.69, B: -0.52 Mean night-time leakage severity at 3 months: A: -0.19, B: -0.02 Mean night-time leakage severity at 6 months: A: -0.40, B: +0.02 Urinary frequency at 3 months: A: 98/120, B: 59/67 Urinary frequency at 6 months: A: 73/89, B: 47/54 Urinary urgency at 3 months: A: 95/121, B: 50/67 Urinary urgency at 6 months: A: 65/91, B: 40/54 Nocturia at 3 months: A: 102/119, B: 60/67 Nocturia at 6 months: A: 77/89, B: 46/53 Not cured of all urinary symptoms at 3 months: A: 75.3%, B: 82.1% Not cured of all urinary symptoms at 6 months: A: 48/89, B: 38/54 Mean number of urinary symptoms at 3 months: A: -0.58, B: -0.41 Mean number of urinary symptoms at 6 months: A: -0.47, B: -0.20 Not satisfied with service: A: 13/109, B: 17/45
Notes	Sub sample of the MRC Incontinence Study 2:1 randomisation. No baseline differences in age and gender between the groups Proportions for which data reported: 82% at 3 months, 63% at 6 months

Brittain 2000a (Continued)

<i>Risk of bias</i>		
Item	Authors' judgement	Description
Allocation concealment?	Yes	A - Adequate

Chu 1997

Methods	RCT
Participants	32 (53%) male and 28 (47%) female adult ischaemic stroke survivors with urinary incontinence/frequency. Subjects described as having multi-focal cerebral infarction. Age: male 50-71, female 45-61.
Interventions	A: 30 people allocated to scalp acupuncture using the "bai hui" channel. Needle in situ for 1-2 days in hot weather, and 3-7 days in cold weather. Needle manipulated every 4 hours. Treatment period: 1-2 weeks. B: 30 people allocated to usual care. No details given.
Outcomes	A: people totally regaining "normal urine" = 16/30, partly regaining "normal urine" = 12/30, not regaining "normal urine" = 2/30. B: no results given.
Notes	Both groups received acupuncture in the channels "guan yuan" and "san yinjiao" combined with nursing care, which was interpreted as usual care

Risk of bias

Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Gelber 1997a

Methods	RCT
Participants	37 adult unilateral stroke survivors with urinary incontinence and normal urodynamic studies. Age and gender mix not given
Interventions	A: 8 people allocated to timed voiding B: 10 people allocated to void on request No further details given
Outcomes	Number of incontinence episodes per day. No numerical results reported
Notes	

Risk of bias

Gelber 1997a (Continued)

Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Gelber 1997b

Methods	RCT
Participants	37 adult unilateral stroke survivors with urinary incontinence and bladder hyperreflexia. Age and gender mix not given
Interventions	A: 9 people allocated to Oxybutinin B: 10 people allocated to timed voiding No further details given
Outcomes	Number of incontinence episodes per day. No numerical results reported
Notes	

Risk of bias

Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Judge 1969

Methods	Cross-over trial Setting: Long stay geriatric hospitals
Participants	13 females with a diagnosis of cerebrovascular accident or "multiple little strokes", 7 with mild incontinence (average 12.3 episodes/week), and 6 with severe incontinence (average 41.1 episodes/week). Inclusion: None stated Exclusion: Faecal impaction, urinary infection Age: mean 82, range 66-92
Interventions	Oestrogen as Quinestradol 0.25mg 4 times a day for one month or placebo with a wash-out period of one month
Outcomes	Number of incontinent episodes per week: Group 1 (mildly incontinent) placebo week 10.14 (SD 4.76), active preparation week 8.43 (SD 2.42) Group 2 (severely incontinent) placebo week 38.80 (SD 5.07), active preparation week 32.4 (SD 2.42)
Notes	Participants in group 1 (defined as 'mildly incontinent') and group 2 (defined as 'severely incontinent') were from two different hospitals

Risk of bias

Judge 1969 (Continued)

Item	Authors' judgement	Description
Allocation concealment?	Yes	A - Adequate

Lewis 1990

Methods	RCT
Participants	23 adults with post-stroke urinary urge incontinence. 5 (22%) haemorrhagic stroke, 18 (78%) with ischaemic stroke. Age and gender mix not given
Interventions	A: 11 people allocated to sensory-motor biofeedback device (Uristop) + timed voiding B: 12 people allocated to timed voiding Treatment time of 2 weeks.
Outcomes	Incontinence after treatment: A: 10/11, B: 10/12 Number of incontinence episodes: A: 5.7 (SD 2.5), B: 3.5 (SD 2.6)
Notes	Continenence defined as zero incontinence episodes in the last two days of the study

Risk of bias

Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Liu 2006

Methods	RCT Setting: inpatient
Participants	35 male (47%) and 40 female (53%) adult stroke survivors with urinary incontinence Inclusion: cerebral infarction or haemorrhage, time between stroke onset and admission to hospital less than 6 months, continent of urine pre-stroke, presence of urinary incontinence, urgent or frequent micturition when condition stabilised after stroke, micturition problems affected quality of life, conscious and able to communicate, normal recognition, aged between 40 and 75 years Exclusion: co-morbidities involving the heart, kidney or other important organs, pre-stroke chronic urinary retention and urinary incontinence, unable to communicate, long-standing chronic urinary tract infection Age: mean 64 years
Interventions	A (39): Ginger-salt-partitioned moxibustion at Sheque (CV 8) and routine acupuncture (e.g. Tsusanli, Yinlingquan). B (36): Routine acupuncture.

Liu 2006 (Continued)

Outcomes	Mean number of urination times each day: A: 7.03, B: 12.60 Mean number of times requesting toileting at night: A: 1.75, B: 4.93 Number of participants with incontinence at 72 hours: A: 16/39; B: 21/36	
Notes	No baseline differences in age, gender, type of stroke, stroke course, frequency symptoms and grade of urinary incontinence between the groups	
Risk of bias		
Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Tiback 2005

Methods	RCT Setting: outpatient	
Participants	26 women adult stroke survivors with mixed stress/urge urinary incontinence Inclusion: women diagnosed with first ever ischaemic stroke according to WHO definition and verified by CAT scan; stroke symptoms in at least 1 month; normal cognitive function (mini-mental state examination score <24); UI according to ICS definition; independent walking ability indoors >100m with/without aids; independence in toilet visits; age between 40 and 85 years Exclusion: urinary tract infection; symptom of vaginal prolapse; chronic respiratory diseases; psychiatric diseases; other neurological diseases; unable to speak Danish Age: median 60 years	
Interventions	A (14): Intervention group: "systematic, controlled, intensive pelvic floor muscle (PFM) training programme". Programme consisted of 1) introduction to theory 2) home exercises, including strength PFM exercise by performing close to maximum contraction (6 sec contraction/6 sec rest) Subjects instructed to repeat exercise programme gradually 6-10 times in supine, standing and sitting positions, 1-2 times daily. 3) group treatment once per week, including isolating PFM contraction (6 sec contraction/6 sec rest) and strength exercises (3 sec contraction/3 sec rest, and 6 sec contraction/6 sec rest). All techniques were repeated 4-8 times while supine, sitting and standing, and also prior to daily activities such as rising, sitting and walking. Vaginal palpation was performed to ensure correct contraction, to evaluate contraction strength and to give feedback to subjects B (12) Comparison group received normal, general rehabilitation with no specific treatment of urinary incontinence	
Outcomes	Two day mean voiding frequency over 24 hours: A: 7.6, B: 9.2 Two day mean daytime voiding frequency: A: 5.9, B: 7.3 Two day mean nighttime voiding frequency: A: 1.7, B: 1.9 Two day mean number of incontinence episodes/24 hours: A: 0.2, B: 1.1 Two day mean number of used pads/24 hours: A: 1.1, B: 2.0 Three day mean voiding frequency over 24 hours: A: 7.6, B: 10.0 Three day mean daytime voiding frequency: A: 5.8, B: 8.1 Three day mean nighttime voiding frequency: A: 1.7, B: 1.8 Three day mean number of incontinence episodes/24 hours: A: 0.2, B: 1.2	

Tiback 2005 (Continued)

	<p>Three day mean number of used pads/24 hours: A: 1.1, B: 1.8 Mean function of pelvic floor muscle: A: 2.3, B: 2.2 Mean strength of pelvic floor muscle: A: 2.6, B: 2.8 Mean static endurance of pelvic floor muscle: A: 23.4, B: 21.3 Mean dynamic endurance of pelvic floor muscle: A:15.7, B: 9.8 Short Form Health Status Questionnaire (SF36) post-intervention: A: 567, B: 595 Short Form Health Status Questionnaire (SF36) at 6 months: A: 550, B: 596 Incontinence Impact Questionnaire (IIQ) post-intervention: A: 44, B:47 Incontinence Impact Questionnaire (IIQ) at 6 months: A: 43, B: 47</p>
--	--

Notes	No baseline differences in age, number of births or mobility between the groups Proportions for which data reported: 92% at post-test and 6 months
-------	---

Risk of bias

Item	Authors' judgement	Description
Allocation concealment?	Yes	A - Adequate

Wikander 1998b

Methods	RCT Setting: hospital rehabilitation ward
Participants	15 (43%) male and 19 (57%) female adults admitted to hospital within 1 month of acute hemispheric stroke Inclusion: New urinary incontinence since stroke Exclusion: none specified Age: mean 74 (SD 6.87)
Interventions	A: 21 people allocated to intervention programme based on assessment using the Functional Independence Measure (FIM). Ward staff educated in use of FIM. Participants assessed on admission and at weekly goal setting meetings. B: 13 people allocated to usual rehabilitation care based on Bobath method
Outcomes	Incontinence after treatment: A: 1/21, B: 3/13 Functional ability (median values of Katz ADL index scale A-G (B = dependent in 1 activity) A: F/B, B: F/F Change in psychological well-being: A: 51.8, B: 12.8 Not discharged to home setting: A: 3/21, B: 8/13 Not independent in transfer from bed to wheelchair: A: 6/21, B: 9/13 Not independent in transfer from wheelchair to toilet: A: 6/21, B: 11/13 Not independent in management of wheelchair: A: 0/21, B: 8/13
Notes	Rehabilitation period 75-91 days No baseline differences in age, gender, side of stroke lesion, time from stroke onset to admission, duration of hospital stay, dysphasia, functional ability, independence in activities of daily living, mobility or psychological well-being between groups.

Wikander 1998b (Continued)

	Participants were randomised to two different wards.	
Risk of bias		
Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Zhang 2002

Methods	RCT Setting: unclear	
Participants	38 male (59%) and 26 female (41%) adult stroke survivors with urinary incontinence Inclusion and exclusion: unclear Age: range 42-62 years	
Interventions	A (36): acupuncture at the following points: Tsusanli, Yinlingquan, Sanyinjiao B (28): general treatment using mannite and other medicines	
Outcomes	Numbers not regaining continence: A: 6/36, B: 26/28	
Notes	No description of baseline comparisons	
Risk of bias		
Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Zhou 1999

Methods	RCT Setting: unclear	
Participants	32 male (40%) and 48 female (60%) adult stroke survivors with urinary incontinence Inclusion and exclusion: unclear	
Interventions	A (40): eye acupuncture and electriferous scalp acupuncture once per day, six days per week for 4 weeks Medication therapy, e.g. Hua Duo Zai shi pill, Xi bi lin, Nao fu kang, vitamins C and E B (40): Medication therapy, e.g. Hua Duo Zai shi pill, Xi bi lin, Nao fu kang, vitamins C and E	
Outcomes	Numbers not regaining continence: A: 18/40, B: 32/40	
Notes	No description of baseline comparisons	
Risk of bias		

Zhou 1999 (Continued)

Item	Authors' judgement	Description
Allocation concealment?	Unclear	B - Unclear

Zhu 2003

Methods	Quasi-experiment Setting: inpatient
Participants	45 male (56%) and 35 female (44%) adult stroke survivors with urinary incontinence Inclusion: Acute cerebral infarction confirmed by CT scan, urinary incontinence Exclusion: Unconscious, mental abnormality Age: mean 63 years
Interventions	A (40): Meclofenoxate 0.3g three times per day for 1 month, in addition to salvia miltirrhiza injection 16 ml, IV, 4 times per day for 2 weeks. B (40%): salvia miltirrhiza injection 16 ml, IV, 4 times per day for 2 weeks
Outcomes	Number whose urinary symptoms did not improve: A: 9/40, B: 27/40 Mean cognitive function: A: 34.5, B: 30.42 Mean activities of daily living: A: 37.4, B: 34.0
Notes	Groups described as 'comparable'

Risk of bias

Item	Authors' judgement	Description
Allocation concealment?	No	C - Inadequate

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Cook 1998	Does not include a measure of continence
Gross 1990	Does not include a measure of continence
Tekeoglu 1998	Does not directly test a method of promoting continence

DATA AND ANALYSES

Comparison 1. INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Number of people with incontinence after treatment	5		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Professional input interventions	2	155	Risk Ratio (M-H, Random, 95% CI)	0.26 [0.01, 4.67]
1.2 Complementary therapy interventions	3	219	Risk Ratio (M-H, Random, 95% CI)	0.44 [0.23, 0.86]
2 Severity of incontinence: mean number of incontinence episodes in 24 hours	1	18	Mean Difference (IV, Fixed, 95% CI)	-1.0 [-2.74, 0.74]
2.1 Behavioural interventions	1	18	Mean Difference (IV, Fixed, 95% CI)	-1.0 [-2.74, 0.74]
3 Urinary symptoms - overall	2		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
3.1 Professional input interventions	1	143	Risk Ratio (M-H, Fixed, 95% CI)	0.77 [0.59, 0.99]
3.2 Pharmacotherapy interventions	1	80	Risk Ratio (M-H, Fixed, 95% CI)	0.33 [0.18, 0.62]
4 Urinary symptoms - frequency (continuous variables)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
4.1 Behavioural interventions	1	18	Mean Difference (IV, Fixed, 95% CI)	-2.3 [-4.14, -0.46]
4.2 Complementary therapy interventions	1	62	Mean Difference (IV, Fixed, 95% CI)	-5.57 [-7.00, -4.14]
5 Urinary symptoms - frequency (dichotomous variables)	1	187	Risk Ratio (M-H, Fixed, 95% CI)	0.93 [0.82, 1.05]
5.1 Professional input interventions	1	187	Risk Ratio (M-H, Fixed, 95% CI)	0.93 [0.82, 1.05]
6 Urinary symptoms - urgency	1	188	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.89, 1.24]
6.1 Professional input interventions	1	188	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.89, 1.24]
7 Urinary symptoms - nocturia (continuous variables)	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
7.1 Behavioural interventions	1	18	Mean Difference (IV, Fixed, 95% CI)	-0.10 [-0.83, 0.63]
7.2 Complementary therapy interventions	1	62	Mean Difference (IV, Fixed, 95% CI)	-3.18 [-3.95, -2.41]
8 Urinary symptoms - nocturia (dichotomous variables)	1	186	Risk Ratio (M-H, Fixed, 95% CI)	0.96 [0.86, 1.07]
8.1 Professional input interventions	1	186	Risk Ratio (M-H, Fixed, 95% CI)	0.96 [0.86, 1.07]
9 Urological measures: mean function of the pelvic floor muscle	1	23	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.48, 0.68]
9.1 Behavioural interventions	1	23	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.48, 0.68]
10 Health status: mean total score SF36	1	24	Mean Difference (IV, Fixed, 95% CI)	-28.0 [-169.66, 113.66]

10.1 Behavioural interventions	1	24	Mean Difference (IV, Fixed, 95% CI)	-28.0 [-169.66, 113.66]
11 Quality of life	2		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
11.1 Professional input interventions	1	34	Mean Difference (IV, Fixed, 95% CI)	-39.0 [-51.19, -26.81]
11.2 Behavioural interventions	1	24	Mean Difference (IV, Fixed, 95% CI)	-3.0 [-47.87, 41.87]
12 Function: mean Barthel score (continuous variables)	1	80	Mean Difference (IV, Fixed, 95% CI)	3.40 [2.85, 3.95]
12.1 Pharmacotherapy interventions	1	80	Mean Difference (IV, Fixed, 95% CI)	3.40 [2.85, 3.95]
13 Function: people independent in transfer from wheelchair to toilet (dichotomous variables)	1	34	Risk Ratio (M-H, Fixed, 95% CI)	0.34 [0.17, 0.69]
13.1 Professional input interventions	1	34	Risk Ratio (M-H, Fixed, 95% CI)	0.34 [0.17, 0.69]
14 Patient satisfaction	1	154	Risk Ratio (M-H, Fixed, 95% CI)	0.32 [0.17, 0.59]
14.1 Professional input interventions	1	154	Risk Ratio (M-H, Fixed, 95% CI)	0.32 [0.17, 0.59]
15 Cost/service use	1	34	Risk Ratio (M-H, Fixed, 95% CI)	0.23 [0.07, 0.72]
15.1 Professional input interventions	1	34	Risk Ratio (M-H, Fixed, 95% CI)	0.23 [0.07, 0.72]

Comparison 3. INTERVENTION VERSUS PLACEBO

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Severity of incontinence: mean number of incontinence episodes per week	1		Paired samples mean (Random, 95% CI)	-3.88 [-8.42, 0.66]
1.1 Oestrogen versus placebo (mild incontinence)	1		Paired samples mean (Random, 95% CI)	-1.75 [-3.31, -0.19]
1.2 Oestrogen versus placebo (severe incontinence)	1		Paired samples mean (Random, 95% CI)	-6.4 [-9.47, -3.33]

Comparison 5. COMBINED INTERVENTION VERSUS SINGLE INTERVENTION

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Number of people with incontinence after treatment	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Sensory-motor biofeedback device + timed voiding versus timed voiding alone	1	23	Risk Ratio (M-H, Fixed, 95% CI)	1.09 [0.80, 1.49]


2 Severity of incontinence: mean number of incontinence episodes	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
2.1 Sensory-motor biofeedback device + timed voiding vs timed voiding alone	1	23	Mean Difference (IV, Fixed, 95% CI)	2.2 [0.12, 4.28]

Analysis 1.1. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 1 Number of people with incontinence after treatment.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 1 Number of people with incontinence after treatment




Analysis 1.2. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 2 Severity of incontinence: mean number of incontinence episodes in 24 hours.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 2 Severity of incontinence: mean number of incontinence episodes in 24 hours




Analysis 1.3. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 3 Urinary symptoms - overall.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 3 Urinary symptoms - overall




Analysis 1.4. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 4 Urinary symptoms - frequency (continuous variables).

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 4 Urinary symptoms - frequency (continuous variables)




Analysis 1.5. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 5 Urinary symptoms - frequency (dichotomous variables).

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 5 Urinary symptoms - frequency (dichotomous variables)




**Analysis 1.6. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 6
Urinary symptoms - urgency.**

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 6 Urinary symptoms - urgency




Analysis 1.7. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 7 Urinary symptoms - nocturia (continuous variables).

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 7 Urinary symptoms - nocturia (continuous variables)




Analysis 1.8. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 8 Urinary symptoms - nocturia (dichotomous variables).

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 8 Urinary symptoms - nocturia (dichotomous variables)




**Analysis 1.9. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 9
Urological measures: mean function of the pelvic floor muscle.**

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 9 Urological measures: mean function of the pelvic floor muscle




Analysis 1.10. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 10 Health status: mean total score SF36.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 10 Health status: mean total score SF36




Analysis 1.11. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 11 Quality of life.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 11 Quality of life




**Analysis 1.12. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 12
Function: mean Barthel score (continuous variables).**

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 12 Function: mean Barthel score (continuous variables)




**Analysis 1.13. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 13
Function: people independent in transfer from wheelchair to toilet (dichotomous variables).**

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 13 Function: people independent in transfer from wheelchair to toilet (dichotomous variables)




Analysis 1.14. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 14 Patient satisfaction.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 14 Patient satisfaction




Analysis 1.15. Comparison 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE, Outcome 15 Cost/service use.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 1 INTERVENTION VERSUS NO INTERVENTION/USUAL CARE

Outcome: 15 Cost/service use




Analysis 3.1. Comparison 3 INTERVENTION VERSUS PLACEBO, Outcome 1 Severity of incontinence: mean number of incontinence episodes per week.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 3 INTERVENTION VERSUS PLACEBO

Outcome: 1 Severity of incontinence: mean number of incontinence episodes per week




Analysis 5.1. Comparison 5 COMBINED INTERVENTION VERSUS SINGLE INTERVENTION, Outcome 1 Number of people with incontinence after treatment.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 5 COMBINED INTERVENTION VERSUS SINGLE INTERVENTION

Outcome: 1 Number of people with incontinence after treatment




Analysis 5.2. Comparison 5 COMBINED INTERVENTION VERSUS SINGLE INTERVENTION, Outcome 2 Severity of incontinence: mean number of incontinence episodes.

Review: Treatment of urinary incontinence after stroke in adults

Comparison: 5 COMBINED INTERVENTION VERSUS SINGLE INTERVENTION

Outcome: 2 Severity of incontinence: mean number of incontinence episodes



APPENDICES

Appendix I. Search methods and terms used for the extra specific searching for this review

Due to the comprehensive nature of the searches already performed by the Stroke Collaborative Review Group, no additional searches were performed other than on CINAHL, combining stroke terms with terms for urinary incontinence without a research methods filter. This was done because of the potentially poor indexing of nursing research. The following strategy was used to search CINAHL (January 1982 to January 2007) on Dialog Dastar:

1. Urination-disorders#.de
2. Urinary-tract-infections#.de
3. Bladder-neurogenic.de
4. Bowel-and-bladder-management#.de
5. Urologic-nursing.de
6. Urologic-care.de
7. Catheters-urinary#.de
8. Catheter-care-urinary#.de
9. Urinary-catheterization#.de
10. Urinary-bladder-irrigation.de
11. Catheter-irrigation-urinary.de
12. Diagnosis-urologic#.de
13. Urinary-tract-physiology#.de
14. Incontinence-aids#.de
15. Urinary-incontinence-and frequency-comfort-questionnaire.de
16. Tube-care-urinary-IOWA-NIC.de
17. Urinary-catheterisation-IOWA-NIC.de
18. Urinary-elimination-management-IOWA-NIC.de
19. Urinary-incontinence-care-IOWA-NIC.de
20. Urinary-retention-care-IOWA-NIC.de
21. Urinary-continenence-IOWA-NOC.de
22. Urinary-elimination-IOWA-NOC.de
23. Altered-urinary-elimination-NANDA#.de
24. Urinary-elimination-component-SABA-HHCC#.de
25. Genito-urinary-function-OMAHA.de
26. Urin\$ or bladder or urethra\$.ti,ab.
27. Catheterization.w..de
28. Catheters.w..de
29. Catheter-care.de
30. Catheter-occlusion.de
31. Catheter-removal.de
32. Tube-removal.de
33. Catheter-placement-determination.de
34. Catheter-related-complications#.de
35. 26 and (27 or 28 or 29 or 30 or 31 or 32 or 33 or 34)
36. Void\$ near (difficult\$ or impair\$ or problem\$ or disorder\$ or impair\$ or control\$)
37. 36.ti or 36.ab
38. Urin\$ with (incontinen\$ or continen\$)
39. 38.ti or 38.ab
40. Bladder with (incontin\$ or continen\$)
41. 40.ti or 40.ab
42. Detrusor with (instability or stability or stable or unstable)
43. 42.ti or 42.ab
44. Bladder with (instability or stability or stable or unstable)
45. 44.ti or 44.ab

46. (stress or urge or pad\$) with (continen\$ or incontinen\$)

47. 46.ti or 46.ab

48. Urodynamic\$.ti or urodynamic\$.ab

49. 37 or 39 or 41 or 43 or 45 or 47 or 48

50. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25

51. 35 or 49 or 50

Key: # = explode, .de = index word, \$ = wild card.

The above search was combined with the CINAHL search from the Cochrane Stroke Group.

Recent unpublished trial data were also searched for on national and international databases i.e. NHS National Research Register, NHS Research Findings Register, US Community of Science NIH Grants, MetaRegister of Controlled Clinical Trials and CRISP, by adapting terms drawn from the existing search strategies of the Incontinence and Stroke Review Groups.

The reference lists of all relevant reviews and trial reports were searched to identify further relevant studies. Major investigators were contacted to ask for any other possible relevant trials, published or unpublished. In addition, contact was made with the authors of other relevant Cochrane reviews to ascertain whether defined subgroups of stroke survivors were identified in trials testing methods of promoting continence in a general population.

The review was publicised on the following websites: Joanna Briggs Institute, Royal College of Nursing Research Society, Royal College of Nursing Continence Interest Group, Association of Continence Advisors, Sigma Theta Tau.

We did not impose any language or other restrictions on any of these searches.

WHAT'S NEW

Last assessed as up-to-date: 13 November 2007.

Date	Event	Description
13 October 2008	Amended	Converted to new review format.

HISTORY

Protocol first published: Issue 4, 2003

Review first published: Issue 3, 2005

Date	Event	Description
14 November 2007	New citation required and conclusions have changed	Substantive amendment. Five new studies added

CONTRIBUTIONS OF AUTHORS

The review was conceived and designed by Lois Thomas. Searching was completed by Beverley French. Trials were considered for inclusion independently by two reviewers (Lois Thomas, Beverley French). Data extraction and review of the methodological quality of the eligible studies was then independently conducted by two reviewers for each study (Chris Sutton, Michael Leathley, Stephen Cross, Beverley French and Lois Thomas). Extracted data and quality assessment were crosschecked and any disagreements discussed and if necessary resolved by Lois Thomas. Lois Thomas and Beverley French wrote the text of the review. James Barrett and Caroline Watkins acted as external reviewers, and contributed to constructing the implications for practice and research.

DECLARATIONS OF INTEREST

None known.

SOURCES OF SUPPORT

Internal sources

- Faculty of Health, University of Central Lancashire, UK.

External sources

- No sources of support supplied

INDEX TERMS

Medical Subject Headings (MeSH)

Acupuncture Therapy [methods]; Randomized Controlled Trials as Topic; Stroke [* complications; rehabilitation]; Urinary Incontinence [drug therapy; etiology; *therapy]

MeSH check words

Adult; Female; Humans; Male