



Article

Some model theory of $SL(2, \mathbb{R})$

Gismatullin, J, Penazzi, D and Pilay, A

Available at <http://clock.uclan.ac.uk/11247/>

Gismatullin, J, Penazzi, D ORCID: 0000-0002-9732-1577 and Pilay, A (2015) Some model theory of $SL(2, \mathbb{R})$. Fundamenta Mathematicae, 229 . pp. 117-128. ISSN 0016-2736

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.4064/fm229-2-2>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

Some model theory of $\mathrm{SL}(2, \mathbb{R})$

Jakub Gismatullin*

University of Leeds and Uniwersytet Wrocławski

Davide Penazzi†

University of Leeds

Anand Pillay‡

University of Leeds

August 2, 2012

Abstract

We study the action of $G = \mathrm{SL}(2, \mathbb{R})$, viewed as a group definable in the structure $M = (\mathbb{R}, +, \times)$, on its type space $S_G(M)$. We identify a minimal closed G -flow I , and an idempotent $r \in I$ (with respect to the Ellis semigroup structure $*$ on $S_G(M)$). We also show that the “ideal group” $(r * I, *)$ is nontrivial (in fact it will be the group with 2 elements), yielding a negative answer to a question of Newelski.

1 Introduction and preliminaries

Abstract topological dynamics concerns the actions of (often discrete) groups G on compact Hausdorff spaces X . Newelski has suggested in a number of papers such as [4], [5], that the notions of topological dynamics may be useful for “generalized stable group theory”, namely the understanding of definable groups in unstable settings, but informed by methods of stable group theory. Given a structure M and a group G definable in M , we have the (left) action of G on its type space $S_G(M)$. When $\mathrm{Th}(M)$ is stable, there is a unique minimal closed G -invariant subset I of $S_G(M)$ which is precisely the set of generic types of G . Moreover (still in the stable case) $S_G(M)$ is equipped with a semigroup structure $*$: $p * q = \mathrm{tp}(a \cdot b/M)$ where a, b are independent realizations of p, q respectively, and $(I, *)$ is a compact Hausdorff topological group which turns out to be isomorphic to $G(\overline{M})/G(\overline{M})^0$ where \overline{M} is a saturated elementary extension of M . In fact this nice situation is more or less characteristic of the stable case, so will not extend as such to unstable settings (other than what we have called in [2] “generically stable groups”). However it was shown in [7] that for the much larger class of so-called *fs* groups definable in *NIP* theories, the situation is not so far from the stable case.

*Supported by the Marie Curie Intra-European Fellowship MODGROUP no. PIEF-GA-2009-254123 and Polish Government MNiSW grant N N201 545938

†Supported by a postdoctoral fellowship on EPSRC grant EP/I002294/1

‡Supported by EPSRC grant EP/I002294/1

In the \mathcal{o} -minimal context the fsG groups are precisely the definably compact groups; for example working in the structure $(\mathbb{R}, +, \times)$, these will be the semialgebraic compact Lie groups. However there is no general model-theoretic machinery (of a stability-theoretic nature) for understanding *simple non compact real Lie groups* (and their interpretations in arbitrary real closed fields). In this paper we try to initiate such a study, focusing on $G = \mathrm{SL}(2, \mathbb{R})$. The reason we work over the standard model $(\mathbb{R}, +, \times)$ rather than an arbitrary or saturated model is that all types over the standard model are definable, hence externally definable sets correspond to definable sets, and the type space is equipped with an “Ellis semigroup structure” $*$. We expect that analogues of our results hold over arbitrary models, expanded by the externally definable sets. In any case the main objective is to identify a minimal closed G -invariant subset I of $S_G(M)$, to identify an idempotent element $r \in I$ and to describe the “ideal group” $r * I$. Now $r * I$ as an abstract group does not depend on the choice of I or r . Newelski asked in [4] whether for groups G definable in NIP theories, $G(\overline{M})/G(\overline{M})^{00}$ is isomorphic, as an abstract group to this $r * I$. In [7] we gave a positive answer for fsG groups in NIP theories. When $G = \mathrm{SL}(2)$ and K is a saturated real closed field then $G(K)$ is simple (modulo its finite centre) as an abstract group, whereby $G(K) = G(K)^{00}$. However we will show that in the case of $\mathrm{SL}(2, \mathbb{R})$ acting on its type space the ideal group $r * I$ is the group with 2 elements, in particular nontrivial, so giving a negative answer to Newelski’s question. Our idempotent will be obtained as an “independent” (with respect to forking) product of realizations of a generic type of T^{00} over \mathbb{R} and an $H(\mathbb{R})$ -invariant type of H where T is a maximal compact and H is the standard Borel subgroups of $\mathrm{SL}(2, \mathbb{R})$. Moreover I will be a “universal minimal” G -flow, from the point of view of “tame” topological dynamics, discussed briefly in the next paragraph.

So before getting on to more detailed preliminaries, let us mention that the third author has recently developed [8] a theory of “tame” or “definable” topological dynamics, concerning roughly the action of (G, \mathcal{B}) on a compact space X , where G is a (discrete) group, and \mathcal{B} is a G -invariant Boolean algebra of subsets of G . We will not give the definition here, but when \mathcal{B} is the Boolean algebra of all subsets of G , this notion reduces to the standard notion of the (discrete) group G acting by homeomorphisms on X . When G is a group definable in a structure M , and \mathcal{B} is the Boolean algebra of definable (with parameters) subsets of G , then the “universal (G, \mathcal{B}) -flow with a distinguished dense orbit” is the type space $S_G(M)$ (rather than the Stone-Cech compactification βG of G as in the standard case). So this makes the study of the action of G on its type space more attractive or intrinsic, although we will not need to know anything about this general theory for the purposes of the current paper.

We will assume a basic knowledge of model theory (types, saturation, definable types, heirs, coheirs,...). References are [9] and [6]. Let us fix a complete 1-sorted theory T , a saturated model \overline{M} of T , and a model M which is elementary substructure of \overline{M} . In the body of the paper T will be RCF , the theory of real closed fields, in the language of rings, and M will be the “standard model” $(\mathbb{R}, +, \times, 0, 1)$. By a definable set in M we mean a subset of M^n definable (with parameters) in M , namely by a formula $\phi(x_1, \dots, x_n, \overline{b})$ where we exhibit the parameters \overline{b} from M . $S_n(M)$ is the space of complete n -types over

M , equivalently, ultrafilters on the Boolean algebra of definable subsets of M^n (which we identify with the Boolean algebra of formulas $\phi(x_1, \dots, x_n)$ with parameters from M , up to equivalence). This is a compact Hausdorff space, under the Stone space topology.

Definition 1.1. (i) A subset $X \subseteq M^n$ is externally definable if there is a formula $\phi(x_1, \dots, x_n, \bar{b})$ where the parameters \bar{b} are from \overline{M} , such that $X = \{\bar{a} \in M^n : \overline{M} \models \phi(\bar{a}, \bar{b})\}$.

(ii) By $S_{ext,n}(M)$ we mean the space of ultrafilters on the Boolean algebra of externally definable subsets of M^n .

Fact 1.2. For any $p(\bar{x}) \in S_{ext,n}(M)$, there is a unique $p'(\bar{x}) \in S_n(\overline{M})$ which is finitely satisfiable in M and such that the “trace on M ” of any formula in p' is in p . This sets up a homeomorphism ι between $S_{ext,n}(M)$ and the closed subspace of $S_n(\overline{M})$ consisting of types finitely satisfiable in M .

Note that if all types over M are definable, then externally definable subsets of M^n are definable and $S_{ext,n}(M)$ coincides with $S_n(M)$.

Lemma 1.3. Suppose that all types over M are definable, and let $p(x) \in S_n(M)$.

(i) For any $B \supseteq M$, p has a unique coheir $p'(x) \in S_n(B)$, namely an extension of p to a complete type over B which is finitely satisfiable in M .

(ii) For any $B \supseteq M$, p has a unique heir over B , which we write as $p|B$. $p|B$ can also be characterized as the unique extension of p to B which is definable over M . Moreover $p|B$ is simply the result of applying the defining schema for p to the set of parameters B .

(iii) For any tuples b, c from \overline{M} , $\text{tp}(b/M, c)$ is definable over M if and only if $\text{tp}(c/M, b)$ is finitely satisfiable in M .

Now suppose G is a group definable over M . We identify G with the group $G(\overline{M})$ and write $G(M)$ for the points in the model M . We have the spaces of types $S_G(M)$, $S_{ext,G}(M)$ and $S_G(\overline{M})$. For $g, h \in G$ we write gh for the product. $G(M)$ acts (on the left) by homeomorphisms on $S_G(M)$ and $S_{ext,G}(M)$.

Definition 1.4. Let $p(x), q(x) \in S_{ext,G}(M)$. Let b realize q in G , and let a realize the unique $p' \in S_G(\overline{M})$ given by 1.2. We define $p * q$ to be the (external) type of ab over M . So in the case when all types over M are definable, this just means: let $b \in G$ realize q and let $a \in G$ realize the unique coheir of p over M , b , then $p * q = \text{tp}(ab/M)$.

The following is contained in [4] and [5]. Everything can be proved directly, but it is a special case of the theory of abstract topological dynamics, as treated in [1] for example.

Lemma 1.5. (i) $(S_{ext,G}(M), *)$ is a semigroup, which we call the Ellis semigroup, and $*$ is continuous in the first coordinate, namely for any $q \in S_{ext,G}(M)$ the map taking $p \in S_{ext,G}(M)$ to $p * q \in S_{ext,G}(M)$ is continuous.

- (ii) Left ideals of $S_{ext,G}(M)$ (with respect to $*$) coincide with subflows, namely closed $G(M)$ -invariant subsets.
- (iii) If $I \subseteq S_{ext,G}(M)$ is a minimal subflow, then I contains an idempotent r such that $r * r = r$, and $(r * I, *)$ is a group, whose isomorphism type does not depend on I or r .

As mentioned earlier, in the stable case there is a unique minimal subflow, the space of generic types of G over M . We will, below, consider the case where T is the theory of real closed fields, $M = (\mathbb{R}, +, \times)$ is the standard model and $G = \text{SL}(2, -)$. Sometimes we write \mathbb{R} for M to be consistent with standard notation. So $G(\mathbb{R})$ is the interpretation of G in M , namely $\text{SL}(2, \mathbb{R})$, and \mathbb{R} as a structure is $(\mathbb{R}, +, \times)$. It is well-known that all types over \mathbb{R} are definable [3], hence Lemma 1.5 applies to $G(\mathbb{R})$ acting on $S_G(\mathbb{R})$.

2 $\text{SL}(2, \mathbb{R})$

We review some basic and well-known facts about $\text{SL}(2, \mathbb{R})$, the group of 2×2 matrices over \mathbb{R} with determinant 1. All the objects, maps etc. we mention will be semialgebraic and so pass over to $\text{SL}(2, K)$ where K is a saturated real closed field. We sometimes write G for $\text{SL}(2)$, so $G(\mathbb{R})$ for $\text{SL}(2, \mathbb{R})$. Write I for the identity matrix. The centre of $\text{SL}(2, \mathbb{R})$ is $\{I, -I\}$. The quotient of $\text{SL}(2, \mathbb{R})$ by this centre is called $\text{PSL}(2, \mathbb{R})$.

$H(\mathbb{R})$ will denote the standard Borel subgroup of $G(\mathbb{R})$, namely the subgroup consisting of matrices $\begin{pmatrix} b & c \\ 0 & b^{-1} \end{pmatrix}$ where $b \in \mathbb{R}_{>0}$ and $c \in \mathbb{R}$. $H(\mathbb{R})$ is precisely the semidirect product of $(\mathbb{R}_{>0}, \times)$ with $(\mathbb{R}, +)$. We let $T(\mathbb{R})$ denote $\text{SO}(2, \mathbb{R})$: the subgroup of $G(\mathbb{R})$ consisting of matrices $\begin{pmatrix} x & -y \\ y & x \end{pmatrix}$ with $x, y \in \mathbb{R}$ and $x^2 + y^2 = 1$. The symbol T here stands for torus. $H(\mathbb{R}) \cap T(\mathbb{R}) = \{I\}$ and any element of G can be written uniquely in the form ht (as well as $t_1 h_1$) for $t, t_1 \in T$ and $h, h_1 \in H$. $T(\mathbb{R})$ is a maximal compact subgroup of $G(\mathbb{R})$. Note that $-I \in T(\mathbb{R})$.

We write $V(\mathbb{R})$ for the homogeneous space $G(\mathbb{R})/H(\mathbb{R})$ (space of left cosets $\{gH(\mathbb{R}) : g \in G(\mathbb{R})\}$), and π (or $\pi(\mathbb{R})$) for the projection $G(\mathbb{R}) \rightarrow V(\mathbb{R})$. Note that $\pi|_{T(\mathbb{R})} : T(\mathbb{R}) \rightarrow V(\mathbb{R})$ is a homeomorphism. We write the action of $G(\mathbb{R})$ on $V(\mathbb{R})$ by \cdot . Understanding this action will be quite important for us. The usual action of $G(\mathbb{R})$ on the real projective line by Mobius transformations factors through the action of $G(\mathbb{R})$ on $V(\mathbb{R})$, and we will try to describe what is going on.

So this standard action of $G(\mathbb{R})$ on $\mathbb{P}^1(\mathbb{R})$ is: $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix}$, where $\begin{pmatrix} x \\ y \end{pmatrix}$ is a representative of an element of $\mathbb{P}^1(\mathbb{R})$. It is well defined because $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ has determinant 1.

We identify $\begin{pmatrix} x \\ 1 \end{pmatrix}$ with $x \in \mathbb{R}$, and treat $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ as the “point at infinity”. It is easy to prove the following fact.

Remark 2.1. (i) $Stab_{G(\mathbb{R})}\left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right) = H_1(\mathbb{R})$, where $H_1(\mathbb{R}) = H(\mathbb{R}) \times \{I, -I\}$.

(ii) $Z(G(\mathbb{R})) = \{I, -I\}$ acts trivially on $\mathbb{P}^1(\mathbb{R})$, and the resulting action of $\mathrm{PSL}(2, \mathbb{R}) = G(\mathbb{R})/Z(G(\mathbb{R}))$ on $\mathbb{P}^1(\mathbb{R})$ is the usual faithful action.

Let π_1 denote the map from $G(\mathbb{R})$ to $\mathbb{P}^1(\mathbb{R})$ taking g to $g \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. So by Remark 2.1(i), π_1 induces an isomorphism of $G(\mathbb{R})$ -homogeneous spaces $G(\mathbb{R})/H_1(\mathbb{R})$ and $\mathbb{P}^1(\mathbb{R})$. Moreover

Remark 2.2. The restriction of π_1 to $T(\mathbb{R})$ induces a homeomorphism between $T(\mathbb{R})/\{I, -I\}$ and $\mathbb{P}^1(\mathbb{R})$, such that the identity of $T(\mathbb{R})/\{I, -I\}$ goes to $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Finally, by virtue of the homeomorphism $\pi|_{T(\mathbb{R})}$ between $T(\mathbb{R})$ and $V(\mathbb{R})$ and the action of $G(\mathbb{R})$ on $V(\mathbb{R})$, we have an action (also written \cdot) of $G(\mathbb{R})$ on $T(\mathbb{R})$. Note that $g \cdot t$ is the unique $t_1 \in T(\mathbb{R})$ such that $gt = t_1 h_1$ for some (unique) $h_1 \in H(\mathbb{R})$. Likewise by virtue of Remark 2.2, and the action of $G(\mathbb{R})$ on $\mathbb{P}^1(\mathbb{R})$, we obtain an action \cdot_1 of $G(\mathbb{R})$ on $T(\mathbb{R})/\{I, -I\}$. We clearly have:

Remark 2.3. The action \cdot_1 of $G(\mathbb{R})$ on $T(\mathbb{R})/\{I, -I\}$ is induced by the action \cdot of $G(\mathbb{R})$ on $T(\mathbb{R})$. In particular, for any $g \in G(\mathbb{R})$ and $t \in T(\mathbb{R})$, $g \cdot t \in g \cdot_1 (t/\{I, -I\})$.

As remarked above all this passes to a saturated model K of RCF in place \mathbb{R} . We write G for $G(K) = \mathrm{SL}(2, K)$, H for $H(K)$ etc, V for $V(K)$ etc. But now our groups and homogeneous spaces contain nonstandard points, and the study of their types and interaction, is what this paper is about.

3 Main results

We follow the conventions at the end of the last section. ($G = \mathrm{SL}(2)$, K a saturated real closed field, etc.) We say that $a \in K$ is *infinite*, if $a > \mathbb{R}$. And call a *negative infinite* if $a < \mathbb{R}$. $\mathrm{Fin}(K)$ denotes the elements of K which are neither infinite nor negative infinite. Any $a \in \mathrm{Fin}(K)$ has a standard part $\mathrm{st}(a) \in \mathbb{R}$. Also given $B \subset K$, a is *infinite (negative infinite) over B* if $a > \mathrm{dcl}(B)$ ($a < \mathrm{dcl}(B)$). Call $a \in K$ *positive infinitesimal* if $a > 0$ and $a < r$ for all positive $r \in \mathbb{R}$. Likewise for *negative infinitesimal* and for *infinitesimal over B* . Note that if for example $a \in K$ is positive infinitesimal, $p(x) = \mathrm{tp}(a/\mathbb{R})$, and $B \subset K$ then $p|_B$ is the type of an element which is positive infinitesimal over B .

We sometimes write g/H for the left coset gH . The projection $\pi: G \rightarrow V = G/H$ induces a surjective continuous map which we also call π from $S_G(\mathbb{R})$ to $S_V(\mathbb{R})$. Both these type spaces are acted on (by homeomorphisms) by $G(\mathbb{R})$, and we clearly have:

Lemma 3.1. π is $G(\mathbb{R})$ -invariant: namely for any $p \in S_G(\mathbb{R})$, and $g \in G(\mathbb{R})$, $\pi(gp) = g \cdot \pi(p)$.

Definition 3.2. Let $p_1 \in S_G(\mathbb{R})$, and $q \in S_V(\mathbb{R})$. Define $p_1 * q$ to be $\text{tp}(g \cdot b)$ where b realizes q , and g realizes the unique coheir of p_1 over M, b .

With above notation, the following extends Lemma 3.1.

Lemma 3.3. For any $p, p_1 \in S_G(\mathbb{R})$, $\pi(p_1 * p) = p_1 * \pi(p)$.

Proof. Fix $p, p_1 \in G$. Then $p_1 * p = \text{tp}(g_1 g / \mathbb{R})$ where g_1 realizes p_1 , g realizes p and $\text{tp}(g_1 / \mathbb{R}, g)$ is finitely satisfiable in \mathbb{R} . But then $\pi(p_1 * p) = \text{tp}((g_1 g / H) / \mathbb{R}) = \text{tp}(g_1 \cdot (g / H) / \mathbb{R})$. Now $\text{tp}(g_1 / \mathbb{R}, g / H)$ is finitely satisfiable in \mathbb{R} and g / H realizes $\pi(p)$. Hence $\text{tp}(g_1 \cdot (g / H) / \mathbb{R}) = p_1 * \pi(p)$, as required. \square

As above the symbol g will range over elements of G . Also h ranges over elements of H , and t over elements of T . If $h = \begin{pmatrix} b & c \\ 0 & b^{-1} \end{pmatrix}$ is in h we identify it with the pair $(b, c) \in \mathbb{R}_{>0} \times \mathbb{R}$. And if $t = \begin{pmatrix} x & -y \\ y & x \end{pmatrix}$ is an element of T we identify it with the pair (x, y) (so T is identified with the unit circle under complex multiplication).

We now fix some canonical types: $p_0 = \text{tp}(b, c / \mathbb{R})$ where b is infinite and c is infinite over b . It is easy to check that p_0 is *left* $H(\mathbb{R})$ -invariant. Namely if h realizes p_0 and $h_1 \in H(\mathbb{R})$, then $h_1 h$ also realizes p_0 .

Note that all nonalgebraic types (over \mathbb{R}) of elements of T are generic in the sense of [2]. In fact T is the simplest possible *fs* group in *RCF*. Let $q_0 = \text{tp}(x, y / \mathbb{R})$ (as the type of an element of T) where y is positive infinitesimal and $x > 0$ (so x is the positive square root of $1 - y^2$). We call q_0 the type of a “positive infinitesimal” of T : it is infinitesimally close to the identity, on the “positive” side.

Likewise, for any $t \in T(\mathbb{R})$ and $t_1 \in T$, we will say that t_1 is “infinitesimally close, on the positive side” to t if $t_1 t^{-1}$ realizes q_0 .

The bijection (homeomorphism) between T and V given by $\pi|_T$ induces a homeomorphism (still called π) between $S_T(\mathbb{R})$ and $S_V(\mathbb{R})$, so we will sometimes identify them below, although we distinguish between q and $\pi(q)$ (for $q \in S_T(\mathbb{R})$).

Definition 3.4. We define r_0 to be $\text{tp}(th / \mathbb{R})$ where $h \in H$ realizes p_0 and $t \in T$ realizes the unique coheir of q_0 over \mathbb{R}, h .

Note that $\pi(r_0) = \pi(q_0)$. Our first aim is to show that $\text{cl}(G(\mathbb{R})r_0) = I$ is a minimal closed $G(\mathbb{R})$ -flow, and that r_0 is an idempotent. Note that $\text{cl}(G(\mathbb{R})r_0)$ is precisely the set of $p * r_0 \in S_G(\mathbb{R})$ for p ranging over $S_G(\mathbb{R})$. Likewise for $\text{cl}(G(\mathbb{R}) \cdot \pi(r_0))$.

Lemma 3.5. For any $p \in S_G(\mathbb{R})$, $p * \pi(q_0) = \pi(q_0)$ if and only if p is of the form $\text{tp}(t_1 h_1 / \mathbb{R})$ with $h_1 \in H$ and $t_1 \in T$ the identity or a realization of q_0 .

Proof. Let $\text{tp}(t_1, h_1 / \mathbb{R}, t)$ be finitely satisfiable in \mathbb{R} with $h_1 \in H$, $t_1 \in T$ and t realizing q_0 (so t / H realizes $\pi(q_0)$). $\text{tp}(t / \mathbb{R}, t_1, h_1)$ is the unique heir of q_0 over (\mathbb{R}, t_1, h_1) . In particular $t \in T$ is positive infinitesimal over (\mathbb{R}, t_1, h_1) as is $t / H \in V$. Now $h_1 \cdot (1 / H) = 1 / H$, hence clearly $h_1 \cdot (t / H)$ is also infinitesimally close (over M, d_1, h_1) to $1 / H$.

Claim. $h_1 \cdot (t/H)$ is on the “positive” side of $1/H$ and realizes the unique heir of $\pi(q_0)$ over \mathbb{R}, t_1, h_1 .

Proof of Claim. When we mention “positive side” we are identifying V and T . Now the map π_1 from T to \mathbb{P}^1 is a “local homeomorphism” taking the identity to $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and taking positive infinitesimals in T to infinite $x \in K$ and (by definition) respects the action of G . Hence it suffices to show that for $h_1 = (b, c) \in H$, and $x \in K$ infinite, such that $tp(h_1/\mathbb{R}, x)$ is finitely satisfiable in \mathbb{R} , then $h_1 \cdot x = b^2x + bc$ is (positive) infinite over \mathbb{R}, b, c . This is clear: Firstly, x is infinite over \mathbb{R}, b, c . Now as $b^2 > 0$, b^2x is positive infinite over \mathbb{R}, b, c , as is $b^2x + bc$. \square

By the claim $tp(h_1 \cdot (t/H)/\mathbb{R}, t_1, h_1) = tp((t/H)/\mathbb{R}, t_1, h_1)$. So without loss of generality $h_1 = 1$. So we are in the situation of $t, t_1 \in T$, t realizes q_0 and $tp(t_1/\mathbb{R}, t)$ is finitely satisfiable over \mathbb{R} . It is then clear that t_1t realizes q_0 if and only if t_1 is the identity, or itself realizes q_0 . As $t_1 \cdot (t/H) = (t_1t/H)$, and by virtue of π inducing a homeomorphism between $S_T(\mathbb{R})$ and $S_V(\mathbb{R})$, we see that $t_1 \cdot (t/H)$ realizes $\pi(q_0)$ if and only if t_1 is the identity or a realization of q_0 . This proves the lemma. \square

Corollary 3.6. *Let $t \in T$ realize q_0 and let $h_1 \in H$ be such that $tp(h_1/\mathbb{R}, t)$ is finitely satisfiable in \mathbb{R} . Then $h_1t = t_1h_2$ for $t_1 \in T$ realizing q_0 .*

Proof. We have just seen in the first part of the proof of 3.5 that $h_1 \cdot (t/H)$ realizes $\pi(q_0)$, which suffices. \square

Lemma 3.7. *For any $p \in S_G(\mathbb{R})$, $p * r_0 = r_0$ if and only if $p = tp(t_1h_1/\mathbb{R})$ with $h_1 \in H$ and $t_1 \in T$ realizing q_0 .*

Proof. If $p * r_0 = r_0$ then by Lemma 3.3, $p * \pi(r_0) = \pi(r_0)$. As $\pi(r_0) = \pi(q_0)$, by Lemma 3.5, p is of the required form.

Now let $p = tp(t_1h_1/M)$ with t_1 realizing q_0 . Suppose th realizes r_0 and $tp(t_1, h_1/\mathbb{R}, t, h)$ is finitely satisfiable in \mathbb{R} . Note that (as $tp(t/M, h)$ is finitely satisfiable in \mathbb{R}) we have that $tp(t_1, h_1, t/M, h)$ is finitely satisfiable in \mathbb{R} , so by Lemma 1.3(iii), $tp(h/M, t_1, h_1, t)$ realizes $p_0|(M, t_1, h_1, t)$. Now by Corollary 3.6, $h_1t = t_2h_2$ for t_2 realizing q_0 and $h_2 \in H$. Note that we still have $tp(h/M, t_1, t_2, h_2) = p_0|(M, t_1, t_2, h_2)$. Now p_0 is a (definable) left $H(\mathbb{R})$ -invariant type of H , so for any model $K' \supset \mathbb{R}$, $p_0|K'$ is also a left $H(K')$ -invariant type of H . Hence $h_3 = h_2h$ realizes $p_0|(\mathbb{R}, t_1, t_2, h_2)$.

Now $t_1h_1th = t_1t_2h_3$. As t_1, t_2 both realize q_0 so does their product t_1t_2 and we have just seen that $tp(t_1t_2/\mathbb{R}, h_3)$ is the unique coheir over (\mathbb{R}, h_3) of q_0 . So $t_1t_2h_3$ realizes r_0 as required. \square

From Lemmas 3.5 and 3.7 we conclude easily:

Corollary 3.8. *The restriction of $\pi: S_G(\mathbb{R}) \rightarrow S_V(\mathbb{R})$ to $\text{cl}(G(\mathbb{R})r_0)$ is a homeomorphism between $\text{cl}(G(\mathbb{R})r_0)$ and $\text{cl}(G(\mathbb{R}) \cdot \pi(r_0))$*

Lemma 3.9. *The set $S_{V,na}(\mathbb{R})$ of nonalgebraic types in $S_V(\mathbb{R})$ is the unique minimal closed $G(\mathbb{R})$ -invariant subset of $S_V(\mathbb{R})$.*

Proof. Let for now S denote the set of nonalgebraic types in $S_V(\mathbb{R})$, a closed subspace. It is obviously $G(\mathbb{R})$ -invariant. To show minimality it is enough, to note that, identifying (via π) S with the space of nonalgebraic types in $S_T(\mathbb{R})$, it is minimal closed $T(\mathbb{R})$ -invariant. If U is a basic open subset of $S_T(\mathbb{R})$ which is not a finite set (of isolated points), then by o -minimality, U contains an “interval” in $S_T(\mathbb{R})$, namely the set of types containing a formula defining an interval, with endpoints in $T(\mathbb{R})$, in T with respect to the circular ordering on T . But clearly if $s \in S$ then for some $g \in T(\mathbb{R})$, $gs \in U$. So S contains no proper $T(\mathbb{R})$ -invariant closed subset, whereby S is minimal closed $T(\mathbb{R})$ -invariant, as required. Uniqueness is clear. \square

From 3.8 and 3.9 we deduce:

Proposition 3.10. *$I = \text{cl}(G(\mathbb{R})r_0)$ is a minimal $G(\mathbb{R})$ -invariant closed subspace of $S_G(\mathbb{R})$ and is homeomorphic as a $G(\mathbb{R})$ -flow to $S_{V,na}(\mathbb{R})$ under π .*

Lemma 3.11. *$r_0 * r_0 = r_0$. Namely r_0 is an idempotent in I .*

Proof. This is special case of Lemma 3.7. \square

So we have so far accomplished the first aim: description of a minimal (closed) subflow I of $S_G(\mathbb{R})$ and an idempotent $r_0 \in I$. We now want to describe the “ideal group” $r_0 * I$. Note first:

Lemma 3.12. *The restriction of π to $r_0 * I$ is a bijection with $r_0 * S_{V,na}(M)$*

Proof. By 3.3 and 3.10. \square

We first consider the action of $H < G$ on \mathbb{P}^1 from Section 2. Note that H fixes $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$. We identify any other element $\begin{pmatrix} x \\ 1 \end{pmatrix}$ of \mathbb{P}^1 with $x \in K$. With this notation:

Lemma 3.13. *Let $x \in \mathbb{P}^1$, let h realize p_0 such that $\text{tp}(h/\mathbb{R}, x)$ is finitely satisfiable in \mathbb{R} . Then $h \cdot x$ is positive infinite or negative infinite (in particular infinitesimally close to $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ in \mathbb{P}^1).*

Proof. So $h = (b, c)$ with b (positive) infinite, and c (positive) infinite over b . And $h \cdot x = b^2x + bc$.

- Case (i), x is finite (positive or negative). Then clearly $b^2x + bc$ is positive infinite (as bc is infinite over $|b^2x|$).
- Case (ii), x is positive infinite. Then clearly $b^2x + bc$ is positive infinite.
- Case (iii), x is negative infinite. Now as $\text{tp}(x/b, c)$ is definable over \mathbb{R} , x is negative infinite over $\{b, c\}$, i.e. $x < \text{dcl}(b, c)$. Hence $b^2x < \text{dcl}(b, c)$, whereby $b^2x + bc$ is still negative infinite.

\square

Now we consider the homeomorphism (induced by π_1) between $T/\{I, -I\}$ and \mathbb{P}^1 given in 2.2 and the corresponding action \cdot_1 of G on $T/\{I, -I\}$. As the identity of $T/\{I, -I\}$ goes to $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ under π_1 , we deduce from Lemma 3.13:

Corollary 3.14. *Let $t/\{I, -I\} \in T/\{I, -I\}$ and let h realize p_0 such that $\text{tp}(h/\mathbb{R}, t)$ is finitely satisfiable in \mathbb{R} . Then $h \cdot_1(t/\{I, -I\})$ is infinitesimal in $T/\{I, -I\}$ (namely infinitesimally close to the identity or equal to the identity).*

We now consider the action \cdot of G on T induced by the action of G on V and the homeomorphism between T and V induced by π . And we use Remark 2.3 to conclude:

Lemma 3.15. *Let $t \in T$ and let h realize p_0 such that $\text{tp}(h/\mathbb{R}, t)$ is finitely satisfiable in \mathbb{R} . Then $h \cdot t$ is infinitesimally close to the identity I (i.e. $(1, 0)$) or to $-I$ (i.e. $(-1, 0)$). Moreover both possibilities happen. Namely if t is infinitesimally close to $-I$ so is $h \cdot t$, and if t is infinitesimally close to I then so is $h \cdot t$.*

Proof. The first part follows from Corollary 3.14. The rest follows by continuity and the fact that $h \cdot I = I$ and $h \cdot -I = -I$ (as $-I$ commutes with h). \square

Remember q_0 is the type of a “positive infinitesimal” in T . We let q_1 denote the type of an element of T infinitesimally close to $-I$ and on the “positive side”. Now we can conclude:

Proposition 3.16. $r_0 * S_{V,na}(\mathbb{R})$ has two elements, $\pi(q_0)$ and $\pi(q_1)$.

Proof. We will work instead with the action \cdot of G on $S_T(\mathbb{R})$ induced by the homeomorphism induced by π between $S_T(\mathbb{R})$ and $S_V(\mathbb{R})$. So for a type q of an element of T , by $r_0 * q$ we mean $\text{tp}(g \cdot t/\mathbb{R})$ where t realizes q and g realizes r_0 such that $\text{tp}(g/\mathbb{R}, t)$ is finitely satisfiable in \mathbb{R} .

So let $t_1 \in T \setminus T(\mathbb{R})$ (i.e. t_1 realizes a nonalgebraic type in $S_T(\mathbb{R})$). And let th realize r_0 such that $\text{tp}(th/\mathbb{R}, t_1)$ is finitely satisfiable in \mathbb{R} . Then $\text{tp}(h/\mathbb{R}, t_1)$ is finitely satisfiable in \mathbb{R} , and we may assume that $\text{tp}(t/\mathbb{R}, h, t_1)$ is finitely satisfiable in \mathbb{R} . (And remember that t realizes q_0 and h realizes p_0). By Lemma 3.15, $h \cdot t_1 = t_2$ say, is infinitesimally close to either I or $-I$ (and each can happen for suitable choice of t_1). Note also that $t \cdot t_2$ is just tt_2 (product in T). Now as t realizes q_0 and its type over (\mathbb{R}, t_2) is finitely satisfiable in \mathbb{R} it is easy to see that tt_2 realizes q_0 if t_2 is infinitesimally close to I , and realizes q_1 if t_2 is infinitesimally close to $-I$. This concludes the proof of Proposition 3.16. \square

Putting together with earlier results we summarize (where r_0 is as in Definition 3.4).

Theorem 3.17. (i) $I = \text{cl}(G(\mathbb{R})r_0)$ is a minimal closed $G(\mathbb{R})$ -invariant subset of $S_G(M)$.

(ii) r_0 is an idempotent, with respect to the Ellis semigroup structure $*$ on $S_G(\mathbb{R})$.

(iii) The ideal group $(r_0 * I, *)$ has two elements.

Proof. (i) is Proposition 3.10. (ii) is Lemma 3.11. And (iii) follows from Proposition 3.16 and Lemma 3.12. \square

We finish the paper with some remarks on extensions of our results. Note the ideal group above is precisely the centre of $\mathrm{SL}(2, \mathbb{R})$. Now any finite cover G of $\mathrm{PSL}(2, \mathbb{R})$ can be realized as a semialgebraic group over \mathbb{R} . The proofs above go over, essentially word-for-word, to show that the ideal group of G coincides with the finite group $Z(G)$. The idempotent r_0 and minimal closed G -invariant subset I of $S_G(\mathbb{R})$ are chosen in exactly the same way, and $Z(G)$ is contained in T .

It is natural to also ask about the case where $G = \widetilde{\mathrm{SL}(2, \mathbb{R})}$, the universal cover of $\mathrm{SL}(2, \mathbb{R})$. Now G can be naturally interpreted (defined) in the two-sorted structure $M = ((\mathbb{Z}, +), (\mathbb{R}, +, \times))$. Again all types over this standard model are definable, the \mathbb{Z} -sort being stable. H will be as before and the role of the maximal compact T is now played by the universal cover of $SO(2, \mathbb{R})$, interpreted naturally on the set $\mathbb{Z} \times SO(2, \mathbb{R})$. The above analysis goes through to show, among other things, that the ideal group is $\widehat{\mathbb{Z}}$, the profinite completion of $(\mathbb{Z}, +)$, which is in fact precisely the set of generic types of \mathbb{Z} .

We expect that all this again goes through, with an arbitrary real closed field K in place of \mathbb{R} , where we expand the structure by all externally definable sets and consider the action of $G(K)$ on $S_{ext, G}(K)$.

Finally we expect that essentially the same results hold for arbitrary simple (modulo discrete centre) noncompact Lie groups.

References

- [1] J. Auslander, *Minimal flows and their extensions*, North Holland, Amsterdam, 1988.
- [2] E. Hrushovski and A. Pillay, On NIP and invariant measures, *J. European Math. Soc.* 13 (2011), 1005 — 1061.
- [3] D. Marker and C. Steinhorn, Definable types in \mathcal{o} -minimal theories, *Journal of Symbolic Logic* 59 (1994), 185-198.
- [4] L. Newelski, Topological dynamics of definable group actions, *Journal of Symbolic Logic*, 74 (2009), 50 — 72.
- [5] L. Newelski, Model theoretic aspects of the Ellis semigroup, *Israel J. Math.*, 2012.
- [6] A. Pillay, *Geometric Stability Theory*, Oxford University Press, 1996.
- [7] A. Pillay, Topological dynamics and definable groups, submitted.
- [8] A. Pillay, Tame topological dynamics, in preparation.
- [9] B. Poizat, *A course in model theory*, Springer-Verlag, NY, 2000.