Article

Effect of Gait Imagery Tasks on Lower Limb Muscle Activity With Respect to Body Posture

Kolarova, B., Krobot, A., Polehlova, K., Hlustik, P. and Richards, Jim

Available at <https://clok.uclan.ac.uk/14576/>

Kolarova, B., Krobot, A., Polehlova, K., Hlustik, P. and Richards, Jim orcid iconORCID: 0000-0002-4004-3115 (2016) Effect of Gait Imagery Tasks on Lower Limb Muscle Activity With Respect to Body Posture. Perceptual and Motor Skills, 122 (2). pp. 411-431. ISSN 0031-5125

It is advisable to refer to the publisher's version if you intend to cite from the work.
<http://dx.doi.org/10.1177/0031512516640377>

For more information about UCLan's research in this area go to
<http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to
<http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

EFFECT OF GAIT IMAGERY TASKS ON LOWER LIMB MUSCLE ACTIVITY WITH RESPECT TO BODY POSTURE

BARBORA KOLÁŘOVÁ, ALOIS KROBOT, AND KAMILA POLEHLOVÁ

Department of Physiotherapy, Faculty of Health Sciences, Palacky University Olomouc, Czech Republic

Department of Rehabilitation, University Hospital Olomouc, Czech Republic

PETR HLUŠTÍK

Department of Neurology, Faculty of general medicine and dentistry, University Hospital Olomouc, Czech Republic

JIM D. RICHARDS

Allied Health Research Unit, University of Central Lancashire, Preston, United Kingdom

Summary. ~~(max 180 words)~~–

The objective of this study was to evaluate the effect of gait imagery tasks on lower limb muscle activity with respect to body posture. The sitting and standing position and lower limb muscle activity was evaluated in 27 healthy female students (24.4 ± 1.3 yrs, 167.2 ± 5.2 cm, 60.10 ± 6.4 kg). Surface electromyography was assessed during rest and in three different experimental conditions using mental imagery. These included; a rhythmic gait, rhythmic gait simultaneously with observation of a model and rhythmic gait after performing rhythmic gait. The normalized rmsEMG values with respect to corresponding rest position were compared using non-parametric statistics. Standing gait imagery tasks had facilitatory effect on proximal lower limb muscle activity. However, EMG activity of distal leg muscles decreased for all gait imagery tasks in the sitting position, when the proprioceptive feedback was less appropriate. For subsequent gait motor imagery tasks the muscle activity decreased, probably as result of habituation. In conclusion the effect of motor imagery on muscle activity appears to depend on relative strength of facilitatory and inhibitory inputs.

Keywords: gait, motor imagery, surface electromyography

33 Motor imagery (MI) represents a pure cognitive process, which positively influences motor
34 performance in healthy subjects which has been shown for sport performance, e.g. gymnastics, ballet
35 and tennis (Guillot, Di Rienzo, Macintyre, Moran, & Collet, 2012). In addition this has been shown
36 in patients following motor impairment and has been used in physical therapy during recovery of
37 function (Lotze & Halsband, 2006; Mizuguchi et al., 2012). Specifically walking skills in
38 neurological patients improved after motor imagery exercise (Dunsky, Dickstein, Marcovitz, Levy,
39 & Deutsch, 2008; Oostra, Oomen, Vanderstraeten, & Vingerhoets, 2015). Home-based motor
40 imagery gait training programs have been shown to improve gait parameters including; walking
41 speed, stride length, cadence, single and double support time in chronic post-stroke subjects (Dunsky,
42 Dickstein, Marcovitz, Levy, & Deutsch, 2008). Motor imagery training includes imagery of walking
43 tasks in combination with physical therapy has been suggested to be more effective for improving
44 gait velocity in sub-acute stroke patients than physical therapy alone (Oostra, Oomen, Vanderstraeten,
45 & Vingerhoets, 2015). In addition videotape-based locomotor imagery training together with regular
46 physical therapy has been shown to improve walking ability in post-stroke and people with
47 Parkinson's disease more than gait training alone (El-Wishy & Fayez, 2013; Hwang et al., 2010).

48 Motor imagery can be described as a conscious mental simulation of an action without actual
49 execution, is accompanied by activity in specific neural substrates (both supraspinal and spinal)
50 similar to those involved in the actual executed movement. Meta-analysis on effect of motor imagery
51 on brain structures conducted by Hetu, *et al.* (2013) provided evidence that motor imagery activates
52 motor related brain networks including large fronto-parietal and subcortical regions involved in motor
53 execution. Several studies provided evidence that motor imagery increases excitability in
54 corticospinal tracts which projects directly to motoneurons and their interneurons controlling the
55 muscles (Clark, Mahato, Nakazawa, Law, & Thomas, 2014; Cowley, Clark, & Ploutz-Snyder, 2008;
56 Oku, Ishida, Okada, & Hiraoka, 2011; Roosink & Zijdwind, 2010). This has been shown to increase
57 the excitability of spinal reflexes (Li, Kamper, Stevens, & Rymer, 2004) and also in muscle
58 proprioceptive structures (muscle spindle Ia afferent fibers) (Bonnet, Decety, Jeannerod, & Requin,
59 1997). So it seems that the motoneuron pool of muscle involved in imaginary movement receives
60 summation of neural inputs via descending and ascending neural pathways in similar way as during
61 real movement. The possibility that mental imagery can have an effect on the muscles that create the
62 movement is supported by the positive influence of motor imagery training on muscle strength (Clark
63 et al., 2014; Yue & Cole, 1992). However the influence of motor imagery on electromyography
64 (EMG) measures is not clear yet. To date several studies have found no significant effect of motor
65 imagery on electromyographic activity during imaginary pointing arm movement for upper limb
66 muscles (Demougeot & Papaxanthis, 2011; Gentili, Papaxanthis, & Pozzo, 2006) during imaginary
67 pointing arm movement for upper limb muscles including anterior deltoid, triceps and biceps brachii,

pectoralis major. In addition, Ranganathan, Siemionow, Liu, Sahgal, & Yue (2004) found no increase in activity of biceps brachii and finger abductor during imaginary isometric little finger abduction and elbow flexion, and Lemos, Rodrigues, & Vargas (2014) who found no increase in activity of the gastrocnemius lateralis during imaginary rising on tiptoes. However, Oku, Ishida, Okada, & Hiraoka (2011) found increased EMG in extensor carpi radialis activity during imaginary wrist extension and Guillot et al. (2007) and Dickstein, Gazit-Grunwald, Plax, Dunskey, & Marcovitz (2005) showed increased EMG activity of nine upper limb muscles in agonists, synergists, fixators and antagonists during imaginary lifting a weighted dumbbell and increased EMG activity of gastrocnemius medialis and rectus femoris when performing imaginary rising on tiptoes respectively.

Surface electromyographic measurements reflect, to some extent, the effort of neural system for movement execution as EMG signal is usually proportional to the level of motor unit activity (Richards, 2008). The muscle activity is altered by variations in the balance between inhibitory and facilitatory input which go in parallel to the motoneuron pool, the terminal part of spinal afferent or efferent sensory/motor pathways (Daroff et al., 2012). So it might be accepted that even during MI the magnitude of EMG activity reflects the summation of facilitatory and inhibitory inputs. This assumption is supported by recent findings, which had shown that the increase of EMG activity during MI mirrors a number of facilitatory inputs including mental effort related to e.g. characteristics of imagined object, the heavier was the object lifted in imagination the showed a greater EMG signal during MI (Bakker, Boschker, & Chung, 1996) and tends to be more pronounced in complex functional movements (Bakker et al., 1996; Guillot et al., 2012; Guillot et al., 2007). The EMG signal during motor imagery is classified mostly as subliminal (Guillot et al., 2012; Guillot et al., 2007) or background muscle activity (Oku, Ishida, Okada, & Hiraoka, 2011) which indicates that detectable muscle activity during MI does not have comparable magnitude and phasic pattern to real movement execution. As the amount of increase in EMG amplitude during motor imagery is positively correlated with the amount of corticospinal excitability (Oku, Ishida, Okada, & Hiraoka, 2011) and with respect to previous findings that corticospinal excitability and brain activity during motor imagery is enhanced with the real sensory feedback generated by holding an object which is imaginary manipulated (Mizuguchi et al., 2012) we speculate that EMG activity during gait imagery may be influenced by character of sensory feedback with respect to sitting (non-default position for walking) or standing (default position for walking) body position during imagination.

With respect to imaginary training protocols in sport or in rehabilitation it has been suggested that simultaneously observing somebody doing the task during motor imagery further positively influences neural activity and enhances motor learning processes (Nedelko, Hassa, Hamzei, Schoenfeld, & Dettmers, 2012; Roosink & Zijdwind, 2010; Wright, Williams, & Holmes, 2014). In similar way with respect to motor learning even previous practice of imaginary movement facilitates

neural activity more than imagery before practice, improves imagination ability of this movement (Wriessnegger, Steyrl, Koschutnig, & Muller-Putz, 2014) and combination of imagination with real practice is more effective for motor recovery than movement imagination or execution alone. Therefore the simultaneous observing of imaginary movement will have facilitatory effect on muscle activity.

It has also been previously suggested that the effectivity of the motor imagery training depends on individual's imaging ability (Gregg, Hall, & Butler, 2010). Subjects with a good motor imagery ability show a greater performance improvement following motor imagery training than do subjects with a poor imagery ability (Mizuguchi, Yamagishi, Nakata, & Kanosue, 2015).

The aim of the present study was to analyze the effect of gait imagery tasks from the first person perspective on both proximal and distal lower limb muscle activity. Based on the prior finding that motor imagery activates neural structures in similar way as movement execution and that muscle activity reflects the summation of neural inputs coming to motoneuron pool via afferent and efferent pathways we hypothesized that: (1) imagination of gait (which is considered as complex functional task) modulates lower limb muscle activity, (2) the magnitude of muscle activity reflects character of peripheral sensory inflow during imagination with respect to body posture and (3) the magnitude of muscle activity is further influenced with respect to additional cognitive and motor task.

Therefore this study aimed to evaluate the electromyographic activity of proximal and distal lower limb muscles, which participate synergically on gait execution, during gait imagery tasks compare to rest conditions. This would potentially further our understanding of influence of gait imagery task on motor system and the effect of imagining or observing gait activity of lower limb muscles. This in turn provides important information for gait imagery rehabilitation protocols and could increase our understanding of gait control mechanisms.

Method

Participants

Twenty seven healthy females participated in this study. Their mean ($\pm$ SD) age, height and weight were 24.4 ± 1.3 yrs, 167.2 ± 5.2 cm and 60.10 ± 6.4 kg. All participants were recruited from students from a Physiotherapy department of Palacky University. All participants had good cognitive function and communicative skills necessary for motor imagery and were able to generate gait motor imagery. Only participants with at least moderate visual and kinesthetic imagery ability, evaluated by Revised Movement Imagery Questionnaire (MIQ-R), were included in the study (Smith & Collins, 2004). MIQ-R represents a reliable tool to assess motor imagery ability in healthy persons. MIQ-R consists of an eight-item self-report questionnaire using two 7-point scales to evaluate ability to form visual and kinaesthetic mental images (Hall & Martin, 1997). The exclusion criteria included

138 psychiatric, neurological or musculoskeletal disorders, balance or walking problems, the use of a
139 walking aid, chronic pain, pregnancy, the use of medication affecting the level of vigilance and
140 uncorrected visual impairments. The dominant lower limb was the right side in all participants,
141 determined as preference for kicking a ball (Seeley, Umberger, & Shapiro, 2008). Testing occurred
142 in a quiet room in one day. All participants signed an informed consent prior to participating in this
143 study. The procedures, which were approved by the local ethics committee, were performed
144 according to the ethical standards of the Declaration of Helsinki.

145 *Motor imagery ability measures*

146 When completing the MIQ-R, participants are asked to perform one of four movement tasks
147 and then rate the ease with which they form visual and kinaesthetic images of this movement (from
148 1 = “very hard to see/feel” to 7 = “very easy to see/feel”). In the study mean MIQ-R scores (SD) were
149 47.7 (5.9) for both subscales, 24.15 (2.94) for the visual subscale, and 23.15 (3.15) for the kinaesthetic
150 subscale. The MIQ-R has demonstrated adequate internal consistency with Cronbach α coefficients
151 0.78 and 0.76 for visual and kinaesthetic subscales respectively. MIQ-R mean scores and consistency
152 were comparable to those observed in previous MI studies (Hall & Martin, 1997; Guillot, et al., 2012).

153 *Electromyography measures*

154 Muscle activity was measured using surface EMG using two self-adhesive electrodes (Ag-
155 AgCl). The electrodes were placed in parallel to the muscle fibers in the midline over the muscle belly
156 with an inter electrode distance of 2 cm. Prior to placing the EMG surface electrodes, the skin was
157 abraded and cleaned. EMG activity was recorded from biarticular lower limb muscles involved with
158 gait execution by synergistic action (Chvatal & Ting, 2012). Three distal muscles of the dominant
159 lower limb: tibialis anterior (TA), gastrocnemius lateralis (GL), gastrocnemius medialis (GM), and
160 three proximal muscles of the dominant lower limb: biceps femoris (BF), semitendinosus (ST) and
161 rectus femoris (RF) were measured. The reference electrode was placed over the fibula head. EMG
162 data were recorded at 1000 Hz using the wireless system TeleMyo 2400T G2 (Noraxon Co., USA)
163 with a system bandwidth was 20-1000 Hz. Real-time EMG signals were sent via telemetry at 1,000
164 Hz to an A-D converter (Noraxon Co., USA). The raw EMG signals were full wave rectified and the
165 root mean squared value of EMG (rmsEMG) signals was calculated using a time averaging period of
166 25 ms (Guillot, et al., 2007). The processing of the signal was performed by using the software
167 MyoResearch XP Master Edition 1.08.17 (Noraxon Co., USA). Raw EMG signal was visually
168 checked prior to processing and analysis to verify the absence of any artifacts.

169 *Procedure*

170 The test protocol was conducted with respect to previous findings such that the imagination
171 ability was enhanced when imagination was done from first person perspective, and is performed

172 with externally given auditive feedback (Guillot, *et al.*, 2007; Heremans, *et al.*, 2012; Koehler, *et al.*,
173 2012; Mizuguchi, *et al.*, 2012; Roosink & Zijdewind, 2010).

174 EMG data were initially collected in two default rest positions, sitting (non-default position
175 for walking) and standing (default position for walking) without performing any voluntary activity or
176 motor imagery, and then within six motor imagery experimental conditions in the following order:

- 177 1. gait imagery in the sitting position, gait imagery in the standing position,
- 178 2. gait imagery and simultaneous gait observation in the sitting position, gait imagery and
179 simultaneous gait observation in the standing position,
- 180 3. gait imagery in the sitting position after gait execution, gait imagery in the standing position
181 after gait execution.

182 Experimental conditions are illustratively demonstrated in Figure 1.

183 FIGURE 1

184 *Figure 1 insert here*

185 Default sitting or standing positions were standardized for all experimental conditions. In the
186 sitting position, the participants were seated upright in a chair that leaned against the back and arm
187 rest. In the standing position, the participants were standing upright with hands along their body. In
188 both default positions, the feet were placed a pelvic width apart. In all experimental situations, the
189 position of the feet was unchanged.

190 For every participant and for all tested conditions, the rhythm of gait was given to the
191 participants using a metronome set at 110 beats per minute, to replicate a normal gait cadence All
192 tested participants reported that they were able to imagine gait well at this step frequency. In the first
193 experimental imaginary gait conditions for sitting and standing, the participants were instructed to
194 imagine a rhythmic gait as vividly as possible, in the first person perspective, the instruction was
195 “*Imagine yourself walking on the pace of the metronome*“ without making any actual movements. In
196 second tested conditions, the participants observed the rhythmic gait of a second person in frontal
197 plane from posterior side on the projection screen (200 x 200 cm) placed 2 meters in front of them.
198 The participants were instructed to watch the gait and to simultaneously imagine a rhythmic gait as
199 if they were walking (the instruction was “*Observe the woman on the screen walking at the pace of*
200 *the metronome and simultaneously imagine yourself walking at the same pace*“). Next, real rhythmic
201 walking at the pace of the metronome in hospital corridor was performed by the participants for a few
202 minutes to enhance further rhythmic gait imagination ability (Wriessnegger, Steyrl, Koschutnig, &
203 Muller-Putz, 2014). Just after real rhythmic walking, third experimental conditions were performed,
204 the instruction within the gait imagery task after gait execution was the same task as in the first
205 experimental conditions “*Imagine yourself walking on the pace of the metronome*“. Each gait imagery

task lasted for approximately 60 seconds. None of participants mentioned feelings of fatigue during the experimental session.

Data processing

The rmsEMG [%] was calculated for every experimental condition in sitting or standing position and then normalized to the rmsEMG of default sitting or standing rest positions. For the rest sitting and standing positions the average rmsEMG values of all tested the muscles were calculated over a 20 seconds interval. These values calculated during the rest condition without any motor imagery were considered as reference values. For all rhythmic gait imagery tasks the mean rmsEMG values were calculated over six gait cycles for the dominant lower limb. The duration of evaluated EMG period was 6.6 seconds which was calculated from the metronome frequency where one gait cycle was 1.1 seconds. This period was selected from the middle part of the measured data for every experimental condition with respect to adaptation on the imagery task. The mean rmsEMG values during experimental gait imagery tasks were expressed as a percentage of reference value. Gait imagery experimental tasks conducted in sitting position were normalized to the respective reference value obtained in rest sitting position and gait imagery tasks conducted in standing position were normalized to the respective reference value obtained in rest standing position for every participant and tested muscle.

Statistical analysis

Data were tested to determine if they were normally distributed using Kolmogorov-Smirnov test. All data were found not to be normally distributed, ($p < 0.05$), therefore non-parametric tests were used throughout the analysis. For the statistical analysis the non-parametric Wilcoxon signed-rank test was performed with the alpha value was set at $p < 0.05$. This allowed the comparison of the reference values for sitting and standing positions and normalized EMG data for experimental conditions in sitting and standing positions respectively (hypothesis 1). And the comparison of normalized EMG data with respect to the default sitting and standing positions (hypothesis 2) alpha value was set at $p < 0.05$. The differences between each of the gait imagery conditions in the sitting or standing position (hypothesis 3) were explored with Friedman tests with post-hoc Wilcoxon tests. As normalized data for three experimental imagery conditions were compared and the alpha value was calculated using Bonferroni's adjustment as $0.05/3$ and set at $p < 0.017$). In addition the effect size for non-parametric data (Fritz, Morris, & Richler, 2012) Z values were computed. All statistical analysis were performed using Statistica 9.0.

Results

For all tested muscles in rest default sitting and standing position the EMG activity was almost silent, the mean and standard deviation reference rest electromyography data [μV] are presented in **Table 1**. All experimental gait imagery conditions were normalized as a percentage of the rest values separately for each posture, muscle and participant, descriptive statistics of these data are presented in **Table 2**. First gait imagery task in standing position had facilitatory effect on proximal lower limb muscle activity (Table 2, Table 3). However, EMG activity of distal leg muscles decreased for all gait imagery tasks in the sitting position, when the proprioceptive feedback was less appropriate.

TABLE 1

Table 1 insert here

TABLE 2

Table 2 insert here

Gait imagery tasks vs. rest (Hypothesis 1)

Conditions using rhythmic gait imagery mostly indicated an inhibitory effect on lower limb muscle activity compared to the rest default positions (**Table 3**). In the sitting position this was apparent for GM and GL and for TA in all experimental conditions, for BF and ST during gait imagery and simultaneous gait observation and gait imagery after gait execution.

In the standing position significant inhibition was only present in GL for second gait imagery condition and in TA for second and third gait imagery condition. In the standing position, the first gait imagery task in the proximal tested muscles (BF, RF) resulted in an increased EMG activity.

TABLE 3

Table 3 insert here

Standing vs. sitting position (Hypothesis 2)

When comparing of the normalized EMG data between experimental conditions and between the sitting and standing positions, muscle activity was mostly higher in the standing position (**Table 3**). This support the hypothesis that standing facilitates muscle activity in comparison to sitting. The difference were significant for GL ($p < 0.01$, $ES > 0.3$) and BF ($p < 0.05$, $ES > 0.3$) in every experimental condition, for GM and TA ($p < 0.05$, $ES > 0.3$) in the first (SI1 $\times$ TI1) and third gait imagery condition (SI3 $\times$ TI3), for ST and RF ($p < 0.05$, $ES > 0.3$) in first gait imagery (SI1 $\times$ TI1) and imagery during gait observation (SI2 $\times$ TI2) conditions.

Subsequent gait imagery tasks (Hypothesis 3)

When comparing experimental conditions, in sitting position the EMG activity was lower during the rhythmic gait imagery after rhythmic gait execution in comparison to the second gait imagery condition for GM ($Z=2.83$, $p=0.005$, $ES=0.36$), GL ($Z=3.24$, $p=0.001$, $ES=0.038$), and TA ($Z=3.73$, $p<0.001$, $r=0.49$) and in comparison to the first gait imagery condition (SI1 $\times$ SI3) for GM ($Z=2.64$, $p=0.01$, $ES=0.39$), GL ($Z=2.79$, $p<0.001$, $ES=0.44$), and TA ($Z=3.63$, $p<0.001$, $ES=0.51$). In the standing position, the muscle activity was lower in the third tested condition compared to the first tested condition for RF ($Z=3.05$, $p<0.001$, $ES=0.42$). For other comparisons the values did not differ significantly.

Discussion

Guillot (2007) showed that MI was accompanied by subliminal EMG activity of muscles participating on imagined movement execution. However the increase of lower limb muscle activity during rhythmic gait imagery was not major finding in our study. Lower limb muscles mostly decreased EMG activity during the experimental tasks using gait imagery compared to the rest conditions, where EMG activity of all muscles was almost silent (Table 1). This was significant especially for distal leg muscles in the sitting position (Table 2 and Table 3). The muscle activity increase during MI compared to rest conditions was previously demonstrated mostly for upper limb tasks (Bakker, *et al.*, 1996; Guillot, Di Rienzo, *et al.*, 2012; Guillot, *et al.*, 2007; Solodkin, *et al.*, 2004) or for non-gait foot tasks (Bakker, *et al.*, 2007; Bonnet, *et al.*, 1997). To follow on from the results of Bakker *et al.* (2008) it could be suggested that during gait imagery compared to imagery of non-gait or postural foot task supraspinal control is suppressed to some extent. Bakker *et al.* (2008) compared corticospinal excitability within motor imagery of simple foot task (dorsiflexion) and MI of gait measured by motor evoked potentials from task-related muscle m. tibialis anterior in sitting position. They found that motor evoked potentials areas increased during motor imagery of simple foot task, however corticospinal excitability within gait imagery increased just in selected group of subjects (5 from 16) who had larger increased during imagined foot dorsiflexion, so compared to the majority of participants this simple task did not show and increase in muscle activity during gait imagery.

As supraspinal control might be suppressed during imagery of postural task we speculate that the less expressed effect of gait imagery on muscle activity could be influenced by neural gait control mechanisms. Rhythmic complex patterns of synergistic muscle activity required for locomotion are to great extent under control of neural autonomy of CPG, neural networks located in lumbosacral spine connected with supraspinal motor regions and with lower limb afferent peripheral sensors (Solopova *et al.*, 2015, Dietz, 2003, 2010; Chvatal & Ting, 2012; Dietz, 2003; MacKay-Lyons, 2002).

309 Motor imagery of lower-limb movements including gait relies mainly on the supplementary motor
310 area, cerebellum, putamen, and parietal regions (Hetu et al., 2013). Activity of these areas is required
311 more for gait planning with respect to changes of external environment rather than for stereotype
312 locomotion which has been shown to be more automatic (Hetu, *et al.*, 2013; la Fougere, *et al.*, 2010).
313 Activity of CPG might be modulated to a great extent by afferent sensory feedback from lower limb
314 receptors even with suppressed supraspinal control than has been previously demonstrated on spinal-
315 cord-injured humans (Bussel et al, 1996, Dietz, 2003, 2010; Harkema, *et al.*, 1997;) or in situations
316 without any extra demands on gait with respect to e.g. additional task or changes in the external
317 environment (Bussel, *et al.*, 1996; Calancie, *et al.*, 1994). Particularly phasic peripheral sensory
318 information associated with lower limb loading during walking evokes lower limb muscle activity
319 (Harkema et al., 1997). Harkema et al. (1997) found that by 70% unloaded body weight stepping (but
320 not 100% unloaded body weight stepping) movements induced by a driven gait orthosis on a treadmill
321 in healthy subjects elicited muscle activity of distal extensor lower limb muscles, namely
322 gastrocnemius medialis and soleus. So the EMG activity of distal lower limb muscles during the gait
323 is to a great extent dependent on phasic peripheral sensory information especially in situations when
324 no extra attention or demands on posture control are needed. The importance of proprioceptive
325 feedback for muscle activity during walking was suggested further McCrea (2001), who found that
326 feedback from extensor proprioceptors induces locomotor dependent reflexes that contribute
327 considerably to extensor muscle activity during real walking. So it is probable that especially distal
328 lower limb motor neurons don't receive enough facilitatory inputs to evoke muscle activity during
329 stereotype rhythmic gait imagery tasks in sitting position. Furthermore it seems that during ~~the~~
330 imagining of gait in a position in which walking is impossible dominate inhibitory effect over possible
331 facilitatory on the muscle activity.

332 The emerging question from these current findings is not only why tested gait imagery
333 conditions do not have facilitatory effect on muscle activity, which was the major focus in previous
334 studies, but why gait imagery tasks resulted in decreased muscle activity compared to the rest
335 condition in our experiment.

336 To date a decrease of EMG activity during imagination of movement execution task has not
337 been described. Decreased excitability of motor neural system during movement imagery compared
338 to rest condition, specifically decreased activity of corticospinal tract, has been previously reported
339 for imagination of muscle relaxation (Kato, Watanabe, Muraoka, & Kanosue, 2015) or during
340 imagination of suppressing movements (Sohn, Dang, & Hallett, 2003) for upper limb tasks. Few
341 studies found decreased corticospinal excitability during imagination of postural tasks in comparison
342 to rest conditions (Hiraoka, 2002; Oishi et al., 1994). Hiraoka (2002) suggested that imagination of
343 stumbling in standing posture lead to decrease excitability of soleus H-reflex and Oishi (1994) found

that imaginary of skating motion in elite skate sprinters led to suppression of soleus H-reflex during whole period of imaginary movement. All these finding are support the previous suggestion that motor commands during motor imagery must be inhibited throughout the neural system to some extent to prevent overt movement execution (Guillot, 2007; Jeanarod, 2001) as EMG activity (if present) is just at subliminal intensity without tonic specific activity as during real movement (Guillot, 2007; Guillot, 2012; Jeanarod, 2001).

Inhibitory processes, which presumably propagate to the spinal motoneurons in parallel with the excitatory inputs might have origin on the cortical, brainstem or either on spinal level (Jeannerod, 2006; Prut & Fetz, 1999). We speculate that the cause of EMG decrease, which occurred mostly in sitting position during gait imagery tasks, presumably mostly took place on spinal level as sitting and standing differs mostly by means of different proprioceptive input. It is probable that muscle spindle afferents is gating the strength of Ia afferent synaptic input onto target motor neurons during gait imagery in the same way as during gait execution (MacKay-Lyons, 2002). One of proposed mechanisms of muscle activity inhibition is presynaptic inhibition according to a previous finding that soleus H-reflex excitability as function of EMG level is decreased during gait (Stein & Capaday, 1998). Presynaptic inhibition reduces the amount of neurotransmitter released at the presynaptic terminal of the Ia axon which lead to decrease in EMG activity (Brooke et al., 1997; Bonnet et al. 1997). Furthermore we speculate that muscle activity decrease during gait imagery task might be influenced by depression of afferent neuronal discharge as has been demonstrated during fictive locomotion in the cat induced by mesencephalic locomotor region stimulation (Perreault et al., 1999). Decrease of muscle and cutaneous afferent-evoked monosynaptic field potentials reflected a reduction of depolarizing synaptic current into spinal neurons during fictive locomotion (Perreault et al., 1999).

The influence of posture

For all tested muscles in most of experimental conditions was muscle activity during gait imaginary tasks significantly lower in sitting position compared to muscle activity during gait imaginary tasks in standing position (see Table 2, 3). Thus, the standing position compared to sitting position had an excitatory effect on muscle activity during rhythmic gait imagery tasks. Standing posture is congruent with walking and thus offer more appropriate somatosensory (tactile, proprioceptive and visual) feedback compared to incongruent positions with walking such as sitting or lying. Presence of real somatosensory feedback facilitates activity of neural structures within motor imagery and motor observation (Mizuguchi et al., 2012; Vargas et al., 2004). Mizuguchi et al. (2012) found that imagination of squeezing the ball and holding the real ball at the same time enhanced the MEPs in comparison to the same situation just without the ball. Vargas et al. (2004) observed that corticospinal

excitability increased in situation when hand posture was compatible with the imagined task compared to incompatible hand posture with the imagined task. Saimpont et al., 2012) proved that posture might influence even accuracy of imagined movement, in their experiment the time duration of gait motor imagery in standing posture (body posture congruent with walking) was more comparable with real gait than gait motor imagery in sitting posture. It has been also previously shown, that standing posture compared to supine posture (the one most used throughout the studies concerning effect of gait observation or gait imagery) has excitatory effect on neural structures (Nakazawa et al., 2003; Shimba et al., 2010). Nakazawa et al. (2003) demonstrated that both stretch reflex and MEP elicited in tibialis anterior were significantly greater in standing compare to supine posture (background EMG was silent in both conditions). Shimba et al. (2010) found that even passive standing posture (accomplished by using gait orthosis) had higher impact on increased stretch reflex of m. soleus compared to supine position. This might reflect facilitatory effect of standing position on muscle spindle Ia afferent fibers. Facilitation of muscle spindle activity with respect to position congruent with imaginary movement found also Bonnet et al. (1997). In their study they showed that mental simulation of pressure on a pedal with the foot in reclined sitting position with their feet on two pedals led to larger changes in T-reflex amplitude compared to H-reflex amplitude (activity of muscle spindle Ia afferent fibres is elicited within the T-reflex, but not by H-reflex) in the leg involved in the simulation. Even the extension of the hip in the standing position might have facilitatory effect on muscle activity compared to sitting position, because also afferent input from hip joints is important for the leg muscle activation during locomotion in dependence on hip position (Dietz and Duysens, 2000; Dietz et al., 2002; Grillner & Rossignol, 1978). Grillner & Rossignol (1978) previously proved that preventing the hip from extension in chronic spinal cats inhibits the flexors muscle activity. As EMG activity depends on level of motoneuron pool excitation it is probable that muscle proprioceptive (muscle spindle) afferents is gating the strength of Ia afferent synaptic input onto target motoneurons during gait imagery, same as during gait execution (MacKay-Lyons, 2002). Then the level of proprioceptors activation might be crucial for the the subthreshold activation of target muscles during gait imagery tasks. This assumption is in accordance with previous studies the appropriate proprioceptive feedback (concretly posture congruent with imaginary task) provided excitatory input to the motor system and facilitates muscle activity.

For the proximal tested muscles (BF and RF) the gait imagery task in the standing position was the only experimental condition when the muscle activity increased compared to the rest position. It has been previously suggested that the proximal leg muscles (e.g., BF) are mostly controlled by the monosynaptic corticospinal pathways compared to mostly polysynaptic corticospinal innervations of the distal leg muscles (e.g., GM) (Brouwer & Ashby, 1991; Cowan, Day, Marsden, & Rothwell, 1986). So presumably during the gait imagery task, the direct neural input from the cortex to the

motoneuron may enhance the ability of the cortex to control the proximal leg muscles (Brouwer & Ashby, 1991). This assumption is in accordance with previous findings that during hand movements dominates monosynaptic cortical-motoneuronal input (Nicolas et al., 2001) and mostly for upper limb movements the presence of EMG activity during imagery tasks has been already demonstrated. It is possible that motor imagery does not provide equivalent neural input to proximal and distal leg muscles, but this has to be further explored. And still just biarticular lower limb muscles were measured. To follow our results it is likely that the imagining of rhythmic gait provides inhibitory input mostly to the distal leg muscles in the default sitting position. In accordance to previously mentioned studies inhibition might reflect the summation of several factors including: decreased supraspinal effort for stereotype gait imagery tasks, spinal inhibitory mechanisms (presynaptic inhibition), different neural drive to the motoneurons of distal and proximal leg muscles, and default sitting posture which does not provide appropriate feedback for real walking. However the results of this study are limited to young woman population with good imagery ability, and to stereotype rhythmic gait imagery task. Therefore, further research is required with respect to different genders and populations.

Comparison of EMG activity during experimental conditions

Combination of motor imagery and observation (Wright, Williams, & Holmes, 2014) or previous imagined movement execution (Wriessnegger, Steyrl, Koschutnig, & Muller-Putz, 2014) enhances activity of neural structures and motor learning processes (Gomes, *et al.*, 2014; Nedelko, Hassa, Hamzei, Schoenfeld, & Dettmers, 2012) compared to motor imagery itself. Based on this assumption we hypothesized, that both simultaneous motor imagery with motor observation and previous execution of imagined movement would have further facilitatory effect on muscle activity compared to gait imagery alone. So we added these “augmented” imagery conditions in given order to the experimental protocol. However in our experiment the second and the third experimental condition mostly led to muscle activity decrease compared to the first tested situation. As the order of first, second and third experimental conditions were not randomized we suggest that the decrease in muscle activity within repeated tested motor imagery tasks in our experiment might reflect to some extent the gradual habituation effect. It has been previously described, that cortical activity is mostly pronounced during initial trials of complex motor imagery tasks (imagery of volleyball spike attack) compared to second and third motor imagery where the short-term habituation effect might be present (Stecklow et al., 2010). None of tested participants reported feelings of tiredness during the experiment the mental fatigue, which has been previously reported for prolonged imagery tasks (Rozand et al., 2016), was not the reason of decreased muscle activity for subsequent imagery tasks.

448 We suggest here that more challenging imagery tasks as part of gait rehabilitation are required, then
449 habituation effect might be avoided (Marchal-Crespo et al., 2014).

450

451 The results of this study potentially further our understanding of influence of rhythmic gait
452 imagination on lower limb muscles with respect to the body posture. This in turn provides important
453 information for gait imagery rehabilitation protocols and could increase our understanding of gait
454 control mechanisms.

455 References

- 456 Bakker, F. C., Boschker, M. S. J., & Chung, T. (1996) Changes in muscular activity while imagining
457 weight lifting using stimulus or response propositions. *Journal of Sport & Exercise*
458 *Psychology*, 18(3), 313-324.
- 459 Bakker, M., de Lange, F. P., Stevens, J. A., Toni, I., & Bloem, B. R. (2007) Motor imagery of gait: a
460 quantitative approach. *Exp Brain Res*, 179(3), 497-504.
- 461 Bakker, M., Overeem, S., Snijders, A. H., Borm, G., van Elswijk, G., Toni, I., & Bloem, B. R. (2008)
462 Motor imagery of foot dorsiflexion and gait: effects on corticospinal excitability. *Clin*
463 *Neurophysiol*, 119(11), 2519-2527.
- 464 Bonnet, M., Decety, J., Jeannerod, M., & Requin, J. (1997) Mental simulation of an action modulates
465 the excitability of spinal reflex pathways in man. *Brain Res Cogn Brain Res*, 5(3), 221-228.
- 466 Brouwer, B., & Ashby, P. (1991) Altered corticospinal projections to lower limb motoneurons in
467 subjects with cerebral palsy. *Brain*, 114 (Pt 3), 1395-1407.
- 468 Bussel, B., Roby-Brami, A., Neris, O. R., & Yakovleff, A. (1996) Evidence for a spinal stepping
469 generator in man. Electrophysiological study. *Acta Neurobiol Exp (Wars)*, 56(1), 465-468.
- 470 Calancie, B., Needham-Shropshire, B., Jacobs, P., Willer, K., Zych, G., & Green, B. A. (1994)
471 Involuntary stepping after chronic spinal cord injury. Evidence for a central rhythm generator
472 for locomotion in man. *Brain*, 117 (Pt 5), 1143-1159.
- 473 Clark, B. C., Mahato, N., Nakazawa, M., Law, T., & Thomas, J. (2014) The Power of the Mind: The
474 Cortex as a Critical Determinant of Muscle Strength/Weakness. *J Neurophysiol*, jn 00386
475 02014.
- 476 Cowan, J. M., Day, B. L., Marsden, C., & Rothwell, J. C. (1986) The effect of percutaneous motor
477 cortex stimulation on H reflexes in muscles of the arm and leg in intact man. *J Physiol*, 377,
478 333-347.
- 479 Cowley, P. M., Clark, B. C., & Ploutz-Snyder, L. L. (2008) Kinesthetic motor imagery and spinal
480 excitability: the effect of contraction intensity and spatial localization. *Clin Neurophysiol*,
481 119(8), 1849-1856.
- 482 Demougeot, L., & Papaxanthis, C. (2011) Muscle fatigue affects mental simulation of action. *J*
483 *Neurosci*, 31(29), 10712-10720.
- 484 Dickstein, R., Gazit-Grunwald, M., Plax, M., Dunskey, A., & Marcovitz, E. (2005) EMG activity in
485 selected target muscles during imagery rising on tiptoes in healthy adults and poststroke
486 hemiparetic patients. *J Mot Behav*, 37(6), 475-483.
- 487 Dietz, V. (2003) Spinal cord pattern generators for locomotion. *Clin Neurophysiol*, 114(8), 1379-
488 1389.
- 489 Dietz, V. (2010) Behavior of spinal neurons deprived of supraspinal input. *Nat Rev Neurol*, 6(3), 167-
490 174.
- 491 Dunskey, A., Dickstein, R., Marcovitz, E., Levy, S., & Deutsch, J. E. (2008) Home-based motor
492 imagery training for gait rehabilitation of people with chronic poststroke hemiparesis. *Arch*
493 *Phys Med Rehabil*, 89(8), 1580-1588.
- 494 El-Wishy, A. A., & Fayez, E. S. (2013) Effect of Locomotor Imagery Training Added to Physical
495 Therapy Program on Gait Performance in Parkinson Patients: A Randomized Controlled
496 Study. *Egyptian Journal of Neurology, Psychiatry & Neurosurgery*, 50(1), 31.
- 497 Gentili, R., Papaxanthis, C., & Pozzo, T. (2006) Improvement and generalization of arm motor
498 performance through motor imagery practice. *Neuroscience*, 137(3), 761-772.
- 499 Gregg, M., Hall, C., & Butler, A. (2010) The MIQ-RS: A Suitable Option for Examining Movement
500 Imagery Ability. *Evid Based Complement Alternat Med*, 7(2), 249-257.
- 501 Guillot, A., Di Rienzo, F., Macintyre, T., Moran, A., & Collet, C. (2012) Imagining is Not Doing but
502 Involves Specific Motor Commands: A Review of Experimental Data Related to Motor
503 Inhibition. *Front Hum Neurosci*, 6, 247.

504 Guillot, A., Lebon, F., Rouffet, D., Champely, S., Doyon, J., & Collet, C. (2007) Muscular responses
505 during motor imagery as a function of muscle contraction types. *Int J Psychophysiol*, 66(1),
506 18-27.

507 Hall, C. R., & Martin, K. A. (1997) Measuring movement imagery abilities: A revision of the
508 Movement Imagery Questionnaire. . *Journal of Mental Imagery*, 21(1-2), 143-154.

509 Harkema, S. J., Hurley, S. L., Patel, U. K., Requejo, P. S., Dobkin, B. H., & Edgerton, V. R. (1997)
510 Human lumbosacral spinal cord interprets loading during stepping. *J Neurophysiol*, 77(2),
511 797-811.

512 Heremans, E., Nieuwboer, A., Spildooren, J., De Bondt, S., D'Hooze A, M., Helsen, W., & Feys, P.
513 (2012) Cued motor imagery in patients with multiple sclerosis. *Neuroscience*, 206, 115-121.

514 Hetu, S., Gregoire, M., Saimpont, A., Coll, M. P., Eugene, F., Michon, P. E., & Jackson, P. L. (2013)
515 The neural network of motor imagery: an ALE meta-analysis. *Neurosci Biobehav Rev*, 37(5),
516 930-949.

517 Hiraoka, K. (2002) Imagining stumbling inhibits motor-evoked potentials in the soleus muscle. *Int J*
518 *Neurosci*, 112(6), 613-622.

519 Hwang, S., Jeon, H. S., Yi, C. H., Kwon, O. Y., Cho, S. H., & You, S. H. (2010) Locomotor imagery
520 training improves gait performance in people with chronic hemiparetic stroke: a controlled
521 clinical trial. *Clin Rehabil*, 24(6), 514-522.

522 Chvatal, S. A., & Ting, L. H. (2012) Voluntary and reactive recruitment of locomotor muscle
523 synergies during perturbed walking. *J Neurosci*, 32(35), 12237-12250.

524 Jeannerod, M. (2006) The origin of voluntary action: history of a physiological concept. *C R Biol*,
525 329(5-6), 354-362.

526 Kato, K., Watanabe, J., Muraoka, T., & Kanosue, K. (2015) Motor imagery of voluntary muscle
527 relaxation induces temporal reduction of corticospinal excitability. *Neurosci Res*, 92, 39-45.

528 Koehler, S., Egetemeir, J., Stenneken, P., Koch, S. P., Pauli, P., Fallgatter, A. J., & Herrmann, M. J.
529 (2012) The human execution/observation matching system investigated with a complex
530 everyday task: a functional near-infrared spectroscopy (fNIRS) study. *Neurosci Lett*, 508(2),
531 73-77.

532 la Fougere, C., Zwergal, A., Rominger, A., Forster, S., Fesl, G., Dieterich, M., . . . Jahn, K. (2010)
533 Real versus imagined locomotion: a [18F]-FDG PET-fMRI comparison. *Neuroimage*, 50(4),
534 1589-1598.

535 Li, S., Kamper, D. G., Stevens, J. A., & Rymer, W. Z. (2004) The effect of motor imagery on spinal
536 segmental excitability. *J Neurosci*, 24(43), 9674-9680.

537 Lotze, M., & Halsband, U. (2006) Motor imagery. *J Physiol Paris*, 99(4-6), 386-395.

538 MacKay-Lyons, M. (2002) Central pattern generation of locomotion: a review of the evidence. *Phys*
539 *Ther*, 82(1), 69-83.

540 McCrea, D. A. (2001) Spinal circuitry of sensorimotor control of locomotion. *J Physiol*, 533(Pt 1),
541 41-50.

542 Mizuguchi, N., Sakamoto, M., Muraoka, T., Moriyama, N., Nakagawa, K., Nakata, H., & Kanosue,
543 K. (2012) Influence of somatosensory input on corticospinal excitability during motor
544 imagery. *Neurosci Lett*, 514(1), 127-130.

545 Mizuguchi, N., Yamagishi, T., Nakata, H., & Kanosue, K. (2015) The effect of somatosensory input
546 on motor imagery depends upon motor imagery capability. *Front Psychol*, 6, 104.

547 Nedelko, V., Hassa, T., Hamzei, F., Schoenfeld, M. A., & Dettmers, C. (2012) Action imagery
548 combined with action observation activates more corticomotor regions than action
549 observation alone. *J Neurol Phys Ther*, 36(4), 182-188.

550 Oku, K., Ishida, H., Okada, Y., & Hiraoka, K. (2011) Facilitation of corticospinal excitability during
551 motor imagery of wrist movement with visual or quantitative inspection of EMG activity.
552 *Percept Mot Skills*, 113(3), 982-994.

553 Oostra, K. M., Oomen, A., Vanderstraeten, G., & Vingerhoets, G. (2015) Influence of motor imagery
554 training on gait rehabilitation in sub-acute stroke: A randomized controlled trial. *J Rehabil*
555 *Med*, 47(3), 204-209.

556 Prut, Y., & Fetz, E. E. (1999) Primate spinal interneurons show pre-movement instructed delay
557 activity. *Nature*, 401(6753), 590-594.

558 Ranganathan, V. K., Siemionow, V., Liu, J. Z., Sahgal, V., & Yue, G. H. (2004) From mental power
559 to muscle power--gaining strength by using the mind. *Neuropsychologia*, 42(7), 944-956.

560 Roosink, M., & Zijdwind, I. (2010) Corticospinal excitability during observation and imagery of
561 simple and complex hand tasks: implications for motor rehabilitation. *Behav Brain Res*,
562 213(1), 35-41.

563 Seeley, M. K., Umberger, B. R., & Shapiro, R. (2008) A test of the functional asymmetry hypothesis
564 in walking. *Gait Posture*, 28(1), 24-28.

565 Smith, D., & Collins, D. (2004) Mental practice, motor performance, and the late CNV. *Journal of*
566 *Sport & Exercise Psychology*, 26(3), 412-426.

567 Sohn, Y. H., Dang, N., & Hallett, M. (2003) Suppression of corticospinal excitability during negative
568 motor imagery. *J Neurophysiol*, 90(4), 2303-2309.

569 Solodkin, A., Hlustik, P., Chen, E. E., & Small, S. L. (2004) Fine modulation in network activation
570 during motor execution and motor imagery. *Cereb Cortex*, 14(11), 1246-1255.

571 Van de Crommert, H. W., Mulder, T., & Duysens, J. (1998) Neural control of locomotion: sensory
572 control of the central pattern generator and its relation to treadmill training. *Gait Posture*, 7(3),
573 251-263.

574 Wriessnegger, S. C., Steyrl, D., Koschutnig, K., & Muller-Putz, G. R. (2014) Short time sports
575 exercise boosts motor imagery patterns: implications of mental practice in rehabilitation
576 programs. *Front Hum Neurosci*, 8, 469.

577 Wright, D. J., Williams, J., & Holmes, P. S. (2014) Combined action observation and imagery
578 facilitates corticospinal excitability. *Front Hum Neurosci*, 8, 951.

579 Yue, G., & Cole, K. J. (1992) Strength increases from the motor program: comparison of training
580 with maximal voluntary and imagined muscle contractions. *J Neurophysiol*, 67(5), 1114-
581 1123.

582

583 References

- 584 Aoki, H., Tsukahara, R., & Yabe, K. (2002) Cortical and spinal motor excitability during the
 585 premovement EMG silent period prior to rapid voluntary movement in humans. *Brain*
 586 *research*, 949, 178-187.
- 587 Bakker, F. C., Boschker, M. S. J., & Chung, T. (1996) Changes in muscular activity while imagining
 588 weight lifting using stimulus or response propositions. *Journal of sport & exercise*
 589 *psychology*, 18, 313-324.
- 590 Bakker, M., de Lange, F. P., Stevens, J. A., Toni, I., & Bloem, B. R. (2007) Motor imagery of gait: a
 591 quantitative approach. *Experimental brain research. Experimentelle Hirnforschung.*
 592 *Experimentation cerebrale*, 179, 497-504.
- 593 Beattie, K., Kenny, I. C., Lyons, M., & Carson, B. P. (2014) The effect of strength training on
 594 performance in endurance athletes. *Sports medicine*, 44, 845-865.
- 595 Bonnet, M., Decety, J., Jeannerod, M., & Requin, J. (1997) Mental simulation of an action modulates
 596 the excitability of spinal reflex pathways in man. *Brain research. Cognitive brain research*, 5,
 597 221-228.
- 598 Brouwer, B., & Ashby, P. (1991) Altered corticospinal projections to lower limb motoneurons in
 599 subjects with cerebral palsy. *Brain : a journal of neurology*, 114 , 1395-1407.
- 600 Bussel, B., Roby-Brami, A., Neris, O. R., & Yakovlev, A. (1996) Evidence for a spinal stepping
 601 generator in man. Electrophysiological study. *Acta neurobiologiae experimentalis*, 56, 465-
 602 468.
- 603 Calancie, B., Needham-Shropshire, B., Jacobs, P., Willer, K., Zych, G., & Green, B. A. (1994)
 604 Involuntary stepping after chronic spinal cord injury. Evidence for a central rhythm generator
 605 for locomotion in man. *Brain : a journal of neurology*, 117, 1143-1159.
- 606 Clark, B. C., Mahato, N., Nakazawa, M., Law, T., & Thomas, J. (2014) The Power of the Mind: The
 607 Cortex as a Critical Determinant of Muscle Strength/Weakness. *Journal of neurophysiology*,
 608 112, 3219-3226.
- 609 Cowan, J. M., Day, B. L., Marsden, C., & Rothwell, J. C. (1986) The effect of percutaneous motor
 610 cortex stimulation on H reflexes in muscles of the arm and leg in intact man. *The Journal of*
 611 *physiology*, 377, 333-347.
- 612 Decety, J. (1996) The neurophysiological basis of motor imagery. *Behavioural brain research*, 77,
 613 45-52.
- 614 Dietz, V. (2003) Spinal cord pattern generators for locomotion. *Clinical neurophysiology : official*
 615 *journal of the International Federation of Clinical Neurophysiology*, 114(8), 1379-1389.
- 616 Dietz, V. (2010) Behavior of spinal neurons deprived of supraspinal input. *Nature reviews.*
 617 *Neurology*, 6, 167-174.

618 Dietz, V., & Duysens, J. (2000) Significance of load receptor input during locomotion: a review. *Gait*
619 *& posture*, 11, 102-110.

620 Dunskey, A., Dickstein, R., Marcovitz, E., Levy, S., & Deutsch, J. E. (2008) Home-based motor
621 imagery training for gait rehabilitation of people with chronic poststroke hemiparesis.
622 *Archives of physical medicine and rehabilitation*, 89, 1580-1588.

623 Fadiga, L., Buccino, G., Craighero, L., Fogassi, L., Gallese, V., & Pavesi, G. (1999) Corticospinal
624 excitability is specifically modulated by motor imagery: a magnetic stimulation study.
625 *Neuropsychologia*, 37, 147-158.

626 Grillner, S., & Rossignol, S. (1978) On the initiation of the swing phase of locomotion in chronic
627 spinal cats. *Brain research*, 146, 269-277.

628 Fritz, C. O., Morris, P.E., Richler, J. J. (2012) Effect size estimates: current use, calculations, and
629 interpretation. *Journal of experimental psychology General*, 141, 2-18.

630 Gomes, T. V., Ugrinowitsch H., Marinho, N., Shea, J. B., Raisbeck, L. D., Benda, R. N. (2014) Effects
631 of mental practice in novice learners in a serial positioning skill acquisition. *Perceptual and*
632 *motor skills*, 119, 397-414.

633 Guillot, A., Desliens, S., Rouyer, C., & Rogowski, I. (2013) Motor imagery and tennis serve
634 performance: the external focus efficacy. *Journal of sports science & medicine* 12, 332-338.

635 Guillot, A., Di Rienzo, F., Macintyre, T., Moran, A., & Collet, C. (2012) Imagining is Not Doing but
636 Involves Specific Motor Commands: A Review of Experimental Data Related to Motor
637 Inhibition. *Frontiers in human neuroscience*, 6, 247.

638 Guillot, A., Genevois, C., Desliens, S., Saieb, S., & Rogowski, I. (2012) Motor imagery and 'placebo-
639 racket effects' in tennis serve performance. *Psychol Sport Exerc*, 13, 533-540.

640 Guillot, A., Lebon, F., Rouffet, D., Champely, S., Doyon, J., & Collet, C. (2007) Muscular responses
641 during motor imagery as a function of muscle contraction types. *International journal of*
642 *psychophysiology : official journal of the International Organization of Psychophysiology*,
643 66, 18-27.

644 Hall, C. R., & Martin, K. A. (1997) Measuring movement imagery abilities: a revision of the
645 Movement Imagery Questionnaire. *Journal of Mental Imagery*, 21, 143-154.

646 Harkema, S. J., Hurley, S. L., Patel, U. K., Requejo, P. S., Dobkin, B. H., & Edgerton, V. R. (1997)
647 Human lumbosacral spinal cord interprets loading during stepping. *Journal of*
648 *neurophysiology*, 77, 797-811.

649 Heremans, E., Nieuwboer, A., Spildooren, J., De Bondt, S., D'Hooge A, M., Helsen, W., & Feys, P.
650 (2012) Cued motor imagery in patients with multiple sclerosis. *Neuroscience*, 206, 115-121.

651 Hetu, S., Gregoire, M., Saimpont, A., Coll, M. P., Eugene, F., Michon, P. E., & Jackson, P. L. (2013)
 652 The neural network of motor imagery: an ALE meta-analysis. *Neuroscience and*
 653 *biobehavioral reviews*, 37, 930-949.

654 Jeannerod, M. (2006) *Motor cognition: What actions tell the Self*. New York: Oxford University
 655 Press.

656 Jeannerod, M. (2001) Neural simulation of action: a unifying mechanism for motor cognition.
 657 *NeuroImage*, 14, S103-109.

658 Jeon, H., Kim, J., Ali, A., & Choi, S. (2014) Noise distraction and mental practice in closed and open
 659 motor skills. *Perceptual and motor skills*, 119, 156-168.

660 Kolarova, B., Krobot, A., Habermannova, P., Kolar, P., & Bastlova P. (2015). Motor imagery and
 661 motor observation in cognitive rehabilitation. *Rehabilitacia*, 52, *in-press*.

662 Kaufman, M. T., Churchland, M. M., Ryu, S. I., & Shenoy, K. V. (2014) Cortical activity in the null
 663 space: permitting preparation without movement. *Nature neuroscience*, 17, 440-448.

664 Kho, A. Y., Liu, K. P., & Chung, R. C. (2014) Meta-analysis on the effect of mental imagery on
 665 motor recovery of the hemiplegic upper extremity function. *Australian occupational therapy*
 666 *journal*, 61, 38-48.

667 Koehler, S., Egetemeir, J., Stenneken, P., Koch, S. P., Pauli, P., Fallgatter, A. J., & Herrmann, M. J.
 668 (2012) The human execution/observation matching system investigated with a complex
 669 everyday task: a functional near-infrared spectroscopy (fNIRS) study. *Neuroscience letters*,
 670 508, 73-77.

671 la Fougere, C., Zwergal, A., Rominger, A., Forster, S., Fesl, G., Dieterich, M., . . . Jahn, K. (2010)
 672 Real versus imagined locomotion: a [18F]-FDG PET-fMRI comparison. *NeuroImage*, 50,
 673 1589-1598.

674 Lemos, T., Rodrigues, E. C., & Vargas, C. D. (2014) Motor imagery modulation of postural sway is
 675 accompanied by changes in the EMG-COP association. *Neuroscience letters*, 577, 101-105.

676 Lotze, M. (2013) Kinesthetic imagery of musical performance. *Frontiers in human neuroscience*, 7,
 677 280.

678 MacKay-Lyons, M. (2002) Central pattern generation of locomotion: a review of the evidence.
 679 *Physical therapy*, 82, 69-83.

680 Mizuguchi, N., Nakata, H., Hayashi, T., Sakamoto, M., Muraoka, T., Uchida, Y., & Kanosue, K.
 681 (2013) Brain activity during motor imagery of an action with an object: a functional magnetic
 682 resonance imaging study. *Neuroscience research*, 76, 150-155.

683 Mizuguchi, N., Sakamoto, M., Muraoka, T., Moriyama, N., Nakagawa, K., Nakata, H., & Kanosue,
 684 K. (2012) Influence of somatosensory input on corticospinal excitability during motor
 685 imagery. *Neuroscience letters*, 514, 127-130.

686 Mulder, T. (2007) Motor imagery and action observation: cognitive tools for rehabilitation. *Journal*
687 *of neural transmission*, 114, 1265-1278.

688 Nakazawa, K., Kawashima, N., Obata, H., Yamanaka, K., Nozaki, D., & Akai, M. (2003) Facilitation
689 of both stretch reflex and corticospinal pathways of the tibialis anterior muscle during standing
690 in humans. *Neuroscience letters*, 338, 53-56.

691 Nedelko, V., Hassa, T., Hamzei, F., Schoenfeld, M. A., Dettmers, C. (2012) Action imagery
692 combined with action observation activates more corticomotor regions than action observation
693 alone. *J Neurol Phys Ther* 36, 182-188.

694 Oku, K., Ishida, H., Okada, Y., & Hiraoka, K. (2011) Facilitation of corticospinal excitability during
695 motor imagery of wrist movement with visual or quantitative inspection of EMG activity.
696 *Perceptual and motor skills*, 113, 982-994.

697 Prut, Y., & Fetz, E. E. (1999) Primate spinal interneurons show pre-movement instructed delay
698 activity. *Nature*, 401,, 590-594.

699 Richards, J. (2008) *Biomechanics in clinic and research*. Churchill Livingstone, London. ISBN 978-
700 0-443-10170-0.

701 Roosink, M., & Zijdewind, I. (2010) Corticospinal excitability during observation and imagery of
702 simple and complex hand tasks: implications for motor rehabilitation. *Behavioural brain*
703 *research*, 213,, 35-41.

704 Roy, F. D., & Gorassini, M. A. (2008) Peripheral sensory activation of cortical circuits in the leg
705 motor cortex of man. *The Journal of physiology*, 586, 4091-4105.

706 Seeley, M. K., Umberger, B. R., & Shapiro, R. (2008) A test of the functional asymmetry hypothesis
707 in walking. *Gait & posture*, 28, 24-28.

708 Shimba, S., Kawashima, N., Ohta, Y., Yamamoto, S., & Nakazawa, K. (2010) Enhanced stretch reflex
709 excitability in the soleus muscle during passive standing posture in humans. *Journal of*
710 *electromyography and kinesiology : official journal of the International Society of*
711 *Electrophysiological Kinesiology*, 20 , 406-412.

712 Smith, D., & Collins, D. (2004) Mental practice, motor performance, and the late CNV. *Journal of*
713 *sport & exercise psychology*, 26, 412-426.

714 Solodkin, A., Hlustik, P., Chen, E. E., & Small, S. L. (2004) Fine modulation in network activation
715 during motor execution and motor imagery. *Cerebral cortex*, 14, 1246-1255.

716 Tsukahara, R., Aoki, H., Yabe, K., & Mano, T. (1995) Effects of premotion silent period on single
717 motor unit firing at initiation of a rapid contraction. *Electroencephalography and clinical*
718 *neurophysiology*, 97, 223-230.

719 Van de Crommert, H. W., Mulder, T., & Duysens, J. (1998) Neural control of locomotion: sensory
 720 control of the central pattern generator and its relation to treadmill training. *Gait & posture*,
 721 7, 251-263.

722 Wriessnegger, S. C., Steyrl, D., Koschutnig, K., Müller-Putz, G. R. (2014) Short time sports exercise
 723 boosts motor imagery patterns: implications of mental practice in rehabilitation programs.
 724 *Frontiers in human neuroscience*, 30, 469.

725 Wright, D. J., Williams, J., Holmes, P. S. (2014) Combined action observation and imagery facilitates
 726 corticospinal excitability. *Frontiers in human neuroscience*, 8, 951.

727 Yue, G., & Cole, K. J. (1992) Strength increases from the motor program: comparison of training
 728 with maximal voluntary and imagined muscle contractions. *Journal of neurophysiology*, 67,
 729 1114-1123.

730

Abeer A. El-Wishy, Eman S. Fayed Effect of Locomotor Imagery Training Added to Physical Therapy
 Program on Gait Performance in Parkinson Patients: A Randomized Controlled Study

731

732 (Chvatal & Ting, 2012 - Voluntary and Reactive Recruitment of Locomotor Muscle Synergies during
 733 Perturbed Walking)

734 Spinal circuitry of sensorimotor control of locomotion

735 David A McCrea Depression of muscle and cutaneous afferent-evoked monosynaptic field
 736 potentials during fictive locomotion in the cat

737 M-C Perreault, S J Shefchyk, I Jimenez, D A McCrea

738 J Physiol. 1999 December 15; 521(Pt 3): 691–703. doi: 10.1111/j.1469-7793.1999.00691.x

739 Marcus Vinicius Stecklow^I; Antonio Fernando Catelli Infantosi^I; Maurício Cagy^{II} EEG changes
 740 during sequences of visual and kinesthetic motor imagery Arquivos de Neuro-Psiquiatria, ISSN
 741 0004-282X, Arq. Neuro-Psiquiatr. vol.68 no.4 São Paulo Aug. 2010
 742 <http://dx.doi.org/10.1590/S0004-282X2010000400015>

743 Hiraoka, 2012. Imagining stumbling inhibits motor-evoked potentials in the soleus muscle"

744 Daroff, 2012, Bradley's Neurology in Clinical Practice

745 Rozand V, Lebon F, Stapley PJ, Papaxanthis C, Lepers R (2016). A prolonged motor imagery session
746 alter imagined and actual movement durations: Potential implications for
747 neurorehabilitation. *Behav Brain Res.* 2016 Jan 15;297:67-75. doi: 10.1016/j.bbr.2015.09.036.
748 Epub 2015 Sep 30.

749

750 . *J Neurosci.* 2011 Jul 20;31(29):10712-20. doi: 10.1523/JNEUROSCI.6032-10.2011.

751 **Muscle fatigue affects mental simulation of** 752 **action.**

753 Demougeot L¹, Papaxanthis C.

754

755 *Neuropsychologia.* 2004;42(7):944-56.

756 **From mental power to muscle power--gaining** 757 **strength by using the mind.**

758 Ranganathan VK¹, Siemionow V, Liu JZ, Sahgal V, Yue GH.

759

760 • G. Nicolas, *et al.* Corticospinal excitation of presumed cervical propriospinal neurones and its
761 reversal to inhibition in humans *J. Physiol. (Lond.)*, 533 (2001), pp. 903–919

762

FIGURE 1
Illustration of tested experimental conditions

TABLE 1
Mean EMG [μ V] reference values ($\pm$ SD) for all tested muscles in default sitting and standing positions

	Gastrocnemius medialis		Gastrocnemius lateralis		Tibialis anterior		Biceps femoris		Semitendinosus		Rectus femoris	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Sitting position	1.35	0.53	1.48	0.52	1.59	0.53	1.3	0.39	1.2	0.4	1.21	0.48
Standing position	6.17	3.72	3.65	1.79	2.45	0.96	2.6	2.57	2.82	3.6	1.72	1.4

771 TABLE 2
772 Normalized elektromyographic activity with respect to reference value for every muscle [%] during
773 gait imagery tasks in sitting and standing position

		Gait imagery		Gait imagery and observation		Gait imagery after gait execution	
		<i>Med</i>	<i>IQR(Q1 – Q3)</i>	<i>Med</i>	<i>IQR (Q1 – Q3)</i>	<i>Med</i>	<i>IQR (Q1 – Q3)</i>
Gastrocnemius lateralis	S	73.15	(58.31–97.48)	69.07	(54.05–92.82)	61.62	(45.73–84.55)
	T	95.33	(85.23–127.63)	87.31	(70.27–95.68)	89.85	(81.15–106.95)
Gastrocnemius medialis	S	80.64	(54.15–97.92)	79.13	(51.56–98.24)	60.22	(45.91–91.24)
	T	97.19	(78.13–129.47)	84.53	(70.58–109.11)	91.09	(75.83–122.77)
Tibialis anterior	S	75.24	(64.25–112.14)	77.7	(62.84–95)	59.53	(50.49–86.9)
	T	96.58	(75.73–119.36)	88.13	(82.11–99.03)	85.34	(70.78–103.78)
Biceps femoris	S	117.9	(91.09–221.63)	101.49	(86.37–151.14)	104.77	(82.31–129.04)
	T	93.5	(88.57–103.43)	91.48	(82.49–102.03)	85.86	(78.97–98.64)
Semitendinosus	S	92.26	(78.35–108.78)	88.40	(76.73–102.7)	87.33	(76.62–107.02)
	T	111.28	(89.03–158.43)	99.1	(87.29–129.14)	98.15	(71.07–148.37)
Rectus femoris	S	91.32	(86.17–106.95)	90.33	(82.06–100.34)	90.83	(75.08–104.5)
	T	111.11	(93.8–270.79)	98.3	(84.09–156.77)	97.24	(78.19–154.44)

S – sitting position, T – standing position, Med – median, (*Q1 – Q3*) – (25th – 75th percentile of data)

774
775
776
777
778
779
780
781

782 TABLE 3
783 Results of statistical analysis (Wilcoxon signed rank test and Effect Size) of changes in the muscle
784 activity during gait imagery tasks

		gait imagery tasks in the sitting position compare to default sitting rest position			gait imagery tasks in the standing position compare to default standing rest position			gait imagery tasks in the sitting position compare to gait imagery tasks in the standing position		
		<i>Wilcoxon's</i>		<i>Effect</i> <i>Size</i>	<i>Wilcoxon's</i>		<i>Effect</i> <i>Size</i>	<i>Wilcoxon's</i>		<i>Effect</i> <i>Size</i>
		<i>Z</i>	<i>p</i>		<i>Z</i>	<i>p</i>		<i>Z</i>	<i>p</i>	
Gastrocnemius medialis	I1	3.00	<.001	0.41	0.29	0.77	0.04	2.21	0.03	0.3
	I2	3.15	<.001	0.43	1.78	0.08	0.24	1.42	0.16	0.19
	I3	4.08	<.001	0.56	0.29	0.77	0.04	3.99	<.001	0.54
Gastrocnemeius lateralis	I1	3.29	<.001	0.45	0.65	0.52	-0.08	4.30	<.001	0.58
	I2	4.04	<.001	0.55	2.71	0.01	0.34	2.79	0.01	0.38
	I3	4.42	<.001	0.6	1.15	0.25	0.16	4.18	<.001	0.59
Tibialis anterior	I1	2.16	0.03	0.29	0.36	0.72	0.05	2.38	0.02	0.32
	I2	2.26	0.02	0.31	2.07	0.04	0.28	1.39	0.16	0.19
	I3	3.89	<.001	0.53	2.81	<.001	0.38	3.08	<.001	0.42
Biceps Femoris	I1	1.71	0.09	0.23	2.16	0.03	-0.29	2.59	0.01	0.35
	I2	3.05	<.001	0.42	1.13	0.26	-0.15	2.64	0.01	0.36
	I3	3.10	<.001	0.42	0.77	0.44	-0.11	1.99	0.05	0.271
Semitendinosus	I1	1.42	0.16	0.19	1.75	0.08	0.24	2.50	0.01	0.34
	I2	3.17	<.001	0.43	0.53	0.60	-0.07	1.80	0.07	0.26
	I3	2.09	0.04	0.28	0,22	0,83	-0.03	1,13	0.26	0.15
Rectus femoris	I1	1.18	0.24	0.16	2.45	0.01	-0.33	3.39	<.001	0.46
	I2	1.95	0.05	0.27	0.86	0.39	-0.12	2.35	0.02	0.32
	I3	1.49	0.14	0.2	0.26	0.79	-0.04	0.96	0.34	0.13

I1 - gait imagery, I2 - gait imagrey and observation, I3 - gait imagery after gait execution