

Central Lancashire Online Knowledge (CLoK)

Title	Quantification of chemotaxis-related alkane accumulation in <i>Acinetobacter baylyi</i> using Raman microspectroscopy
Type	Article
URL	https://clock.uclan.ac.uk/17225/
DOI	https://doi.org/10.1021/acs.analchem.6b02297
Date	2017
Citation	Li, Hangbing, Martin, Francis L and Zhang, Dayi (2017) Quantification of chemotaxis-related alkane accumulation in <i>Acinetobacter baylyi</i> using Raman microspectroscopy. <i>Analytical Chemistry (including News & Features)</i> , 89 (7). pp. 3909-3918. ISSN 0003-2700
Creators	Li, Hangbing, Martin, Francis L and Zhang, Dayi

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1021/acs.analchem.6b02297>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Quantification of chemotaxis-related alkane accumulation in *Acinetobacter baylyi* using Raman microspectroscopy

Hanbing Li, Francis Luke L Martin, and Dayi Zhang

Anal. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.analchem.6b02297 • Publication Date (Web): 03 Mar 2017

Downloaded from <http://pubs.acs.org> on March 3, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



**Quantification of chemotaxis-related alkane accumulation in
Acinetobacter baylyi using Raman microspectroscopy**

Hanbing Li^a, Francis Luke L Martin^{a,b}, Dayi Zhang^{a,*}

^a Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

^b School of Pharmacy and Biomedical Sciences, University of Central Lancashire,
Preston PR1 2HE, UK

***Corresponding author** Dr Dayi Zhang, Lancaster Environment Centre, Lancaster
University, Lancaster, LA1 4YQ, UK; Tel.: +44(0)1524510288 ; Fax:
+44(0)1524510082 ; Email: d.zhang@lancaster.ac.uk

Abstract

Alkanes are one of the most widespread contaminants in the natural environment, primarily as a consequence of biological synthesis and oil spills. Many indigenous microbes metabolize alkanes, and the chemotaxis and accumulation in some strains has been identified. For the first time, we apply Raman microspectroscopy to identify such chemotaxis-related affinity, and quantify the alkane concentrations *via* spectral alterations. Raman spectral alterations were only found for the alkane chemo-attractant bacteria *Acinetobacter baylyi* ADP1, not for *Pseudomonas fluorescense*, which exhibits limited chemotaxis towards alkane. The significant alterations were attributed to the strong chemotactic ability of *A. baylyi* enhancing the affinity and accumulation of alkane molecules on cell membranes or cellular internalization. Spectral fingerprints of *A. baylyi* significantly altered after 1-h exposure to pure alkanes (dodecane or tetradecane) and alkane mixtures (mineral oil or crude oil), but not monocyclic aromatic hydrocarbons (MAHs) or polycyclic aromatic hydrocarbons (PAHs). A semi-log linear regression relationship between Raman spectral alterations and alkane concentrations showed its feasibility in quantifying alkane concentration in environmental samples. Pure alkanes or alkane mixtures exhibited different limits of detection and regression slopes, indicating that the chemotaxis-related alkane accumulation in *A. baylyi* is dependent on the carbon chain length. This work provides a novel biospectroscopy approach to characterize the chemotaxis-related alkane bioaccumulation, and has immense potential for fast and high-throughput screening bacterial chemotaxis.

Keywords: *Acinetobacter baylyi*; Chemotaxis; Alkane; Affinity; Bioaccumulation; Raman microspectroscopy; Spectral fingerprints

Introduction

Many hazardous chemicals have been released into the environment through various industrial activities. Particularly, with industrial development and urbanization processes, the increasing usage of crude oil has consequently been associated with numerous oil spill accidents and contaminated sites. Since 1969, there have been >40 large oil spill incidents worldwide, including the Exxon Valdez oil spill in Prince William Sound in 1989^{1,2}, the Deepwater Horizon oil spill in Gulf of Mexico³ and the Xingang oil spill in Dalian⁴ in 2010. These incidents have resulted in large areas of oil-contaminated sites, damaged ecological systems and threats to human health^{5,6}. Chemical analysis of alkanes predominantly includes gravimetric⁷⁻⁹, fluorophotometry^{10,11}, infrared (IR) absorption¹², gas chromatography flame ionization detector (GC-FID)¹³ or gas chromatography mass spectrometry (GC-MS) methods¹⁴, but samples require pre-treatment using solvent extraction or solid-phase microextraction (SPME)¹⁵. To evaluate alkane bioavailability or bioaccessibility, some whole-cell bioreporters have been developed¹⁶ and even applied in real-world cases^{4,17}, but they suffer from relatively low reproducibility. Considering the rapid response required to oil spill incidents, it is necessary to develop some real-time and non-destructive screening tools to achieve fast and reliable detection of crude oil contamination.

In the natural environment, alkanes are the main components of crude oil and carbon sources for many bacteria, and alkane chemotaxis has been found in some species¹⁸. Chemotaxis-related affinity to alkanes facilitates bacterial movement in alkane gradients and contributes to their elevated abilities to access and utilize alkane substrates¹⁹. In *Pseudomonas aeruginosa* PAO1, the *tlpS* gene, which is located downstream of the alkane monooxygenase *alkB1* gene, is believed to encode a methyl-accepting chemotaxis protein (MCP) that confers the chemotactic response to hexadecane²⁰. Another *Pseudomonas* strain, *P. putida* GPo1, has the *alkN* gene encoding an MCP for alkane chemotaxis²¹. *Flavimonas oryzae* is reported to exhibit weak chemotactic responses to hexadecane but its mechanisms remain unknown²². The alkane chemotaxis machinery of marine alkane degrader *Alcanivorax dieselolei* consists of eight cytoplasmic chemotaxis proteins, including MCP and flagellar proteins²³. *Acinetobacter baylyi* is the bacterial species with the strongest alkane chemotaxis and accumulation^{16,24-26}. In *A. baylyi*, the putative gene encoding

an MCP is *chpA* which has high similarity to *cheY* (fused chemotactic sensory histidine kinase) and *ompR* (two-component response regulator) in *Escherichia coli*, suggesting potentially similar chemotaxis machinery. *A. baylyi* cells have high affinity only to alkane droplets, but not to other hydrophobic molecules¹⁶, suggesting that the MCP has high recognition specificity to linear alkanes. Meanwhile, alkane chemotaxis further helps microbial accumulation, internalization and degradation of alkanes by providing enhanced opportunities to seek out linear alkanes. The fimbriae on the surface of *Acinetobacter* cells are postulated to drive the access to alkane droplets for cellular uptake²⁷⁻²⁹. Trapped alkane droplets on the cell surface are subsequently transported into bacterial cells after being emulsified by biosurfactant mediation³⁰⁻³².

Raman microspectroscopy has been widely applied in biological molecule detection, as a fast, reproducible and non-destructive approach generating a chemical fingerprint³³. It has been well developed to identify bacteria and study their responses to their environment³⁴⁻³⁷. Most spectrally-identified biological molecules are well characterized, such as purine, pyrimidine, amino acids and proteins³⁸⁻⁴². Though many approaches, such as swarm plates⁴³, capillary assays⁴⁴, temporal stimulation of tethered cells⁴⁵ and automated tracking of swimming cells⁴⁶, have been applied to test the bacterial chemotaxis towards specific chemicals, there is still no study exploring Raman spectral alterations during chemotaxis-related affinity and bioaccumulation processes. The chemotactic behaviour of alkanes hints at the possibility that alkane molecules will accumulate on the membrane and/or in the cytoplasm of bacterial cells, potentially allowing them to be identified in Raman spectra.

In the present study, we used the Raman microspectroscopy to distinguish the spectral alterations and quantify alkane concentrations based on the chemotaxis-related alkane accumulation of *A. baylyi*. For the first time, we demonstrate that chemotaxis-related alkane affinity and accumulation is measurable by Raman microspectroscopy and the MCP of *A. baylyi* is highly specific to alkane molecules, instead of monocyclic aromatic hydrocarbons (MAHs) or polycyclic aromatic hydrocarbons (PAHs). Not only contributing to fast quantification of aqueous alkane concentrations, this method also has the potential as a screening tool in characterising chemotaxis-related affinity and accumulation towards various chemicals.

Experimental section

Strains and Growth Conditions

The two bacterial strains used in this study were the alkane chemotactic *Acinetobacter baylyi* ADP1 and chemotaxis-negative *Pseudomonas fluorescence*. Both strains were grown in minimal medium with 20 mM sodium succinate as the sole carbon source, at 30°C and 150 rpm for 16 h. The 1.0 litre minimal medium contained 1.0 g (NH₄)₂SO₄, 2.5 g KH₂PO₄, 0.1 g MgSO₄·7H₂O, 0.005 g FeSO₄·7H₂O, 0.25 g nitrilotriacetic acid (NTA), 0.55 g NaOH, and 1 mL Bauchop and Elsdén solution. Strains were further centrifuged at 4,000 rpm for 5 min and washed three times using sterile deionized water, and finally suspended in minimal medium for further experimentation.

Hydrocarbon exposure

Unless specifically stated otherwise, all chemicals in this study were of analytical grade and purchased from Sigma Aldrich (UK). The hydrocarbons included six pure chemicals: dodecane, tetradecane, toluene, xylene, phenanthrene and naphthalene; and, two hydrocarbon mixtures: crude oil and mineral oil. They were dissolved in dimethyl sulfoxide (DMSO) to prepare the 10 g/L stock solution, and serially diluted to 0.1-5.0 g/L. The hydrocarbon stock solution was separately mixed with bacterial suspensions (1:100, v/v) to reach a final concentration of 100 mg/L. After 1-h exposure at 30°C, the medium was removed by 5 min centrifugation at 5,000 rpm, and the cell pellets were re-suspended in deionized water and washed twice.

The time-dependence of chemotaxis-related chemo-association was measured by exposing *A. baylyi* suspensions to dodecane, tetradecane, crude oil or mineral oil for 0.5, 1, 3 and 5 h at 30°C, respectively. The final alkane concentrations were 100 mg/L. The cells were further centrifuged at 5,000 rpm for 7 min, and washed twice by deionized water.

To establish the dose response of alkane accumulation, different volumes of dodecane, tetradecane, crude oil or mineral oil stock solution were added to 1 mL washed *A. baylyi* suspensions. The final concentrations were: 1, 2, 5, 10, 20, 50 and 100 mg/L. After 1-h exposure at 30°C, the mixtures were centrifuged at 5,000 rpm for 7 min, and washed twice by deionized water.

Hydrocarbon chemotaxis capillary test

To validate chemotaxis effects, conventional capillary tests were applied according to Alder's protocol⁴⁷ with some modifications. Briefly, the capillary tubes (internal diameter of 0.2 mm and length of 10 cm) were plunged into 1 mL chemotaxis medium with attractant (alkanes, PAHs, MAHs and oils). After approximately 10 min, the liquid was drawn up to approximately 1 cm the length of the tube, the capillary was then inserted (without rinsing) into the bacterial suspension (*A. baylyi* or *P. fluorescence*) and incubated for 1 h at 30°C. To test the time-dependency of chemotaxis, the capillary tubes with tetradecane or crude oil attractants were incubated within the *A. baylyi* suspension for 0.5, 1, 3 or 5 h at 30°C, similar to the hydrocarbon exposure experiment described above.

Instead of plate counting, we used quantitative chain polymerase reaction (qPCR) to quantify the 16S rRNA copy numbers of chemotactic bacterial in duplicate. After 1-h incubation, the capillary tube was removed and the exterior of the open end was directly plunged into qPCR buffer. The 10 µL qPCR buffer consisted of 1 µL of primer 314F (5'-CCTACGGGNGGCWGCAG-3'), 1 µL of primer 802R (5'-TACNVGGGTATCTAATCC-3'), 3 µL molecular water and 5 µL iTaq™ Universal SYBR® Green Supermix (BioRad, USA). The thermos cycling program was run complete except for the extra fluorescence data acquisition at 80°C for 15 s in each cycle. This was: initial denaturation at 94°C for 3 min; 34 amplification cycles of 94°C for 45 s, 52°C for 45 s, 72°C for 45 s, and fluorescence data acquisition at 80°C for 15 s. Standard curves were obtained with serial dilutions of quantified plasmid DNA (*via* nanodrop) containing the fragment of 16S rRNA, as described previously⁴⁸.

Chemical analysis of hydrocarbons

After exposing chemotactic *A. baylyi* suspensions to hydrocarbon solutions, the determination of residual hydrocarbons at different time points was achieved by following the hexane extraction method⁴⁸. At 0, 0.5, 1, 3 or 5 h, the 50 mL cell-hydrocarbon suspensions were added with 50 mL hexane and subsequently ultrasonically homogenized for 2 min (40 kHz). The supernatant was fractionized by column chromatography, in which the hexane supernatant was loaded onto a glass column (Φ10 mm × 100 mm, consisting of 2 cm anhydrous Al₂O₃ and 0.3 cm anhydrous Na₂SO₄ from the bottom to the top) and washed with 50 mL hexane. The

elution was evaporated in 40°C water bath and the hydrocarbons were measured by the gravimetric method.

Raman Microspectroscopy Analysis

Ten μL of washed cells were transferred onto a slide covered with aluminium foil and air-dried before Raman microspectroscopy analysis⁴⁹. Raman spectra were obtained using an InVia confocal micro-Raman system (Renishaw, Gloucestershire, UK) equipped with a 100 mW 785 nm laser diode with a Rayleigh holographic edge filter. The spectrometer's entrance slit of 50 μm combined with a 1200 lines per mm (1 cm^{-1} spatial resolution) diffraction grating allowed dispersion of Raman signals onto a Master Renishaw Pelletier cooled charged couple detector (CCD). A white light camera mounted on the microscope allowed the use of dark-field images to visualize locations for spectral acquisition. An attached microscope (Leica Microsystems, Milton Keynes, UK) with $\times 50$ objective (0.75 numerical aperture; $\approx 1\text{ }\mu\text{m}$ spatial resolution) was utilized for sample detection and acquisition. All sample spectra were obtained using 100% laser power (26 mW at sample), 15 sec acquisition time and one accumulation within a spectral range from 500 to 2000 cm^{-1} . The Renishaw system was calibrated with a Renishaw silicon calibration source for wavenumber shifts. All the treatments were carried out in triplicates, and at least twenty biological replicates were performed and analysed for each sample.

Raman Spectra Analysis

Unless otherwise stated, all data were normalized and analysed using the IRootLab toolbox for Matlab (version R2013b, MathWorks, USA)⁵⁰. Principal component analysis (PCA) was carried out to reduce the dimensionality of the multivariate data and allow visualization of the natural variance within the dataset⁵¹. Post-exposure to different alkane or oil concentrations, the distance of individual Raman spectra to that of comparator negative controls (alkane/oil concentration = 0 mg/L) was calculated based on the values of principal component (PC) 1 (PC1) and PC2. The dispersions of individual group to negative control and pure alkane/oil groups were evaluated by output data derived from PCA and linear discriminant analysis (LDA) and were visualized as dispersion indicator (D_I) scores plots, in which increasing D_I between two categories is proportional to dissimilarity [details see Electronic Supplementary Information (ESI)]. The statistical significance of differences and variance analysis

(P -value <0.05) of LDA among the different treatments was performed using a one-way ANOVA and least significant difference (LSD) test.

Results and Discussion

Raman spectral characterization of A. baylyi exposed to alkanes

Exposed to alkanes or oil droplets, *A. baylyi* bacterial cells were predominantly found on the oil-water interface and could emulsify hydrocarbons into small droplets (see ESI Figure S1). The majority of *A. baylyi* cells were found on the alkane-water interface, not in the aqueous phase, demonstrating their high affinity to alkane droplets. In contrast, the strains negative for or with minor alkane chemotactic properties, like *P. fluorescence*, exhibited a very weak oil droplets affinity and most bacterial cells remained in the water phase (see ESI Figure S1). Similarly, the droplets of alkane mixtures (mineral oil) or hydrocarbon mixtures (crude oil) also showed strong attraction to *A. baylyi*, but no adhesion with *P. fluorescence*. *A. baylyi* ADP1 has been reported with high affinity to alkane droplets¹⁶, explained as alkane chemotaxis^{43,52}. From the results of hydrocarbon capillary assay (Figure. 2A), we confirmed the chemotaxis-related affinity of *A. baylyi* to alkanes. The accumulation of 16S rRNA copies of *A. baylyi* in capillaries in oil mixture and pure alkane treatments ranged from 3.9×10^8 to 1.07×10^9 per capillary, significantly higher than those in PAHs and MAHs treatments (8.7×10^5 to 3.6×10^7 16S rRNA copies per capillary). In contrast, the low 16S rRNA copy numbers of *P. fluorescence* in capillary tubes post-exposure to oil mixtures and pure alkanes (2.9×10^4 to 9.7×10^6 16S rRNA copies per capillary) suggested that *P. fluorescence* is not an alkane chemotactic strain.

The hydrocarbon capillary assay proves that *A. baylyi* ADP1 cells can swim down the attractant gradient towards alkanes. From another perspective, MCP on the cell membranes of *A. baylyi* ADP1 effectively recognizes and binds the alkane molecules, eventually trapping and accumulating the soluble fraction of alkane molecules in the aqueous phase. Considering the low solubility of alkane in water, it generates the alkane gradient and contributes to the access of *A. baylyi* ADP1 cells to alkane molecules. This is strongly suggested by the significant accumulation of *A. baylyi* cells in capillary following exposure to alkane mixtures or pure alkanes (Figure 2A). Alkane chemotaxis significantly helps the internalization and utilization of the insoluble alkanes by *A. baylyi* ADP1¹⁶. The accumulation of alkane molecules either

on cell membranes or internalized within cells provides the opportunity to employ Raman microspectroscopy to investigate the chemotaxis-related affinity and accumulation of alkanes.

Since dodecane and tetradecane are both linear alkane molecules with carbon chain lengths of 12 and 14 respectively, their Raman spectra are similar (Figure 1A and Table 1, see ESI Figure S2A). The predominant bands include 890 cm^{-1} (terminal methyl CH_3 rock), 1133 cm^{-1} (C-C asymmetric stretch and CH_2 wag), 1297 cm^{-1} (CH_2 twist) and 1439 cm^{-1} (CH_2 bend)⁵³. The only difference between dodecane and tetradecane is the bands at 961 cm^{-1} and 1033 cm^{-1} in dodecane, which are attributed to CH_2 rock⁵⁴. As the distillation product of crude oil, the Raman spectra of mineral oil (see ESI Figure S2C) show distinct peaks at bands 1058 cm^{-1} and 890 cm^{-1} , assigned to CH_2 twist-rock and C-C stretch, respectively⁵⁵. Although the CH_2 twist peak at 1297 cm^{-1} is observed for both crude oil and mineral oil, crude oil also contains MAHs and PAHs, and therefore possesses three unique peaks (Figure 1B): benzene ring breathing band 1060 cm^{-1} , symmetric and antisymmetric CH_2 scissors vibration bands 1437 cm^{-1} and 1459 cm^{-1} ⁵⁶.

Table 1. Variations and assignments of Raman bands of alkanes and *A. baylyi* before and after exposure.

Band (cm ⁻¹)	Tentative bands assignment	Origin	Alkane	<i>A. baylyi</i>	Alkane + <i>A. baylyi</i>
890	Terminal methyl CH ₃ rock	Dodecane, tetradecane, mineral oil	++	-	+ ↑
961	CH ₂ rock	Dodecane	++	-	+ ↑
1033	CH ₂ rock	Dodecane	+	-	-
1058	C-C stretch	Mineral oil	++	+	+
1133	C-C asymmetric stretch and CH ₂ wag	Dodecane, tetradecane, crude oil, mineral oil	+++	+	+
1297	CH ₂ twist	Dodecane, tetradecane, crude oil	+++	+	+++ ↑
1437	Symmetric CH ₂ scissors vibration	Crude oil	+++	-	+++ ↑
1439	CH ₂ bend	Dodecane, tetradecane, mineral oil	+++	+++	+++ ↑
1459	Antisymmetric CH ₂ scissors vibration	Crude oil	+++	-	+++ ↑
723	Ring breathing vibration	Bacterial nucleic acids	-	++	+ ↓
777	Ring breathing vibration	Bacterial nucleic acids	-	++	+ ↓
1002	Benzene ring breathing	Bacterial phenylalanine	-	+++	++ ↓
1238	/	Bacterial amino acids	-	+	- ↓
1311	/	Bacterial amino acids	-	+	- ↓
1441	Deformations of CH ₂	Bacterial glycine	-	+++	+++ ↑
1663	C=C stretching	Bacterial proteins	-	+++	+++

“-” represents no peak identified; “+” represents weak peak; “++” represents medium peak; “+++” represents strong peak. Brown or green symbols indicate characteristic peaks belonging to alkanes and *A. baylyi*, respectively. Red or blue arrows represent the increases or decreases in respective Raman intensity of *A. baylyi* post-exposure to alkanes.




Figure 1. Raman spectra of *A. baylyi* and *P. fluorescence*, pre- and post-exposure to 100 mg/L tetradecane (A) or crude oil (B). PCA of Raman spectra shows the significant segregation of *A. baylyi* categories when exposed to 100 mg/L tetradecane (C) or crude oil (D) due to chemotaxis-related alkane affinity and accumulation, but not *P. fluorescence*. Twenty Raman spectra were randomly obtained per treatment.

The Raman spectra of original *A. baylyi* and *P. fluorescence* cells are similar, due to their close relationship to the order *Pseudomonadales* from genomic and proteomic analyses^{25,26,57}. From Figure 1A, 1B and Table 1, both 777 cm⁻¹ and 723 cm⁻¹ bands refer to nucleic acids⁵⁸. The band at 1441 cm⁻¹ is assigned to glycine in bacterial cells⁴⁰. Amino acids bands are also observed in the Raman spectra at 1238 cm⁻¹ and 1311 cm⁻¹⁵⁸. The 1663 cm⁻¹ band represents protein and the 1002 cm⁻¹ band points to

phenylalanine. Post-exposure to tetradecane or crude oil, entirely different Raman spectra are identified for the two strains. The Raman spectra of *P. fluorescence* remain unchanged, explained by the fact that alkanes exhibit no chemo-attraction and are not accumulated in *P. fluorescence*. Accordingly, all the *P. fluorescence* cells fall into one category after PCA (Figure 1C and 1D) and are indistinguishable pre- and post-exposure to tetradecane or crude oil. The Raman spectra of *A. baylyi* exhibit significant alterations post-exposure to alkanes (Figure 1A, 1B and Table 1). Post-exposure bacterial spectra are coupled with some key alkane bands. Two strong alkane bands, 1297 cm^{-1} (CH_2 twist) and 1439 cm^{-1} (CH_2 bend), are observed in all Raman spectra of *A. baylyi* exposed to tetradecane or crude oil.

In dodecane treatment, the specific 961 cm^{-1} and 1033 cm^{-1} bands are attributed to CH_2 rock (Table 1). The bands 1058 cm^{-1} (CH_2 twist-rock) and 890 cm^{-1} (C-C stretch) only appear in *A. baylyi* treated with mineral oil (see ESI Figure S2D), and the bands 1060 cm^{-1} (ring breathing of benzene) and $1430\text{--}1470\text{ cm}^{-1}$ (symmetric and antisymmetric CH_2 scissors vibrations) are found in the spectra of *A. baylyi* exposed to crude oil (Figure 1B). It is worth noting that both chemotaxis-related affinity and accumulation of alkane might contribute to the appearance of alkane characteristic bands, where alkane affinity significantly accelerates the access, accumulation and potential internalization of alkanes by *A. baylyi* and enhances Raman signals. PCA of Raman spectra (Figure 1C and 1D) further proves that alkane affinity and accumulation result in the significant differentiation of *A. baylyi* post-exposure to tetradecane or crude oil. The category comprising *A. baylyi* treated with tetradecane is very close to that of pure tetradecane, whereas the crude oil exposure leads to similarity between *A. baylyi* and crude oil. They are both significantly separated from the original *A. baylyi* group. The results show that the significant Raman spectral alterations are caused by the alkane affinity and accumulation of *A. baylyi*. Moreover, the strong terminal methyl CH_3 rock (890 cm^{-1}) is only found in *A. baylyi* spectra post-exposed to tetradecane, readily discriminated from those in crude oil treatment. Therefore, our findings reveal that chemotaxis-related alkane affinity contributes to the adhesion of alkane molecules onto the membrane of *A. baylyi* cells, allowing further alkane accumulation or cellular internalization. Raman spectral alterations highlight such chemotaxis-related affinity and accumulation, and even distinguish the different chemo-attracted molecules quickly and with high reproducibility.

Specific chemotaxis-related alkane affinity and accumulation in A. baylyi

To test the specificity of the hydrocarbons affinity and accumulation in *A. baylyi*, various hydrocarbon molecules were tested, including pure alkanes (dodecane or tetradecane), alkane mixtures (mineral oil or crude oil), MAHs (toluene or xylene) and PAHs (naphthalene or phenanthrene). Raman spectral alterations are only found for pure alkanes or alkane mixtures (Figure 2B, see ESI Figure S2). PCA differentiation illustrates that the categories for post-exposure to MAHs or PAHs co-cluster due to significant Raman bands for benzene ring breathing peaks at 1001 cm^{-1} and 1027 cm^{-1} for toluene or xylene, 760 cm^{-1} for naphthalene and 705 cm^{-1} for phenanthrene (see ESI Figure S2)⁵⁹⁻⁶¹. The category for crude oil is located between pure alkanes and MAHs/PAHs, because it consists of various hydrocarbons, including alkanes, MAHs and PAHs. As a by-product from crude oil distillation, mineral oil is a mixture of n-alkanes and located very close to the category for pure alkanes⁶². Post-exposure to MAHs (toluene, xylene) or PAHs (phenanthrene and naphthalene), PCA of *A. baylyi* fails to segregate these categories from unexposed comparator cells. The results were consistent with hydrocarbon capillary assay (Figure 2A) in which the accumulated 16S rRNA copy numbers of *A. baylyi* were much higher in oil mixtures and pure alkanes treatments than those in PAHs and MAHs treatments, indicating that *A. baylyi* has chemo-attraction towards neither MAHs nor PAHs, which has been proved previously^{16,24}. After PCA, the categories for *A. baylyi* exposed to pure alkane or alkane mixtures co-cluster with those of the pure chemicals respectively, and are completely separated from the unexposed comparator cells. These results prove that the chemotactic association between alkane and *A. baylyi* is the key factor in altering Raman spectra. Such chemotaxis-related affinity is specific to alkanes and helps further alkane accumulation or internalization within *A. baylyi* cells, resulting in significant spectral alterations compared to those induced by other hydrophobic molecules (MAHs or PAHs). Thus, we conclude from the Raman spectral discrimination that the alterations in *A. baylyi* exposed to mineral oil or crude oil is attributable to chemotaxis-related affinity towards alkanes; these are the main components of mineral oil and crude oil.

Previous research has examined the mechanisms of alkane chemotaxis. In the presence of alkanes, OmpS is most likely the first sensor protein to transmit signals back to bacterial cells⁶³. The chemotactic signals are stimulated by a complex protein

system that is comprised of coupling CheW and methyl-accepting chemotaxis (MCP) to the histidine kinase CheA⁶⁴⁻⁶⁶. It is also reported that the MCP is captured firmly in this protein complex, whereas a substantial part of CheW and CheA is free in cells⁶⁷. The signals from OmsP are likely received by the MCP protein embedded on CheW/CheA complex. The coupling protein CheW2 is found in *Alcanivorax dieselolei* as the chemotaxis complex only induced by long-chain alkanes⁶⁸⁻⁷¹. Hence, bacterial chemotaxis towards alkanes is hypothesized to exhibit high specificity, and the chemotaxis-related alkane affinity and accumulation therefore follows the same behaviour. Our results provide direct evidence that Raman spectral alterations of *A. baylyi* are only observed for linear n-alkanes, not MAHs or PAHs.




Figure 2. (A) Validation of alkane chemotaxis *via* capillary assay. High amount of bacterial 16S rRNA in capillary indicates significant chemotaxis of *A. baylyi* (red bar) towards dodecane, tetradecane, mineral oil or crude oil, whereas *P. fluorescence* (blue bar) does not show chemotaxis to these hydrocarbons. (B) Raman spectral segregation by PCA of *A. baylyi* exposed to different hydrocarbons. The tested chemicals with alkane-chemotaxis effect include: dodecane, tetradecane, mineral oil or crude oil. The tested chemicals without chemotaxis effect were: monocyclic aromatic hydrocarbons (toluene or xylene) and polycyclic aromatic hydrocarbons (naphthalene or phenanthrene). Twenty Raman spectra were randomly obtained per treatment.

Time-dependent Raman spectra alteration

To identify the optimal exposure time of *A. baylyi* to alkanes, bacterial cells were treated with 100 mg/L tetradecane or crude oil for 0.5, 1.0, 3.0 or 5.0 h. The results from PCA (Figure 3) indicate that the Raman spectral alterations of *A. baylyi* alkane association are time-dependent. With the sensitive chemotactic affinity of *A. baylyi* towards alkanes, significant spectral alterations are observed within 0.5 h for both tetradecane or crude oil, and the most discrimination is at 1.0 h. Interestingly, after 0.5 h exposure, the cluster of *A. baylyi* cells treated with tetradecane exhibits most discrimination compared to the untreated category (Figure 3A) *versus* cells exposed to crude oil (Figure 3B). Thus affinity of tetradecane appears stronger than crude oil. This is very likely explained by the complex composition of crude oil, which consists of alkanes, MAHs and PAHs. On the surface of crude oil droplets, all these molecules are evenly distributed, leaving fewer active sites of chemo-attractive alkanes for the effective recognition by the MCP of *A. baylyi*. The diverse hydrophobic components in crude oil droplets reduce alkanes bioavailability, challenge the access of bacteria to alkane molecules⁷² and slow down alkanes utilization by *A. baylyi*⁷³. The accessibility of *A. baylyi* to tetradecane is therefore faster and such stronger adhesion to pure alkane droplets has been reported⁷⁴⁻⁷⁶.




Figure 3. PCA segregation of *A. baylyi* Raman spectra against exposure time in tetradecane (A) and crude oil (B) treatments. Twenty Raman spectra were randomly obtained per treatment.

A. baylyi has the alkane monooxygenase encoding *alkM* gene and can metabolize n-alkanes¹⁶. The activation of *alkM* is reported to occur within 30 min post-exposure to alkanes, when significant alkane degradation occurs^{66,74}. The alkane molecules accumulated on cell membrane or internalized inside *A. baylyi* cells were therefore hypothesized to be rapidly consumed. Our test on residual hydrocarbon concentrations (see ESI Figure S3) proves this hypothesis that the dramatic alkane degradation was only observed within the first 1 hour and its concentration remained stable thereafter. Thus, the amount of alkanes captured and accumulated by *A. baylyi* cells peaked at 1 hour and resulted in the most discrimination in Raman spectral fingerprints (Figure 3). This explains the diminishing spectral alterations after 1-h exposure. As a consequence, the Raman spectral categories of 0.5, 3 and 5 h exposure are clustered close to the original *A. baylyi* category in either tetradecane or crude oil treatments (Figure 3, see ESI Figure S4). Thus, the highest alkane affinity and accumulation of alkanes was at 1.0 h, at which time the strongest spectral alterations are found. Therefore, 1-h exposure was applied for all the subsequent experiments. Compared to the conventional hydrocarbon chemotaxis capillary assay (see ESI Figure S5) in which the maximum bacterial enrichment was 0.5 h for crude oil and 3 h for tetradecane, our Raman assay had different response time (1.0 h for both crude oil and tetradecane). It might be explained by the different chemotaxis-related behaviour quantified by the two methods. The chemotaxis capillary assay measures the chemotaxis-driven bacterial migration in alkane gradients, whereas the Raman spectral alteration represents the chemotaxis-related affinity and accumulation of alkane molecules on *A. baylyi* cells. It is worth highlighting that the Raman assay is a faster and higher throughput interrogation of specific chemical affinity and accumulation than the conventional capillary assay. Though the chemotactic response is around 1 hour in both methods, the Raman assay takes only 20 min for wash/dry and 1-2 min spectral measurement, much less than that required in determining bacterial enrichment by the capillary assay (3-4 hours by qPCR or 18 hours by plate counting). Additionally, the attractant only needs to be mixed together with targeting bacterial suspension, easier than the plunge and transfer of attractant medium in numerous capillaries in the conventional assay, allowing high-throughput detection for many chemicals in 96-well plates.

Quantification of alkane affinity and accumulation by Raman spectral alterations

Raman spectral alterations of *A. baylyi* chemotaxis-related affinity and accumulation towards alkane allow evaluation of the sensitivity of alkane affinity and to quantify alkane concentration in aquatic phase. Figure 4 illustrates the PCA categorisation of *A. baylyi* Raman spectra following different exposures (1, 2, 5, 10, 20, 50, 100 mg/L) of dodecane, tetradecane, mineral oil or crude oil, respectively. After 1-h exposure to the targeted alkane molecules or mixtures, all the characteristic peaks of *A. baylyi* remained the same, and the new altered peaks originated from alkanes (see ESI Figure S6). The intensity of alkane characteristic peaks increased with increasing dodecane (or tetradecane) exposure, especially for the band at 1297 cm^{-1} which is annotated to CH_2 twist and dominates in alkane molecules. The Raman peak of CH_2 bend at 1493 cm^{-1} was positively related to the alkane concentrations. The terminal methyl CH_3 Raman signal of alkane at 890 cm^{-1} was detected when the concentrations of dodecane and tetradecane was 20 mg/L or higher. Besides pure alkanes, similar Raman spectral alterations were found for treatments with alkane mixtures (mineral oil or crude oil; Figure 4C and 4D). Spectral alternations of *A. baylyi* are separated post-exposure to different concentrations of mineral oil or crude oil. The key altered peaks of mineral oil and crude oil include the 1297 cm^{-1} and 1439 cm^{-1} bands representing CH_2 twist and bend, respectively (see ESI Figure S7). The consistent alteration (1297 cm^{-1} and 1439 cm^{-1}) in all bacterial spectra exposed to alkanes and alkane mixtures prove that *A. baylyi* has specific affinity towards alkane molecules^{16,77}.

The results of LDA and dispersion indicator (D_I) scores plots show the specific chemotaxis-related affinity and accumulation of *A. baylyi* towards different alkanes (Figure 5). There is no significant discrimination between the categories of *A. baylyi* exposed to 0, 1 or 2 mg/L dodecane (Figure 5A), while *A. baylyi* treated with 1 mg/L tetradecane is completely segregated from the negative control (0 mg/L, Figure 5B). This indicates that the limit of alkane detection *via* Raman microspectroscopy is 5 mg/L for dodecane and 1 mg/L for tetradecane. Although *A. baylyi* can respond to alkane levels as low as 0.1 mg/L by bioreporter data¹⁶, the lower sensitivity of Raman spectra is possibly attributed to the weak signal of accumulated or internalized alkane molecules. At low alkane concentrations, only partial MCP recognizes and captures

alkane molecules. Similar behaviour is observed for alkane mixtures, with a limit of detection of 1 mg/L for mineral oil and 2 mg/L for crude oil.




Figure 4. PCA segregation of *A. baylyi* Raman spectra at different concentrations of dodecane (A), tetradecane (B), mineral oil (C) or crude oil (D). For each treatment, twenty replicates were performed.

With increasing alkane levels, the more alkane molecules accumulate on or inside *A. baylyi* cells and consequently result in increasing alkane characteristic peaks and more significant Raman spectral alterations. Above the limit of detection, all the alkanes or alkane mixtures gave rise to a semi-log linear regression relationship with D_I (see ESI Figure S8). The semi-log linear slope of pure alkanes and alkane mixtures followed the order: mineral oil (0.583) > crude oil (0.532) > tetradecane (0.478) >

dodecane (0.399). The quantification range was from 1 mg/L to 100 mg/L for tetradecane and mineral oil, 2 mg/L to 100 mg/L for crude oil and 5 mg/L to 100 mg/L for dodecane. The values of semi-log linear slopes were consistent with the limits of detection for all the pure alkanes and alkane mixtures. It indicates the selectivity of alkane affinity and accumulation in *A. baylyi* varies with different carbon chain lengths. For instance, the *cheY1* and *cheY2* genes in *Alcanivorax dieselolei* are triggered by C₈-C₂₄ n-alkanes and alkanes longer than C₂₄, respectively⁷¹. In this study, *A. baylyi* had higher affinity towards and subsequently accumulation of tetradecane than dodecane, due to a lower limit of detection (1 mg/L vs. 5 mg/L) and higher slopes (0.478 vs. 0.399). The expression of genes related to alkane affinity and internalization in *A. baylyi* is therefore dependent on alkane carbon chain length, and it can be evaluated and quantified by Raman spectra.




Figure 5. D_I scores plots of *A. baylyi* Raman spectra following different concentrations of dodecane (A), tetradecane (B), mineral oil (C) or crude oil (D). For each treatment, twenty replicates were performed. Boxes with different letters indicated a significant difference based on LSD (P -value <0.05).

The differing spectral alteration in mineral oil and crude oil treatments suggests that *A. baylyi* exhibits the stronger sensitivity of chemotaxis-related affinity and accumulation for mineral oil than crude oil. Considering the composition of the two alkane mixtures, mineral oil is fully composed of alkanes with different carbon chain lengths, whereas alkanes only account for 20-25% of the total weight of crude oil. Not all the molecules of crude oil can be chemotactically sensed by *A. baylyi* cells, consequently resulting in the lowest affinity and accumulation. More interestingly from linear slopes, both tetradecane and dodecane have lower affinity than mineral oil and crude oil. From the discussion above, the MCP recognition and expression is dependent on the carbon chain length of alkanes. For instance, the expression of *alkM* alkane monooxygenase gene in *A. baylyi* is more highly induced by dodecane than tetradecane, suggesting that the *alkR* encoding AraC-XylS-like transcriptional regulators has unique sensitivity towards alkanes with different carbon chain length^{29,36,75,78}. Thus, *A. baylyi* is hypothesized to have higher affinity towards some other alkane molecules with different carbon chain lengths in mineral oil or crude oil, and Raman spectral alterations of *A. baylyi* are enhanced post-exposure to mineral oil and crude oil.

Though the current detection limit of Raman assay is still above the natural hydrocarbon contamination level (0.1-1 mg/L) and does not meet with the monitoring requirement for environmental hydrocarbons, it can be improved by optimizing attractant medium, such as pH and ionic strength. Meanwhile, alkane affinity and accumulation has been extensively investigated, but there is a lack of effective approaches to quantify its sensitivity, which is one of restriction factors in bioremediation at oil contaminated sites. Our results successfully prove the concept that Raman spectral alterations provide a rapid, non-destructive tool towards understanding alkane affinity and accumulation. More research is required to address the mechanisms of chemotaxis-related alkane affinity and accumulation to a wider range of alkanes with differing carbon chain lengths.

Conclusion

Herein, we developed a novel alkane quantification method based on chemotaxis-related alkane affinity and accumulation of *A. baylyi* and Raman spectral alterations. To the best of our knowledge, this is the first study to show that Raman microspectroscopy can be used to identify and quantify the microbial chemotaxis-

related affinity and accumulation of alkanes. It opens up new possibilities for the rapid biological detection of aquatic alkane concentrations with natural microorganisms, instead of gene-modified bioreporters. Importantly, alkane chemo-affinity facilitates bacterial access to alkanes, and internalization determines alkane metabolism inside bacterial cells, both playing a key role in alkane degradation in the natural environment. The effective quantification of alkane affinity and accumulation *via* Raman spectra contributes to new insights into the microbial utilization of alkanes. This technique is also feasible to identify and characterize bacterial affinity and accumulation towards other molecules.

Acknowledgement

The authors would like to thank National Natural Science Foundation of China (No. 41301331) and Lancaster University FST research grant for financial support. Hanbing Li is supported by China Scholarship Council (CSC).

Supporting Information

Dispersion indicator model; Chemotaxis of *A. baylyi* and *P. fluorescence* to mineral oil, tetradecane, xylene and phenanthrene (Figure S1); Raman spectra of *A. baylyi* exposure to different hydrocarbons (Figure S2); Relationship between dispersion indicator of Raman spectra and residual hydrocarbon contents (Figure S3); Raman spectra of *A. baylyi* against exposure time (Figure S4); Alkane chemotaxis against time via chemotaxis capillary assay (Figure S5); Raman spectra of *A. baylyi* exposure to different concentrations of dodecane and tetradecane (Figure S6); Raman spectra of *A. baylyi* exposure to different concentrations of mineral oil and crude oil (Figure S7); Semi-log linear regression of the n-alkane concentrations and dispersion indicator within the quantification range (Figure S8).


Reference

- (1) Bence, A. E.; Kvenvolden, K. A.; Kennicutt, M. C. *Org. Geochem.* **1996**, *24*, 7-42.
- (2) Bragg, J. R.; Prince, R. C.; Harner, E. J.; Atlas, R. M. *Nature* **1994**, *368*, 413-418.
- (3) Camilli, R.; Reddy, C. M.; Yoerger, D. R.; Van Mooy, B. A. S.; Jakuba, M. V.; Kinsey, J. C.; McIntyre, C. P.; Sylva, S. P.; Maloney, J. V. *Science* **2010**, *330*, 201-204.
- (4) Zhang, D.; Ding, A.; Cui, S.; Hu, C.; Thornton, S. F.; Dou, J.; Sun, Y.; Huang, W. E. *Water Res.* **2013**, *47*, 1191-1200.
- (5) Peterson, C. H.; Rice, S. D.; Short, J. W.; Esler, D.; Bodkin, J. L.; Ballachey, B. E.; Irons, D. B. *Science* **2003**, *302*, 2082-2086.
- (6) Piatt, J. F.; Lensink, C. J.; Butler, W.; Kendziorek, M.; Nysewander, D. R. *Auk* **1990**, *107*, 387-397.
- (7) USEPA. **1998**.
- (8) USEPA. **1998**.
- (9) USEPA. **1999**.
- (10) Zhou, Z.; Guo, L.; Shiller, A. M.; Lohrenz, S. E.; Asper, V. L.; Osburn, C. L. *Mar. Chem.* **2013**, *148*, 10-21.
- (11) Wang, C.; Shi, X.; Li, W.; Wang, L.; Zhang, J.; Yang, C.; Wang, Z. *Mar. Pollut. Bull.* **2016**.
- (12) Lay-Ekuakille, A.; Palamara, I.; Caratelli, D.; Morabito, F. C. *Rev. Sci. Instrum.* **2013**, *84*, 015103.
- (13) Krupcik, J.; Gorovenko, R.; Spanik, I.; Bockova, I.; Sandra, P.; Armstrong, D. W. *J. Chromatogr. A* **2013**, *1301*, 225-36.
- (14) Liu, W.; Chen, J.; Lin, X.; Fan, Y.; Tao, S. *Environ. Pollut.* **2007**, *146*, 470-7.
- (15) D'Auria, M.; Racioppi, R.; Velluzzi, V. *J. Chromatogr. Sci.* **2008**, *46*, 339-344.
- (16) Zhang, D.; He, Y.; Wang, Y.; Wang, H.; Wu, L.; Aries, E.; Huang, W. E. *Microbial Biotech.* **2012**, *5*, 87-97.
- (17) Sticher, P.; Jaspers, M. C. M.; Stemmler, K.; Harms, H.; Zehnder, A. J. B.; vanderMeer, J. R. *Appl. Environ. Microbiol.* **1997**, *63*, 4053-4060.
- (18) Wang, W.; Shao, Z. *Front. Microbiol.* **2013**, *4*.
- (19) Parales, R. E.; Harwood, C. S. *Curr. Opin. Microbiol.* **2002**, *5*, 266-273.
- (20) Smits, T. H. M.; Witholt, B.; van Beilen, J. B. *Anton. Leeuw. Int. J. G.* **2003**, *84*, 193-200.
- (21) van Beilen, J. B.; Panke, S.; Lucchini, S.; Franchini, A. G.; Rothlisberger, M.; Witholt, B. *Microbiol. SGM* **2001**, *147*, 1621-1630.
- (22) Lanfranchi, M. P.; Alvarex, H. M.; Studdert, C. A. *Environ. Microbiol.* **2003**, *5*, 1002-1008.
- (23) Lai, Q.; Li, W.; Shao, Z. *J. Bacteriol.* **2012**, *194*, 6674-6674.
- (24) Huang, W. E.; Singer, A. C.; Spiers, A. J.; Preston, G. M.; Whiteley, A. S. *Environ Microbiol* **2008**, *10*, 1668-80.
- (25) Vaneechoutte, M.; Young, D. M.; Ornston, L. N.; De Baere, T.; Nemec, A.; Van Der Reijden, T.; Carr, E.; Tjernberg, I.; Dijkshoorn, L. *Appl. Environ. Microbiol.* **2006**, *72*, 932-936.
- (26) Young, D. M.; Parke, D.; Ornston, L. N. *Annu. Rev. Microbiol.* **2005**, *59*, 519-551.
- (27) Foster, J. *Oxygenases. Academic, New York* **1962**, 241-261.
- (28) Rosenberg, E. *Trends Biotechnol.* **1993**, *11*, 419-424.
- (29) Wentzel, A.; Ellingsen, T. E.; Kotlar, H.-K.; Zotchev, S. B.; Throne-Holst, M. *Appl. Microbiol. Biotechnol.* **2007**, *76*, 1209-1221.

- (30) Boulton, C.; Ratledge, C. *Top. Enzyme Ferment. Biotechnol.* **1984**, *9*, 11-77.
- (31) Haferburg, D.; Hommel, R.; Claus, R.; Kleber, H.-P. In *Bioproducts*; Springer, 1986, pp 53-93.
- (32) Singer, M.; Finnerty, W. **1984**.
- (33) Efrima, S.; Zeiri, L. *J. Raman Spectrosc.* **2009**, *40*, 277-288.
- (34) Efrima, S.; Bronk, B. *J. Phys. Chem. B* **1998**, *102*, 5947-5950.
- (35) Guzelian, A. A.; Sylvia, J. M.; Janni, J. A.; Clauson, S. L.; Spencer, K. M. In *Environmental and Industrial Sensing*; International Society for Optics and Photonics, 2002, pp 182-192.
- (36) Wang, Y.; Ravindranath, S.; Irudayaraj, J. *Anal. Bioanal. Chem.* **2011**, *399*, 1271-1278.
- (37) Zhou, H.; Yang, D.; Ivleva, N. P.; Mircescu, N. E.; Schubert, S. r.; Niessner, R.; Wieser, A.; Haisch, C. *Anal. Chem.* **2015**, *87*, 6553-6561.
- (38) Cheng, H.-W.; Huan, S.-Y.; Yu, R.-Q. *The Analyst* **2012**, *137*, 3601-3608.
- (39) Cheng, H.-W.; Luo, W.-Q.; Wen, G.-L.; Huan, S.-Y.; Shen, G.-L.; Yu, R.-Q. *The Analyst* **2010**, *135*, 2993-3001.
- (40) Cui, L.; Chen, P.; Chen, S.; Yuan, Z.; Yu, C.; Ren, B.; Zhang, K. *Anal. Chem.* **2013**, *85*, 5436-43.
- (41) Liu, Y.; He, L.; Mustapha, A.; Li, H.; Hu, Z.; Lin, M. *J. Appl. Microbiol.* **2009**, *107*, 1193-1201.
- (42) Schuster, K. C.; Reese, I.; Urlaub, E.; Gapes, J. R.; Lendl, B. *Anal. Chem.* **2000**, *72*, 5529-5534.
- (43) Kearns, D. B. *Nat. Rev. Microbiol.* **2010**, *8*, 634-44.
- (44) Berg, H. C.; Turner, L. *Biophys. J.* **1990**, *58*, 919-930.
- (45) Block, S. M.; Segall, J. E.; Berg, H. C. *J. Bacteriol.* **1983**, *154*, 312-323.
- (46) Shi, L. Z.; Nascimento, J.; Chandsawangbhuwana, C.; Berns, M. W.; Botvinick, E. L. *Microsc Res Tech* **2006**, *69*, 894-902.
- (47) Adler, J. *Microbiology* **1973**, *74*, 77-91.
- (48) Wang, X.; Zhao, X.; Li, H.; Jia, J.; Liu, Y.; Ejenavi, O.; Ding, A.; Sun, Y.; Zhang, D. *Res. Microbiol.* **2016**, DOI: 10.1016/j.resmic.2016.07.004.
- (49) Cui, L.; Butler, H. J.; Martin-Hirsch, P. L.; Martin, F. L. *Anal. Methods* **2016**, *8*, 481-487.
- (50) Trevisan, J.; Angelov, P. P.; Scott, A. D.; Carmichael, P. L.; Martin, F. L. *Bioinformatics* **2013**, *29*, 1095-1097.
- (51) Butler, H. J.; Ashton, L.; Bird, B.; Cinque, G.; Curtis, K.; Dorney, J.; Esmonde-White, K.; Fullwood, N. J.; Gardner, B.; Martin-Hirsch, P. L.; Walsh, M. J.; McAinsh, M. R.; Stone, N.; Martin, F. L. *Nat. Protoc.* **2016**, *11*, 664-687.
- (52) Mao, H.; Cremer, P. S.; Manson, M. D. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, *100*, 5449-54.
- (53) Mizushima, S.-i.; Simanouti, T. *J. Am. Chem. Soc.* **1949**, *71*, 1320-1324.
- (54) Kalyanasundaram, K.; Thomas, J. *J. Phys. Chem.* **1976**, *80*, 1462-1473.
- (55) Graham, S. F.; Haughey, S. A.; Ervin, R. M.; Cancouët, E.; Bell, S.; Elliott, C. T. *Food Chem.* **2012**, *132*, 1614-1619.
- (56) Orange, D.; Knittle, E.; Farber, D.; Williams, Q. *Geol. Soc. Spec. Pub.* **1996**, *5*, 65-81.
- (57) Barbe, V.; Vallenet, D.; Fonknechten, N.; Kreimeyer, A.; Oztas, S.; Labarre, L.; Cruveiller, S.; Robert, C.; Duprat, S.; Wincker, P. *Nucleic Acids Res.* **2004**, *32*, 5766-5779.
- (58) De Gelder, J.; De Gussem, K.; Vandenabeele, P.; Moens, L. *J. Raman Spectrosc.* **2007**, *38*, 1133-1147.

- (59) Cloutis, E.; Szymanski, P.; Applin, D.; Goltz, D. *Icarus* **2016**, 274, 211-230.
- (60) Konradi, J.; Singh, A. K.; Materny, A. *Phys. Chem. Chem. Phys.* **2005**, 7, 3574-9.
- (61) Biedermann, M.; Grob, K. *J. Chromatogr. A* **2012**, 1255, 56-75.
- (62) Tarnow, P.; Hutzler, C.; Grabiger, S.; Schon, K.; Tralau, T.; Luch, A. *PLoS One* **2016**, 11, e0147239.
- (63) Wang, S.-K.; Wang, F.; Hu, Y.-R.; Stiles, A. R.; Guo, C.; Liu, C.-Z. *ACS Appl. Mater. Interfaces* **2014**, 6, 109-115.
- (64) Briegel, A.; Li, X.; Bilwes, A. M.; Hughes, K. T.; Jensen, G. J.; Crane, B. R. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, 109, 3766-3771.
- (65) Hazelbauer, G. L.; Falke, J. J.; Parkinson, J. S. *Trends Biochem. Sci.* **2008**, 33, 9-19.
- (66) Xu, L.; Guo, C.; Wang, F.; Zheng, S.; Liu, C.-Z. *Bioresour. Technol.* **2011**, 102, 10047-10051.
- (67) Liu, J.; Hu, B.; Morado, D. R.; Jani, S.; Manson, M. D.; Margolin, W. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, 109, E1481-8.
- (68) Lanfranconi, M. P.; Alvarez, H. M.; Studdert, C. A. *Environ. Microbiol.* **2003**, 5, 1002-1008.
- (69) Smits, T. H.; Witholt, B.; van Beilen, J. B. *Anton. Leeuw. Int. J. G.* **2003**, 84, 193-200.
- (70) Sourjik, V.; Armitage, J. P. *EMBO J.* **2010**, 29, 2724-2733.
- (71) Wang, W.; Shao, Z. *Nat. Commun.* **2014**, 5.
- (72) Prince, R. C.; McFarlin, K. M.; Butler, J. D.; Febbo, E. J.; Wang, F. C.; Nedwed, T. J. *Chemosphere* **2013**, 90, 521-6.
- (73) Lal, B.; Khanna, S. *J. Appl. Bacteriol.* **1996**, 81, 355-362.
- (74) Maeng, J. H.; Sakai, Y.; Tani, Y.; Kato, N. *J. Bacteriol.* **1996**, 178, 3695-3700.
- (75) Ratajczak, A.; Geißdörfer, W.; Hillen, W. *J. Bacteriol.* **1998**, 180, 5822-5827.
- (76) Throne-Holst, M.; Wentzel, A.; Ellingsen, T. E.; Kotlar, H.-K.; Zotchev, S. B. *Appl. Environ. Microbiol.* **2007**, 73, 3327-3332.
- (77) Tanaka, D.; Takashima, M.; Mizuta, A.; Tanaka, S.; Sakatoku, A.; Nishikawa, A.; Osawa, T.; Noguchi, M.; Aizawa, S.-I.; Nakamura, S. *Curr. Microbiol.* **2010**, 60, 203-209.
- (78) Wang, X. B.; Nie, Y.; Tang, Y. Q.; Wu, G.; Wu, X. L. *Appl. Environ. Microbiol.* **2013**, 79, 400-2.

Table of Content Graphic



For TOC only.