Article

Biochemical alterations in duckweed and algae induced by carrier solvents: Selection of an appropriate solvent in toxicity testing

Hu, Li-Xin, Tian, Fei, Martin, Francis L and Ying, Guang-Gou

Available at <http://clock.uclan.ac.uk/17392/>

Hu, Li-Xin, Tian, Fei, Martin, Francis L ORCID: 0000-0001-8562-4944 and Ying, Guang-Gou (2017) Biochemical alterations in duckweed and algae induced by carrier solvents: Selection of an appropriate solvent in toxicity testing. Environmental Toxicology and Chemistry . ISSN 0730-7268

It is advisable to refer to the publisher's version if you intend to cite from the work.
<http://dx.doi.org/10.1002/etc.3804>

For more information about UCLan's research in this area go to
<http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to
<http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

**BIOCHEMICAL ALTERATIONS IN DUCKWEED AND ALGAE INDUCED BY CARRIER
SOLVENTS: SELECTION OF AN APPROPRIATE SOLVENT IN TOXICITY TESTING**

LI-XIN HU, FEI TIAN, FRANCIS L. MARTIN, and GUANG-GUO YING

Environ Toxicol Chem., **Accepted Article** • DOI: 10.1002/etc.3804

Accepted Article

"Accepted Articles" are peer-reviewed, accepted manuscripts that have not been edited, formatted, or in any way altered by the authors since acceptance. They are citable by the Digital Object Identifier (DOI). After the manuscript is edited and formatted, it will be removed from the "Accepted Articles" Web site and published as an Early View article. Note that editing may introduce changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. SETAC cannot be held responsible for errors or consequences arising from the use of information contained in these manuscripts.

BIOCHEMICAL ALTERATIONS IN DUCKWEED AND ALGAE INDUCED BY CARRIER
SOLVENTS: SELECTION OF AN APPROPRIATE SOLVENT IN TOXICITY TESTING

Running title: Selection of carrier solvents in toxicity testing

LI-XIN HU,^{a,b} FEI TIAN,^a FRANCIS L. MARTIN,^c and GUANG-GUO YING^{a,d,*}

^aState Key Laboratory of Organic Geochemistry, Guangzhou Institute of Geochemistry, Chinese
Academy of Sciences, Guangzhou, China

^bUniversity of Chinese Academy of Sciences, Beijing, China

^cSchool of Pharmacy and Biomedical Sciences, University of Central Lancashire, Preston, UK

^dThe Environmental Research Institute, MOE Key Laboratory of Environmental Theoretical Chemistry,
South China Normal University, Guangzhou, China

* Address correspondence to guang-guo.ying@gig.ac.cn

This article contains online-only Supplemental Data

This article is protected by copyright. All rights reserved

Submitted 8 December 2016; Returned for Revision 5 March 2017; Accepted 22 March 2017

This article is protected by copyright. All rights reserved

Abstract: Carrier solvents are often used in aquatic toxicity testing for test chemicals with hydrophobic properties. However, the knowledge of solvent effects on test organisms remains limited. This study aimed to understand biochemical effects of the four common solvents ((methanol, ethanol, acetone and dimethylsulfoxide (DMSO)) on two test species *Lemna minor* and *Raphidocelis subcapitata* by applying Fourier transform infrared spectroscopy (FTIR) coupled with multivariate analysis in order to select appropriate solvents in the toxicity testing. The results showed biochemical variations associated with solvent treatments at different doses on test species. From the obtained infrared spectra, the structures of lipid membrane and protein phosphorylation in the test species were found sensitive to the solvents. Methanol and ethanol mainly affected the protein secondary structure, while acetone and DMSO primarily induced the alterations of carbohydrate and proteins in the test species. The FTIR results demonstrated that methanol and ethanol showed higher biochemical alterations in the test species than acetone and DMSO, especially at the high doses (0.1% and 1% v/v). Based on the growth inhibition and FTIR spectroscopy, acetone and DMSO can be used as carrier solvent in the toxicity testing when their doses are lower than 0.1% v/v. This article is protected by copyright. All rights reserved

Keywords: Toxicity test; Carrier solvent; FTIR spectroscopy; Biochemical fingerprint

INTRODUCTION

Considering the insoluble or sparingly soluble property of some chemicals, it becomes crucial to use a carrier solvent in toxicity testing. Methanol, ethanol, acetone and dimethylsulfoxide (DMSO) are the most commonly used solvents in toxicity tests [1]. However, their potential effects on test organisms have become a crucial issue. OECD has recommended a maximum acceptable solvent concentration of 0.01% v/v, and the United States environmental protection agency (US EPA) has set 0.05% solvent as the maximum concentration for acute toxicity tests [1, 2]. The solvent doses in the aquatic assays are often higher than these limits of recommendation because of the low solubility of test chemicals, small test volume and other limitations [3]. Although the carrier solvents are often used for the toxicity testing of hydrophobic chemicals, the toxic effects and toxicity data of these solvents are still very limited, especially for those subtle changes at molecular level. Thus there is a need for running concurrent solvent and negative controls to assess carrier influence in these assays.

There have been some reports on toxicity of some carrier solvents to alga, fish, invertebrate and other organisms [4-7]. Syed et al. [8] studied the effect of acetone, acetonitrile, dimethyl sulfoxide (DMSO), ethanol and methanol on oxidative phosphorylation (ATP synthesis) in rat liver mitochondria, and found that all the organic solvents inhibited the oxidative phosphorylation in a concentration dependent manner, but with differences in potencies. Among the tested organic solvents, acetonitrile and acetone were more potent than ethanol, methanol, and DMSO [8]. Ma et al. [5] assessed the toxicity of seven solvents (acetone, ethanol, methanol, DMSO, *N,N*-dimethyl-formamide, furanidine and acetidin) to green alga, and indicate that acetone and DMSO would be suitable solvents to use in toxicity bioassays involving algae. As a higher plant, *Lemna gibba* was found to be much more

This article is protected by copyright. All rights reserved

sensitive to methanol when compared with some algae, and the inhibitory effect was strongly associated with the increase in the nonphotochemical energy dissipation [9]. Chen et al. [10] investigated the behavior effects of ethanol and DMSO to zebrafish embryo-larvae, and observed that the solvents at low concentrations could alter locomotor activity of larval zebrafish without causing any observable developmental defects. Therefore, solvent effects in toxicity bioassays should be further investigated, especially at the biochemical level.

Duckweed (*Lemna minor*) and alga (*Raphidocelis subcapitata*) are common test species in aquatic toxicity tests of chemicals [11]. Their effects can be further explored at the subcellular level using Fourier transform infrared spectroscopy (FTIR). FTIR is a useful tool to provide detailed biochemical information according to the vibration of the chemical bonds [12], and it can be used as endpoints in toxicity testing [13]. For instance, the vibration in $3000\text{-}2800\text{ cm}^{-1}$ are often assigned to the stretching vibration of C-H of CH_3 and CH_2 , and the vibrations of $1750\text{-}1700\text{ cm}^{-1}$ are associated with C=O stretching vibration of lipid, vibrations in these regions can indicate the structure and degree of oxidation of the lipid [13]. $1700\text{-}1580\text{ cm}^{-1}$ mainly represents the C=O stretching vibrations coupled with C-N stretching vibrations of amide I, this region can indicate the secondary structure of protein [14]. The amide II is associated with the vibration of C-N stretching vibrations along with N-H in-plane bending vibration at $1580\text{-}1480\text{ cm}^{-1}$ region [15]. The biochemical alterations at 1250 and 1080 cm^{-1} are often related with asymmetric and symmetric phosphate stretching vibrations, the alterations at these bands may be an indicator of the nucleic acids changes [16]. The vibrations in $1200\text{-}900\text{ cm}^{-1}$ are often assigned to carbohydrate [17]. In order to extract the biomarkers to evaluate the biochemical effects induced by the reagents, a sophisticated unsupervised approach like principle component analysis and a powerful classification tool like linear discriminant analysis are applied to analyze the

This article is protected by copyright. All rights reserved

complicated IR data [18].

The aims of this study was to assess the alterations of biochemical information induced by four common carrier solvents (methanol, ethanol, acetone and DMSO), and to select an appropriate solvent and concentration in duckweed and alga toxicity testing for hydrophobic chemicals.

MATERIALS AND METHODS

Reagents

The solvents tested in this study included ethanol (99.9%), methanol (99.99%), acetone (99.9%), dimethylsulfoxide (DMSO, 99.99%). Ethanol and acetone were purchased from FUYU Chemical Co. (China), while methanol and DMSO were purchased from Merck Corporation (Germany). All solvent doses in the toxicity tests were given as percent (%) volume/volume.

Test species

The duckweed *Lemna minor* used in this study was cultured in laboratory in the Swedish standard (SIS) medium (Table S1) under the following conditions: 2000 lx; 25±1 °C; Humidity, 60%; day: night 14 h: 10 h. The alga *Raphidocelis subcapitata* was cultured in the Blue-Green (BG11) medium (Table S1) under the following conditions: 2000 lx; 25±1 °C; Humidity, 60%; day: night 14 h: 10 h.

Toxicity testing

Duckweed toxicity test. Exposure tests for duckweeds were conducted in a number of six well plates. Each well contained 10 mL of the SIS medium and 4 three-frond colonies of approximately the same size. Four doses (1%, 0.1%, 0.01%, and 0.001% v/v) for each carrier solvent were used, and each treatment was replicated three times. All fronds were placed under the same conditions shown as above. The number of fronds in each well was counted every two days and then the solution was renewed. At the end of the exposure, two fronds from each treatment were picked out for FTIR and biomass

This article is protected by copyright. All rights reserved

analysis. The fronds for FTIR analysis were fixed with 10% formalin in phosphate buffer saline (PBS) for 12 h [13].

Alga toxicity test. Exposure tests for algae were conducted in a series of 24 well plates. Each well contained 2 mL of BG11 medium and 1.0×10^4 cells/mL of algae. Four doses (1%, 0.1%, 0.01%, and 0.001% v/v) for each carrier solvent were used, and each treatment was replicated four times. The algal biomass in each well was measured at the absorbance of 437 nm each day using a microplate reader (BMG Labtech FLUOstar Omega) [19]. At the end of the exposure, the algal cells were centrifuged at 1000 g for 5 min, washed with PBS and then fixed with 10% formalin in PBS for 12 h.

FTIR spectroscopy and pre-processing of spectra data

Duckweed fronds in 10% formalin in PBS were washed three times with milli-Q water, applied to BaF₂ slides (30×3 mm, BTD, Tianjin), and then dried in the desiccator for 24 h. For algal cells, the fixed cells were centrifuged at 1000 g for 5 min, washed three times with milli-Q water and suspended with 100 µL water, and 10 µL of the mixture was dropped onto a BaF₂ slide for infrared measurement. IR spectra were obtained using a Bruker vertex 70 FTIR spectrometer (Bruker Optics Ltd., Germany) equipped with a Hyperion 1000 microscope, which contained a liquid nitrogen cooled detector.

Instrument parameters were set at 64 scans, and 8 cm⁻¹ resolution. Prior to starting each new slide, a background was taken. For each replication, approximately 50 IR spectra were acquired at different positions across each sample.

Raw data obtained from tested fronds were analyzed in the irootlab toolbox (<http://irootlab.googlecode.com/>) running on Matlab r2010a [20]. Each spectrum was cut at the C-H stretching vibration region (3000-2800 cm⁻¹) and the biochemical fingerprint region (1800-900 cm⁻¹) for further analysis. These two regions were then baseline corrected and normalized to the maximum

This article is protected by copyright. All rights reserved

peak, 1650 cm^{-1} for the biochemical region and 2920 cm^{-1} for the C-H stretching region [13]. Following these preprocessing, a multivariate analysis, principal component analysis (PCA) and linear discriminant analysis (PCA-LDA) was applied to identify biochemical alterations [21]. As a powerful dimensionality reduction tools, PCA is applied to reduce numbers of wavenumbers to 10 PCs [22].

LDA was applied to discriminate groups with different treatments, and it could also minimize the within-class variances and maximize the inter-class variances [21]. In this study, the first LDA factor (LD_1) was used to represent the primary biochemical alterations induced by reagents, which was shown in scores plots. Cluster vectors were also used to describe the difference between treatment groups (color line) and the control (black straight line) [23].

Statistical significance test

To determine whether treatment groups were significantly different from the reference control, one-way analysis of variance (ANOVA) with Dunnett's post hoc tests was applied. A p value below 0.05 is regarded to be statistically significant. All ANOVA analyses were performed in GraphPad Prism 5 (GraphPad Software, USA).

RESULTS

Growth inhibition effects

The growth inhibitions of duckweed fronds and algal cells under different test concentrations of the solvents are shown in Figure 1. For duckweeds, with the increasing concentrations of the tested solvents, the growth inhibition rates were gradually increasing. Compared with the reference control, the four solvents at the low doses (0.001% and 0.01% v/v) had little difference in growth inhibition. While their concentrations increased up to 0.1%, the fronds treated with methanol and ethanol were significantly inhibited, with the inhibition rate exceeding 50%, while the fronds treated with acetone

This article is protected by copyright. All rights reserved

and DMSO shown little effect on inhibition. When the fronds were exposed to the reagents at 1% v/v, they all showed different degrees of inhibition. In addition to the growth inhibition effect, some morphological differences of fronds were noticed, and the differences included the area, transparency and color of the fronds. Compared with the low doses (0.001% and 0.01% v/v), the fronds treated with the high doses (0.1% and 1% v/v) showed smaller, more transparent and pale features. Besides, the fronds exposed to methanol and ethanol at the high doses (0.1% and 1% v/v) emerged with leaking cytosol.

For algae, the dose related effects were not evident, except for the ethanol group. The growth inhibition for the methanol group showed little variations with the inhibition rate remaining at about 10% with the increasing doses. The acetone group and DMSO group showed little change in growth inhibition at the low doses, and a slight stimulatory effect at the high doses.

Biochemical Effects

Duckweeds. To visualize the biochemical effects in duckweeds, integrated FTIR data obtained from the exposure experiment were analyzed with multivariate analysis. Throughout the spectral region, C-H stretching vibration region (3000 to 2800 cm^{-1}) and biochemical fingerprint region (1800 to 900 cm^{-1}), there were no clear differences between different treatment groups. Considering the large amounts of spectra obtained and its multidimensional character, PCA – LDA was applied to discriminate different categories and identify biomarkers associated with biochemical alterations.

Scores plots in two-dimensions (Figure 2) showed differences between different classes, nearness in spectral points of classes implies similarity, while segregation of classes indicates dissimilar. Clusters vectors indicated the distinguishing features induced by the treatments compared to the reference control. The effects associated with different treatments were achieved by calculating the distance

along LD₁ from the mean spectra of treatment classes and the reference control.

Compared with the control, the methanol group induced distinct, statistically significant ($p < 0.001$) dose related alterations at the high doses (0.1% and 1% v/v) in biochemical fingerprint (Figure 2a).

Following exposure of fronds to the low doses (0.001% and 0.01% v/v) of the solvent, a slight separation from the reference control along LD₂ but not LD₁ was observed. This phenomenon shows that the low dose treatments induced small alterations when compared to the significant biochemical effects caused by the high dose treatments. For the high dose treatments, there was a clear separation along LD₁ in C-H stretching vibration region (Figure 3a).

In clusters vectors, we can identify wavenumbers distinguishing the control from the methanol treatments. The main alterations induced with the high methanol doses (1% and 0.1% v/v) included vibrations in absorbance spectra associated with lignin ($\sim 1600\text{ cm}^{-1}$), lipids ($\sim 1740\text{ cm}^{-1}$), Amide II (1550 cm^{-1}), carbohydrate ($\sim 1150\text{ cm}^{-1}$) and protein phosphorylation (980 cm^{-1}). Only slight biochemical alterations were induced by the low doses (0.01% and 0.001% v/v) when compared with the control. The main alterations associated with 0.01% v/v methanol treatment included vibrations in absorbance spectra associated with lipids ($\sim 1700\text{ cm}^{-1}$), protein (amino acid residue (1420 cm^{-1})), carbohydrate ($\sim 1100\text{ cm}^{-1}$) and starch (1000 cm^{-1}). The main alterations associated with 0.001% v/v methanol treatment included vibrations in absorbance spectra associated with starch ($\sim 1025\text{ cm}^{-1}$) and protein phosphorylation (980 cm^{-1}) (Figure 2b).

The dose-related alterations in C-H stretching region by methanol treatments were observed in LD₁ (Figure 3a). There was no overlap of spectral points in 1% v/v methanol treatment, compared to the other doses. Clusters vectors in C-H region showed that the main difference in LD₁ was caused by C-H stretching vibration of CH₃ ($\sim 2960\text{ cm}^{-1}$) and CH₂ ($\sim 2920\text{ cm}^{-1}$), and this spectral alteration was

This article is protected by copyright. All rights reserved

assigned to lipid (Figure 3b).

Exposure of fronds to ethanol also resulted in a dose-related response in biochemical information (Figure 2c). Compared with the reference control, there are two distinct dose related alterations in 2-D scores plots showed distinct dose-related patterns. The overlapping at the high doses indicated similarity in biochemical alterations; and a small separation of the low doses from the control along LD₂ indicated slight biochemical alterations at the low doses. As shown in CV plots, the main alterations at the high ethanol doses included amide I (1627 cm⁻¹), lipid (1740 cm⁻¹), and amide II (1542 cm⁻¹) (Figure 2d). Subtle alterations at the low doses included carbohydrate (1164 cm⁻¹), amide I (1608 cm⁻¹) and lipid (1735 cm⁻¹).

Complete separation of the high ethanol doses from the low doses and control was observed in C-H region (Figure 3c). The alterations associated with ethanol exposure included symmetric and asymmetric C-H stretching vibrations of CH₃ and CH₂ (Figure 3d).

For acetone treatments, there were incomplete segregations between each treatment and the control in both biochemical fingerprint region and C-H region (Figure 2e). But not much overlap in spectral points was observed between the high doses and low doses, so their spectral points were divided at LD₁ in biochemical region. The alterations induced by the high acetone doses were mainly found on starch (~1050 cm⁻¹), nucleic acids (1064 cm⁻¹) and protein (1625 cm⁻¹), while for the low acetone doses only subtle changes were found on protein (amide I and amide II), carbohydrate (1164 cm⁻¹) and nucleic acids (1083 cm⁻¹) (Figure 2f). Little separation was observed along LD₂ in C-H stretching vibrations (Figure 3e), and the CV plots showed dose-related biochemical variations on CH₂ (2850 cm⁻¹) (Figure 3f).

For DMSO treatments, incomplete segregations were observed between each treatment and the

control (Figure 2g). No overlap was found between the 1% v/v dose and the other doses in biochemical region and C-H stretching region. Spectral alterations induced by DMSO treatments were mainly associated with amide I ($\sim 1600\text{ cm}^{-1}$), amide II ($\sim 1520\text{ cm}^{-1}$) and starch ($\sim 1000\text{ cm}^{-1}$) (Figure 2h). The main alterations for the 0.001% v/v DMSO treatment included vibrations associated with amide I (1608 cm^{-1}), amide II ($\sim 1520\text{ cm}^{-1}$), while those for the 1% v/v DMSO treatment included vibrations associated with starch (1030 cm^{-1}), and carbohydrate (1126 cm^{-1}). In C-H stretching vibrations region, the highest dose induced marked spectral alterations, with all kinds of C-H stretching vibrations being changed (Figure 3g and 3h).

Algae. The methanol treatments induced dose-related alterations in algal cells. The spectra of methanol treatments were partially overlapped with the control (Figure 4a and 5a), suggesting only slight alterations in algal cells induced by the methanol doses. In CV plots (Figure 4b), the distinguishing wavenumbers were mainly focused on amide I, carbohydrate and DNA/RNA at the high doses, and proteins and starch at the low doses.

For ethanol treatments, clear segregations were observed between the treatment groups and the control and the separation from the control increased with the doses along LD₁ (Figure 4c and 5c). The alterations obtained at CH stretching region were the same as the biochemical fingerprint region, with the principal alterations on protein (amide I and amide II) and lipid related vibrations (Figure 4d and 5d).

For acetone treatments, no clear separation was found between the treatments and the control at both biochemical fingerprint region and CH stretching region (Figure 4e and 5e). Spectra alterations induced by acetone were generally associated with starch, amide I and protein phosphorylation (Figure 4f).

For the DMSO treatments, the differences between different treatments and the control were very subtle, and the coefficient values were very small at the biochemical region (Figure 4g and 4h). Spectra alterations at CH stretching region shown a dose related effects, and the ratio of vibration on CH₂ and CH₃ were also changed (Figure 5g and 5h).

DISCUSSION

The results from the present study showed species specific dose-related effects from the carrier solvents based on the traditional endpoints and biochemical endpoints. For the two test species (*Lemna minor* and *Raphidocelis subcapitata*), acetone and DMSO clearly showed less growth effects and biochemical alterations than methanol and ethanol. The result from the present study is also consistent with a previous study [3], which indicated that the solvent toxicity was ethanol, methanol > acetone > DMSO. Compared with the reference control, the distances between the treatment groups and the reference control on duckweed (Figure 2) were greater than the distances on alga (Figure 4). As the clusters vectors plots shown, the height of the peaks of duckweed ranged from 0.1 to 0.2, while the height of peaks of alga was limited to 0.015. So the solvents had less toxic effects on algae than duckweeds, especially at the high doses.

This may be due to fast algal cell reproduction. The results based on the biomass and FTIR analysis showed consistency to some degree (Figure S1), but FTIR analysis gave us more information on specific biochemical alterations.

Since the cell is mainly constituted of lipid and protein, alterations of these two elements, especially the lipid, could disrupt the structure and function of cells [24]. The present study showed dose-related alterations of C-H stretching region due to the exposure to the solvents, which could lead to disorder of the C-H chain in the phospholipid, and further affecting cell membrane permeability and

This article is protected by copyright. All rights reserved

fluidity [25, 26]. Thus these solvents especially methanol and ethanol at the high doses could alter the structure of the membrane lipids, and induce the peroxidation of the membrane lipids. In accordance with the previous studies [27], our findings also provided the direct evidence to the relation between the biochemical mechanism of ethanol toxicity and the lipid peroxidation.

The change of protein structure would affect the normal function of organism and metabolic materials. The relative content of α -helix, β -sheet and β -turn represents the stabilization of protein structure [28]. The vibrations associated with 1400 cm^{-1} are assigned to the amino acids residues, and the damage of these residues could affect the secondary structure [29]. Besides, the vibrations at 970 cm^{-1} are related to protein phosphorylation, which is associated with the structure of β -sheet [30]. The results from the present study showed alterations of β -sheet structure by the methanol, ethanol and acetone treatments, and changes of α -helix by the methanol treatment. The alterations of amino acids residues were associated with the acetone and DMSO groups. Protein phosphorylation was observed for the treatments of all four solvents, suggesting sensitive response to the carrier solvents.

The alterations of DNA/RNA are always associated with the bands at 1250 cm^{-1} and 1080 cm^{-1} , and these two bands are assigned to asymmetric and symmetric stretching vibration of phosphodiester linkage [31]. The alterations of phosphodiester linkage were detected in the present study for the four solvents, even at the doses below the OECD recommend limit. The alterations of carbohydrates are mainly reflected on the band of 1000 cm^{-1} to 1200 cm^{-1} [32]. These characteristic peaks are related to the photosynthesis of plant cells. According to the present study, acetone and DMSO at the high doses might result in the changes of carbohydrates.

Based on the biochemical alterations and PCA - LDA scores plots, appropriate solvents can be selected for use in duckweed and alga toxicity testing. For duckweed, a clear separation of the high

This article is protected by copyright. All rights reserved

doses (1% and 0.1% v/v) from the low doses (0.01% and 0.001% v/v) and the control was observed for methanol and ethanol treatments in biochemical region, suggesting significant effects of methanol and ethanol on algal cells (Figure 2). In contrast, overlapping for acetone and DMSO treatments was observed, indicating less biochemical effects from these two solvents. Much higher growth inhibition rates were found for both methanol and ethanol than acetone and DMSO at the high doses (Figure 1). This suggests that methanol and ethanol are not suitable to be used as carrier solvents at the high doses (1% and 0.1% v/v) in duckweed toxicity testing. Both the present study and previous study [9] showed sensitivity of duckweeds to methanol when compared with algae.

For algae, a clear segregation of different doses was only observed for ethanol, while overlapping for the other three solvents (Figure 4). Previous studies [4, 5] investigated toxicity of chemical solvents on algae based on growth inhibition. They found that methanol and ethanol are unsuitable for toxicity testing, while acetone and DMSO would be suitable solvents to use in toxicity bioassays involving algae.

CONCLUSION

The results from this study showed that biochemical alterations in the test organisms duckweeds and algae could be induced by the carrier solvents even at the low doses. The effects to the two test species varied among the four solvents, and slight differences were found between the two species. Methanol and ethanol generally caused more severe toxic effects than acetone and DMSO; thus acetone and DMSO are better carrier solvents for hydrophobic chemicals in duckweed and alga toxicity testing. For other toxicity tests, a preliminary experiment is essential to find out what solvent and dose should be used before testing. It is also crucial to ensure that the solvent used in the testing has no toxic effects, whether in the growth and development, or in the subcellular biochemical information. The solvent

This article is protected by copyright. All rights reserved

control should be included together with the toxicity test, while the interaction between the test chemical and carrier solvent should also be avoided.

Supplemental Data—The Supplemental Data are available on the Wiley Online Library at DOI: 10.1002/etc.xxxx.

Acknowledgment—The authors would like to acknowledge the financial support from Guangdong Provincial Applied Science and Technology Research Program (2015B020235012), National Natural Science Foundation of China (U1401235) and National Water Pollution Control Program of China (2014ZX07206-005). Thanks to J.Y. Li at Lancaster University for his assistance in bio-spectroscopy analysis.

Data Availability—Data is included in the supporting materials.

REFERENCES

1. Chen TH, Wang YH, Wu YH. 2011. Developmental exposures to ethanol or dimethylsulfoxide at low concentrations alter locomotor activity in larval zebrafish: implications for behavioral toxicity bioassays. *Aquat Toxicol* 102:162-166.
2. Hutchinson T, Shillabeer N, Winter M, Pickford D. 2006. Acute and chronic effects of carrier solvents in aquatic organisms: a critical review. *Aquat Toxicol* 76:69-92.
3. Okumura Y, Koyama J, Takaku H, Satoh H. 2001. Influence of organic solvents on the growth of marine microalgae. *Arch Environ Con Tox* 41:123-128.
4. El Jay A. 1996. Toxic effects of organic solvents on the growth of *Chlorella vulgaris* and *Selenastrum capricornutum*. *B Environ Contam Tox* 57:191-198.
5. Ma J, Chen J. 2005. How to accurately assay the algal toxicity of pesticides with low water solubility. *Environ Pollut* 136:267-273.
6. David RM, Jones HS, Panter GH, Winter MJ, Hutchinson TH, Chipman JK. 2012. Interference with xenobiotic metabolic activity by the commonly used vehicle solvents dimethylsulfoxide and methanol in zebrafish (*Danio rerio*) larvae but not *Daphnia magna*. *Chemosphere* 88:912-917.
7. Stevens AS, Pirotte N, Plusquin M, Willems M, Neyens T, Artois T, Smeets K. 2015. Toxicity profiles and solvent–toxicant interference in the planarian *Schmidtea mediterranea* after dimethylsulfoxide (DMSO) exposure. *J Appl Toxicol* 35:319-326
8. Syed M, Skonberg C, Hansen SH. 2013. Effect of some organic solvents on oxidative phosphorylation in rat liver mitochondria: Choice of organic solvents. *Toxicol in Vitro* 27:2135-2141.
9. Dewez D, Dautremepuits C, Jeandet P, Vernet G, Popovic R. 2003. Effects of methanol on

photosynthetic processes and growth of *Lemna gibba*. *Photochem Photobi* 78:420-424.

10. Chai J, Yan X, Liu RT, Wang M. 2011. Study on the toxic interaction of methanol, ethanol and propanol against the bovine hemoglobin (BHb) on molecular level. *Spectrochim Acta A* 79:1406-1410.

11. Mkandawire M, Teixeira da Silva JA, Dudel EG. 2014. The *Lemna* bioassay: contemporary issues as the most standardized plant bioassay for aquatic ecotoxicology. *Crit Rev Env Sci Tec* 44:154-197.

12. Martin FL, Kelly JG, Llabjani V, Martin-Hirsch PL, Patel II, Trevisan J, Fullwood NJ, Walsh MJ. 2010. Distinguishing cell types or populations based on the computational analysis of their infrared spectra. *Nat Protoc* 5:1748-1760.

13. Hu LX, Ying GG, Chen XW, Huang GY, Liu YS, Jiang YX, Pan CG, Tian F, Martin FL. 2016. FTIR spectroscopy as a novel approach towards providing effect-based endpoints in duckweed toxicity testing. *Environ Toxicol Chem* 36(2):346-353

14. Baker MJ, Trevisan J, Bassan P, Bhargava R, Butler HJ, Dorling KM, Fielden PR, Fogarty SW, Fullwood NJ, Heys KA. 2014. Using Fourier transform IR spectroscopy to analyze biological materials. *Nat Protoc* 9:1771-1791.

15. Lewis PD, Lewis KE, Ghosal R, Bayliss S, Lloyd AJ, Wills J, Godfrey R, Kloer P, Mur LA. 2010. Evaluation of FTIR spectroscopy as a diagnostic tool for lung cancer using sputum. *BMC Cancer* 10:1.

16. Li J, Strong R, Trevisan J, Fogarty SW, Fullwood NJ, Jones KC, Martin FL. 2013. Dose-related alterations of carbon nanoparticles in mammalian cells detected using biospectroscopy: potential for real-world effects. *Environ Sci Technol* 47:10005-10011.

17. Li J, Ying GG, Jones KC, Martin FL. 2015. Real-world carbon nanoparticle exposures induce brain and gonadal alterations in zebrafish (*Danio rerio*) as determined by biospectroscopy techniques.

Analyst 140:2687-2695.

This article is protected by copyright. All rights reserved

18. Llabjani V, Trevisan JI, Jones KC, Shore RF, Martin FL. 2011. Derivation by infrared spectroscopy with multivariate analysis of bimodal contaminant-induced dose-response effects in MCF-7 cells. *Environ Sci Technol* 45:6129-6135.
19. Gitchaiwat, A., Likhitlert, S., Kositchaiyong, A., Israngkura, K., Taptim, K., & Sombatsompop, N. 2015. Uses of 2-hydroxypropyl-3-piperazinyl-quinoline carboxylic acid methacrylate and Terbutryn as algaecides in low-density polyethylene mulching films for agricultural applications. *J Plast Film Sheet*, 8756087915578467.
20. Trevisan J, Angelov PP, Scott AD, Carmichael PL, Martin FL. 2013. IRootLab: a free and open-source MATLAB toolbox for vibrational biospectroscopy data analysis. *Bioinformatics*:btt084.
21. Llabjani V, Trevisan J, Jones KC, Shore RF, Martin FL. 2010. Binary mixture effects by PBDE congeners (47, 153, 183, or 209) and PCB congeners (126 or 153) in MCF-7 cells: biochemical alterations assessed by IR spectroscopy and multivariate analysis. *Environ Sci Technol* 44:3992-3998.
22. Wagner H, Liu Z, Langner U, Stehfest K, Wilhelm C. 2010. The use of FTIR spectroscopy to assess quantitative changes in the biochemical composition of microalgae. *J Biophotonics* 3:557-566.
23. Li J, Tian M, Cui L, Dwyer J, Fullwood NJ, Shen H, Martin FL. 2016. Low-dose carbon-based nanoparticle-induced effects in A549 lung cells determined by biospectroscopy are associated with increases in genomic methylation. *Sci Rep-UK* 6.
24. Michael P, Salonen E, Terama E, Vattulainen I, Faller R, Lee BW, Holopainen J, Karttunen M. 2006. Under the influence of alcohol: the effect of ethanol and methanol on lipid bilayers. *Biophys J* 90:1121-1135.
25. Ozek NS, Tuna S, Erson-Bensan AE, Severcan F. 2010. Characterization of microRNA-125b expression in MCF7 breast cancer cells by ATR-FTIR spectroscopy. *Analyst* 135:3094-3102.

26. Vilen B, Jeney S, Sienkiewicz A, Marcoux P, Miller L, Forró L. 2010. Evidence of lipid peroxidation and protein phosphorylation in cells upon oxidative stress photo-generated by fullerenes. *Biophys Chem* 152:164-169.
27. Corongiu F P, Lai M, Milia A. 1983. Carbon tetrachloride, bromotrichloromethane and ethanol acute intoxication. New chemical evidence for lipid peroxidation in rat tissue microsomes. *Biochemical Journal* 212(3): 625-631.
28. Roccatano D, Colombo G, Fioroni M, Mark AE. 2002. Mechanism by which 2, 2, 2-trifluoroethanol/water mixtures stabilize secondary-structure formation in peptides: a molecular dynamics study. *P Natl A Sci* 99:12179-12184.
29. Rost B, Sander C. 1994. Combining evolutionary information and neural networks to predict protein secondary structure. *Proteins* 19:55-72.
30. Bah A, Vernon RM, Siddiqui Z, Krzeminski M, Muhandiram R, Zhao C, Sonenberg N, Kay LE, Forman-Kay JD. 2015. Folding of an intrinsically disordered protein by phosphorylation as a regulatory switch. *Nature* 519:106-109.
31. Obinaju BE, Fullwood NJ, Martin FL. 2015. Distinguishing nuclei-specific benzo [a] pyrene-induced effects from whole-cell alterations in MCF-7 cells using Fourier-transform infrared spectroscopy. *Toxicology* 335:27-34.
32. Butler HJ, McAinsh MR, Adams S, Martin FL. 2015. Application of vibrational spectroscopy techniques to non-destructively monitor plant health and development. *Anal Methods-UK* 7:4059-4070.

Figure 1. Dose-response curves showing the effects of different solvents on growth inhibition of duckweeds (A) and algae (B) at 96 hours. Standard deviations are represented by error bars.

Figure 2. Spectra of duckweeds in the region of biochemical fingerprint ($1800\text{-}900\text{ cm}^{-1}$). Principle component analysis coupled with linear discriminant analysis (PCA-LDA) datasets was applied for the duckweeds exposed to different carrier solvents. PCA-LDA scores plots in two-dimensions (90% confidence ellipsoids) were acquired from fronds of duckweed, the distance between different groups suggested the difference. Clusters vectors plots show the wavenumbers discriminating effects corresponding to the reference control. The heights of the clusters vectors peaks were proportional to the alterations of biochemical information compared to the reference control. The black horizontal line represents the control. The hint line represents a typical duckweed infrared spectrum of the biochemical fingerprint region ($1800\text{-}900\text{ cm}^{-1}$). These scores plots are on the left. Panels represent: (a, b) methanol; (c, d) ethanol; (e, f) acetone; and (g, h) DMSO.

Figure 3. Spectra of duckweeds in the region of C-H stretching vibration region ($3000\text{-}2800\text{ cm}^{-1}$).

Principle component analysis coupled with linear discriminant analysis (PCA-LDA) datasets was obtained from the duckweeds exposed to different carrier solvents. PCA-LDA scores plots in two-dimensions (90% confidence ellipsoids) were acquired from fronds of duckweed, the distance between different groups suggested the difference. Clusters vectors plots show the wavenumbers discriminating effects corresponding to the reference control. The black horizontal line represents the control. The hint line represents a typical duckweed infrared spectrum of C-H stretching vibration region ($3000\text{-}2800\text{ cm}^{-1}$). These scores plots are on the left. Panels represent: (a, b) methanol; (c, d) ethanol; (e, f) acetone; and (g, h) DMSO.

Figure 4. Spectra of algae in the region of biochemical fingerprint ($1800-900\text{ cm}^{-1}$). Principle component analysis coupled with linear discriminant analysis (PCA-LDA) datasets was obtained from the duckweed exposed to different kinds of carrier solvents. PCA-LDA scores plots in two-dimensions (90% confidence ellipsoids) were acquired from fronds of duckweeds; the distance between different groups suggested the difference. Clusters vectors plots show the wavenumbers discriminating effects corresponding to the reference control. The heights of the clusters vectors peaks were proportional to the alterations of biochemical information compared to the reference control. The black horizontal line represents the control. The hint line represents a typical algal infrared spectrum of the biochemical fingerprint region ($1800-900\text{ cm}^{-1}$). These scores plots are on the left. Panels represent: (a, b) methanol; (c, d) ethanol; (e, f) acetone; and (g, h) DMSO.

Figure 5. Spectra of algae in the region of C-H stretching vibration region ($3000-2800\text{ cm}^{-1}$). Principle component analysis coupled with linear discriminant analysis (PCA-LDA) datasets was obtained from the duckweeds exposed to different carrier solvents. PCA-LDA scores plots in two-dimensions (90% confidence ellipsoids) were acquired from fronds of duckweeds; the distance between different groups suggested the difference. Clusters vectors plots show the wavenumbers discriminating effects corresponding to the reference control. The black horizontal line represents the control. The hint line represents a typical algae infrared spectrum of C-H stretching vibration region ($3000-2800\text{ cm}^{-1}$). These scores plots are on the left. Panels represent: (a, b) methanol; (c, d) ethanol; (e, f) acetone; and (g, h) DMSO.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.