

Central Lancashire Online Knowledge (CLoK)

Title	The effects of Ankle Protectors on Lower Limb Kinematics in male football players: A comparison to Braced and Unbraced Ankles
Type	Article
URL	https://clock.uclan.ac.uk/17482/
DOI	https://doi.org/10.3920/CEP160031
Date	2017
Citation	Graydon, Robert William, Fewtrell, David John, Atkins, Stephen and Sinclair, Jonathan Kenneth (2017) The effects of Ankle Protectors on Lower Limb Kinematics in male football players: A comparison to Braced and Unbraced Ankles. <i>Comparative Exercise Physiology</i> , 13 (4). pp. 251-258. ISSN 1755-2540
Creators	Graydon, Robert William, Fewtrell, David John, Atkins, Stephen and Sinclair, Jonathan Kenneth

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.3920/CEP160031>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Comparative Exercise Physiology

The effects of Ankle Protectors on Lower Limb Kinematics in male football players. A comparison to Braced and Unbraced Ankles.

--Manuscript Draft--

Manuscript Number:	CEP-160031R2
Full Title:	The effects of Ankle Protectors on Lower Limb Kinematics in male football players. A comparison to Braced and Unbraced Ankles.
Article Type:	Research article
Corresponding Author:	Robert Graydon Preston, UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	
Corresponding Author's Secondary Institution:	
First Author:	Robert Graydon
First Author Secondary Information:	
Order of Authors:	Robert Graydon David Fewtrell Stephen Atkins Jonathan Sinclair
Order of Authors Secondary Information:	
Abstract:	<p>Football (Soccer) players have a high risk of injuring the lower extremities. To reduce the risk of ankle inversion injuries ankle braces can be worn. To reduce the risk of ankle contusion injuries ankle protectors can be utilized. However, athletes can only wear one of these devices at a time. The effects of ankle braces on stance limb kinematics has been extensively researched, however ankle protectors have had little attention. Therefore, the current study aimed to investigate the effects of ankle protectors on lower extremity kinematics during the stance phase of jogging and compare them with braced and uncovered ankles. Twelve male participants ran at 3.4 m.s⁻¹ in three test conditions; ankle braces (BRACE), ankle protectors (PROTECTOR) and with uncovered ankles (WITHOUT). Stance phase kinematics were collected using an eight-camera motion capture system. Kinematic data between conditions were analysed using one-way repeated measures ANOVA. The results showed that BRACE (absolute range of motion (ROM) =10.72° & relative ROM =10.26°) significantly (P<0.05) restricted the ankle in the coronal plane when compared to PROTECTOR (absolute ROM =13.44° & relative ROM =12.82°) and WITHOUT (absolute ROM =13.64° & relative ROM =13.10°). It was also found that both BRACE (peak dorsiflexion =17.02° & absolute ROM =38.34°) and PROTECTOR (peak dorsiflexion =18.46° & absolute ROM =40.15°) significantly (P<0.05) reduced sagittal plane motion when compared to WITHOUT (peak dorsiflexion =19.20° & absolute ROM =42.66°). Ankle protectors' effects on lower limb kinematics closely resemble that of an unbraced ankle. Therefore, ankle protectors should only be used as a means to reduce risk of ankle contusion injuries and not implemented as a method to reduce the risk of ankle inversion injuries. Furthermore, the reductions found in sagittal plane motion of the ankle could possibly increase the bodies energy demand needed for locomotion when ankle protectors are utilised.</p>

1 **The effects of Ankle Protectors on lower limb kinematics in male football players. A**
2 **comparison to Braced and Unbraced Ankles.**

3 *Graydon, R¹, Fewtrell, D¹, Atkins, S², and Sinclair, J¹*

4 1. *Centre for Applied Sport & Exercise Sciences, School of Sport & Wellbeing,*
5 *University of Central Lancashire, UK*

6 2. *School of Health Sciences, University of Salford, UK*

7
8 **Corresponding author contact details:**

9 Robert Graydon

10 *Centre for Applied Sport & Exercise Sciences*

11 University of Central Lancashire

12 Preston

13 Lancashire

14 PR1 2HE

15 E-mail: RWGraydon@uclan.ac.uk

16 **Funding Declaration**

17 No external funding was provided.

18 **Conflict of Interest Declaration**

19 No conflict of interest.

20

21 **Keywords:** Biomechanics, motion analysis, Ankle Braces, Ankle Protectors, Football,
22 Soccer.

23 **Abstract**

24 Football (Soccer) players have a high risk of injuring the lower extremities. To reduce the risk
25 of ankle inversion injuries ankle braces can be worn. To reduce the risk of ankle contusion
26 injuries ankle protectors can be utilized. However, athletes can only wear one of these devices
27 at a time. The effects of ankle braces on stance limb kinematics has been extensively
28 researched, however ankle protectors have had little attention. Therefore, the current study
29 aimed to investigate the effects of ankle protectors on lower extremity kinematics during the
30 stance phase of **jogging** and compare them with braced and uncovered ankles. Twelve male
31 participants ran at $3.4 \text{ m}\cdot\text{s}^{-1}$ in three test conditions; ankle braces (BRACE), ankle protectors
32 (PROTECTOR) and with uncovered ankles (WITHOUT). Stance phase kinematics were
33 collected using an eight-camera motion capture system. Kinematic data between conditions
34 were analysed using one-way repeated measures ANOVA. The results showed that BRACE
35 (absolute range of motion (ROM) = 10.72° & relative ROM = 10.26°) significantly ($P<0.05$)
36 restricted the ankle in the coronal plane when compared to PROTECTOR (absolute ROM
37 = 13.44° & relative ROM = 12.82°) and WITHOUT (absolute ROM = 13.64° & relative ROM
38 = 13.10°). It was also found that both BRACE (peak dorsiflexion = 17.02° & absolute ROM
39 = 38.34°) and PROTECTOR (peak dorsiflexion = 18.46° & absolute ROM = 40.15°)
40 significantly ($P<0.05$) reduced sagittal plane motion when compared to WITHOUT (peak
41 dorsiflexion = 19.20° & absolute ROM = 42.66°). Ankle protectors' effects on lower limb
42 kinematics closely resemble that of an unbraced ankle. Therefore, **ankle protectors should only**
43 **be used as a means to reduce risk of ankle contusion injuries and not implemented as a method**
44 **to reduce the risk of ankle inversion injuries. Furthermore, the reductions found in sagittal plane**

45 motion of the ankle could possibly increase the bodies energy demand needed for locomotion
46 when ankle protectors are utilised.

47

48 **Introduction**

49 Football (Soccer) is an immensely popular sport with an estimated 265 million participants
50 worldwide (FIFA Communications Division, 2007). Unfortunately, as with any sport, there is
51 an inherent risk of injury to participants and football is no exception. Figures for injury
52 incidences vary among studies due to differing methodologies, time frames observed, ability
53 of participants and competitions observed but conclude there are approximately 25 to 43.53
54 injuries per 1000 hours of competitive match play (Andersen, et al., 2004; Hägglund, et al.,
55 2013; Hawkins & Fuller, 1999; Salces, et al., 2014). Losing an integral team member can lead
56 to a reduced chance of winning competitive matches and further more lead to loss of major
57 trophies (Hägglund, et al., 2013). Therefore, an understanding of the common types of injury
58 sustained by players and also methods to reduce the occurrence of injury is a high priority for
59 football clubs.

60

61 Footballing injuries mainly occur to the lower extremities (Ekstrand, et al., 2011) with the ankle
62 being one of the most commonly injured sites amongst players (Junge & Dvorak, 2013). Ankle
63 inversion injuries and contusion injuries account for a large proportion of the total amount of
64 ankle injuries (Waldén, et al., 2013). Once a player has suffered an ankle inversion injury they
65 have an increased risk of reinjuring the ankle (Thacker, et al., 1999). To reduce the risk of ankle
66 inversion injuries ankle braces can be worn (Kaplan, 2011), the ankles can be taped (Verhagen,
67 et al., 2000), or a neuromuscular training program can be utilised (McGuine & Keene, 2006).
68 Using tape to support the ankle has been found to be ineffective after approximately fifteen

69 minutes of use (Lohkamp, et al., 2009) and expensive (Olmsted, et al., 2004), whereas
70 neuromuscular training programs have been found to be effective but take long periods of time
71 to implement (Emery & Meeuwisse, 2010). This makes ankle braces an attractive alternative
72 because they are easy to put on, do not need to be regularly replaced, and have been found to
73 reduce the risk of ankle inversion injury by restricting the range of motion of the ankle (Farwell,
74 et al., 2013; Janssen, et al., 2014; Pedowitz, et al., 2008). To reduce the risk of contusion
75 injuries ankle protectors can be worn which utilise foam constructs to reduce forces being
76 transferred to the ankle (Ankrah & Mills, 2002; Ankrah & Mills, 2004). Unfortunately, due to
77 ankle braces and ankle protectors aiming to reduce differing injuries at the same location only
78 one of these devices can be used at any one time. This selection is dependent on whether the
79 wearer wants to reduce **the risk of** acute or chronic injuries.

80 Ankle braces effects on ankle kinematics have been well established and have been found to
81 reduce the amount of movement of the ankle (Tang, et al., 2010; DiStefano, et al., 2008) whilst
82 having little effect on running performance (Locke, et al., 1997; Gross, et al., 1997;
83 Bocchinfuso, et al., 1994). **The effects of ankle braces on knee and hip kinematics has also**
84 **been previously studied and found to, in some sporting tasks, increase knee axial rotation which**
85 **could indicate a higher risk of knee injury (Santos, et al., 2004).** However, the effects of ankle
86 protectors' on ankle kinematics during running has, to the author's best knowledge, had no
87 attention. As the location of ankle protectors are the same as ankle braces there is a possibility
88 that they inadvertently act like ankle braces by reducing the amount of movement of the ankle
89 whilst running. **If ankle protectors are found to produce similar ankle kinematics to braced**
90 **ankles, health care professionals could potentially recommend ankle protectors to reduce the**
91 **risk of both ankle inversion injuries and ankle contusion injuries.** Therefore, the current study
92 aims to investigate; firstly, the effects of ankle protectors on ankle kinematics during the stance
93 phase of a wearers running gait, secondly, compare the effects of ankle protectors on ankle

94 kinematics with braced and unbraced ankles to establish which it more closely resembles, **and**
95 **thirdly, investigate the effects of ankle protectors on knee and hip kinematics.**

96

97 **Method**

98 *Participants*

99 Twelve male participants took part in this study. Participants were recruited from local and
100 university football teams using poster adverts. The inclusion criteria for the study was that the
101 participant were aged between 18 and 35, currently playing for a football team, and were injury
102 free at the time of testing. All participants provided written consent in line with the University
103 of Central Lancashire's ethical panel (STEMH 309).

104

105 *Ankle Braces and Ankle Protectors*

106 The ankle protectors used for the current investigation were a pair of Nike ankle shield 10 (Nike
107 Inc, Washington County, Oregon, USA) and the ankle braces used were a pair of Aircast A60
108 (DJO, Vista, CA, USA).

109

110 *****Figure 1 here*****

111

112 *Procedure*

113 Participants performed running trials across a 22m biomechanics laboratory in three test
114 conditions; wearing ankle braces (BRACE), wearing ankle protectors (PROTECTOR) and
115 with uncovered ankles (WITHOUT). Five successful trials were recorded for each test

116 condition. A successful trial was determined as one in which the participant landed with the
117 whole of their right foot on an embedded force platform (Kistler Instruments Ltd., Alton,
118 Hampshire) located in the centre of the laboratory, did not focus on the force plate as to alter
119 their natural gait pattern (Sinclair, et al., 2014), and kept within a speed tolerance of $3.4 \text{ m}\cdot\text{s}^{-1}$
120 $\pm 5\%$. The force plate sampled at 1000 Hz and was used to determine the start and end of the
121 stance phase during the running trials. These points were determined as the point where the
122 force plate first recorded a vertical ground reaction force (VGRF) that exceeded 20N and ended
123 when the VGRF dropped back down below 20N (Sinclair, et al., 2011).

124

125 Kinematic data were recorded using an eight camera motion capture system (Qualisys Medical
126 AB, Goteburg, Sweden) tracking retro-reflective markers at a sampling rate of 250 Hz. Using
127 the calibrated anatomical system technique (CAST) (Cappozzo, et al., 1995) the retro-reflective
128 markers were attached to the 1st and 5th metatarsal heads, calcaneus, medial and lateral
129 malleoli, the medial and lateral femoral epicondyles, the greater trochanter, Left and right
130 anterior superior iliac spine, and left and right posterior superior iliac spine. These markers
131 were used to model the right foot, shank, thigh, and pelvis segments in six degrees of freedom.
132 Rigid plastic mounts with four markers on each were also attached to the shank and thigh and
133 were secured using elasticated bandage. These were used as tracking markers for the shank and
134 thigh segments. To track the foot the 1st and 5th metatarsal heads and the calcaneus were used
135 and to track the pelvis the left and right anterior superior iliac spine and left and right posterior
136 superior iliac spine were used. In the BRACE condition the medial and lateral malleoli
137 locations were found by placing the index finger under the rigid construct of the brace to locate
138 the anatomical landmark then matching the location to the exterior of the Brace where the
139 marker was then fixed to. In the PROTECTOR condition the medial and lateral malleoli
140 locations were located by palpating the soft foam construct to find the underlying anatomical

141 landmarks. To assess the speed of the participant a single marker was attached to the xiphoid
142 process and was checked for velocity using the QTM software after each trial was recorded.
143 Before dynamic trials were captured a static trial of the participant stood in the anatomical
144 position was captured which was used to identify the location of the tracking makers with
145 reference to the anatomical markers. To define each plane of motion firstly the Z (transverse)
146 axis follows the segment from distal to proximal and denotes internal/external rotation,
147 secondly the Y (coronal) axis is orientated from anterior to posterior of the segment and denotes
148 adduction/abduction, and thirdly the X (sagittal) axis is orientated from medial to lateral of the
149 segment and denotes flexion/extension.

150

151 *Data Processing*

152 Anatomical and tracking markers were identified within the Qualisys Track Manager software
153 and then exported as C3D files to be analysed using Visual 3-D software (C-Motion,
154 Germantown, MD, USA). To define the centre points of the ankle and knee segments the two
155 marker methods were utilised for both. These methods calculate the centre of the joint using
156 the positioning of the malleoli markers for the ankle centre and the femoral epicondyle markers
157 for the knee centre (Graydon, et al., 2015; Sinclair, et al., 2015). To calculate the hip joint
158 centre a regression equation which uses the position of the ASIS markers was utilised (Sinclair,
159 et al., 2014). The running trials were filtered at 12Hz using a low pass 4th order zero-lag filter
160 Butterworth filter. Data were normalized to 100% of the stance phase then processed trials
161 were used to produce means of the five trials for each test condition for each participant. 3D
162 kinematics of the ankle, knee and hip joints of the right leg were calculated using an XYZ
163 cardan sequence of rotations. The 3D joint kinematic measures which were extracted for further
164 analysis were 1) angle at footstrike, 2) angle at toe-off, 3) peak angle during the stance phase,

165 4) Absolute range of motion (Absolute ROM) calculated by taking the maximum angle from
166 the minimum angle during stance, 5), Relative range of motion (Relative ROM) calculated
167 using the angle at footstrike and the first peak value after footstrike.

168

169 *Statistical analyses*

170 Data analysis was conducted using SPSS v22.0 (SPSS Inc., Chicago, IL, USA). **The means of**
171 **the five trials for each of** the three test conditions were compared using one-way repeated
172 measures ANOVA with significant findings, accepted at $P < 0.05$ level, being further explored
173 using post-hoc pairwise comparisons. Effect sizes were determined using partial Eta² (η^2).

174

175 **Results**

176 The demographic of the participants of the current study were; age 24.8 ± 4.8 years, height
177 174.8 ± 5.8 cm, body mass 73.4 ± 10.5 kg and BMI 24.0 ± 2.7 .

178 Tables 1, 2, and 3 present the key parameters of interest for each condition and Figures 1, 2,
179 and 3 display the 3D kinematic waveforms recorded for each condition in each plane of motion.

180

181 *****Tables 1-3 close to here*****

182

183 For the ankle joint, in the Sagittal plane, significant main effects were found for the Angle at
184 footstrike $F_{(2, 22)} = 5.04$, $P < 0.05$, $\eta^2 = 0.31$, Angle at toe-off $F_{(2, 22)} = 11.95$, $P < 0.05$, $\eta^2 = 0.52$,
185 Peak dorsiflexion angle $F_{(2, 22)} = 23.27$, $P < 0.05$, $\eta^2 = 0.68$, and Absolute ROM $F_{(2, 22)} = 31.12$,
186 $P < 0.05$, $\eta^2 = 0.74$. Post-hoc analysis revealed that the BRACE condition exhibited significantly

187 (P<0.05) lower angle at footstrike than the PROTECTOR condition. It also revealed the
188 BRACE and PROTECTOR conditions had a significant (P<0.05) reduction in angle at toe off
189 than the WITHOUT condition. The BRACE condition significantly (P<0.05) reduced peak
190 dorsiflexion when compared to the other groups and all three conditions were significantly
191 (P<0.05) different from each other for Absolute range of motion with the WITHOUT condition
192 having the most ROM and BRACE condition having the least ROM.

193 For the ankle joint, in the coronal plane, significant main effects were found for the Angle at
194 footstrike $F_{(2, 22)} = 7.34$, $P < 0.05$, $\eta^2 = 0.40$, Angle at toe-off $F_{(2, 22)} = 6.02$, $P < 0.05$, $\eta^2 = 0.35$, Peak
195 Inversion angle $F_{(2, 22)} = 10.22$, $P < 0.05$, $\eta^2 = 0.48$, Peak Eversion angle $F_{(1.19, 13.14)} = 6.80$,
196 $P < 0.05$, $\eta^2 = 0.38$, Relative ROM $F_{(2, 22)} = 18.40$, $P < 0.05$, $\eta^2 = 0.63$, and Absolute ROM $F_{(2, 22)}$
197 $= 25.19$, $P < 0.05$, $\eta^2 = 0.70$. Post-hoc analysis revealed that the BRACE condition significantly
198 (P<0.05) reduced angle at footstrike, angle at toe off, and peak inversion angle when compared
199 with the WITHOUT condition. The BRACE condition also exhibited significantly (P<0.05)
200 lower peak eversion angle when compared to the PROTECTOR condition. It was also revealed
201 that the BRACE condition had significantly (P<0.05) lower Absolute and Relative ROM's
202 when compared to both the WITHOUT and PROTECTOR conditions.

203

204 No significant differences ($P > 0.05$) were found in the transverse plane for the ankle or in any
205 of the planes of motion for both the knee joint and the hip joint.

206

207 *****Figures 2, 3, and 4 close to here*****

208

209

210 Discussion

211 The aim of the current study was to investigate the effects of ankle protectors on ankle
212 kinematics during the stance phase of a wearers running gait, compare the effects of ankle
213 protectors with braced and unbraced ankles to establish which it more closely resembles, **and**
214 **investigate the effects of ankle protectors on knee and hip kinematics.**

215

216 Previous research reviewing the effectiveness of ankle braces has found them to reduce the risk
217 of inversion injury (Farwell, et al., 2013) and it is a reduction in coronal plane kinematics which
218 is likely the main contributor to the reduction in risk of inversion injuries (Tang, et al., 2010).
219 Ankle protectors aim to reduce contusion injuries and have previously been found to be
220 effective at this (Ankrah & Mills, 2004). However, **it was previously unknown whether** an
221 ankle protector inadvertently restricts the ankle, due to its location, which may cause
222 restrictions similar to ankle braces. It is evident from the results from the current study that
223 ankle protectors do not significantly restrict the ankle in the coronal plane and replicate similar
224 movement to that of an ankle free of orthotic support. The lack of restriction is due to the soft
225 foam construct of the ankle protector which is far less rigid than the plastic polymer contained
226 within the brace. It is this rigidity that is the main contributor to the ankle braces efficiency
227 at restricting the ankle. **Therefore, ankle protectors do not offer the benefits of protecting**
228 **against ankle inversion injuries like ankle braces.**

229

230 The sagittal plane results produced some interesting observations. The angle at toe off was
231 significantly reduced in the BRACED & PROTECTOR conditions when compared to the
232 WITHOUT condition. Also Absolute ROM was reduced in these conditions too, these results
233 suggest that there is an impedance on the ankle when wearing an ankle protector. The reduction

234 in movement in this plane might be due to the way both the ankle braces and ankle protectors
235 sit on the ankle. The ankle braces have a support strap that runs around the front and rear of the
236 ankle which allows the brace to be tightened. The tightening of this strap is likely to reduce the
237 movement of the ankle by restricting the ankle in the sagittal plane. As for the ankle protector,
238 although the soft foam is designed not to come all the way over the front of the foot, on many
239 of the participants the foam did encroach on the front of the foot due to its “one size fits all”
240 design. The location of the foam at the front of the ankle joint could possibly explain the
241 reduction of sagittal plane movement when wearing the ankle protector. **Reductions in ankle
242 motion in the sagittal plane have been shown to increase energy expenditure (Huang, et al.,
243 2015). The reductions in ankle ROM seen in the current study could suggest that ankle
244 protectors could cause earlier onset of fatigue for a wearer during prolong use such as during
245 competitive match play. This is beyond the scope of the current study but should be
246 investigated further.**

247

248 Although no restrictions of the ankle in the coronal plane were observed for the ankle protectors
249 there is a possibility they might provide proprioceptive cues to the wearer, which may be
250 beneficial to reduce the overall risk of inversion injury. This has been seen with ankle taping
251 where the effectiveness of the tape does not exceed more than approximately fifteen minutes
252 of use (Lohkamp, et al., 2009) but has been found to significantly reduce the risk of ankle injury
253 when compared to not wearing any tape (Verhagen, et al., 2000). Again this is beyond the
254 scope of the current investigation but one that should be researched in the future to compare
255 inversion injury rates of players wearing ankle protectors’ verses players who do not wear ankle
256 protectors.

257

258 Previous research has shown some ankle devices alter knee and hip kinematics which could
259 increase the likelihood of sustaining an injury higher up the kinematic chain (Santos, et al.,
260 2004). Looking at the results of the current study it can be seen that the knee and hip kinematics
261 were found to not be significantly different between the test conditions. The implementation of
262 the ankle braces and ankle protectors used in the current study do not increase the risk of
263 injuring the knee or hip by altering the kinematics of these locations.

264

265 The current study has limited applicability due to the relatively comfortable jogging pace the
266 participants ran at and further research is required to investigate the effects of ankle protectors
267 during nonlinear motion, during jumping, during kicking a football, and also how they affect
268 female footballers. Furthermore, some of the kinematic data show large standard deviations.
269 These large deviations may be due to differing running styles exhibited by the participants, and
270 in some cases such as the hip, due to the movement of the tightly fitted sports shorts worn by
271 participants. Also although markers affixed to the malleoli were not used to track the dynamic
272 movement there is still a possibility that error in their application may cause errors within the
273 data collected as they were used for defining segments in the static model.

274 The current study has established that ankle protectors provide very little restriction to the ankle
275 when jogging and do not restrict the ankle like ankle braces. Therefore, ankle protectors should
276 only be used as a means to reduce risk of ankle contusion injuries and not implemented as a
277 method to reduce the risk of ankle inversion injuries. It must be noted that although no
278 restrictions were seen in the coronal plane there were reductions in sagittal plane motion for
279 the ankle which could possibly increase energy demand needed for locomotion.

280

281 **References**

- 282 Ambegaonkar, J. P. et al., 2011. Ankle stabilizers affect agility but not vertical jump or dynamic
283 balance performance.. *Foot & Ankle Specialist*, 4(6), pp. 354-360.
- 284 Andersen, T. E., Floerenes, T. W., Arnason, A. & Bahr, R., 2004. Video Analysis of the
285 Mechanisms for Ankle Injuries in Football. *The American Journal of Sports Medicine*, 32(1),
286 pp. 69-79.
- 287 Ankrah, S. & Mills, N., 2002. Ankle Protection in Football Shin Guards. *The Engineering of*
288 *Sport*, pp. 128-135.
- 289 Ankrah, S. & Mills, N., 2004. Analysis of ankle protection in Association football.. *Sports*
290 *Engineering*, 7(1), pp. 41-52.
- 291 Bocchinfuso, C., Sitler, M. R. & Kimur, I. F., 1994. Effects of two semi-rigid prophylactic
292 ankle stabilizers on speed, agility, and vertical jump. *Journal of Sport Rehabilitation*, 3(2), pp.
293 125-134.
- 294 Burks, R. T., Bean, B. G., Marcus, R. & Barker, H. B., 1991. Analysis of athletic performance
295 with prophylactic ankle devices. *The American Journal of Sports Medicine*, 19(2), pp. 104-
296 106.
- 297 Cappozzo, A., Catani, F., Croce, U. D. & Leardini, A., 1995. Position and orientation in space
298 of bones during movement: anatomical frame definition and determination.. *Clin Biomech*,
299 10(4), pp. 171-178.
- 300 DiStefano, L. J., Padua, D. A., Brown, C. N. & Guskiewicz, K. M., 2008. Lower Extremity
301 Kinematics and Ground Reaction Forces After Prophylactic Lace-Up Ankle Bracing. *Journal*
302 *of Athletic Training*, 43(3), pp. 234-241.

303 Ekstrand, J., Hägglund, M. & Waldén, M., 2011. Injury incidence and injury patterns in
304 professional football: the UEFA injury study. *British Journal of Sports Medicine*, Volume 45,
305 pp. 553-558.

306 Emery, C. A. & Meeuwisse, W. H., 2010. The effectiveness of a neuromuscular prevention
307 strategy to reduce injuries in youth soccer: a cluster-randomised controlled trial. *British Journal*
308 *of Sports Medicine*, 44(8), pp. 555-562.

309 Farwell, K. E. et al., 2013. The Effectiveness of Prophylactic Ankle Braces in Reducing the
310 Incidence of Acute Ankle Injuries in Adolescent Athletes: A Critically Appraised Topic.
311 *Journal of Sport Rehabilitation*, Volume 22, pp. 137-142.

312 FIFA Communications Division, 2007. [Online]
313 Available at:
314 http://www.fifa.com/mm/document/fifafacts/bcoffsurv/bigcount.statspackage_7024.pdf
315 [Accessed 20 01 2017].

316 Graydon, R., Fewtrell, D., Atkins, S. & Sinclair, J., 2015. The test-retest reliability of different
317 ankle joint center location techniques. *The Foot and Ankle Online Journal*, 8(1), p. 11.

318 Gross, M. T. et al., 1997. Effect of ankle orthoses on functional performance for individuals
319 with recurrent lateral ankle sprains.. *J Orthop Sports Phys Ther.*, 25(4), pp. 245-252.

320 Hägglund, M. et al., 2013. Injuries affect team performance negatively in professional football:
321 an 11-year follow-up of the UEFA Champions League injury study. *British Journal of Sports*
322 *Medicine*, Volume 47, pp. 738-742.

323 Hawkins, R. D. & Fuller, C. W., 1999. A prospective epidemiological study of injuries in four
324 English professional football clubs. *British Journal Of Sports Medicine*, 33(3), pp. 196-203.

325 Huang, T. P., Shorter, K. A., Adamczyk, P. G. & Kuo, A. D., 2015. Mechanical and energetic
326 consequences of reduced ankle plantar-flexion in human walking. *Journal of Experimental*
327 *Biology*, Volume 218, pp. 3541-3550.

328 Janssen, K. W., Mechelen, W. V. & Verhagen, E. A. L. M., 2014. Bracing superior to
329 neuromuscular training for the prevention of self-reported recurrent ankle sprains: a three-arm
330 randomised controlled trial. *British Journal of Sports Medicine*, Volume 48, pp. 1235-1239.

331 Jones, D., Rawnsley, P. & Switzer, A., 2014. *Deloitte*. [Online]
332 Available at: [http://www2.deloitte.com/content/dam/Deloitte/uk/Documents/sports-business-](http://www2.deloitte.com/content/dam/Deloitte/uk/Documents/sports-business-group/deloitte-uk-annual-review-football-finance.pdf)
333 [group/deloitte-uk-annual-review-football-finance.pdf](http://www2.deloitte.com/content/dam/Deloitte/uk/Documents/sports-business-group/deloitte-uk-annual-review-football-finance.pdf)
334 [Accessed 28 03 2015].

335 Junge, A. & Dvorak, J., 2013. Injury surveillance in the World Football Tournaments 1998 -
336 2012. *British Journal of Sports Medicine*, Volume 47, pp. 782-788.

337 Kaplan, Y., 2011. Prevention of ankle sprains in sport: a systematic literature review. *British*
338 *Journal Of Sports Medicine*, 45(4), p. 355.

339 Locke, A., Sitler, M., Aland, C. & Kimura, I., 1997. Long-Term use of a Softshell Prophylactic
340 Ankle Stabilizer on speed, agility, and vertical jump performance. *Journal of Sport*
341 *Rehabilitation*, Volume 6, pp. 235-245.

342 Lohkamp, M., Craven, S., Walker-Johnson, C. & Greig, M., 2009. The Influence of Ankle
343 Taping on Changes in Postural Stability During Soccer-Specific Activity. *Journal of Sport*
344 *Rehabilitation*, Volume 18, pp. 482-492.

345 MacKean, L. C., Bell, G. & Burnham, R. S., 1995. Prophylactic ankle bracing vs. taping:
346 effects on functional performance in female basketball players. *J Orthop Sports Phys Ther*,
347 22(2), pp. 77-81.

348 McGuine, T. A. & Keene, J. S., 2006. The Effect of a Balance Training Program on the Risk
349 of Ankle Sprains in High School Athletes. *The American Journal of Sports Medicine*, 34(7),
350 pp. 1103-1111.

351 Olmsted, L. C., Vela, L. I., Denegar, C. R. & Hertel, J., 2004. Prophylactic Ankle Taping and
352 Bracing: A Numbers-Needed-to-Treat and Cost-Benefit Analysis. *Journal of Athletic Training*,
353 39(1), pp. 95-100.

354 Pedowitz, D. I. et al., 2008. Prophylactic Bracing Decreases Ankle Injuries in Collegiate
355 Female Volleyball Players. *The American Journal of Sports Medicine*, 36(2), pp. 324-327.

356 Salces, J. N. et al., 2014. Epidemiology of injuries in First Division Spanish Football. *Journal*
357 *of Sports Sciences*, 32(13), pp. 1263-1270.

358 Santos, M. J., McIntire, K., Foecking, J. & Liu, W., 2004. The effects of ankle bracing on
359 motion of the knee and hip joint during trunk rotation tasks.. *Clinical Biomechanics*, 19(9),
360 pp. 964-971.

361 Sinclair, J., Edmundson, C. J., Brooks, D. & Hobbs, S. J., 2011. Evaluation of kinematic
362 methods of identifying gait events during running. *Int J Sp Sci Eng*, Volume 5, pp. 188-192.

363 Sinclair, J., Hebron, J. & Taylor, P. J., 2015. The test-retest reliability of knee joint center
364 location techniques.. *Journal of Applied Biomechanics*, Volume 31, pp. 117-121.

365 Sinclair, J. et al., 2014. The Influence of Different Force and Pressure Measuring Transducers
366 on Lower Extremity Kinematics Measured During Running. *Journal of applied biomechanics*,
367 30(1), pp. 166-172.

368 Sinclair, J., Taylor, P. J., Currigan, G. & Hobbs, S. J., 2014. The test-retest reliability of three
369 different hip joint centre location techniques.. *Movement & Sport Sciences*, Volume 83, pp. 31-
370 39.

371 Tang, Y. M., Wu, Z. H., Liao, W. H. & Chan, K. M., 2010. A study of semi-rigid support on
372 ankle supination sprain kinematics. *Scandinavian Journal of Medicine & Science in Sports*,
373 Volume 20, pp. 822-826.

374 Thacker, S. B. et al., 1999. The Prevention of Ankle Sprains in Sports. *The American Journal*
375 *of Sports Medicine*, 27(6), pp. 753-760.

376 Verhagen, E. A. L. M., van Mechelen, W. & de Vente, W., 2000. The Effect of Preventive
377 Measures on the Incidence of Ankle Sprains. *Clinical Journal of Sport Medicine*, 10(4), pp.
378 291-296.

379 Waldén, M., Häggglund, M. & Ekstrand, J., 2013. Time-trends and circumstances surrounding
380 ankle injuries in men's professional football: an 11-year follow-up of the UEFA Champions
381 League injury study. *British Journal of Sports Medicine*, Volume 47, pp. 748-753.

382 Yaggie, J. A. & Kinzey, S. J., 2001. A comparative analysis of selected ankle orthoses during
383 functional tasks. *Journal of Sport Rehabilitation*, 10(3), pp. 174-183.

384

385 **List of figures**

386 **Figure 1.** On the left a pair of Nike ankle shield 10 ankle protectors and on the right an Aircast
387 A60 ankle brace.

388 **Figure 2.** Ankle joint kinematics during the stance phase of locomotion a. sagittal, b. coronal
389 and c. transverse planes (PROTECTOR = black, BRACE = grey, WITHOUT = dash) (DF =
390 dorsiflexion, IN = inversion, EXT = external rotation).

391 **Figure 3.** Knee joint kinematics during the stance phase of locomotion a. sagittal, b. coronal
392 and c. transverse planes (PROTECTOR = black, BRACE = grey, WITHOUT = dash) (FL =
393 flexion, AD = adduction, INT = internal rotation).

394 **Figure 4.** Hip joint kinematics during the stance phase of locomotion a. sagittal, b. coronal and
 395 c. transverse planes (PROTECTOR = black, BRACE = grey, WITHOUT = dash) (FL = flexion,
 396 AD = adduction, INT = internal rotation).

397 **Tables**

398 **Table 1.** Kinematic data (means and stand deviations) for the ankle obtained during stance
 399 phase of the running gait.

	WITHOUT	PROTECTOR	BRACE	
Sagittal plane (+ = dorsiflexion/ - = plantarflexion)				
Angle at footstrike (°)	6.20 ± 7.42	6.05 ± 6.82	4.15 ± 5.64	B
Angle at toe-off (°)	-23.65 ± 4.13	-21.69 ± 3.85	-21.32 ± 3.22	A
Peak dorsiflexion (°)	19.20 ± 3.21	18.46 ± 2.41	17.02 ± 2.09	AB
Absolute ROM (°)	42.66 ± 3.29	40.15 ± 3.73	38.34 ± 2.99	AB
Relative ROM (°)	13.00 ± 6.45	12.41 ± 5.96	12.87 ± 5.41	
Coronal plane (+ = inversion/ - = eversion)				
Angle at footstrike (°)	3.32 ± 2.86	2.54 ± 3.07	1.46 ± 2.55	A
Angle at toe-off (°)	0.02 ± 3.41	-1.06 ± 3.59	-1.24 ± 3.05	A
Peak Inversion (°)	3.87 ± 2.79	3.16 ± 3.07	1.92 ± 2.74	A
Peak Eversion (°)	-9.78 ± 3.70	-10.28 ± 3.78	-8.80 ± 3.74	B
Absolute ROM (°)	13.64 ± 3.23	13.44 ± 3.20	10.72 ± 2.30	AB
Relative ROM (°)	13.10 ± 3.94	12.82 ± 3.69	10.26 ± 2.87	AB
Transverse plane (+ = external/ - = internal)				
Angle at footstrike (°)	-1.15 ± 2.10	-0.56 ± 2.66	-0.43 ± 2.91	
Angle at toe-off (°)	5.06 ± 3.87	5.61 ± 3.95	4.87 ± 4.42	
Peak Internal rotation (°)	-8.82 ± 4.44	-8.33 ± 4.53	-8.06 ± 4.38	
Absolute ROM (°)	13.94 ± 4.18	14.02 ± 4.02	13.12 ± 3.43	
Relative ROM (°)	7.67 ± 3.13	7.78 ± 2.83	7.63 ± 2.47	

400 **Note.** A = significant difference from WITHOUT condition, B = Significant difference from PROTECTOR
 401 condition.

402

403

404

405 **Table 2.** Kinematic data (means and stand deviations) for the Knee obtained during stance
 406 phase of the running gait.

	WITHOUT	PROTECTOR	BRACE
Sagittal plane (+ = Flexion / - = Extension)			
Angle at footstrike (°)	11.99 ± 4.35	12.58 ± 4.36	12.83 ± 3.81
Angle at toe-off (°)	12.49 ± 4.62	14.32 ± 6.05	14.12 ± 5.50
Peak Flexion (°)	40.09 ± 3.97	40.55 ± 3.70	40.17 ± 3.98
Absolute ROM (°)	30.56 ± 4.43	30.31 ± 3.42	29.54 ± 3.54
Relative ROM (°)	28.10 ± 4.96	27.97 ± 4.96	27.34 ± 4.08
Coronal plane (+ = Adduction / - = Abduction)			
Angle at footstrike (°)	0.14 ± 4.18	-0.6 ± 4.24	-0.43 ± 4.50
Angle at toe-off (°)	-3.16 ± 2.78	-3.14 ± 2.92	-3.15 ± 3.00
Peak Adduction (°)	2.92 ± 4.66	2.73 ± 4.66	2.56 ± 4.38
Absolute ROM (°)	6.52 ± 2.40	6.65 ± 2.30	6.42 ± 1.76
Relative ROM (°)	2.79 ± 2.65	2.79 ± 2.76	2.99 ± 2.60
Transverse plane (+ = Internal / - = External)			
Angle at footstrike (°)	-12.96 ± 6.03	-12.18 ± 7.46	-11.94 ± 7.23
Angle at toe-off (°)	-8.37 ± 4.39	-7.52 ± 4.98	-7.17 ± 5.00
Peak Internal Rotation (°)	0.20 ± 6.72	0.62 ± 7.67	0.31 ± 7.22
Absolute ROM (°)	14.07 ± 5.89	13.84 ± 6.32	13.12 ± 6.30
Relative ROM (°)	13.16 ± 6.49	12.25 ± 6.90	12.25 ± 6.69

407 **Note.** A = significant difference from WITHOUT condition, B = Significant difference from PROTECTOR
 408 condition.

409

410

411

412

413

414

415

416

417 **Table 3.** Kinematic data (means and stand deviations) for the Hip obtained during stance
 418 phase of the running gait.

	WITHOUT	PROTECTOR	BRACE
Sagittal plane (+ = Flexion / - = Extension)			
Angle at footstrike (°)	36.72 ± 9.56	37.78 ± 8.34	36.82 ± 8.95
Angle at toe-off (°)	-3.61 ± 8.28	-2.72 ± 7.14	-3.11 ± 7.23
Peak Flexion (°)	39.64 ± 9.24	39.81 ± 9.10	38.70 ± 9.38
Absolute ROM (°)	43.27 ± 9.48	42.45 ± 9.76	41.81 ± 9.64
Relative ROM (°)	40.35 ± 10.18	40.41 ± 9.86	39.93 ± 9.90
Coronal plane (+ = Adduction / - = Abduction)			
Angle at footstrike (°)	4.41 ± 4.87	3.99 ± 4.70	4.55 ± 5.30
Angle at toe-off (°)	0.37 ± 2.36	0.38 ± 3.33	0.46 ± 3.63
Peak Adduction (°)	10.51 ± 5.10	10.75 ± 5.30	10.79 ± 5.81
Absolute ROM (°)	10.86 ± 2.63	11.07 ± 2.53	11.09 ± 2.38
Relative ROM (°)	6.10 ± 3.28	6.76 ± 3.56	6.24 ± 3.76
Transverse plane (+ = Internal / - = External)			
Angle at footstrike (°)	2.48 ± 7.76	2.45 ± 7.50	2.61 ± 8.57
Angle at toe-off (°)	-7.32 ± 6.56	-7.47 ± 7.21	-6.91 ± 6.74
Peak External Rotation (°)	-8.20 ± 6.71	-8.18 ± 7.01	-7.61 ± 6.59
Absolute ROM (°)	11.48 ± 4.24	11.56 ± 4.57	11.14 ± 4.59
Relative ROM (°)	10.68 ± 4.52	10.63 ± 4.83	10.22 ± 4.57

419 **Note.** A = significant difference from WITHOUT condition, B = Significant difference from PROTECTOR
 420 condition.

421

422

Dear Reviewers and Editor,

I have made further revisions based on your feedback and I hope that I have addressed all the concerns that you have. Below I have detailed the changes I have made and I have highlighted the changes in red on the revised version of the paper.

Editor: The referees have reviewed the revision of this interesting paper and both find it much improved. They both have minor concerns that need to be addressed in a second revision. Please make the changes in a red font and also provide a point by point response to the comments. The editor will review the revision.

Reviewer #1: Thank you very much for the re-submission of the paper which in my opinion has improved a lot.

There are three general points before I write some more detailed comments:

1. Why is it so important to know if protectors restrict ankle movement? Please include in the introduction and discussion some answers to this question.

I have added a statement about why ankle protectors restricting movement might be important.

2. You present the kinematic data from the knee and hip but you don't give a reason in the introduction why you collect those data and also in the discussion section you don't write something about this.

I have added a statement explaining the inclusion of knee and hip data in the introduction and have added a section to the discussion too.

3. Be a bit more precise with your conclusion that the movement closely resembles that of an unbraced ankle: don't forget that in sagittal plane there are significant differences

I have changed the conclusion so it is more precise and includes the differences found in the sagittal plane.

Abstract:

Lines 41-44: Adapt the conclusion

Adapted the conclusion to be more precise

Introduction:

Line 76: It's not really true to say that acute or chronic injuries can be prevented depending on the device. Please amend.

I have added the phrase "the risk of..." to this sentence

Discussion:

This section is the weakest section of the article. I am missing more discussion on implications of the changes kinematics on performance/injury... Also, I think you could be a bit more critical with your results: For example include a discussion on the accuracy of data especially when placing the markers on top of the devices. Your data have very large standard deviations - please also include this in the discussion or/and refer to the Effect size. As I said above, I am missing the discussion of knee and hip data.

I have added more information which I hope covers all the points above.

Line 210: Why is it imperative to understand the kinematics? What influence has a restriction on performance/ injury?

I think the sentence I used was a poor choice so I have changed this so the paragraph reads better

Line 213-214: Yes, there is no significant difference if you look at the parameters you have analysed, however, if you look at Figure 2b - it looks like especially in this plane there is a big difference. Is this the graph of one person or the mean of all?

It is the mean of all participants. I have just spotted that the ankle graphs have been mislabelled in the version I have sent in and should be PROTECTOR = dash, BRACE = black, WITHOUT = grey on that one. This is different to the labelling of the knee and hip so I have redone the ankle graphs so the lines match the same style as the knee and hip and are now correctly labelled in the version I have sent in with this submission.

Line 225-229: again, what is the implication on function?

Added information to this paragraph to discuss implication

Line 241: this is not true for sagittal plane.

Lines 245-247: Why is this important? What are the implications for practice?

Added sentences to answer this.

Reviewer #2: This version of the manuscript is much improved, and I appreciate the authors incorporating the reviewers' comments into the revised manuscript. I have some further suggestions to improve the current version.

The abstract refers to both running and jogging. I believe jogging is probably the better word choice here, and throughout the manuscript in any reference to applicability, given that the speed is 3.4m/s, and this would be a relatively slow speed during a soccer match. I feel most would associate the term "running" during soccer with more of a sprint speed. You have generally used jogging throughout the manuscript, which I agree with. I think the few mentions of "running" within the paper are fine as-is, but the tables could be updated to use the term "jogging" for consistency.

Line 105. It should be specified that the 22m distance is part of a runway. It should also be stated what material it is (i.e., rubber track, concrete, artificial turf, etc.).

Line 117. Data are plural, thus data "were" recorded

Changed to "were recorded"

Line 162-163. While I realize you mentioned the means were computed in the previous paragraph, please state "The means of the five trials for each of the three test conditions were compared..."

Added "The means of the five trials for each of..." to the sentence

Line 169. No need to report height, body mass, and BMI beyond one decimal point.

Changed to 1 decimal place

Line 220-221. Awkwardly worded, run-on sentence

I am sorry I did not mention this previously, but it should also be mentioned that the rigid/semi-rigid materials used for both the ankle protector and the ankle brace do not necessarily represent actual movement at the ankle. Markers placed on the skin are subject to some movement, which is a limitation in 3D motion analysis, but markers placed on the surfaces of the protector/brace may differ from the true location of the malleolus during dynamic movements, and this may differ

between the brace and protector. The authors have done a good job describing how they did the best job possible in ensuring proper static position, but the possibility of ankle markers having some error during movement, and the error differing between conditions, must be included in the limitations.

I appreciate that this is an issue however the markers affixed to the malleoli were not used to track the dynamic movement only for defining the segment in the static. The shank was tracked using a rigid plastic mount located on the shank itself and the foot was tracked using the 1st and 5th metatarsal heads and the calcaneus. This allowed me to compute the ankle movement without using the malleoli markers in the dynamic trials.

I appreciate the inclusion of the other various limitations for the final paragraph.

Figure 1- Photos



Figure 2- Ankle




Figure 3- Knee




Figure 4- Hip

