

Central Lancashire Online Knowledge (CLoK)

Title	First-Order Convergence and Roots
Type	Article
URL	https://clock.uclan.ac.uk/17855/
DOI	https://doi.org/10.1017/S0963548315000048
Date	2016
Citation	Christofides, Demetres and Kral, Daniel (2016) First-Order Convergence and Roots. <i>Combinatorics, Probability and Computing</i> , 25 (02). pp. 213-221. ISSN 0963-5483
Creators	Christofides, Demetres and Kral, Daniel

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1017/S0963548315000048>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

First order convergence and roots*

Demetres Christofides[†] Daniel Král'[‡]

Abstract

Nešetřil and Ossona de Mendez introduced the notion of first order convergence, which unifies the notions of convergence for sparse and dense graphs. They asked whether if $(G_i)_{i \in \mathbb{N}}$ is a sequence of graphs with M being their first order limit and v is a vertex of M , then there exists a sequence $(v_i)_{i \in \mathbb{N}}$ of vertices such that the graphs G_i rooted at v_i converge to M rooted at v . We show that this holds for almost all vertices v of M and we give an example showing that the statement need not hold for all vertices.

2010 *Mathematics Subject Classification.* 03C98 (Primary), 05C63, 28E15.

1 Introduction

The theory of limits of combinatorial objects keeps attracting more and more attention and its applications in various areas such as extremal combinatorics, computer science and many others grow. The most understood is the case of dense graph convergence which originated in the series of papers by Borgs, Chayes, Lovász, Sós, Szegedy and Vesztergombi [3–5, 12, 13]. This development is also reflected in a recent monograph by Lovász [11]. Another line of research concentrated around the convergence of sparse graphs (such as those with bounded maximum degree) known as the Benjamini-Schramm convergence [1, 2, 8, 9]. Nešetřil and Ossona de Mendez [14, 15] proposed a notion of first order convergence to unify the two notions for the dense and sparse settings.

First order convergence is a notion of convergence for all relational structures. For simplicity, we limit our exposition to graphs and rooted graphs only but all

*This work was done during a visit to the Institut Mittag-Leffler (Djursholm, Sweden).

[†]School of Sciences, UCLan Cyprus, 7080 Pyla, Cyprus. E-mail: dchristofides@uclan.ac.uk.

[‡]Mathematics Institute, DIMAP and Department of Computer Science, University of Warwick, Coventry CV4 7AL, UK. E-mail: d.kral@warwick.ac.uk. The work leading to this invention has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no. 259385.

our arguments extend to the general setting naturally. If ψ is a first order formula with k free variables and G is a finite graph, then the *Stone pairing* $\langle \psi, G \rangle$ is the probability that a uniformly chosen k -tuple of vertices of G satisfies ψ . A sequence $(G_n)_{n \in \mathbb{N}}$ of graphs is called *first order convergent* if the limit $\lim_{n \rightarrow \infty} \langle \psi, G_n \rangle$ exists for every first order formula ψ . A *modeling* M is a (finite or infinite) graph whose vertex set is equipped with a probability measure such that the set of all k -tuples of vertices of M satisfying a formula ψ is measurable in the product measure for every first order formula ψ with k free variables. In the analogy to the graph case, the *Stone pairing* $\langle \psi, M \rangle$ is the probability that a randomly chosen k -tuple of vertices satisfies ψ . If a finite graph is viewed as a modeling with a uniform discrete probability measure on its vertex set, then the stone pairings for the graph and the modeling obtained in this way coincide.

A modeling M is a limit of a first order convergent sequence $(G_n)_{n \in \mathbb{N}}$ if

$$\lim_{n \rightarrow \infty} \langle \psi, G_n \rangle = \langle \psi, M \rangle$$

for every first order formula ψ . It is not true that every first order convergent sequence of graphs has a limit modeling [15] but it can be shown, e.g., that first order convergent sequences of trees do [10, 16].

Nešetřil and Ossona de Mendez [15, Problem 1] raised the following problem, which we formulate here for graphs only.

Problem 1. *Let M be a modeling that is a limit of a first order convergent sequence $(G_n)_{n \in \mathbb{N}}$ and let v be a vertex of M . Does there exist a sequence $(v_n)_{n \in \mathbb{N}}$ of vertices of the graphs $(G_n)_{n \in \mathbb{N}}$ such that the modeling M rooted at v is a limit of the sequence $(G'_n)_{n \in \mathbb{N}}$ where G'_n is obtained from G_n by rooting at v_n ?*

We prove that the statement from Problem 1 is true for almost every vertex v of M .

Theorem 1. *Let M be a modeling that is a limit of a first order convergent sequence $(G_n)_{n \in \mathbb{N}}$. It holds with probability one that if M' is a modeling obtained from M by rooting at a random vertex v of M , then there exist a sequence $(v_n)_{n \in \mathbb{N}}$ of vertices of $(G_n)_{n \in \mathbb{N}}$ such that M' is a limit of the sequence $(G'_n)_{n \in \mathbb{N}}$ where G'_n is obtained from G_n by rooting at v_n .*

Theorem 1 follows from a more general Theorem 5 which we prove in Section 3. In Section 4, we present an example that the statement of Theorem 1 cannot be strengthened to all vertices v of M , i.e., the answer to Problem 1 is negative. This also answers a more general problem [15, Problem 2] in the negative way.

2 Notation

We assume that the reader is familiar with standard graph theory and logic terminology as it can be found, e.g., in [6, 7]. We briefly review here less standard

terminology and notation only. Throughout the paper, we write $[k]$ for the set of positive integers between 1 and k (inclusively).

There is a close connection between the first order logic and the so-called Ehrenfeucht-Fraïssé games. The p -round Ehrenfeucht-Fraïssé game is played by two players, the spoiler and the duplicator, on two relational structures. We explain the game when played on two graphs G and H . At the beginning of each round, the spoiler chooses a vertex in any one of the two graphs and the duplicator responds with choosing a vertex in the other. One vertex can be chosen several times in different rounds of the game. Let v_i be the vertex chosen in the i -th round in G and w_i the vertex chosen in the i -th round in H . The duplicator wins the game if the subgraph of G induced by v_1, \dots, v_p and the subgraph of H induced by w_1, \dots, w_p are isomorphic through the isomorphism mapping v_i to w_i .

It can be shown that the duplicator has a winning strategy for the p -round Ehrenfeucht-Fraïssé game played on G and H if and only if G and H satisfy the same first order sentences with quantifier depth at most p . More generally, suppose that $\psi(x_1, \dots, x_k)$ is a first order formula with k free variables and with quantifier depth d , G and H are two graphs, and v_1, \dots, v_k and w_1, \dots, w_k are (not necessarily distinct) vertices of G and H , respectively. If the duplicator has a winning strategy for the $(k + d)$ -round Ehrenfeucht-Fraïssé game when played on G and H with the vertices v_1, \dots, v_k and w_1, \dots, w_k played in the first k rounds (so, it remains to play d rounds of the game), then G satisfies $\psi(v_1, \dots, v_k)$ if and only if H satisfies $\psi(w_1, \dots, w_k)$. This correspondence can be used to show [7] that the set $F_{p,q}^m$ of all non-equivalent first order formulas with p free variables and quantifier depth at most q for m -rooted graphs is finite for all positive integers m , p and q (the language for m -rooted graphs consists of a single binary relation representing the adjacency and m constants representing the roots).

3 Almost every rooting is good

In this section, we prove our main result which provides a positive answer to Problem 1 in the almost every sense. As preparation for the proof, we need to establish several technical lemmas.

Lemma 2. *Let ψ be a first order formula for m -rooted graphs and let $[a, b] \subseteq [0, 1]$ be a non-empty interval. For every $\varepsilon > 0$, there exists a first order formula ψ' such that the following holds for every m -rooted modeling M :*

- if $\langle \psi, M \rangle \in [a, b]$, then $\langle \psi', M \rangle > 1 - \varepsilon$, and
- if $\langle \psi, M \rangle \notin (a - \varepsilon, b + \varepsilon)$, then $\langle \psi', M \rangle < \varepsilon$.

Proof. If ψ is a sentence, i.e., it has no free variables, then the statement is trivial. Suppose that ψ has k free variables. Let ψ_n be the first order formula with nk

free variables grouped in n k -tuples such that ψ_n is true if and only if at least $an - n^{2/3}$ and at most $bn + n^{2/3}$ of these k -tuples do satisfy ψ . Formally,

$$\begin{aligned} & \psi_n(x_1^1, \dots, x_k^1, \dots, x_1^n, \dots, x_k^n) \\ = & \bigvee_{i=\lceil an - n^{2/3} \rceil}^{\lfloor bn + n^{2/3} \rfloor} \bigvee_{A \in \binom{[n]}{i}} \left(\bigwedge_{j \in A} \psi(x_1^j, \dots, x_k^j) \wedge \bigwedge_{j \notin A} \neg \psi(x_1^j, \dots, x_k^j) \right). \end{aligned}$$

The Chernoff bound implies that the formula ψ' can be chosen to be the formula ψ_n for n sufficiently large. \square

An interval is a *dyadic interval of order* $k \in \mathbb{N}$ if it is of the form $[a2^{-k}, (a+1)2^{-k}]$ for some integer a . A point x is ε -far from an interval J if $|x - y| \geq \varepsilon$ for every $y \in J$. Otherwise, we say that x is ε -close to J . A *multidimensional interval* is a subset of $[0, 1]^d$ that is the product of d intervals; if J is a multidimensional interval, then J_i denotes the i -th term in the product. A multidimensional interval J is *dyadic of order* $k \in \mathbb{N}$ if every J_i is dyadic of order k .

The next lemma is a direct consequence of Lemma 2. Recall that $F_{p,q}^m$ is the set of all non-equivalent first order formulas with p free variables and quantifier depth at most q , and the set $F_{p,q}^m$ is finite for all m, p and q .

Lemma 3. *Let m, p and q be integers and let $J \subseteq [0, 1]^{F_{p,q}^m}$ be a multidimensional interval. For every $\varepsilon > 0$, there exists a first order formula $\psi_{p,q}^{m,J,\varepsilon}$ such that the following holds for every m -rooted modeling M :*

- if $\langle \psi, M \rangle \in J_\psi$ for every $\psi \in F_{p,q}^m$, then $\langle \psi_{p,q}^{m,J,\varepsilon}, M \rangle > 1 - \varepsilon$, and
- if $\langle \psi, M \rangle$ is ε -far from J_ψ for at least one $\psi \in F_{p,q}^m$, then $\langle \psi_{p,q}^{m,J,\varepsilon}, M \rangle < \varepsilon$.

If $\psi_{p,q}^{m,J,\varepsilon}$ is the formula from Lemma 3, then $\widehat{\psi}_{p,q}^{m,J,\varepsilon}$ is the formula obtained from $\psi_{p,q}^{m,J,\varepsilon}$ by adding m new free variables such that $\widehat{\psi}_{p,q}^{m,J,\varepsilon}$ is satisfied if and only if $\psi_{p,q}^{m,J,\varepsilon}$ is satisfied for the modeling obtained from M by rooting at the m -tuple specified by the new free variables, i.e., the m constants in $\psi_{p,q}^{m,J,\varepsilon}$ are replaced with the new m free variables of $\widehat{\psi}_{p,q}^{m,J,\varepsilon}$. An m -tuple of vertices v_1, \dots, v_m of a modeling M is *negligible* if there exist integers p and q and a dyadic multidimensional interval $J \subseteq [0, 1]^{F_{p,q}^m}$ such that

- $\langle \psi, M' \rangle_{\psi \in F_{p,q}^m} \in J$ where M' is the m -rooted modeling obtained from M by rooting at v_1, \dots, v_m , and
- there exists $\varepsilon_0 > 0$ such that $\langle \widehat{\psi}_{p,q}^{m,J,\varepsilon}, M \rangle \leq \varepsilon$ for every $0 < \varepsilon < \varepsilon_0$.

The next lemma asserts that very few m -tuples can be negligible.

Lemma 4. *If M is a modeling and m is an integer, then the set of negligible m -tuples of M is a subset of a set of measure zero.*

Proof. Note that there are countably many triples p, q and $J \subseteq [0, 1]^{F_{p,q}^m}$ where J is dyadic. Hence, it is enough to show for every p, q and J , that if there exists $\varepsilon_0 > 0$ such that $\langle \widehat{\psi}_{p,q}^{m,J,\varepsilon}, M \rangle < \varepsilon$ for every $0 < \varepsilon < \varepsilon_0$, then there exists a set of measure zero containing all m -tuples v_1, \dots, v_m such that $\langle \psi, M' \rangle_{\psi \in F_{p,q}^m} \in J$ where M' is obtained from M by rooting at v_1, \dots, v_m . Fix p, q and J for the rest of the proof. Let X be the set of all such m -tuples, and let k_0 be an integer such that $2^{-k_0} < \varepsilon_0$.

Let $F_k(v_1, \dots, v_m)$ for $k \in \mathbb{N}$ be the function from M^m to $[0, 1]$ defined to be $\langle \psi_{p,q}^{m,J,2^{-k}}, M' \rangle$ where M' is the modeling obtained from M by rooting at v_1, \dots, v_m . Since the set of tuples satisfying $\widehat{\psi}_{p,q}^{m,J,2^{-k}}$ is measurable, the function F_k is measurable in the corresponding product space. Moreover, it holds that

$$\int F_k(v_1, \dots, v_m) dv_1 \cdots dv_m = \langle \widehat{\psi}_{p,q}^{m,J,2^{-k}}, M \rangle < 2^{-k}$$

for every $k \geq k_0$. Observe that Lemma 3 yields that

$$X \subseteq \bigcap_{k=k_0}^{\infty} F_k^{-1}([1 - 2^{-k}, 1]).$$

Since the function F_k takes values between 0 and 1 (inclusively), the measure of $F_k^{-1}([1 - 2^{-k}, 1])$ is less than $2^{-k}/(1 - 2^{-k})$. It follows that X is a subset of a set of measure zero. \square

We are now ready to prove our main theorem.

Theorem 5. *Let M be a modeling that is a limit of a first order convergent sequence $(G_n)_{n \in \mathbb{N}}$ and let m be a positive integer. It holds with probability one that if M' is a modeling obtained from M by rooting at a random m -tuple of vertices of M , then there exist a sequence $(v_{n,1}, \dots, v_{n,m})_{n \in \mathbb{N}}$ of m -tuples such that the graphs $(G_n)_{n \in \mathbb{N}}$ rooted at these m -tuples first order converge to M' .*

Proof. By Lemma 4, we can assume that the randomly chosen m -tuple w_1, \dots, w_m of the vertices of M is not negligible. It is enough to show for every p, q and $\delta > 0$ that there exists n_0 such that every graph G_n , $n \geq n_0$, contains an m -tuple $v_{n,1}, \dots, v_{n,m}$ of vertices such that the graph G'_n obtained from G_n by rooting at the vertices $v_{n,1}, \dots, v_{n,m}$ satisfies that

$$|\langle \psi, G'_n \rangle - \langle \psi, M' \rangle| \leq \delta \tag{1}$$

for every $\psi \in F_{p,q}^m$. Note that (1) implies that

$$|\langle \psi, G'_n \rangle - \langle \psi, M' \rangle| \leq \delta$$

for every $\psi \in F_{p',q'}^m$ for $p' \in [p]$ and $q' \in [q]$.

Fix the integers p and q and the real $\delta > 0$ for the rest of the proof. Choose an integer k such that $2^{-k} < \delta$ and a real $\varepsilon > 0$ such that $2^{-k} + \varepsilon < \delta$. Further, let $J \subseteq [0, 1]^{F_{p,q}^m}$ be the dyadic multidimensional interval of order k containing the point $\langle \psi, M' \rangle_{\psi \in F_{p,q}^m}$. Since the m -tuple $v_{n,1}, \dots, v_{n,m}$ is not negligible, there exists $\varepsilon' < \varepsilon$ such that

$$\langle \widehat{\psi}_{p,q}^{m,J,\varepsilon'}, M \rangle > \varepsilon'.$$

Since the sequence $(G_n)_{n \in \mathbb{N}}$ converges to M , there exists n_0 such that

$$\langle \widehat{\psi}_{p,q}^{m,J,\varepsilon'}, G_n \rangle > \varepsilon' \quad (2)$$

for every $n \geq n_0$. By the definition of the formula $\widehat{\psi}_{p,q}^{m,J,\varepsilon'}$, the inequality (2) implies that every graph G_n , $n \geq n_0$, contains an m -tuple $v_{n,1}, \dots, v_{n,m}$ of vertices such that

$$\langle \psi_{p,q}^{m,J,\varepsilon'}, G'_n \rangle > \varepsilon' \quad (3)$$

where G'_n is obtained from G_n by rooting at $v_{n,1}, \dots, v_{n,m}$. By Lemma 3, the Stone pairing $\langle \psi, G'_n \rangle$ is ε' -close to J_ψ for every $\psi \in F_{p,q}^m$. It follows that

$$|\langle \psi, G'_n \rangle - \langle \psi, M \rangle| < 2^{-k} + \varepsilon' < \delta$$

for every $\psi \in F_{p,q}^m$. The proof of the theorem is now finished. \square

4 Counterexample

We now show that the statement of Theorem 1 cannot be strengthened to all vertices. Before doing so, we need to introduce some additional notation.

If a (finite or infinite) graph G is bipartite, we write $G(A, B)$ where A and B are the two parts of G . The adjacency matrix M of G is the matrix with rows indexed by A and columns indexed by B such that M_{ab} is equal to 1 if the vertices $a \in A$ and $b \in B$ are adjacent, and it is equal to zero, otherwise. If $G(A, B)$ is a bipartite graph and W is a subset of its vertices, then W_A is $A \cap W$ and W_B is $B \cap W$. The adjacency matrix of G restricted to W is the submatrix with rows and columns indexed by W_A and W_B , respectively. Suppose that W is a subset of vertices of $G(A, B)$, W' is a subset of vertices of $G'(A', B')$ and there is a one-to-one correspondence between the vertices of W and W' . When we say that the adjacency matrices of G and G' restricted to W and W' are the same, we mean that they are the same in the stronger sense that the rows/columns for the corresponding vertices are the same.

A bipartite graph $G(A, B)$ is ℓ -universal, if every vector from $\{0, 1\}^B$ appears at least ℓ times among the rows of the adjacency matrix of G . If W is a subset of vertices of $G(A, B)$, then the ℓ -shadow of W is the multiset S such that each of the vectors $u \in \{0, 1\}^{W_A}$ is included to S exactly $\min\{k, \ell\}$ times where k is the number of times u appears among the columns of the adjacency matrix of

G restricted to $W_A \times (B \setminus W_B)$. If $W_A = \emptyset$, then the ℓ -shadow of W consists of $\min\{|B|, \ell\}$ null vectors (i.e. vectors of dimension zero).

The following is the key lemma in our construction.

Lemma 6. *Let p and q be two non-negative integers. Suppose that $G(A, B)$ and $G'(A', B')$ are two $(p + q)$ -universal graphs and that w_1, \dots, w_q and w'_1, \dots, w'_q are two sequences of the vertices of $G(A, B)$ and $G'(A', B')$, respectively. Let $W = \{w_1, \dots, w_q\}$ and $W' = \{w'_1, \dots, w'_q\}$. If the adjacency matrices of $G(A, B)$ and $G'(A', B')$ restricted to $W_A \times W_B$ and to $W'_{A'} \times W'_{B'}$, respectively, are the same (with the row/column corresponding to w_i being the same as that of w'_i), and the 2^p -shadows of W and W' are also the same, then the duplicator has a winning strategy for the $(p + q)$ -round Ehrenfeucht-Fraïssé game where the vertices chosen in the first q rounds are w_1, \dots, w_q and w'_1, \dots, w'_q .*

Proof. We proceed by induction on p . If $p = 0$, then the graphs induced by the vertices of W and W' are isomorphic since the adjacency matrices of G and G' restricted to W and W' are the same.

Suppose that $p > 0$. By symmetry, we can assume that the spoiler chooses a vertex of G in the next round. Let w_{q+1} be the chosen vertex. If $w_{q+1} = w_i$ for some $1 \leq i \leq q$, the duplicator responds with w'_i . So, we can now assume that w_{q+1} is different from all the vertices w_1, \dots, w_q and we distinguish two cases based on whether w_{q+1} belongs to A or B .

Let us start with the analysis of the case when $w_{q+1} \in A \setminus A_W$. Let x be the row of the adjacency matrix of G corresponding to w_{q+1} . We will construct a vector $x' \in \{0, 1\}^{B'}$ which will determine the response of the duplicator.

Set $x'_{w'_i} = x_{w_i}$ for $w'_i \in W'_{B'}$. Fix a vector $u \in \{0, 1\}^{W_A}$. Let $u_0, u_1 \in \{0, 1\}^{W_A \cup \{w_{q+1}\}}$ be the two extensions of u , and let m_0 and m_1 be the multiplicities of u_0 and u_1 , respectively, in the 2^{p-1} -shadow of $W \cup \{w_{q+1}\}$. Finally, let W_u be the set of the vertices v of $B' \setminus W'_{B'}$ such that the column of v restricted to $W'_{A'}$ is u . If $m_0 + m_1 < 2^p$, then the 2^p -shadow of W' contains the vector u exactly $m_0 + m_1$ times. Set x'_v to 0 for m_0 of the vertices $v \in W_u$ and to 1 for m_1 of such vertices. If $m_0 + m_1 \geq 2^p$, at least one of the numbers m_0 or m_1 is at least 2^{p-1} . If $m_0 \geq 2^{p-1}$, set x'_v to 1 for $\min\{m_1, 2^{p-1}\}$ of vertices $v \in W_u$ and to 0 for all other $v \in W_u$. If $m_0 < 2^{p-1}$ and $m_1 \geq 2^{p-1}$, set x'_v to 0 for m_0 of vertices $v \in W_u$ and to 1 for all other $v \in W_u$. Performing this for every vector $u \in \{0, 1\}^{W_A}$, the entire vector $x \in \{0, 1\}^{B'}$ is defined.

Since the graph G' is $(p + q)$ -universal, there exists a vertex $w'_{q+1} \in A'$ different from the vertices w'_1, \dots, w'_q such that the row of w'_{q+1} in the adjacency matrix of G' is equal to x' . The duplicator responds with the vertex w'_{q+1} . Observe that the choice of x' implies that the adjacency matrices of G and G' restricted to $W \cup \{w_{q+1}\}$ and $W' \cup \{w'_{q+1}\}$, respectively, are the same and that the 2^{p-1} -shadows of $W \cup \{w_{q+1}\}$ and $W' \cup \{w'_{q+1}\}$ are also the same. The existence of the winning strategy for the duplicator now follows by induction.

It remains to consider the case that $w_{q+1} \in B \setminus B_W$. Let u be the column of w_{q+1} in the adjacency matrix of G restricted to W_A . Clearly, u is contained in the 2^p -shadow of W . Consequently, there is a vertex $w'_{q+1} \in B' \setminus B'_{W'}$ such that the column of w'_{q+1} in the adjacency matrix of G' restricted to $W'_{A'}$ is u . The duplicator responds with the vertex w'_{q+1} . The adjacency matrices of G and G' restricted to $W \cup \{w_{q+1}\}$ and $W' \cup \{w'_{q+1}\}$, respectively, are the same. The 2^{p-1} -shadow of $W \cup \{w_{q+1}\}$ in G is obtained from the 2^p -shadow of W by removing u from the shadow and restricting the multiplicity of each vector to be at most 2^{p-1} . Likewise, the 2^{p-1} -shadow of $W' \cup \{w'_{q+1}\}$ in G' is obtained from the 2^p -shadow of W' by removing u from the shadow and restricting the multiplicity of each vector to be at most 2^{p-1} . Note that if $A = A' = \emptyset$, each of the 2^{p-1} -shadows consists of 2^{p-1} null vectors. Since the 2^p -shadows of W and W' are the same, the 2^{p-1} -shadows of $W \cup \{w_{q+1}\}$ and $W' \cup \{w'_{q+1}\}$ are also the same. The existence of the winning strategy for the duplicator now follows by induction. \square

Let $\mathbf{s} = (s_n)_{n \in \mathbb{N}}$ be a sequence of integers such that $s_n \geq 2$ for every $n \in \mathbb{N}$. For each $x \in [0, 1]$, there exists a unique sequence $(x_n)_{n \in \mathbb{N}}$ of integers such that

$$x = \sum_{n=1}^{\infty} \frac{x_n}{\prod_{k=1}^n s_k},$$

$0 \leq x_n < s_n$ for every n , and there is no $n \in \mathbb{N}$ such that $x_n \neq s_n$ and $x_{n'} = s_{n'}$ for every $n' \geq n$. We define $M_{\mathbf{s}}$ to be the following modeling. The vertex set of $M_{\mathbf{s}}$ is the unit square $[0, 1]^2$ with the uniform measure on its Borel subsets. Fix a sequence $(z_n)_{n \in \mathbb{N}}$ of distinct vertices, say $z = (2^{-n}, 0)$, and let $Z = \{z_n, n \in \mathbb{N}\}$. The modeling $M_{\mathbf{s}}(A, B)$ is the bipartite graph with $A = [0, 1]^2 \setminus Z$ and $B = Z$ such that a vertex $(x, y) \in A = [0, 1]^2 \setminus Z$ is adjacent to a vertex $z_n \in B = Z$ if and only if $x_n \neq 0$.

We next verify that every first order definable subset of $M_{\mathbf{s}}^k$ is Borel. A subset X of $M_{\mathbf{s}}^{\ell}$ is *basic* if there exist $v_1, \dots, v_p \in B$ (we allow $p = 0$), a matrix $M \in \{0, 1\}^{\ell \times p}$, an integer q and a multiset $S \subseteq \{0, 1\}^p$ such that the set X is formed by all ℓ -tuples $w_1, \dots, w_{\ell} \in A$ such that the adjacency matrix restricted to $\{v_1, \dots, v_p, w_1, \dots, w_{\ell}\}$ is M and the 2^q -shadow of $\{v_1, \dots, v_p, w_1, \dots, w_{\ell}\}$ is S . In particular, if X is basic, then $X \subseteq A^{\ell}$.

Let $X(\ell, B', M, T)$ for a non-negative integer ℓ , a finite subset $B' \subseteq B$, a matrix $M \in \{0, 1\}^{\ell \times B'}$ and a subset $T \subseteq \{0, 1\}^{\ell}$, be the set of ℓ -tuples $w_1, \dots, w_{\ell} \in A$ such that the the adjacency matrix of $M_{\mathbf{s}}$ restricted to $\{w_1, \dots, w_{\ell}\} \times B'$ is M and all the columns of the adjacency matrix not associated with vertices of B' belong to T after restricting to w_1, \dots, w_{ℓ} . Observe that the set $X(\ell, B', M, T) \subseteq A^{\ell}$ is Borel for all ℓ, B', M and T . Since every basic set is a countable union of sets $X(\ell, B', M, T)$, every basic set is Borel.

Fix a first order formula ψ with k free variables and quantifier depth d . By Lemma 6, the set of k -tuples of $M_{\mathbf{s}}^k$ satisfying ψ can be partitioned into countably

many subsets such that each of them after a suitable permutation of coordinates is either a basic set or a product of a basic set and one or more single element subsets of B . Consequently, every first order definable subset of $M_{\mathbf{s}}^k$ is Borel.

The next lemma directly follows from the definition of a modeling $M_{\mathbf{s}}$.

Lemma 7. *Let $\mathbf{s} = (s_n)_{n \in \mathbb{N}}$ be a sequence of integers such that $s_n \geq 2$ for every $n \in \mathbb{N}$. The modeling $M_{\mathbf{s}}(A, B)$ is ℓ -universal for every $\ell \in \mathbb{N}$. For all integers p and ℓ , it holds with probability one that a random p -tuple of vertices of $M_{\mathbf{s}}$ contains p different vertices from A and the ℓ -shadow of the p -tuple is the multiset containing each vector $\{0, 1\}^p$ with multiplicity ℓ .*

Observe that Lemmas 6 and 7 yield that $\langle \psi, M_{\mathbf{s}} \rangle = \langle \psi, M_{\mathbf{s}'} \rangle$ for every first order formula ψ and any two sequences \mathbf{s} and \mathbf{s}' .

We now define the graph $H_n(A, B)$ to be the graph with $A = [2^n] \times [n]$ and $B = [n]$ such that $(a, a') \in A$ is adjacent to $b \in B$ iff the b -th bit of a when written in binary is 1. We summarize the properties of the graphs H_n in the next lemma.

Lemma 8. *Let p and ℓ be two integers. The graph $H_n(A, B)$ is ℓ -universal if $n \geq \ell$, and the probability that a random p -tuple of vertices of H_n contains p different vertices from A and the ℓ -shadow of them is the multiset containing each vector $\{0, 1\}^p$ with multiplicity ℓ tends to one as n tends to infinity.*

The next theorem follows directly from Lemmas 6–8.

Theorem 9. *Let $\mathbf{s} = (s_n)_{n \in \mathbb{N}}$ be a sequence of integers such that $s_n \geq 2$ for every $n \in \mathbb{N}$. It holds for every first order formula ψ that*

$$\lim_{n \rightarrow \infty} \langle \psi, H_n \rangle = \langle \psi, M_{\mathbf{s}} \rangle .$$

In particular, the modeling $M_{\mathbf{s}}$ is a limit of $(H_n)_{n \in \mathbb{N}}$.

Let $\psi_0(x)$ be the first order formula that is true if x is adjacent to the root. If $s_n = 3$ for every $n \in \mathbb{N}$, then the set of neighbors of every vertex of B in the modeling $M_{\mathbf{s}}(A, B)$ has measure $2/3$. Hence, if $M'_{\mathbf{s}}$ is the modeling obtained from $M_{\mathbf{s}}$ by rooting at an arbitrary vertex of B , then $\langle \psi_0, M'_{\mathbf{s}} \rangle = 2/3$. Since no vertex of H_n is adjacent to more than $2^{n-1}n$ vertices (the vertices of A are adjacent to at most n vertices each and each vertex of B is adjacent to $2^{n-1}n$), it holds that

$$\limsup_{n \rightarrow \infty} \langle \psi_0, H'_n \rangle \leq \frac{1}{2}$$

for any sequence $(H'_n)_{n \in \mathbb{N}}$ of rooted graphs obtained from H_n . We conclude that the sequence $(H_n)_{n \in \mathbb{N}}$, the modeling $M_{\mathbf{s}}(A, B)$ with $\mathbf{s} = (3)_{n \in \mathbb{N}}$ and rooting $M_{\mathbf{s}}$ at any vertex of B provide a counterexample to Problem 1.

References

- [1] D. Aldous, R. Lyons: Processes on unimodular random networks, *Electron. J. Probab.* 12 (2007), no. 54, 1454–1508.
- [2] I. Benjamini, O. Schramm: Recurrence of distributional limits of finite planar graphs, *Electron. J. Probab.* 6 (2001), no. 23, 1–13.
- [3] C. Borgs, J.T. Chayes, L. Lovász, V.T. Sós, K. Vesztergombi: Convergent sequences of dense graphs I: Subgraph frequencies, metric properties and testing, *Adv. Math.* 219 (2008), 1801–1851.
- [4] C. Borgs, J.T. Chayes, L. Lovász, V.T. Sós, K. Vesztergombi: Convergent sequences of dense graphs II. Multiway cuts and statistical physics, *Ann. of Math.* 176 (2012), 151–219.
- [5] C. Borgs, J. Chayes, L. Lovász, V.T. Sós, B. Szegedy, K. Vesztergombi: Graph limits and parameter testing, in: *Proc. 38rd Annual ACM Symposium on the Theory of Computing (STOC)*, ACM, New York, 2006, 261–270.
- [6] R. Diestel: *Graph theory*, Springer-Verlag, Heidelberg, 2010.
- [7] H.-D. Ebbinghaus, J. Flum: *Finite model theory*, Springer-Verlag, Heidelberg, 2005.
- [8] G. Elek: Note on limits of finite graphs, *Combinatorica* 27 (2007), 503–507.
- [9] H. Hatami, L. Lovász, B. Szegedy: Limits of local-global convergent graph sequences, to appear in *Geometric and Functional Analysis*.
- [10] D. Král’, M. Kupec, V. Tůma: Modelings of trees, in preparation.
- [11] L. Lovász: *Large networks and graph limits*, AMS, Providence, RI, 2012.
- [12] L. Lovász, B. Szegedy: Limits of dense graph sequences, *J. Combin. Theory Ser. B* 96 (2006), 933–957.
- [13] L. Lovász, B. Szegedy: Testing properties of graphs and functions, *Israel J. Math.* 178 (2010), 113–156.
- [14] J. Nešetřil, P. Ossona de Mendez: A model theory approach to structural limits, available as [arXiv:1303.2865](#).
- [15] J. Nešetřil, P. Ossona de Mendez: A unified approach to structural limits, and limits of graphs with bounded tree-depth, available as [arXiv:1303.6471](#).
- [16] J. Nešetřil, P. Ossona de Mendez: Modeling limits in hereditary classes: reduction and application to trees, available as [arXiv:1312.0441](#).