

**PROFILING HEAD ACCELERATIONS USING TRIAXIAL ACCELEROMETRY DURING ‘BODIES IN FRONT’ TRAINING PLAY: A PRELIMINARY STUDY OF THE SPORT OF RUGBY LEAGUE**

1 Stephen J Atkins\*, 2 Ian Bentley, 2 Howard T Hurst, 2 Mark E Mark E, 3 Cassandra Richardson, 2 Jonathan K Sinclair.

1 School of Health Sciences, University of Salford, England; 2 Centre for Applied Sport and Exercise Sciences, University of Central Lancashire, England; 3 School of Psychology, University of Central Lancashire, England

10.1136/bjsports-2016-097270.19

Objective; To provide unique descriptive information on head accelerations, both linear and rotational, during ‘bodies in front’ training play.

Design A convenience sample were tested using a cross-sectional design. Players wore a triaxial accelerometer (XPatch, X2 Biosystems, Seattle) during two sessions of training gameplay (session duration  $142 \pm 26$  min). Accelerometers were affixed on the left side of the head, at a point approximate to the external occipital protuberance. Setting Testing was undertaken at an elite performance training centre.

Participants 20 elite, professional, rugby league players.

Main outcome measures; Minimum recordable accelerations were set at 10 g (linear acceleration) and 25 rads s<sup>-2</sup> (rotational acceleration).

Results; A total of 1256 recorded ‘hits’ were analysed, excluding evident artefacts. No player was reported to have suffered a concussion. Player ‘loading’ was  $46 \pm 33$  hits per session (range 7– 170). Average linear ( $14.9 \pm 8.8$  g) and rotational ( $532 \pm 347$  rads s<sup>-2</sup>) accelerations were not significantly different between training sessions ( $P > 0.05$ ). Frequency of hits are summarised in the table below

Linear acceleration (g)	Cumulative%	Rotational acceleration (rads s <sup>-2</sup> )	Cumulative%
10–20	92.6	25–500	64.7
21–40	5.4	501–1000	27.5
41–60	0.7	1001–2000	5.4
61–80	0.5	2001–3000	1.6
81–100	0.4	3001–4000	0.8
>101	0.4	>4001	0

Conclusions; The majority of head accelerations occurred within lower ‘sub-concussive’ zones. This confirms existing research findings from other sports. Whilst few accelerations occurred in higher zones, there were a number of very high loading impacts observed in ‘hit up forward’ positions. Our research confirms the suitability of using portable accelerometers in determining head accelerations in full-contact, ‘armourless’, invasion games. Competing interests None.