

Article

Experiences of patients with mental illness' interactions with medical students: a systematic review

Joiner, Adam, Dearman, Samuel, Gordon, Morris and Vince, Gill

Available at <http://clock.uclan.ac.uk/19466/>

Joiner, Adam, Dearman, Samuel, Gordon, Morris and Vince, Gill (2018) Experiences of patients with mental illness' interactions with medical students: a systematic review. The Canadian Journal of Psychiatry, 63 (1). pp. 4-11. ISSN 0706-7437

It is advisable to refer to the publisher's version if you intend to cite from the work.

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

EXPERIENCES OF PATIENTS WITH MENTAL ILLNESS' INTERACTIONS WITH MEDICAL STUDENTS: A SYSTEMATIC REVIEW

Abstract

Objectives

Mental health is a key area for learning within undergraduate medical education. Given the nature of mental illness, interactions may have the potential to uniquely impact patients. This study set out to systematically review studies reporting experiences and perceptions of patients with mental illness' clinical interactions with medical students. This includes which factors encourage patients to interact with medical students, and if patients perceive negative and positive effects from these interactions.

Method

Studies reporting patient experiences of involvement in undergraduate medical ~~ineal~~ were included. A standardized search of online databases was carried out independently by two authors and consensus reached on the inclusion of studies. Data extraction and quality assessment were also completed independently, after which a content analysis of interventions was conducted and key themes extracted. Studies were included from peer-reviewed journals, in any language.

Results

Eight studies from five countries were included, totaling 1088 patients. The majority of patients regard interacting with medical students as a positive experience. Patients described feeling comfortable with medical students, and the majority believes it is important for students to "see real patients". Patients describe benefits to them as enjoyment, being involved in student education, and developing illness narrative.

Conclusions

Results suggest that most patients with mental illness want to interact with medical students, and this should be encouraged during student placements. Further research

however is required to understand in more depth what else can be done to improve the comfort and willingness for patients to interact with students, including barriers to this.

Declaration of interests

None

Given that quality and safety of medical care is dependent on the education of medical students,¹ it is important to understand how patients experience the interaction with medical students. Experiential learning is an important, evidence-based part of undergraduate medical development² and interaction with patients remains a requirement in order that medical students learn and develop. Learning occurs by observing and reflecting on experiences, then forming abstract concepts and generalizations based on reflections, and finally testing the concepts in new situations.³ Medical students can interact with patients at four progressive levels: passive observer, active observer, actor in rehearsal, actor in performance.^{4,5} The progression through these levels occurs as students become more senior, which is imperative to an evolving sense of participation and contributing to ongoing learning and development.⁵ Learning is enhanced when the student experience of participation is simultaneously challenging and supportive,^{4,5,6} such as increasing independence alongside quality supervision.⁶ Workplace experience including patient interaction promotes the development of professional identity,⁷ as well as the skills and knowledge required for clinical practice.

With respect to mental illness, it is regrettable that medical students can sometimes experience discouragement from clinical staff when interacting with patients ~~with certain symptoms such as self-harm~~, including in psychiatry placements. ~~A lack of understanding by clinical staff as to how people with mental illness feel about interacting with medical students might explain this observation. This is an area of particular importance to explore~~ It is important to understand this further given the importance of experiential learning, ~~given as that~~ patients are potentially vulnerable, and medical student attitudes towards psychiatry^{8,9} and people with mental illness¹⁰ may be negative and stigmatizing, compared to physical illness. Negative attitudes towards mental illness may become increasingly apparent as students progress through medical school¹⁰ and such negative attitudes can become harmful.¹¹ ~~It is of particular interest to understand how patients experience interaction~~ ing with medical students. By definition the interaction between patients and students is a dyadic process and whilst the literature exploring the effects of experiential and workplace learning richly describes the perspective of medical students,¹² it is equally important to understand the perspective of patients. There is a professional duty to

understand the experience of patients for clinical reasons, but this information is also of undoubted value in improving the educational quality of the interaction.

We investigated the experiences of patients with mental illness with respect to' clinical interactions with medical students, addressing two key objectives. Firstly, to explore the nature of the experiences of interacting with medical students. This included~~s~~ which factors encourage, or discourage, patients to interact with medical students (Descriptive). Secondly, to explore whether patients with mental illness perceived contact with medical students as having positive or negative outcomes, either for themselves or the students (Justification).¹³

Methods

We sought to understand the impact of interacting with medical students on patients with mental illness and how patients perceive this interaction as impacting on the students. Whilst this work is not directly situated within an educational environment, ~~as~~ the focus is ~~very much~~ within the context of medical education, the review has been reported in line with the methodology of the Best Evidence Medical Education Collaborative¹⁴ and STORIES guidelines.¹⁵⁴

As each patient can reasonably be expected to experience interactions with medical students differently, an absolute single truth as to this experience is not possible. Instead, we wanted to establish a 'probable' truth,¹⁶⁵ aligning with a research paradigm of post-positivism^{165,176} supporting evidence--informed translation to practice of the results.¹⁸⁷ Ethical review was not sought as it is not required for systematic reviews.

All study designs were considered for this review, as long as they reported experiences and attitudes of patients involved with medical students. For the purpose of this review, 'experience' was defined as observations made through contact with medical students, 'perception' as the way in which patients regard, understand, feel or interpret interaction with medical students, 'view' as a particular way of considering or regarding medical students, 'attitude' as an established way of thinking, feeling about, and behaving towards medical students and 'opinion' as a judgement formed about medical students.

We accepted studies with any patient with mental illness aged 18 or over who has interacted with medical students for the purpose of an assessment. All settings were acceptable, such as inpatients and outpatients, and all psychiatric diagnoses were included. Studies needed to include empirical data for selection with respect to patient experience, perceptions, views, attitudes or opinions on either (a) how patients believed/felt clinical interactions with medical students impacts on them or other patients, and/or (b) how patients believed/felt students' interaction with 'real' patients impacts on medical students. We excluded studies which explored medical students' experiences/views,

healthcare professionals experiences/views, and healthcare students who are not medical students.

Commentary pieces, ~~surveys,~~ audits ~~and/or~~ review articles were not included. We excluded from this review studies exploring the experiences of patients whose primary diagnosis is not a mental illness and the experiences of 'professional' patients (patients who are trained to provide specific interactions for students with specific objectives, for example in an examination), the medical students themselves, simulated patients or involvement of patients who have had any medical education training. The focus was the experiences of patients with mental illness who are interacting with medical students in the 'real' clinical environment, i.e. an assessment by a medical student. We included any empirical studies, but excluded opinion pieces. Studies from all countries published in all languages were included. There was no time limit on the search.

We ran the following exploded search terms combination using Boolean operations in ~~July~~ ~~March~~ 2017: [psychiatr* OR "mental" OR Patient OR Patients OR "service user*"] AND ["medical student*" OR "student*"] AND [experience* OR Attitude OR Attitudes OR Opinion OR Opinions OR View OR Views OR perception*]. The databases searched using these terms were Medline and PsycInfo. A grey literature search was not performed due to resource limitations, however hand searching of selected article references was performed.

Citations were reviewed independently by each of the authors. Potentially relevant abstracts were independently reviewed and full papers obtained for any studies that appeared to meet the inclusion criteria. Disputes were resolved by consensus. The full manuscripts for all included studies were assessed independently by three of the authors. A data extraction summary was completed for each selected study. The descriptive data included: study setting, sample size, country, year of study, methods and measures, key findings and outcomes, and study limitations. Given the nature of the objectives, the data were analyzed using a descriptive and narrative approach. We assessed quality of the studies by scoring five key areas: student sample define, patient sample defined, the study design is appropriate to answer the research question, implementation of the methods,

appropriateness of the analysis. Each of these five items a score was applied where 0=no; 1=to some extent; 2=yes. Therefore the maximum quality score was 10 and minimum 0. The quality scores for each study is in Table 1.

Results

There were ~~10,482~~11,103 potentially relevant citations retrieved through searching. Of these, ~~3029~~ articles were identified as potentially relevant and abstracts were reviewed. After review of these studies, ~~1645~~ were excluded as they did not meet the inclusion criteria. In the case of four further studies it was unclear if they included patients with mental illness. After communications were made with the authors, these four studies were excluded as no patients with mental illness were included. This left ten articles that met the inclusion criteria.¹⁹⁸⁻²⁸⁷ However, two of these ten studies were excluded as they involved patients in primary care²¹⁹ and another study described ~~an~~ idiosyncratic practice which is not transferable to most clinical or educational environments.²⁷⁶ This left eight studies included ~~for in the~~ data extraction and synthesis. No further articles were found as a result of hand searching of references (Figure 1). A total of 1088 patients from five different countries were included in the accepted studies for the data extraction and synthesis. None of the studies were conducted in Canada.

Table 1 summarizes the included studies. All studies used questionnaires/surveys with Likert scales plus open-ended questions, except that by Dogra *et al*,²⁸⁷ which was a qualitative study using focus groups. The surveys used by Lynoe *et al*²⁰¹⁹ and Santulli²³² did not use open-ended questions. Black & Church²⁴³ also included qualitative analysis of interviews with patients, with a mean length of interview of five minutes.

The nature of the experiences of interacting with medical students from the patients' perspective

A “positive” experience of being assessed by medical students is described by 60%¹⁹⁸ and 85%²⁰⁴⁹ of patients, with one study finding that 97% of patients experience interactions with students as “valuable”.²¹²⁰ Öster *et al*²²⁴ found strong agreement from patients that they should have clinical interaction with medical students, an opinion with which patients were more likely to agree with if they had previously interacted with a medical student (mean of 4.53 on a five-point Likert scale) than not (mean of 4.34), $p=0.024$.

Öster *et al*²²⁴ found that 76% of patients felt “comfortable” interacting with medical students and that inpatients and outpatients were equally “comfortable”. In this study, reported comfort reduced slightly when the degree of supervision was specified, dropping from 73% under supervision to 61% without. In the same study, male and female patients were equally comfortable with female students, but female patients were comparatively less comfortable with male student (mean reported comfort level reducing from 3.89 to 3.38 on five-point Likert scales). Female patients were less comfortable interacting with “younger” medical students than male patient, with means of 3.14 and 3.47, respectively, $p=0.01$. When provided with statements in terms of dis-comfort, Santulli found patients generally disagreed (mean of 2.49 on a five-point Likert scale) that interacting with medical student was an uncomfortable experience.²³²

Two studies referred to how the patients experienced the medical students as professionals. Santulli²³² found agreement from patients that students were “kind”, with a mean of 4.51 on a five-point Likert scale. Black and Church²⁴³ found strong agreement that students were “warm and caring” (mean of 4.59 on a five-point Likert Scale) as well as reporting disagreement that they felt “talked down to” (mean of 1.57) or that interacting with medical students felt “unnatural” (mean of 1.63).

Four studies explored patients’ willingness to interact with medical students again in the future. Santulli²³² found agreement that patients are willing to interact with medical students again in the future (mean of 4.31). Rates of willingness to interact with medical students again were reported in three other studies, with rates reported at 91%²¹⁰, 79%²⁵⁴ and 70%²⁶⁵.

Three studies reported on consent. Doshi *et al*²⁵⁴ ~~also~~ reported that 30% did not feel they received an appropriate explanation as to the student's task but that, 93% of patients felt they had been properly consented. Tunde-Ayinmode²⁶⁵ reported this at 83%. Lynoe⁴⁹²⁰ didn't report on consent rates, but found that 40% of patients stated that they would interact with medical students if put "under duress".

Öster *et al*²²⁴ qualitatively explored patient's wishes when interacting with medical students. They found that patients wanted prior notice about age, gender, seniority of students and clarity as to the purpose of interacting with the students.

Patients' perception of positive and negative outcomes from interacting with medical students

Regarding positive outcomes, two studies report rates of "enjoyment" of patients interacting with medical students at 56%²⁷⁶ and 87%²⁶⁵. Santulli²³² found patients typically agreed that student interviews were enjoyable (mean of 3.95 on a five-point Likert scale). Black & Church²⁴³ found that as a result of interacting with medical students, 89.7% of patients felt that the medical students were helpful to their overall treatment, as well as patients agreeing they experienced greater explanations about their treatment (mean of 4.11), involvement in decisions about their care (mean of 3.94), and useful answers to their questions (mean of 4.26). Tunde-Ayinmode²⁶⁵ found that 37% of patients felt that interacting with patients aided their recovery. Gundel & Wefelmaier¹⁹⁸ found 35% of patients reported deeper insight into their illnesses and 24.5% felt they had learnt about themselves. While these numbers appear low for sense of recovery, deeper insight, and learning about themselves, the remainder of patients did not notice any change in these domains, suggesting this may be a benefit for some patients only. Interacting with medical students did not negatively affect these domains. However, Öster *et al*²²⁴ found patients were neutral (mean of 3.33) about whether interaction with students improved quality of care.

Qualitative results described patients feeling they benefited from student interactions by being the *provider* of learning,²⁶⁵ enjoyment,^{224,254} and having a sense of occupation.²⁵⁴

Patients perceived positive outcomes not only for themselves, but for the medical students also. Tunde-Ayinmode *et al*²⁶⁵ reported that 83% of patients felt students' learning was "enhanced" by interacting with "real" patients, with a similar finding by Doshi *et al*²⁵⁴ where the mean score on a six-point Likert scale was 5.01 on a scale of agreement with the statement "seeing real patients is valuable training for medical students".

From qualitative data Doshi *et al*²⁵⁴ found that patients felt interacting with medical students "helps them [students] learn" and makes medical students "good doctors". Öster *et al*²²⁴ found patients felt students learned aspects of empathy and professionalism. Dogra *et al*²⁸⁷ found that patients felt that students learned about "humanity" and the "whole person" as well as developing a sense of "professional identity". They also suggested that students benefited as the experience reduces stigma and engenders an attitude of "hope" and "recovery".

Four studies reported on dissatisfaction experienced by patients. Gundel & Wefelmaier¹⁹⁸ reported 10% of patients describing the experience as "unpleasant" and "uncomfortable". Öster *et al*²²⁴ found that nearly 25% of patients were "not comfortable" interacting with students. Tunde-Ayinmode²⁶⁵ found that 23% of patients were "worried" about interacting with medical students, and 13% of patients felt interacting with medical students was a "waste of time". It is concerning that 53% of patients saw students because they didn't want care to be compromised, and 13% withheld clinical information because they were being assessed by medical students.²⁶⁵ Doshi *et al*²⁵⁴ reported 12% of patients felt that they could not decline interactions with students, 21% that they could not terminate interviews, and 14% felt that such interviews were not confidential. Doshi *et al*²⁵⁴ also explored the chronology of anxiety which patients experienced; they found that 11% were "nervous" at the prospect of being asked to interact with a medical student (although this reduced overtime), and 21% became distressed during interview and 7% after the interview had concluded. None of these studies attempted to establish why some patients had these negative experiences or opinions.

Discussion

The published evidence demonstrates that the experience of the large majority of patients with mental illness is positive when interacting with medical students. The students were similarly described in a positive light, seen as caring, kind and competent and the majority of patients would interact with medical students again. This is similar to the experiences of patients with physical illness when interacting with medical students, who also found students to be “polite” and “kind”.²⁹⁸ This trend continues when patients interact with first and second year medical students, with almost all patients very satisfied with the encounter and 85% prepared to interact with students again.²⁹³⁰ These findings seem to be consistent regardless of the physical health specialty the patients are being treated in.³¹⁹

The evidence suggests patients benefit from interacting with medical students, ranging from a sense of occupation, being a provider of learning, and feeling more involved in their own care. Patients also described improved self-esteem and insights, although the level of agreement for this appears weaker. Studies exploring the perceptions of patients with physical illness interacting with medical students have found patients feel that the quality of care improves, when comparing a hospital before and after the introduction of medical students.^{324,332} Interestingly, patients with physical illness perceived the care they received from physicians improved following the introduction of medical students to a hospital in domains such as friendliness, competence and time spent with patients.³²⁴

The data from the studies reviewed suggests that the majority of patients feel strongly that they should interact with students for a variety of reasons. This would appear to counter the position often encountered in clinical practice that patients with mental illness are often felt as “unsuitable” for clinical interaction with medical students. This review suggests that most people with mental illness can and should interact with medical students. There is less evidence from patients with mental illness that interacting with medical students directly improves quality of care in terms of health outcomes, however the subjective experience of care does appear to improve.

The literature also suggests there are potential negative outcomes for patients, although in a lower order of magnitude than positive outcomes. For example, about one fifth of patients will not be comfortable interacting with medical students. This appears to relate to feelings of vulnerability such as fears of loss of confidentiality, not being able to decline, as well as inconvenience. It is notable that in primary care nearly 70% of patients seeing their GP for a mental health problem don't believe the quality of their care will be positively affected if a medical student is involved, contrary to those patients with physical illness in the hospital setting.³²⁴ In physical health patient populations, the level of perceived negative outcomes appears dependent on the setting, for example there were no perceived negative outcomes in a study in USA,²⁹³⁰ while non-Caucasian populations in the UK are twice as likely to feel uncomfortable interacting with medical students as the Caucasian population.³⁴³ In Kuwait 20% to 25% of patients feel uncomfortable enough to refuse student participation.³⁵⁴ For people with mental illness receiving care in Canada, the literature has yet to explore experience clinical interactions with medical students, and whilst findings from USA and Europe are likely to be at least partially transferable further confirmatory research is required.

Although not a specific objective of this systematic review, it is encouraging that the results serve to challenge stigma against those with mental health problems in a number of ways. Patients with mental health problems appear equally willing to engage with medical students when compared to patients with physical health problems. Patients with mental health problems shouldn't be automatically viewed as frail and not able to interact with students, in fact a number of benefits are described. It is also important to note that patients themselves report that interacting with medical students positively impacts on attitude formation of tomorrow's doctors and reduces stigma.

There is less evidence in terms of patients' perceptions of benefits to students. However in the small number of studies in this review that consider this, the majority of patients view interacting with medical students as an enhancement to learning. Patients appear to view this benefit in terms of learning of empathy and humanity, as well as developing

professionalism and professional identity. Encouragingly, patients with mental illness have also reported that interacting with medical students promotes a sense of hope and the possibility of recovery.

-Lessons for improved practice in medical education can be drawn from these findings. It is useful for those tasked with managing student placement numbers that within the population of patients around 4 in 5 patients will consent to see medical students, and thus the educational capacity of the services can be better understood and patients not be overwhelmed. More detailed educational improvements can be made when noting the clear and recurrent theme in use of language both in the qualitative and quantitative studies, that of “comfort” of patients in relation to interacting with medical students. This appears to be the preferred term or outcome. This being the case, effort should be made jointly between clinical services and medical education departments to have policies and procedures in place that aim to ensure the comfort of their patients. The aim should be to promote and increase levels of comfort. More specifically, three points of good practice are suggested. Firstly, patients should be provided with specific information as to the reported benefits of seeing medical students. Secondly, specific patient briefings occur prior to interacting with medical students and should include the purpose of their interaction, the level of supervision being provided, the number of students, student gender(s) and year of study. Lastly, specific support should be available during interview and afterwards, being mindful of the reported potential for delayed distress.

Strengths and Limitations

This study captures the known international body of evidence, without limits on language or date of study followed by a robust approach to selection, quality assessment and data extraction. The resulting synthesis supports clinical interactions between medical students and people with mental illness. Therefore, this study provides a foundation to challenge any practice which is discouraging medical students from interacting with people with mental illness. This study also supports any psychiatrist who wants to incorporate medical students’ involvement in enhancing patient care.

The foremost limitation of this study arises by virtue of the nature of the primary evidence base. The majority of studies described summary descriptive statistics of Likert scales or alternatively individual comments or responses. There were no studies that undertook a deeper clarification analysis,¹⁴³ aiming to generate new theory or conceptual frameworks to aid understanding in this area. Such understanding could further explore the experiences of patients that have been raised as well as the mechanisms of positive or negative outcomes that have been synthesized. There may be a response bias in the primary research with patients who have enjoyed or benefitted from interaction with students being more likely to participate in the studies. Additionally, a lack of precision as to the terms of the metrics being considered in the primary studies is likely to put limits on what can be concluded. Although as a part of this review we defined our terms relating to patients' experiences and views, this was not seen in the primary literature and in many cases terms appear to be used somewhat interchangeably. Due to resource limitations, a variety of search terms in different combinations was not used, but may have uncovered further literature, given the nature of the terms used.

A range of patients from different settings have contributed to the studies considered. Universally, the description of the patients themselves was quite limited and did not extend to the details of their current diagnoses, for example. As such this review can conclude that, at the level of care setting, the majority of patients are comfortable interacting with medical students and report positive experiences and benefits, but there can be no comment as to variation by other factors such as the effect of suffering from a particular mental health problem.

Conclusions

This review suggests that most patients with mental illness have positive experiences interacting with medical students, and report perceived benefits to themselves and to students. Patients are more likely to be willing to interact with medical students than not, but it is important to enhance the patient experience through proper consent, which includes informing patients of the student's gender, year of study, task expected of the student and level of supervision the student will be receiving. This may contribute to increasing the number of patients who have a positive experience interacting with students.

However, future research is required to understand what barriers patients perceive to interacting with students, and what aspects of the interaction enhance the experience. This includes exploring the effects of diagnosis on this experience.

References

1. General Medical Council. Recognising and approving trainers: the implementation plan. 2012. http://www.gmc-uk.org/Approving_trainers_implementation_plan_Aug_12.pdf 56452109.pdf (accessed 29th September 2015).
2. Taylor DCM, Hamdy H. Adult learning theories: Implications for teaching and learning in medical education: AMEE guide No. 83. *Medical Teacher*. 2013; 35: e1561-e1572.
3. Kolb DA (1984) *Experiential learning: experience as the source of learning and development*. Prentice Hall: New Jersey.
4. Dornan T, Boshuizen H, King N, Scherpbeier A. Experience-based learning: a model linking the process and outcomes of medical students' workplace learning. *Medical Education*. 2007; 41: 84-91.
5. Dornan T, Scherpbeier A, Boshuizen H. Supporting medical students' workplace learning: experience-based learning (ExBL). *Clinical Teacher*. 2009; 6(3): 167-171.


6. Van der Zwet J, Zwietering P, Teunissen P, van der Vleuten C, Scherpbeier A (2010) Workplace learning in general practice: supervision, patient mix and independence emerge from the black box once again. *Medical Teacher*. 32: e294-299.
7. Hay A, Smithson S, Mann K, Dornan T. Medical students' reactions to an experience-based learning model of clinical education. *Perspectives in Medical Education*. 2013; 2(2): 58-71.
8. Wilkinson DG, Greer S, Toone BK. Medical students' attitudes to psychiatry. *Psychological Medicine*. 1983; 13(1): 185-192.
9. Feldman TB. Medical students' attitudes towards psychiatry and mental illness. *Academic Medicine*. 2005; 29(4): 354-356.
10. Korszun A, Dinos S, Ahmed K, Bhui K. Medical student attitudes about mental illness: does medical-school education reduce stigma? *Academic Psychiatry*. 2012; 36(3): 197-204.
11. National Institute for Health and Clinical Excellence. Commissioning Guides: Support for commissioning for self-harm. 2013.
<http://publications.nice.org.uk/support-for-commissioning-for-selfharm-cmg50.pdf> (accessed on 29th November 2013).
12. Wigney T, Parker G. Factors encouraging medical students to a career in psychiatry: Qualitative analysis. *Australian and New Zealand Journal of Psychiatry*. 2008; 42(6): 520-525.
- ~~13.~~ 13. Cook DA, Bordage G, Schmidt HS. Description, justification and clarification: a framework for classifying the purposes of research in medical education. *Medical Education*. 2008 42:128-33.
- ~~13-14.~~ 13-14. <https://www.bemecollaboration.org/> (accessed 20th July 2017)
- ~~14-15.~~ 14-15. Gordon M, Gibbs T. STORIES statement: Publication standards for healthcare education evidence synthesis. *BMC Medicine*. 2014; 12:143 DOI: 10.1186/s12916-014-0143-
- ~~15-16.~~ 15-16. Bunniss S, Kelly DR. Research paradigms in medical education research. *Medical Education*. 2010; 44: 358-366.

- ~~16-17.~~ Gordon M. 2016. Are we talking the same paradigm? Considering methodological choices in health education systematic review. *Medical Teacher*. 2016; 23: 1-5 [Epub ahead of print].
- ~~17-18.~~ Tractenberg R, Gordon M. 2017. Supporting Evidence-Informed Teaching in Biomedical and Health Professions Education Through Knowledge Translation: An Interdisciplinary Literature Review, *Teaching and Learning in Medicine*, DOI:10.1080/10401334.2017.1287572
- ~~18-19.~~ Gundel H, Wefelmaier T. Effects of the clinical interview within a psychiatric teaching group on participating patients and medical students. An empirical study. *Nervenarzt*. 1998; 69(9): 806-810.
- ~~19-20.~~ Lynoe, N , Sandlund, M, Westberg, K & Duchek, M. Informed consent in clinical training - patient experiences and motives for participating. *Medical Education*. 1998; 32(5): 465-471.
- ~~20-21.~~ Walters K, Raven P, Rosenthal J, Russell J, Humphrey C, Buszewicz M. Teaching undergraduate psychiatry in primary care: the impact on student learning and attitudes. *Medical Education*. 2003; 326: 740-743
- ~~21-22.~~ Öster C, Backstrom S, Lantz I, Ramklint M. Psychiatric patients' perspectives of student involvement in their care. *BMC Medical Education*. 2015; 15: 69. DOI: 10.1186/s12909-015-0352-z
- ~~22-23.~~ Santulli RB. Reactions of psychiatric-inpatients to medical-student interviews. *Academic Medicine*. 1993; 68(10suppl):S40-S42.
- ~~23-24.~~ Black, AE & Church, M. Assessing medical student effectiveness from the psychiatric patient's perspective: The medical student interviewing performance questionnaire. *Medical Education*. 1998; 32(5): 472-478.
- ~~24-25.~~ Doshi M, Acharya S, Wall D. Mentally ill inpatients' experiences and opinions on seeing medical students: a questionnaire study. *Medical Teacher*. 2006; 28(6): 568-570.
- ~~25-26.~~ Tunde-Ayinmode M, Rajagopalan M, Little J. Attitudes of psychiatric inpatients to medical student interviews. *Australasian Psychiatry*. 2002; 10(3): 275-278.

- ~~26-27.~~ [27.](#) Greenberg D, Cohen R. Interviewing and case presentations in the presence of patients in undergraduate psychiatry: an experiment. *Medical Teacher*. 2003; 25(2): 167-170
- ~~27-28.~~ [28.](#) Dogra N, Anderson J, Edwards R, Cavendish S. Service user perspectives about their roles in undergraduate medical training about mental health. *Medical Teacher*. 2008; 30(6): 152-156
- ~~28-29.~~ [29.](#) Murata K, Sakuma M, Seki S, Morimoto T. Public attitudes towards practice by medical students: a nationwide survey in Japan. *Teaching and Learning in Medicine*. 2014; 26(4): 335-343
- ~~29-30.~~ [30.](#) Isaacson JH, Neides D, Mayer M, Nottingham K. Patient perceptions of having 1st- and 2nd-year medical students involved in their care. *Teaching and Learning in Medicine*. 2014; 26(2): 164-167.
- ~~30-31.~~ [31.](#) Passaperuma K, Higgins J, Power S, Taylor T. Do patients' comfort levels and attitudes regarding medical student involvement vary across specialties? *Medical Teacher*. 2008; 30(1): 48-54.
- ~~31-32.~~ [32.](#) Esguerra R, Toro J, Ospina JM, Porras A, Diaz C, Reyes S. The transition to a teaching hospital: patient satisfaction before and after the introduction of medical students. *Medical Teacher*. 2014; 36: 710-714.
- ~~32-33.~~ [33.](#) Scheffer C, Edelhauser F, Tauschel D, Reichman M, Tekian A. Can final year medical students significantly contribute to patient care? A pilot study about the perception of patients and clinical staff. *Medical Teacher*. 2010; 32(7): 552-557.
- ~~33-34.~~ [34.](#) Choudhury TR, Moosa AA, Cushing A, Bestwick J. Patients' attitudes towards the presence of medical students during consultations. *Medical Teacher*. 28(7): e198-e203.
- ~~34-35.~~ [35.](#) Marwan Y, Al-Saddique M, Hassan A, Karim J, Al-Saleh M. Are medical students accepted by patients in teaching hospitals? *Med Educ Online*. 2012; 17: 17172

Figures and Tables:

Figure 1. Articles found meeting search criteria






Table 1. Summary of included studies.

Author	Year	Location	Setting	Data type	Patient sample size	Quality Rating	Key findings
Öster <i>et al</i>	2015	Sweden	Adults + older adults	Quantitative & qualitative	655	<u>8</u>	76% comfortable seeing medical students, 73% under supervision dropping to 61% Strong agreement that patients should see medical students.
Dogra <i>et al</i>	2008	UK	Community adults + older adults (+ experience as children)	Quantitative	28	<u>5</u>	Students learn humanity, professional identity, appreciate stigma and treat the wh
Doshi <i>et al</i>	2006	UK	Inpatients (Adults)	Quantitative & qualitative	42	<u>3</u>	Whilst patients reported enjoyment and a sense of occupation in seeing patients, t seeing the student
Tunde- Ayinmode <i>et al</i>	2002	Australia	Inpatients (adults)	Quantitative	30	<u>6</u>	87% of patients enjoyed seeing medical students and 37% felt the experience ben were seeing a student as opposed to a qualified doctor

Black & Church	1998	USA	Inpatients (adult + older adult)	Quantitative	102	<u>4</u>	Medical student involvement can lead to a greater sense of involvement in decision making
Gundel & Wefelmaier	1998	Germany	Inpatients	Quantitative & qualitative	93	<u>3</u>	35% of patients felt they developed deeper insight as a result of seeing students
Lynoe <i>et al</i>	1998	Sweden	Community and inpatients	Quantitative	71	<u>2</u>	85% of patients are positive about seeing medical students
Santulli	1993	USA	Inpatients	Quantitative	67	<u>3</u>	Patients agree that seeing medical students is enjoyable and there is some disagreement about the amount of involvement