

## Central Lancashire Online Knowledge (CLOK)

Title	Effects of barefoot and shod running on lower extremity joint loading, a musculoskeletal simulation study
Type	Article
URL	<a href="https://clock.uclan.ac.uk/id/eprint/21974/">https://clock.uclan.ac.uk/id/eprint/21974/</a>
DOI	<a href="https://doi.org/10.1007/s11332-018-0446-1">https://doi.org/10.1007/s11332-018-0446-1</a>
Date	2018
Citation	Sinclair, Jonathan Kenneth (2018) Effects of barefoot and shod running on lower extremity joint loading, a musculoskeletal simulation study. Sport Sciences for Health. ISSN 1824-7490
Creators	Sinclair, Jonathan Kenneth

It is advisable to refer to the publisher's version if you intend to cite from the work.  
<https://doi.org/10.1007/s11332-018-0446-1>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

**Effects of barefoot and shod running on lower extremity joint loading, a  
musculoskeletal simulation study.**

*Jonathan Sinclair<sup>1</sup>*

*1. Centre for Applied Sport and Exercise Sciences, Faculty of Health and Wellbeing,  
University of Central Lancashire, Lancashire, UK.*

**Correspondence Address:**

Dr. Jonathan Sinclair

Centre for Applied Sport & Exercise Sciences

Faculty of Health and Wellbeing

University of Central Lancashire

Preston

Lancashire

PR1 2HE.

**e-mail:** jksinclair@uclan.ac.uk

**Keywords:** Biomechanics; barefoot; musculoskeletal; joint

**Abstract**

*PURPOSE:* The aim of the current investigation was to utilize a musculoskeletal simulation based approach, to examine the effects of barefoot and shod running on lower extremity joint loading during the stance phase.

*METHODS:* Twelve male runners, ran over an embedded force plate at 4.0 m/s, in both barefoot and shod conditions. Kinematics of the lower extremities were collected using an eight camera motion capture system. Lower extremity joint loading was also explored using a musculoskeletal simulation and mathematical modelling approach, and differences between footwear conditions were examined using paired samples t-tests.

*RESULTS:* Peak Achilles tendon force was significantly larger ( $P=0.039$ ) when running barefoot (6.85 BW) compared to shod (6.07 BW). In addition, both medial ( $P=0.013$ ) and lateral ( $P=0.007$ ) tibiofemoral instantaneous load rates were significantly larger in the barefoot (medial = 289.17 BW/s & lateral = 179.59 BW/s) in relation to the shod (medial = 167.57 BW/s & lateral = 116.40 BW/s) condition. Finally, the barefoot condition (9.70 BW) was associated with a significantly larger ( $P=0.037$ ) peak hip force compared to running shod (8.51 BW).

*CONCLUSIONS:* The current investigation indicates that running barefoot may place runners at increased risk from the biomechanical factors linked to the aetiology of chronic lower extremity pathologies. However, future analyses using habitual barefoot runners, are required before more definitive affirmations regarding injury predisposition can be made.

## **Introduction**

Running is an extremely popular exercise modality. It has been projected that as many as 2 million people in the UK utilize running as a mode of exercise (1). There is an overwhelming body of evidence, which has emphasized the physiological and psychological benefits of physical activity and exercise (2). However, despite the plethora of physical benefits

associated with regular running, it is also associated with a high incidence of chronic pathologies. Each year, up to 80 % of runners will suffer an overuse injury (3).

The knee joint is most susceptible to chronic pathology in runners (3). Specifically, patellofemoral pain syndrome is the most frequent overuse injury encountered in runners (4), characterized by pain at or anterior to the patella aggravated by physical activities that load the patellofemoral joint (5). Pain symptoms are related to excessive patellofemoral loading and typically persist for many years (6). A recent epidemiological analysis has shown that there may be a link between patellofemoral pain in younger adults and subsequent osteoarthritis at this joint (7). Furthermore, tibiofemoral pathologies are also common chronic running injuries; associated with up to 16.8% of all knee injuries (8). The medial aspect of the tibiofemoral joint is known to be significantly more prone to osteoarthritic degeneration than the lateral compartment (9). The causes of tibiofemoral chronic pathologies relate to the magnitude of the stress loading of the joint (10), which is considered to be the mechanical parameter most strongly associated with the onset and progression of knee osteoarthritis. The mechanism responsible for this is thought to be the increased joint contact forces experienced by the medial compartment of the tibiofemoral joint during locomotion (11). Finally, Achilles tendinopathies are also frequently occurring chronic musculoskeletal disorders in runners, accounting for approximately 8–15% of all injuries (12). The pathogenesis of Achilles tendinopathy is considered to be associated with habitual and excessive mechanical loading of the tendon itself, which creates microscopic tears in the tendons' collagen fibres (13).

An array of different treatment/ preventative modalities, have therefore been investigated in an attempt to attenuate the risk of running injuries. An extremely popular conservative

strategy is to select running trainers with appropriate biomechanical properties, as running shoes are proposed as a mechanism by which the rate of chronic injuries can be controlled (14). Recently however, it has been proposed that running using traditional running shoes may place runners at increased risk from the biomechanical factors linked to the aetiology of chronic running injuries (15). This led to a new proposal in footwear research, that running barefoot footwear may be associated with a reduced incidence of chronic running injuries (15). Based on this hypothesis, a number of runners are now choosing to run barefoot or in minimalist footwear (16, 17).

In recent years, barefoot running has received considerable research attention in biomechanical literature. Using a mathematical modelling approach driven by sagittal plane external joint torques and knee kinematics, both Bonacci et al., (18) and Sinclair, (19) showed that running barefoot significantly reduced patellofemoral joint loading during the stance phase of running. Furthermore, using external joint torques and ankle joint kinematics, Sinclair, (19) revealed that barefoot running was associated with significantly increased Achilles tendon forces in comparison to running shod. Finally, Sinclair et al., (16) and Sinclair et al., (17) found that barefoot running significantly increased the loading rate of the external vertical ground reaction force. Previous analyses concerning the biomechanical differences between barefoot and shod running, have utilized either the external ground reaction force or joint torque driven mathematical modelling approaches to explore the loads experienced by the musculoskeletal system. However, the external ground reaction force and joint torques represent global indices of joint loading, and therefore are not representative of localized joint loading (20). Herzog et al., (21) showed that muscles are the primary contributors to lower extremity joint loading. Yet the complex role of muscles in controlling

joint biomechanics during human movement has received insufficient attention within the literature, possibly due to difficulties in calculating muscle kinetics.

However, advances in musculoskeletal modelling have led to the development of bespoke software which allows skeletal muscle force distributions to be simulated during movement using motion capture based data (22). To date, such approaches have not yet been utilized to explore biomechanical differences between barefoot and shod running. Therefore, the aim of the current investigation was to examine the effects of barefoot and shod running on lower extremity joint loading using a musculoskeletal simulation based approach. A study of this nature may provide further insight into the biomechanical differences between barefoot and shod running; particularly with regards to runners' susceptibility to chronic pathologies.

## Methods

### *Participants*

Twelve healthy male runners, volunteered to take part in this study. All were identified as recreational runners who trained 3 times/week, completing a minimum of 35 km. The participants provided written informed consent in accordance with the principles outlined in the Declaration of Helsinki. The mean characteristics of the participants were; age  $24.33 \pm 4.09$  years, height  $1.77 \pm 0.09$  m and body mass  $75.44 \pm 6.58$  kg. The procedure utilized for this investigation was approved by the University of Central Lancashire, Science, Technology, Engineering and Mathematics, ethical committee.

112 *Procedure*

113 Participants ran at 4.0 m/s ( $\pm 5\%$ ), striking an embedded piezoelectric force platform (Kistler,  
114 Kistler Instruments Ltd., Alton, Hampshire) with their right foot. Running velocity was  
115 monitored using infrared timing gates (Newtest, Oy Koulukatu, Finland). The stance phase  
116 was delineated as the duration over which 20 N or greater of vertical force was applied to the  
117 force platform (23). Runners completed a minimum of five successful trials in both barefoot  
118 and shod conditions. The shod condition (New Balance 1260 v2) had an average mass of  
119 0.285 kg, heel thickness of 25 mm and a heel drop of 14 mm. The order that participants ran  
120 in each footwear condition was counterbalanced. Kinematics and ground reaction forces data  
121 were synchronously collected. Kinematic data was captured at 250 Hz via an eight camera  
122 motion analysis system (Qualisys Medical AB, Goteburg, Sweden). Dynamic calibration of  
123 the motion capture system was performed before each data collection session.

124

125 To define the anatomical frames of the thorax, pelvis, thighs, shanks and feet retroreflective  
126 markers were placed at the C7, T12 and xiphoid process landmarks and also positioned  
127 bilaterally onto the acromion process, iliac crest, anterior superior iliac spine (ASIS),  
128 posterior superior iliac spine (PSIS), medial and lateral malleoli, medial and lateral femoral  
129 epicondyles, greater trochanter, calcaneus, first metatarsal and fifth metatarsal. Carbon-fibre  
130 tracking clusters comprising of four non-linear retroreflective markers were positioned onto  
131 the thigh and shank segments. In addition to these the foot segments were tracked via the  
132 calcaneus, first metatarsal and fifth metatarsal, the pelvic segment was tracked using the PSIS  
133 and ASIS markers and the thorax segment was tracked using the T12, C7 and xiphoid  
134 markers. The shod condition was modified by cutting windows into the experimental  
135 footwear at the calcaneus, first metatarsal and fifth metatarsal locations in accordance with

Shultz & Jenkyn (24). This allowed the anatomical markers at these positions to be placed onto the skin in order to match the barefoot condition (25). Static calibration trials were obtained with the participant in the anatomical position in order for the positions of the anatomical markers to be referenced in relation to the tracking clusters/markers. A static trial was conducted with the participant in the anatomical position in order for the anatomical positions to be referenced in relation to the tracking markers, following which those not required for dynamic data were removed.

#### *Processing*

Dynamic trials were digitized using Qualisys Track Manager in order to identify anatomical and tracking markers then exported as C3D files to Visual 3D (C-Motion, Germantown, MD, USA). All data were normalized to 100 % of the stance phase. Ground reaction force and kinematic data were smoothed using cut-off frequencies of 50 and 12 Hz with a low-pass Butterworth 4th order zero lag filter (26). All net joint force parameters throughout were normalized by dividing by bodyweight (BW). Kinematic measures from the hip, knee, ankle which were extracted for statistical analysis were 1) angle at footstrike, 2) peak flexion/dorsiflexion during the stance phase and 3) angular range of motion (ROM) from footstrike to peak angle.

Data during the stance phase were exported from Visual 3D into OpenSim 3.3 software (Simtk.org). A validated musculoskeletal model with 12 segments, 19 degrees of freedom and 92 musculotendon actuators (27) was used to estimate extremity joint forces. The model was scaled for each participant to account for the anthropometrics of each athlete. As muscle



forces are the main determinant of joint compressive forces (21), muscle kinetics were quantified using a static optimization in accordance with Steele et al., (28). Compressive medial/ lateral tibiofemoral and hip joint forces were calculated via the joint reaction analyses function using the muscle forces generated from the static optimization process as inputs. Furthermore, medial and lateral tibiofemoral contact stresses (MPa) were quantified by dividing the tibiofemoral force by the medial and lateral contact areas estimated using the data of Kettelkamp and Jacobs, (29). From the above processing, peak medial tibiofemoral force, peak lateral tibiofemoral force, peak hip force, peak medial tibiofemoral stress and peak lateral tibiofemoral stress were extracted for statistical analyses. In addition medial/ lateral tibiofemoral and hip instantaneous load rates (BW/s) were also extracted by obtaining the peak increase in force between adjacent data points.

Patellofemoral loading during the stance phase of running was quantified using a model adapted from van Eijden et al., (30) in accordance with the protocol of Willson et al., (31). A key drawback of this model is that co-contraction of the knee flexor musculature is not accounted for. Taking this into account, summed hamstring and gastrocnemius forces derived from the static optimization procedure were multiplied by their estimated knee joint muscle moment arms as a function of knee flexion angle (32), and then added together to determine the knee flexor torque during the stance phase. In addition to this, the knee extensor torque was also calculated by dividing the summed quadriceps forces by this muscle groups' knee joint muscle moment arms as a function of knee flexion angle (30). The knee flexor and extensor torques were then summed and subsequently divided by the quadriceps muscle moment arm to obtain quadriceps force adjusted for co-contraction of the knee flexor musculature. Patellofemoral force was quantified by multiplying the derived quadriceps force by a constant which was obtained by using the data of Eijden et al., (30). Finally,

patellofemoral joint stress (MPa) was quantified by dividing the patellofemoral force by the patellofemoral contact area. Patellofemoral contact areas were obtained by fitting a polynomial curve to the sex specific data of Besier et al., (33), who estimated patellofemoral contact areas as a function of the knee flexion angle using MRI. From the above processing, peak patellofemoral force and peak patellofemoral stress were extracted for statistical analyses. In addition, patellofemoral instantaneous load rate (BW/s) was also extracted by obtaining the peak increase in force between adjacent data points.

Finally, Achilles tendon forces were estimated in accordance with the protocol of Almonroeder et al., (34), by summing the muscle forces of the medial gastrocnemius, lateral, gastrocnemius, and soleus muscles. From the above processing, peak Achilles tendon force and Achilles tendon instantaneous load rate (BW/s) were extracted for statistical analyses.

Running barefoot has been shown to alter the step length/ stance time during running (35), which may affect the number of footfalls required to complete a set distance. We therefore firstly calculated integral of the hip, tibiofemoral, patellofemoral and Achilles tendon forces during the stance phase, using a trapezoidal function. In addition to this, we also estimated the total force per mile (BW) by multiplying these parameters by the number of steps required to run a mile. The number of steps required to complete one mile was quantified using the step length (m), which was determined by taking the difference in the horizontal position of the foot centre of mass between the right and left legs at footstrike.

*Statistical analyses*

Means, standard deviations (SD) and 95 % confidence intervals (95% CI) were calculated for each outcome measure for both footwear conditions. The data was screened for normality using Shapiro-Wilk tests which confirmed that the normality assumption was met. Differences between footwear conditions were examined using paired samples t-tests, and effect sizes were calculated using partial eta<sup>2</sup> (pη<sup>2</sup>). Statistical actions were conducted using SPSS v23.0 (SPSS, USA).

## Results

### *Joint kinematics*

The hip was significantly (P=0.017, pη<sup>2</sup> = 0.42) more flexed at footstrike in the shod condition. In addition, peak hip flexion was significantly (P=0.018, pη<sup>2</sup> = 0.41) greater in the shod condition.

The ankle was significantly (P=0.001, pη<sup>2</sup> = 0.66) more dorsiflexed at footstrike in the shod condition. In addition, peak dorsiflexion was significantly (P=0.0004, pη<sup>2</sup> = 0.69) larger in the shod condition, and ankle ROM was significantly (P=0.032, pη<sup>2</sup> = 0.35) greater in the barefoot condition.

@@@ TABLE 1 NEAR HERE @@@

@@@ FIGURE 1 NEAR HERE @@@

### *Temporal parameters*

@@@ **TABLE 2 NEAR HERE** @@@

Step length was significantly ( $P=0.001$ ,  $\eta^2 = 0.65$ ) greater during shod running (Table 2). In addition, the number of steps per mile was significantly ( $P=0.001$ ,  $\eta^2 = 0.65$ ) lower in the shod condition (Table 2).

*Tibiofemoral kinetics*

Medial tibiofemoral force instantaneous load rate was significantly larger ( $P=0.013$ ,  $\eta^2 = 0.33$ ) in the barefoot condition (Table 3). In addition, lateral tibiofemoral force instantaneous load rate was significantly larger ( $P=0.007$ ,  $\eta^2 = 0.50$ ) in the barefoot condition (Table 3).

*Hip kinetics*

Peak hip force was significantly larger ( $P=0.037$ ,  $\eta^2 = 0.34$ ) in the barefoot condition (Table 3; Figure 3e). In addition, hip instantaneous load rate was significantly larger ( $P=0.002$ ,  $\eta^2 = 0.59$ ) in the barefoot condition (Table 3).

*Patellofemoral kinetics*

No differences ( $P>0.05$ ) in patellofemoral loading were observed (Table 3-4; Figure 2ab).

*Achilles tendon kinetics*

Peak Achilles tendon force was significantly larger ( $P=0.039$ ,  $\eta^2 = 0.33$ ) in the barefoot condition (Table 3; Figure 2c). In addition, Achilles tendon force per mile was significantly larger ( $P=0.028$ ,  $\eta^2 = 0.37$ ) in the barefoot condition (Table 4).

@@@ TABLE 3 NEAR HERE @@@

@@@ TABLE 4 NEAR HERE @@@

@@@ FIGURE 2 NEAR HERE @@@

@@@ FIGURE 3 NEAR HERE @@@

## Discussion

The aim of the current examination, was to examine the effects of barefoot and shod running on lower extremity joint loading using a musculoskeletal simulation approach. To the authors knowledge, this represents the first investigation to explore the biomechanical differences between barefoot and shod running using this methodology. This investigation provides further insight into the biomechanical differences between barefoot and shod running.

A key observation from the current analysis, is that patellofemoral loading parameters were not statistically different between barefoot and shod running. This finding opposes those of Bonacci et al., (18) and Sinclair, (19) who showed significant reductions in patellofemoral loading when running barefoot. It is proposed that this observation may relate to the specific

kinematic adjustments that runners made in the current investigation. Typically, when running barefoot the ankle is in a plantarflexed position at footstrike (15-17), and the knee ROM is significantly reduced (19), which effectively attenuates the role of the knee as a shock absorber (19). However, the current investigation showed no differences in knee kinematics when running barefoot, and whilst the ankle angle at footstrike was significantly altered in the barefoot condition, it was still in a dorsiflexed position. As such, it appears that the kinematic adaptations that runners typically make in the absence of footwear were less pronounced in this investigation, which may consequently explain the lack of differences in patellofemoral loading. Additionally, this may relate to the manner in which patellofemoral loading was calculated in the current study, as previous analyses have used mathematical models which do not account for co-contraction of the knee flexors (18, 19). Nonetheless, the current investigation indicates that running barefoot may not always attenuate the patellofemoral loading parameters linked to the aetiology of patellofemoral pain in runners.

The current investigation also revealed that the rate at which both the medial and lateral aspects of the tibiofemoral joint were loaded, was significantly larger in the barefoot condition. This finding is supported by those of Sinclair et al., (36) who found that the tibiofemoral rate of loading measured using an inverse dynamics based approach was significantly larger when running barefoot, in relation to traditional running trainers. This finding may be important, as increased compressive loading at the tibiofemoral joint, is a risk factor for the onset and progression of osteoarthritis (37). Therefore, the current analysis indicates that running barefoot may increase susceptibility to the risk factors associated with tibiofemoral osteoarthritis.

A further important observation from the current investigation was that Achilles tendon loading parameters were shown to be significantly larger in the barefoot condition. This observation concurs with those of Sinclair, (19), who similarly showed that Achilles tendon loading was greater when running barefoot. This observation may provide important clinical information in regards to the initiation and progression of Achilles tendinopathy (38). The aetiology of Achilles tendinopathy is mediated through repeated and excessive mechanical loading of the tendon during activities such as running. Repetitive tendon loads such as those initiate collagen and extracellular matrix synthesis and tissue degradation (39). Therefore, the current investigation shows that running barefoot may place runners at increased risk from the biomechanical parameters linked to Achilles tendinopathy.

In addition, this investigation also showed that peak compressive hip joint loading was significantly larger when running barefoot, in comparison to the shod condition. This study represents the first investigation to contrast hip joint loading during barefoot and shod running using musculoskeletal simulation, therefore comparisons against previous analyses are difficult. However, our findings are partially supported by those of Rooney & Derrick, (40) who showed that non-rearfoot strike runners experienced significantly greater compressive hip joint loading during running. However, in their prospective investigation of running injuries in barefoot and shod runners Altman & Davis, (41) found that hip injuries were statistically more frequent in shod runners. This appears to be contradictory as hip joint pathologies are strongly influenced by compressive hip joint loading (42). It is clear from this observation that further epidemiological research is required concerning the potential clinical influence of running barefoot.

A potential drawback to the current study is that it examined only habitual shod runners, who do not customarily run barefoot. Previous work examining the biomechanics of running barefoot has drawn conflicting observations, often on the basis of the barefoot running experience of their participants (15-17, 43). It can therefore, be speculated that the results from the current analysis may have been different had a sample of habitual barefoot runners been examined. Therefore, repeating the current investigation using habitual barefoot runners is advisable for future research, which may allow more definitive assertions with regards to injury predisposition to be made. That this study utilized a simulation based procedure to quantify muscles forces and joint loading may also serve as a limitation. Whilst this procedure is considered an improvement over previous approaches, in that joint reaction analyses are representative of localized joint loading and muscular co-contraction is accounted for. Musculoskeletal simulations depend on the underlying mathematical model and numerous mechanical assumptions are made in the construction of musculoskeletal simulation models (22). These predominately relate to the constrained rotational degrees of freedom at the knee and ankle joints and the lack of key muscles such as rectus abdominis, which may lead to incorrectly predicted muscle forces. However, as direct quantification of muscle forces are not possible at this time, the current procedure is the most practicable method in dynamic movements.

In conclusion, although the biomechanics of barefoot running have received extensive research attention; there has yet to be a quantitative comparison of lower extremity joint loading during barefoot and shod running using a musculoskeletal simulation based approach. The present investigation therefore adds to the current knowledge, by providing a comprehensive evaluation of lower extremity joint loading during barefoot and shod running conditions. On the basis that hip, tibiofemoral and Achilles tendon loading parameters were



significantly greater when running barefoot, the findings from the current investigation indicate that barefoot running may place runners at increased risk from the biomechanical risk factors linked to the aetiology of chronic lower extremity pathologies. However, future analyses using habitual barefoot runners, are required before more definitive affirmations regarding injury predisposition can be made.

## **Acknowledgements**

We thank Gareth Shadwell for his technical assistance.

## **Compliance with ethical standards**

### ***Conflict of interest***

We declare that we have no conflict of interest.

### ***Ethical approval***

The current research project was approved by an institutional ethical panel. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and the declaration of Helsinki.

### ***Informed consent***

All of the subjects provided written consent.

## **References**

1. <http://www.sportengland.org/media-centre/news/2014/september/05/sport-england-joins-the-great-north-run-team/>
2. Warburton DE, Nicol CW, Bredin SS (2006). Health benefits of physical activity: the evidence. CMAJ 174: 801-809. DOI: 10.1503/cmaj.051351

3. van Gent RN, Siem D, van Middelkoop M, van Os AG, Bierma-Zeinstra SMA, Koes BW (2007). Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med* 41: 469-480. DOI: 10.1136/bjism.2006.033548
4. Crossley KM (2014). Is patellofemoral osteoarthritis a common sequela of patellofemoral pain? *Br J Sports Med* 48: 409–410. DOI: <http://dx.doi.org/10.1136/bjsports-2014-093445>
5. Crossley KM, Stefanik JJ, Selfe J, Collins NJ, Davis IS, Powers CM, McConnell J, Vicenzino B, Bazett-Jones BM, Esculier J-F, Morrissey D, Callaghan MJ (2016). Patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester. Part 1: Terminology, definitions, clinical examination, natural history, patellofemoral osteoarthritis and patient-reported outcome measures. *Br J Sports Med* 50: 839–843. DOI: <http://dx.doi.org/10.1136/bjsports-2016-096384>
6. Collins NJ, Bierma-Zeinstra SM, Crossley KM, van Linschoten RL, Vicenzino B, van Middelkoop M. (2012). Prognostic factors for patellofemoral pain: a multicentre observational analysis. *Br J Sports Med* 47: 227-233. DOI: 10.1136/bjsports-2012-091696
7. Thomas MJ, Wood L, Selfe J, Peat G. (2010). Anterior knee pain in younger adults as a precursor to subsequent patellofemoral osteoarthritis: a systematic review. *BMC Musculoskelet Disord*, 11, 201-211. DOI: 10.1186/1471-2474-11-201.
8. Taunton JE, Ryan MB, Clement DB, McKenzie DC, Lloyd-Smith DR, Zumbo BD (2002). A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med* 36: 95-101. DOI: <http://dx.doi.org/10.1136/bjism.36.2.95>

9. Wise BL, Niu J, Yang M, Lane NE, Harvey W, Felson DT, Lewis CE (2012). Patterns of compartment involvement in tibiofemoral osteoarthritis in men and women and in whites and African Americans. *Arthritis Care Res* 64: 847-852. DOI: 10.1002/acr.21606.
10. Morgenroth DC, Medverd JR, Seyedali M, Czerniecki JM. (2014). The relationship between knee joint loading rate during walking and degenerative changes on magnetic resonance imaging. *Clin Biomech* 29: 664-670. DOI: 10.1016/j.clinbiomech.2014.04.008
11. Schipplein OD, Andriacchi TP (1991). Interaction between active and passive knee stabilizers during level walking. *J Orthop Res* 9: 113-119.
12. Van Ginckel A, Thijs Y, Hesar NGZ, Mahieu N, De Clercq D, Roosen P, Witvrouw E. (2008). Intrinsic gait-related risk factors for Achilles tendinopathy in novice runners: a prospective study. *Gait Posture* 29: 387-391. DOI: 10.1016/j.gaitpost.2008.10.058
13. Cook JL, Purdam CR. (2009). Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy. *Br J Sports Med* 43: 409-416. DOI: <http://dx.doi.org/10.1136/bjsm.2008.051193>
14. Shorten, MA. Running shoe design: protection and performance. pp 159-169 in *Marathon Medicine* (Ed. D. Tunstall Pedoe). 2000; London, Royal Society of Medicine.
15. Lieberman DE, Venkadesan M, Werbel WA, Daoud AI, D'Andrea S, Davis IS, Mang'eni RO, Pitsiladis Y. (2010). Foot strike patterns and collision forces in habitually barefoot versus shod runners. *Nature* 463: 531-535. DOI:10.1038/nature08723

16. Sinclair J, Greenhalgh A, Edmundson CJ, Brooks D, Hobbs SJ (2013). The influence of barefoot and barefoot-inspired footwear on the kinetics and kinematics of running in comparison to conventional running shoes. *Footwear Sci.* 5: 45–53. DOI: 10.1080/19424280.2012.693543
17. Sinclair J, Hobbs SJ, Currigan G, Taylor PJ (2013). A comparison of several barefoot inspired footwear models in relation to barefoot and conventional running footwear. *Comp Ex Phys* 9: 13-21. DOI: <http://dx.doi.org/10.3920/CEP13004>
18. Bonacci J, Vicenzino B, Spratford W, Collins P (2013). Take your shoes off to reduce patellofemoral joint stress during running. *Br J Sports Med* 48: 425-428. DOI: <http://dx.doi.org/10.1136/bjsports-2013-092160>
19. Sinclair J (2014). Effects of barefoot and barefoot inspired footwear on knee and ankle loading during running. *Clin Biomech* 29: 395-399. DOI: 10.1016/j.clinbiomech.2014.02.004
20. Herzog W, Longino D, Clark A (2003). The role of muscles in joint adaptation and degeneration. *Langenbecks Arch Surg* 388: 305-315. DOI:10.1007/s00423-003-0402-6
21. Herzog W, Clark A, Wu J. (2003). Resultant and local loading in models of joint disease. *Arthritis Care Res* 49: 239-247. DOI: 10.1002/art.11004
22. Delp SL, Anderson FC, Arnold AS, Loan P, Habib A, John CT, Thelen DG (2007). OpenSim: open-source software to create and analyze dynamic simulations of movement. *IEEE Trans Biomed Eng* 54:1940-1950. DOI: 10.1109/TBME.2007.901024
23. Sinclair J, Edmundson CJ, Brooks D, Hobbs SJ (2011). Evaluation of kinematic methods of identifying gait Events during running. *International J Sp Sci Eng* 5: 188-192.

24. Shultz R, Birmingham T, Jenkyn TR (2008). Validation of windows for examining kinematics of the foot with respect to the shoe using a multi-segment foot model. *J Foot Ankle Res* 1: 29-34. DOI: 10.1186/1757-1146-1-S1-O29
25. Sinclair J, Greenhalgh A, Taylor PJ, Edmundson CJ, Brooks D, Hobbs SJ (2013). Differences in tibiocalcaneal kinematics measured with skin-and shoe-mounted markers. *Hum Mov* 14: 64-69. DOI: <https://doi.org/10.2478/humo-2013-0005>
26. Sinclair J, Taylor PJ, Atkins S (2015). Influence of running shoes and cross-trainers on Achilles tendon forces during running compared with military boots. *J R Army Med Corps* 161: 140-143. DOI: 10.1136/jramc-2014-000308. Epub 2014 Nov 26.
27. Lerner ZF, DeMers MS, Delp SL, Browning RC (2015). How tibiofemoral alignment and contact locations affect predictions of medial and lateral tibiofemoral contact forces. *J Biomech* 48: 644-650. DOI: 10.1016/j.jbiomech.2014
28. Steele KM, DeMers MS, Schwartz MH, Delp SL (2012). Compressive tibiofemoral force during crouch gait. *Gait Posture* 35: 556-560. DOI: 10.1016/j.gaitpost.2011
29. Kettelkamp DB, Jacobs AW (1972). Tibiofemoral Contact Area-Determination and Implications. *JBJS* 54: 349-356.
30. van Eijden TM, Kouwenhoven E, Verburg J, Weijs WA (1986). A mathematical model of the patellofemoral joint. *J Biomech* 19: 219-229. [https://doi.org/10.1016/0021-9290\(86\)90154-5](https://doi.org/10.1016/0021-9290(86)90154-5)
31. Willson JD, Ratcliff OM, Meardon SA, Willy RW (2015). Influence of step length and landing pattern on patellofemoral joint kinetics during running. *Scand J Med Sci Sports* 25: 736-743. DOI: 10.1111/sms.12383
32. Spoor CW, van Leeuwen JL (1992). Knee muscle moment arms from MRI and from tendon travel. *J Biomech* 25: 201-206. DOI: [https://doi.org/10.1016/0021-9290\(92\)90276-7](https://doi.org/10.1016/0021-9290(92)90276-7)

33. Besier TF, Draper CE, Gold GE, Beaupre GS, Delp SL. (2005). Patellofemoral joint contact area increases with knee flexion and weight-bearing. *J Orthop Res* 23: 345–350. DOI: 10.1016/j.orthres.2004.08.003
34. Almonroeder T, Willson JD, Kernozek TW. (2013). The effect of foot strike pattern on Achilles tendon load during running. *Ann Biomed Eng* 41: 1758-1766. DOI: 10.1007/s10439-013-0819-1.
35. Thompson M, Gutmann A, Seegmiller J, McGowan C. (2014). The effect of stride length on the dynamics of barefoot and shod running. *J Biomech* 47: 2745-2750. DOI: 10.1016/j.jbiomech.2014.04.043
36. Sinclair J. (2016). The Effects of Barefoot and Barefoot Inspired Footwear Running on Tibiofemoral Kinetics. *Hum Mov* 17: 176-180. DOI: <https://doi.org/10.1515/humo-2016-0022>
37. Dabiri Y, Li LP (2013). Altered knee joint mechanics in simple compression associated with early cartilage degeneration. *Comput Math Methods Med* 11: 1-12. DOI: <http://dx.doi.org/10.1155/2013/862903>
38. Selvanetti A, Cipolla M, Puddu G. (1997). Overuse tendon injuries: basic science and classification. *Op Tech Sports Med* 5: 110–117. DOI: [https://doi.org/10.1016/S1060-1872\(97\)80031-7](https://doi.org/10.1016/S1060-1872(97)80031-7)
39. Magnusson SP, Langberg, H, Kjaer M. (2010). The pathogenesis of tendinopathy: balancing the response to loading. *Nat Rev Rheumatol* 6: 262–268. DOI: 10.1038/nrrheum.2010.43
40. Rooney BD, Derrick TR (2013). Joint contact loading in forefoot and rearfoot strike patterns during running. *J Biomech* 46: 2201-2206. DOI: 10.1016/j.jbiomech.2013.06.022

41. Altman AR, Davis IS (2016). Prospective comparison of running injuries between shod and barefoot runners. *Br J Sports Med* 50: 476-480. DOI: <http://dx.doi.org/10.1136/bjsports-2014-094482>
42. Johnson VL, Hunter DJ (2014). The epidemiology of osteoarthritis. *Best Pract Res Clin Rheumatol* 28: 5–15. DOI: 10.1016/j.berh.2014.01.004
43. Squadrone R, Gallozzi, C (2009). Biomechanical and physiological comparison of barefoot and two shod conditions in experienced barefoot runners. *J Sports Med Phys Fitness* 49: 6-13.

## Figures

Figure 1: Joint kinematics as a function of footwear a. = hip, b. = knee and c. = ankle (black = barefoot and grey = shod).

Figure 2: Patellofemoral and Achilles tendon kinetics as a function of footwear a. = patellofemoral force, b. = patellofemoral stress and c. Achilles tendon force (black = barefoot and grey = shod).

Figure 3: Tibiofemoral and hip kinetics as a function of footwear a. = medial tibiofemoral force, b. = medial tibiofemoral stress, c. = lateral tibiofemoral force, d. = lateral tibiofemoral stress and e. = hip force (black = barefoot and grey = shod).

Table 1: Hip, knee and ankle kinematics (Mean, SD and 95% CI's) as a function of footwear.

	Barefoot				Shod				
	Mean	SD	95% CI Lower	95% CI Upper	Mean	SD	95% CI Lower	95% CI Upper	
Hip angle at footstrike (°)	34.29	12.38	26.42	42.15	42.27	7.77	37.34	47.21	*
Peak hip flexion (°)	34.84	12.03	27.20	42.49	42.76	7.24	38.16	47.35	*
Hip ROM (°)	0.56	1.26	0.24	1.36	0.48	1.13	0.22	1.20	
Knee angle at footstrike (°)	25.05	5.45	21.59	28.52	24.67	9.12	18.88	30.47	
Peak knee flexion (°)	45.90	4.48	43.05	48.75	47.90	6.41	43.82	51.97	
Knee ROM (°)	20.85	7.38	16.16	25.54	23.22	8.54	17.80	28.65	
Ankle angle at footstrike (°)	4.56	6.93	0.15	8.96	12.74	2.62	11.07	14.40	*
Peak dorsiflexion (°)	18.35	4.17	15.70	21.00	22.82	3.85	20.37	25.26	*
Ankle ROM (°)	13.80	7.70	8.90	18.69	10.08	4.08	7.49	12.67	*

Key: \* = significant difference



Table 2: Peak hip, knee and ankle loading parameters (Mean, SD and 95% CI's) as a function of footwear.

	Barefoot				Shod				
	Mean	SD	95% CI Lower	95% CI Upper	Mean	SD	95% CI Lower	95% CI Upper	
Peak patellofemoral force (BW)	4.32	0.93	3.73	4.91	4.51	1.07	3.83	5.19	
Peak patellofemoral stress (MPa)	5.05	0.93	4.46	5.64	5.14	0.78	4.65	5.63	
Patellofemoral instantaneous load rate (BW/s)	159.55	56.26	123.81	195.29	149.80	56.60	113.84	185.76	
Peak Achilles tendon force (BW)	6.85	1.95	5.61	8.09	6.07	1.22	5.29	6.84	*
Achilles tendon instantaneous load rate (BW/s)	174.17	85.71	119.71	228.63	142.16	32.01	121.83	162.50	
Peak medial tibiofemoral force (BW)	6.53	1.64	5.49	7.57	6.23	1.25	5.44	7.03	
Peak medial tibiofemoral stress (MPa)	12.51	2.75	10.76	14.26	11.77	2.04	10.47	13.07	
Medial tibiofemoral instantaneous load rate (BW/s)	289.17	142.69	198.50	379.83	167.57	77.16	118.54	216.59	*
Peak lateral tibiofemoral force (BW)	4.17	1.09	3.48	4.87	3.94	0.75	3.47	4.42	
Peak lateral tibiofemoral stress (MPa)	13.15	3.56	10.89	15.41	12.32	2.17	10.94	13.70	
Lateral tibiofemoral instantaneous load rate (BW/s)	179.59	60.90	140.89	218.28	116.40	30.13	97.25	135.54	*
Peak hip force (BW)	9.70	1.32	8.86	10.53	8.51	0.94	7.92	9.11	*
Hip instantaneous load rate (BW/s)	377.38	140.49	288.12	466.64	167.25	78.35	117.47	217.03	*

Key: \* = significant difference

Table 3: Step characteristics (Mean, SD and 95% CI's) as a function of footwear.

	Barefoot				Shod				
	Mean	SD	95% CI Lower	95% CI Upper	Mean	SD	95% CI Lower	95% CI Upper	
Step length (m)	1.27	0.05	1.24	1.31	1.38	0.06	1.34	1.42	*
Steps per mile	632.42	26.41	615.64	649.19	583.20	24.32	567.75	598.65	*

Key: \* = significant difference

Table 4: Joint loading per mile (Mean, SD and 95% CI's) of hip, knee and ankle loading.

	Barefoot				Shod				
	Mean	SD	95% CI Lower	95% CI Upper	Mean	SD	95% CI Lower	95% CI Upper	
Patellofemoral force per mile (BW)	321.49	52.39	288.20	354.77	322.16	84.85	268.25	376.07	
Achilles tendon force per mile (BW)	402.47	93.60	343.00	461.94	356.31	79.19	306.00	406.62	*
Medial tibiofemoral force per mile (BW)	464.62	110.98	394.11	535.14	441.14	81.48	389.38	492.91	
Lateral tibiofemoral force per mile (BW)	283.32	56.09	247.68	318.96	290.12	58.62	252.87	327.37	
Hip force per mile (BW)	854.05	187.03	735.22	972.88	781.19	109.56	711.58	850.80	

Key: \* = significant difference