

Central Lancashire Online Knowledge (CLoK)

Title	Increased exposure to pesticides and colon cancer: Early evidence in Brazil
Type	Article
URL	https://clock.uclan.ac.uk/23305/
DOI	https://doi.org/10.1016/j.chemosphere.2018.06.118
Date	2018
Citation	Martin, Francis L, Martinez, Edson Z., Stopper, Helga, Garcia, Sergio Britto, Uyemura, Sergio Akira and Kannen, Vinicius (2018) Increased exposure to pesticides and colon cancer: Early evidence in Brazil. Chemosphere, 209. pp. 623-631. ISSN 0045-6535
Creators	Martin, Francis L, Martinez, Edson Z., Stopper, Helga, Garcia, Sergio Britto, Uyemura, Sergio Akira and Kannen, Vinicius

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.chemosphere.2018.06.118>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Accepted Manuscript

Increased exposure to pesticides and colon cancer: Early evidence in Brazil

Francis L. Martin, Edson Z. Martinez, Helga Stopper, Sergio Britto Garcia, Sergio Akira Uyemura, Vinicius Kannen



PII: S0045-6535(18)31193-7

DOI: [10.1016/j.chemosphere.2018.06.118](https://doi.org/10.1016/j.chemosphere.2018.06.118)

Reference: CHEM 21650

To appear in: *ECSN*


Received Date: 15 April 2018

Revised Date: 14 June 2018

Accepted Date: 17 June 2018

Please cite this article as: Martin, F.L., Martinez, E.Z., Stopper, H., Garcia, S.B., Uyemura, S.A., Kannen, V., Increased exposure to pesticides and colon cancer: Early evidence in Brazil, *Chemosphere* (2018), doi: 10.1016/j.chemosphere.2018.06.118.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Increased exposure to pesticides and colon cancer: early evidence in Brazil

Francis L. Martin¹, Edson Z. Martinez², Helga Stopper³, Sergio Britto Garcia⁴,
Sergio Akira Uyemura⁵, Vinicius Kannen⁵

¹*School of Pharmacy and Biomedical Sciences, University of Central Lancashire,
Preston PR1 2HE, United Kingdom;*

²*Department of Social Medicine, University of Sao Paulo, Ribeirao Preto, Brazil;*

³*Department of Toxicology, University of Wuerzburg, Wuerzburg, Germany;*

⁴*Department of Pathology, University of Sao Paulo, Ribeirao Preto, Brazil;*

⁵*Department of Toxicology, Bromatology, and Clinical Analysis, University of Sao
Paulo, Ribeirao Preto, Brazil.*

Running title: Pesticide and colon cancer

Corresponding Author:

Tel +55 (16) 3315.9133; Fax: +55 (16) 33154725;

E-mail: vinicius.kannen@fcfrp.usp.br.

ABSTRACT

Environmental factors may increase colon cancer (CC) risk. It has been suggested that pesticides could play a significant role in the etiology of this malignancy. As agriculture is one of the mainstays of the Brazilian economy, this country has been the largest pesticides consumer worldwide. The CC burden is also increasing in Brazil. Herein, we examined data from the Brazilian Federal Government to determine whether CC mortality and pesticide consumption may be associated. Database of the Ministry of Health provided CC mortality data in Brazil, while pesticides use was accessed at the website of Brazilian Institute of Environment and Renewable Natural Resources. The CC mortality in the Brazilian states was calculated as standard mortality rates (SMR). All Bayesian analysis was performed using a Markov chain Monte Carlo method in WinBUGS software. We observed that colon cancer mortality has exhibited a steady increase for more than a decade, which correlated with the amount of sold pesticides in the country. Both observations are concentrated in the Southern and the Southeast regions of Brazil. Although ecological studies like ours have methodological limitations, the current dataset suggests the possibility that pesticide exposure may be a risk factor for colon cancer. It warrants further investigation.

Keywords: Xenobiotics; carcinogens; environment; tumors; intestines

1. Introduction

Colon cancer (CC) has afflicted humans for millennia. Chronic exposure to certain environmental factors appears to be the key to better understanding the etiology of this malignancy (David and Zimmerman, 2010). Over-nutrition and sedentary lifestyle may also be responsible for up to 75% of cancers today (Nebert and Dalton, 2006; David and Zimmerman, 2010). Notably, CC is one of the leading cause of cancer-related deaths (Torre et al., 2015). By 2030, developing countries are expected to exhibit a sharp increase in CC cases (Arnold et al., 2016). Also, it should be pointed out that recent epidemiological trends highlight that the CC burden is shifting towards a younger population (de Magalhaes, 2013; Siegel et al., 2014).

Cancer risk, including CC, appears to be profoundly influenced by environmental factors (Wu et al., 2016). Thus, CC etiology is complex, meaning that a multiple of environmental factors may cause this disease. One of many hazardous and carcinogenic factors promoting malignancies, pesticides have been suggested by the International Agency for Research on Cancer (IARC) to increase cancer risk in humans (Guyton et al., 2015; Guyton et al., 2016). Extensive epidemiological studies support the idea that pesticides are a risk factor for solid tumors (Parron et al., 2014). There has also been some evidence that pesticides promote CC in both humans and rodents (Soliman et al., 1997; Tellez-Banuelos et al., 2016; Hong et al., 2017). It seems feasible that pesticides contaminate human food sources (Nagao and Sugimura, 1993; Lodovici et al., 1997; Sakita et al., 2017), a fact that may be related to increased cancer risk (Arrebola et al., 2015). Another point underlying to study the relationship between pesticides and cancer

must be considered: disease incidence is increasing dramatically (Lodovici et al., 1997; Soliman et al., 1997; Agudo et al., 2009; Andreotti et al., 2010; Boccolini Pde et al., 2013; Parron et al., 2014; Arrebola et al., 2015; Carnero et al., 2015; Coggon et al., 2015; Guyton et al., 2015; Guyton et al., 2016; Tellez-Banuelos et al., 2016; Hong et al., 2017).

Furthermore, the lack of epidemiological and experimental data that accurately correlate CC incidence with detection of individual cancer initiators impairs our current ability to determine the impact of environmental factors on the CC development in humans (Tomasetti and Vogelstein, 2015). For instance, various environmental pollutants were reported to induce DNA damage and adducts, but the precise evolution of such genomic damages into mutations that promote CC remains unknown (Tomasetti and Vogelstein, 2015; Poirier, 2016). Then, it should be considered that instead of those DNA-damaging effects induced by initiators, endogenous and exogenous cancer promoters are classically determined to lead mutated cells towards clonal expansion, enabling them to collect further genomic changes by either high proliferative activity or new carcinogenic hits (Irigaray and Belpomme, 2010). Rather than binding to DNA, a cancer promoter usually activates transcriptional and epigenetic mechanisms that induce proliferation but inhibit apoptosis (Irigaray and Belpomme, 2010; Engstrom et al., 2015). Such mechanistic activity has for long been known to induce proliferation intrinsic errors leading to mutations and the development of CC (Ames and Gold, 1990; Bartkova et al., 2005; Gorgoulis et al., 2005). Interestingly, pesticides may act either as carcinogens or cancer promoters (Agudo et al., 2009; Andreotti et al., 2010; Arrebola et al., 2015; Carnero et al., 2015; Coggon et al.,

2015). Of note is the fact that Brazil has been the most significant consumer of pesticides worldwide for years (Boccolini Pde et al., 2013). Recently, we have hypothesized that pesticides could impact on the CC risk (Uyemura et al., 2017).

Herein, we propose an association between increased CC mortality and pesticide consumption in Brazil. This could suggest that pesticides alter the risk of CC in a human population.

2. Materials and methods

2.1. Collection of public data

CC mortality (<http://www2.datasus.gov.br/DATASUS/index.php?area=0205>) was collected from the database of the Ministry of Health. The quantity of pesticides (tonnes) sold within the country was downloaded from the website of the Brazilian Institute of Environment and Renewable Natural Resources (http://dados.contraosagrototoxicos.org/pt_PT/dataset/comercializacao-ibama-2014; <http://www.ibge.gov.br/>). Complementary data on pesticides and farmed land area for each Brazilian state (Km²) were collected from the Brazilian Institute of Geography and Statistics (<http://www.ibge.gov.br/>).

2.2. Statistical analyses

The CC mortality in the Brazilian states was calculated as standard mortality rates (SMR). Further information on SMR can be found in a previous report authored by Ulm (Ulm, 1990). We determined SMR to be the ratio of observed mortality to expected mortality adjusted for age and gender group. An SMR value >1 indicates excessive mortality. Expected numbers of death were calculated using age and

gender-specific mortality rates for the Brazilian general population (assumed to be the standard population). Within this approach, $w(s,t,f)$ was the death rate for the Brazilian population at the year t ($t = 1$ if 2000, $t = 2$ if 2001, and so on) considering gender s ($s = 1$ if women and $s = 2$ if man) and age group f ($f = 1$ if <50 y old, $f = 2$ if 50 to 59 y, $f = 3$ if 60 to 69 y, $f = 4$ if 70 to 79 y and $f = 5$ if ≥ 80 y). The expected number of death for each Brazilian state p ($p = 1, \dots, 27$) in the year (t) according to the gender (s) is given by:

$$E(p, s, t) = \sum_{f=1}^5 w(s, t, f) \times m(p, s, t, f),$$

where $m(p,s,t,f)$ is the number of inhabitants of the state (p) with gender (s) at the year (t) and group age (f). The SMR is thus given by:

$$SMR(p, s, t) = \frac{Y(p, s, t)}{E(p, s, t)},$$

where $Y(p,s,t)$ is corresponding observed mortality. Spatio-temporally smoothed SMR values were obtained from a Bayesian model based on the Poisson distribution. This statistical model is given by:

$$Y(p,s,t) \mid \mu(p,s,t), E(p,s,t) \sim \text{Poisson} [E(p,s,t) \times \mu(p,s,t)],$$

where $\mu(p,s,t) = \exp[\alpha_0 + \alpha_{sp} + \omega(p,s,t)]$ is the parameter that describes the SMR, α_0 is an intercept, α_{sp} are bivariate random effects that capture spatial dependence in the data ($s = 1,2$, $p = 1,\dots,27$) and $\omega(p,s,t)$ models the longitudinal trend of annual mortality rate for the federation unit p and gender s , considering a multivariate Gaussian process with a mean vector 5×1 with all components equal to zero and a given covariance function. In the Bayesian analysis, it was assumed that α_{sp} follows a conditionally bivariate autoregressive (CAR) structure and α_0 follows a non-informative normal distribution with mean zero and a large variance.

Then, we verified the association between the HDI of each Brazilian state and the corresponding SMR, for which a Bayesian model was fitted to the data.

Thus, $\mu(p,s,t)$ was replaced by:

$$\mu(p,s,t) = \exp[\alpha_0 + \alpha_{sp} + \beta_{st}x(p)],$$

where $x(p)$ is the amount of sold pesticide (measured in tonnes) recorded in each Brazilian state (p) at the year 2000, divided by its respective total cultivated area in hectares (including permanent and temporary crops) and multiplied by 1,000, and β_{st} is the corresponding effect. Credible intervals for β_{st} that do not include zero indicates a significant correlation between the amount of sold pesticide and the mortality rate. Credible intervals are the Bayesian analogues to the traditional 95% confidence intervals. In all Bayesian analysis, the posterior distributions were simulated using a Markov chain Monte Carlo (MCMC) method in WinBUGS software.

3. Results

CC has not only been suggested to be one of the commonest malignancy types in Western countries (Torre et al., 2015) but also that its incidence and mortality may increase throughout the next decade in developing countries (Arnold et al., 2016). This notion inspired us to apply the Bayesian model to calculate SMR values for CC mortality in the Brazilian population. Heatmaps revealed that mortality by CC mainly occurred in the Southern Brazilian states (Figure 1 and 2).

Environmental factors are well-known able of increasing cancer risk (Wu et al., 2016). In addition, the IARC has suggested that pesticide can promote human risk of developing different types of cancer (Guyton et al., 2015; Guyton et al., 2016). In developing countries, some research groups report that pesticides may increase cancer incidence (Soliman et al., 1997; Fonnum and Mariussen, 2009; Yi, 2013; Arrebola et al., 2015). Herein, we analyzed the quantity of pesticide sold in Brazil. We should note that these records were reported by the Federal Government in tonnes for each state, and are the most accurate dataset available to the public. To provide a better perspective of pesticide distribution in each Federal unit, we rated pesticide values by the total cultivated area that was officially reported for each of those Brazilian states. We observed a dramatic increase in pesticide usage from 2000 to 2012 within the country, mainly in the Southern, Southeast and Central-West regions of Brazil (Figure 3).

Next, we examined whether both events were correlated in the Brazilian population. We found an increase in SMR values correlating with the amount of pesticide sold by 2000 in Brazil (Figure 4 and 5). Smoothed curves fitted by loess were added on each graph. Moreover, it shows 95% credible intervals for the

effects (β_{st}) of the amount of sold pesticide on the SMR values for each year (t) and gender (s), obtained from the Bayesian spatiotemporal regression models. From 2000 to 2007, the credible intervals do not contain zero, thus suggesting a significant effect of the amount of sold pesticide recorded in each Brazilian state on their corresponding SMR for CC (Figure 6).

4. Discussion

We should initially consider that some environmental chemicals damage the DNA, whereas other promote the expansion of mutated cells during the development of CC (Lawrence et al., 2013; Tomasetti and Vogelstein, 2015; Poirier, 2016), meaning that we can no longer hypothesize that only DNA damaging compounds impact on the cancer risk in humans. Indeed, it seems that the mutation rate intrinsic to mitosis might be sufficient in invoking oncogenic changes in the rapidly dividing colonic epithelial cell population (Bartkova et al., 2005; Gorgoulis et al., 2005). This was initially observed in classical experiments of rodents exposed to cancer promoters (Ames and Gold, 1990). Persistent epithelial self-renewal requires precise molecular regulation of proliferation in component cells that is, consequently, prey to corruption by environmental and mutational factors. It is, therefore, no surprise that the majority of cancers originated in epithelial tissue are due to somatic mutations that deregulate the molecular constraints on cell pluripotency and proliferation (Lawrence et al., 2013; Tomasetti and Vogelstein, 2015; Vogelstein and Kinzler, 2015).

Manmade compounds (xenobiotics) can access the human body *via* multiple routes, each modifying the risk of cancer (Sakita et al., 2017; Uyemura et al.,

2017). This requires that the increasingly large number of chemicals whose cancer-causing effects remain unknown should be taken into account while discussing the impact of environmental factors on CC risk (Guha et al., 2016). Indeed, most pesticides might have endocrine-disrupting and metabolic effects, as well as bio-accumulating in the human body (Irigaray and Belpomme, 2010; Soto and Sonnenschein, 2010; Walker and Gore, 2011; Ellsworth et al., 2015; Espin Perez et al., 2015; Maqbool et al., 2016). It means that whether pesticides interact at low levels and may increase the risk of cancer, their activity does not need to be simultaneous or continuous. Combining several exposures to different pesticides at multiple time-points could, thus, induce far greater cancer-related effects than single compounds in humans (Goodson et al., 2015).

The massive number of modern xenobiotics has made almost impossible to determine what their precise impact on human cancer risk is (Bouvard et al., 2015; Goodson et al., 2015). For instance, a research group analyzed 6000 human-made compounds and found that 16.3% of those chemicals were pesticides, from which less than 1% had been investigated in the context of cancer (Guha et al., 2016). Alavanja and colleagues studied the effects of 50 commonly used pesticides in 56,813 pesticide applicators and found a potential relationship between exposure to chlorpyrifos and aldicarb with the incidence of colorectal cancer (CRC) (Lee et al., 2007). A meta-analysis study suggested that aldicarb could increase the CC risk, imazethapyr may promote the cancer risk in the proximal colon region, and CRC risk was probably enhanced by exposure to pendimethalin, chlorpyrifos, chlordane, and toxaphene (Alexander et al., 2012). Considering the complex CC etiology together with the little number of epidemiological and experimental data

correlating CC development with the environmental pollution by pesticides becomes clear that further efforts are required to clarify this matter.

In Tunisia, foodstuffs containing pesticides were suggested to increase the risk of breast cancer in women (Arrebola et al., 2015). Different Brazilian research groups have reported high-pesticide levels in human milk in the country (Matuo et al., 1992; Beretta and Dick, 1994; Dorea et al., 1997). Pesticide levels in bovine milk have been reported to exceed safety standards in the Midwest region of Brazil (Avancini et al., 2013). Then, public data from the Brazilian National Health Surveillance Agency (Anvisa; <http://portal.anvisa.gov.br/en/programa-de-analise-de-registro-de-agrotoxicos-para>) show that 20% of food samples analyzed between 2013 and 2015 were unsafe for human use. In 2013, Meyer and colleagues revealed that pesticides could be related to increased non-Hodgkin's lymphoma mortality found in Brazil (Boccolini Pde et al., 2013). Koifman and colleagues hypothesized that cancer-related mortality in Brazilian farm workers could be related to their exposure to pesticides from 1979 to 1998 (Meyer et al., 2003). Meyer and colleagues suggested that the amount of pesticides selected in 1985 could be related to breast, prostate, and ovarian cancer mortality ten years later (Koifman S., 2002). In Martinique, pesticides increased the risk of prostate cancer (Landau-Ossondo et al., 2009). In South-Korea, pesticides increased CC risk (Fonnum and Mariussen, 2009; Yi, 2013). Indeed, high-pesticide serum levels were detected in CC patients in Egypt (Soliman et al., 1997). In rats, pesticides increased the risk of CC (Hong et al., 2017). Then, another research group suggested that pesticides might increase CC risk by promoting inflammation in the colon (Tellez-Banuelos et al., 2016).

Although there has been some evidence that pesticides could be a risk factor for CC (Soliman et al., 1997; Lee et al., 2007; Fonnum and Mariussen, 2009; Alexander et al., 2012; Yi, 2013; Hong et al., 2017), other limitations in studying the effects of these chemicals in cancer have to be considered. Carcinogenic effects of human-made pollutants usually require protracted exposure to be detectable. For instance, asbestos-related effects increasing lung mesothelioma have been reported to take over 63 years to develop (Hodgson et al., 2005). However, we should also consider that asbestos has an established effect in promoting this type of malignancy in the lungs (Hodgson et al., 2005), while the complex activity of multiple pesticides in different types of cancer makes almost impossible to suggest which pesticide directly increases the CC risk in the human population. Moreover, other confounding factors could also have similar effects promoting CC risk. For instance, dietary factors seemed to be one of the main risk factors promoting this disease in humans (Sakita et al., 2017). Notably, a 10% increase in the intake of ultra-processed food furthered by 10% the cancer risk in humans (Fiolet et al., 2018). In Brazil, the risk of developing CRC was related to the high consumption of meat (Angelo et al., 2016). Here, we should also consider that human food sources have been suggested to be contaminated by pesticides in Brazil (Matuo et al., 1992; Beretta and Dick, 1994; Dorea et al., 1997; Avancini et al., 2013; Uyemura et al., 2017). This scenario is quite severe since some types of food with known carcinogenic potential could have a more hazardous effect if they contained pesticides in their composition. Indeed, we do not claim to have found that pesticides cause CC mortality in Brazil, but current evidence should not be ignored and requires further study.

Nevertheless, from our perspective, the CC mortality rates in the Brazilian state Amapá, located at the North region of the country, seems to be an outlier. CC-related death numbers varied from the lowest to the highest rates in the country by 2005. This increase reversed over the subsequent period. Lima and Queiroz analyzed the Brazilian death registry system and found that completeness of death registration in this state was one of the poorest in the country (Lima and Queiroz, 2014). Hence, we advise future studies to have careful consideration on this matter while investigating mortality rates during this period in that Brazilian state.

5. Conclusion

We believe that protracted exposure to pesticide may be a potential risk factor for CC. This fact requires urgent attention from the Federal Government monitoring the exposure of Brazilians to such chemicals. Whereas authorities must oversee the activity of multinational agrochemical and agricultural biotechnology corporations, as well as pesticide usage in agriculture, farmers should be informed by awareness programs to improve their product quality without harming the human population with high pesticide residue levels in the environment and food.

Conflict of interest statement

The authors disclose that no competing interests exist.

Acknowledgements

The authors disclose receipt of the following financial support for the development of this investigation: Sao Paulo Research Foundation (FAPESP; 2014/06428-5; 2015/01723-1). The funder had no role in the study design, data collection, analysis, decision to publish, or preparation of the manuscript.

Authors' role

Study concept and design: VK; Acquisition of data: VK and EZM; Statistical analysis: EZM; Analysis and interpretation of data: All; Drafting the first version of the manuscript: FLM and VK; Critical revision of the manuscript: All; Obtained funding: VK; Study supervision: FLM, EZM and VK.

References

- Agudo, A., Goni, F., Etxeandia, A., Vives, A., Millan, E., Lopez, R., Amiano, P., Ardanaz, E., Barricarte, A., Chirlaque, M.D., Dorronsoro, M., Jakszyn, P., Larranaga, N., Martinez, C., Navarro, C., Rodriguez, L., Sanchez, M.J., Tormo, M.J., Gonzalez, C.A., 2009. Polychlorinated biphenyls in Spanish adults: determinants of serum concentrations. *Environmental research* 109, 620-628.
- Alexander, D.D., Weed, D.L., Mink, P.J., Mitchell, M.E., 2012. A weight-of-evidence review of colorectal cancer in pesticide applicators: the agricultural health study and other epidemiologic studies. *International archives of occupational and environmental health* 85, 715-745.
- Ames, B.N., Gold, L.S., 1990. Too many rodent carcinogens: mitogenesis increases mutagenesis. *Science* 249, 970-971.

- 330 Andreotti, G., Hou, L., Beane Freeman, L.E., Mahajan, R., Koutros, S., Coble, J.,
331 Lubin, J., Blair, A., Hoppin, J.A., Alavanja, M., 2010. Body mass index, agricultural
332 pesticide use, and cancer incidence in the Agricultural Health Study cohort. *Cancer*
333 *causes & control* : CCC 21, 1759-1775.
- 334 Angelo, S.N., Lourenco, G.J., Magro, D.O., Nascimento, H., Oliveira, R.A., Leal,
335 R.F., Ayrizono Mde, L., Fagundes, J.J., Coy, C.S., Lima, C.S., 2016. Dietary risk
336 factors for colorectal cancer in Brazil: a case control study. *Nutrition journal* 15, 20.
- 337 Arnold, M., Sierra, M.S., Laversanne, M., Soerjomataram, I., Jemal, A., Bray, F.,
338 2016. Global patterns and trends in colorectal cancer incidence and mortality. *Gut*.
- 339 Arrebola, J.P., Belhassen, H., Artacho-Cordon, F., Ghali, R., Ghorbel, H., Boussen,
340 H., Perez-Carrascosa, F.M., Exposito, J., Hedhili, A., Olea, N., 2015. Risk of
341 female breast cancer and serum concentrations of organochlorine pesticides and
342 polychlorinated biphenyls: a case-control study in Tunisia. *The Science of the total*
343 *environment* 520, 106-113.
- 344 Avancini, R.M., Silva, I.S., Rosa, A.C., Sarcinelli Pde, N., de Mesquita, S.A., 2013.
345 Organochlorine compounds in bovine milk from the state of Mato Grosso do Sul-
346 Brazil. *Chemosphere* 90, 2408-2413.
- 347 Bartkova, J., Horejsi, Z., Koed, K., Kramer, A., Tort, F., Zieger, K., Guldberg, P.,
348 Sehested, M., Nesland, J.M., Lukas, C., Orntoft, T., Lukas, J., Bartek, J., 2005.
349 DNA damage response as a candidate anti-cancer barrier in early human
350 tumorigenesis. *Nature* 434, 864-870.
- 351 Beretta, M., Dick, T., 1994. Organochlorine compounds in human milk, Porto
352 Alegre, Brazil. *Bulletin of environmental contamination and toxicology* 53, 357-360.

- 353 Boccolini Pde, M., Boccolini, C.S., Chrisman Jde, R., Markowitz, S.B., Koifman, S.,
354 Koifman, R.J., Meyer, A., 2013. Pesticide use and non-Hodgkin's lymphoma
355 mortality in Brazil. *International journal of hygiene and environmental health* 216,
356 461-466.
- 357 Bouvard, V., Loomis, D., Guyton, K.Z., Grosse, Y., Ghissassi, F.E., Benbrahim-
358 Tallaa, L., Guha, N., Mattock, H., Straif, K., International Agency for Research on
359 Cancer Monograph Working, G., 2015. Carcinogenicity of consumption of red and
360 processed meat. *Lancet Oncol.*
- 361 Carnero, A., Blanco-Aparicio, C., Kondoh, H., Lleonart, M.E., Martinez-Leal, J.F.,
362 Mondello, C., Ivana Scovassi, A., Bisson, W.H., Amedei, A., Roy, R., Woodrick, J.,
363 Colacci, A., Vaccari, M., Raju, J., Al-Mulla, F., Al-Temaimi, R., Salem, H.K.,
364 Memeo, L., Forte, S., Singh, N., Hamid, R.A., Ryan, E.P., Brown, D.G., Wise, J.P.,
365 Sr., Wise, S.S., Yasaei, H., 2015. Disruptive chemicals, senescence and
366 immortality. *Carcinogenesis* 36 Suppl 1, S19-37.
- 367 Coggon, D., Ntani, G., Harris, E.C., Jayakody, N., Palmer, K.T., 2015. Soft tissue
368 sarcoma, non-Hodgkin's lymphoma and chronic lymphocytic leukaemia in workers
369 exposed to phenoxy herbicides: extended follow-up of a UK cohort. *Occupational
370 and environmental medicine* 72, 435-441.
- 371 David, A.R., Zimmerman, M.R., 2010. Cancer: an old disease, a new disease or
372 something in between? *Nat Rev Cancer* 10, 728-733.
- 373 de Magalhaes, J.P., 2013. How ageing processes influence cancer. *Nat Rev
374 Cancer* 13, 357-365.

- 375 Dorea, J.G., Granja, A.C., Romero, M.L., 1997. Pregnancy-related changes in fat
376 mass and total DDT in breast milk and maternal adipose tissue. *Annals of nutrition
377 & metabolism* 41, 250-254.
- 378 Ellsworth, R.E., Mamula, K.A., Costantino, N.S., Deyarmin, B., Kostyniak, P.J., Chi,
379 L.H., Shriver, C.D., Ellsworth, D.L., 2015. Abundance and distribution of
380 polychlorinated biphenyls (PCBs) in breast tissue. *Environmental research* 138,
381 291-297.
- 382 Engstrom, W., Darbre, P., Eriksson, S., Gulliver, L., Hultman, T., Karamouzis,
383 M.V., Klaunig, J.E., Mehta, R., Moorwood, K., Sanderson, T., Sone, H., Vadgama,
384 P., Wagemaker, G., Ward, A., Singh, N., Al-Mulla, F., Al-Temaimi, R., Amedei, A.,
385 Colacci, A.M., Vaccari, M., Mondello, C., Scovassi, A.I., Raju, J., Hamid, R.A.,
386 Memeo, L., Forte, S., Roy, R., Woodrick, J., Salem, H.K., Ryan, E.P., Brown, D.G.,
387 Bisson, W.H., 2015. The potential for chemical mixtures from the environment to
388 enable the cancer hallmark of sustained proliferative signalling. *Carcinogenesis* 36
389 Suppl 1, S38-60.
- 390 Espin Perez, A., de Kok, T.M., Jennen, D.G., Hendrickx, D.M., De Coster, S.,
391 Schoeters, G., Baeyens, W., van Larebeke, N., Kleinjans, J.C., 2015. Distinct
392 genotype-dependent differences in transcriptome responses in humans exposed to
393 environmental carcinogens. *Carcinogenesis*.
- 394 Fiolet, T., Srour, B., Sellem, L., Kesse-Guyot, E., Alles, B., Mejean, C.,
395 Deschasaux, M., Fassier, P., Latino-Martel, P., Beslay, M., Hercberg, S., Lavalette,
396 C., Monteiro, C.A., Julia, C., Touvier, M., 2018. Consumption of ultra-processed
397 foods and cancer risk: results from NutriNet-Sante prospective cohort. *Bmj* 360,
398 k322.

399 Fonnum, F., Mariussen, E., 2009. Mechanisms involved in the neurotoxic effects of
400 environmental toxicants such as polychlorinated biphenyls and brominated flame
401 retardants. *J Neurochem* 111, 1327-1347.

402 Goodson, W.H., 3rd, Lowe, L., Carpenter, D.O., Gilbertson, M., Manaf Ali, A.,
403 Lopez de Cerain Salsamendi, A., Lasfar, A., Carnero, A., Azqueta, A., Amedei, A.,
404 Charles, A.K., Collins, A.R., Ward, A., Salzberg, A.C., Colacci, A., Olsen, A.K.,
405 Berg, A., Barclay, B.J., Zhou, B.P., Blanco-Aparicio, C., Baglione, C.J., Dong, C.,
406 Mondello, C., Hsu, C.W., Naus, C.C., Yedjou, C., Curran, C.S., Laird, D.W., Koch,
407 D.C., Carlin, D.J., Felsher, D.W., Roy, D., Brown, D.G., Ratovitski, E., Ryan, E.P.,
408 Corsini, E., Rojas, E., Moon, E.Y., Laconi, E., Marongiu, F., Al-Mulla, F.,
409 Chiaradonna, F., Darroudi, F., Martin, F.L., Van Schooten, F.J., Goldberg, G.S.,
410 Wagemaker, G., Nangami, G.N., Calaf, G.M., Williams, G., Wolf, G.T., Koppen, G.,
411 Brunborg, G., Lyster, H.K., Krishnan, H., Ab Hamid, H., Yasaei, H., Sone, H.,
412 Kondoh, H., Salem, H.K., Hsu, H.Y., Park, H.H., Koturbash, I., Miousse, I.R.,
413 Scovassi, A.I., Klaunig, J.E., Vondracek, J., Raju, J., Roman, J., Wise, J.P., Sr.,
414 Whitfield, J.R., Woodrick, J., Christopher, J.A., Ochieng, J., Martinez-Leal, J.F.,
415 Weisz, J., Kravchenko, J., Sun, J., Prudhomme, K.R., Narayanan, K.B., Cohen-
416 Solal, K.A., Moorwood, K., Gonzalez, L., Soucek, L., Jian, L., D'Abronzio, L.S., Lin,
417 L.T., Li, L., Gulliver, L., McCawley, L.J., Memeo, L., Vermeulen, L., Leyns, L.,
418 Zhang, L., Valverde, M., Khatami, M., Romano, M.F., Chapellier, M., Williams,
419 M.A., Wade, M., Manjili, M.H., Leonart, M.E., Xia, M., Gonzalez, M.J., Karamouzis,
420 M.V., Kirsch-Volders, M., Vaccari, M., Kuemmerle, N.B., Singh, N., Cruickshanks,
421 N., Kleinstreuer, N., van Larebeke, N., Ahmed, N., Ogunkua, O., Krishnakumar,
422 P.K., Vadgama, P., Marignani, P.A., Ghosh, P.M., Ostrosky-Wegman, P.,

423 Thompson, P.A., Dent, P., Heneberg, P., Darbre, P., Sing Leung, P., Nangia-
424 Makker, P., Cheng, Q.S., Robey, R.B., Al-Temaimi, R., Roy, R., Andrade-Vieira,
425 R., Sinha, R.K., Mehta, R., Vento, R., Di Fiore, R., Ponce-Cusi, R., Dornetshuber-
426 Fleiss, R., Nahta, R., Castellino, R.C., Palorini, R., Abd Hamid, R., Langie, S.A.,
427 Eltom, S.E., Brooks, S.A., Ryeom, S., Wise, S.S., Bay, S.N., Harris, S.A.,
428 Papagerakis, S., Romano, S., Pavanello, S., Eriksson, S., Forte, S., Casey, S.C.,
429 Luanpitpong, S., Lee, T.J., Otsuki, T., Chen, T., Massfelder, T., Sanderson, T.,
430 Guarnieri, T., Hultman, T., Dormoy, V., Odero-Marah, V., Sabbisetti, V., Maguer-
431 Satta, V., Rathmell, W.K., Engstrom, W., Decker, W.K., Bisson, W.H., Rojanasakul,
432 Y., Luqmani, Y., Chen, Z., Hu, Z., 2015. Assessing the carcinogenic potential of
433 low-dose exposures to chemical mixtures in the environment: the challenge ahead.
434 *Carcinogenesis* 36 Suppl 1, S254-296.

435 Gorgoulis, V.G., Vassiliou, L.V., Karakaidos, P., Zacharatos, P., Kotsinas, A.,
436 Liloglou, T., Venere, M., Ditullio, R.A., Jr., Kastrinakis, N.G., Levy, B., Kletsas, D.,
437 Yoneta, A., Herlyn, M., Kittas, C., Halazonetis, T.D., 2005. Activation of the DNA
438 damage checkpoint and genomic instability in human precancerous lesions. *Nature*
439 434, 907-913.

440 Guha, N., Guyton, K.Z., Loomis, D., Barupal, D.K., 2016. Prioritizing Chemicals for
441 Risk Assessment Using Chemoinformatics: Examples from the IARC Monographs
442 on Pesticides. *Environ Health Perspect.*

443 Guyton, K.Z., Loomis, D., Grosse, Y., El Ghissassi, F., Benbrahim-Tallaa, L.,
444 Guha, N., Scoccianti, C., Mattock, H., Straif, K., International Agency for Research
445 on Cancer Monograph Working Group, I.L.F., 2015. Carcinogenicity of

tetrachlorvinphos, parathion, malathion, diazinon, and glyphosate. *Lancet Oncol* 16, 490-491.

Guyton, K.Z., Loomis, D., Grosse, Y., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L., Guha, N., Mattock, H., Straif, K., International Agency for Research on Cancer Monograph Working, G., 2016. Carcinogenicity of pentachlorophenol and some related compounds. *Lancet Oncol* 17, 1637-1638.

Hodgson, J.T., McElvenny, D.M., Darnton, A.J., Price, M.J., Peto, J., 2005. The expected burden of mesothelioma mortality in Great Britain from 2002 to 2050. *Br J Cancer* 92, 587-593.

Hong, M.Y., Hoh, E., Kang, B., DeHamer, R., Kim, J.Y., Lumibao, J., 2017. Fish Oil Contaminated with Persistent Organic Pollutants Induces Colonic Aberrant Crypt Foci Formation and Reduces Antioxidant Enzyme Gene Expression in Rats. *J Nutr.*

Irigaray, P., Belpomme, D., 2010. Basic properties and molecular mechanisms of exogenous chemical carcinogens. *Carcinogenesis* 31, 135-148.

Koifman S., K.R.J., Meyer A., 2002. Human reproductive system disturbances and pesticide exposure in Brazil. *Cad Saud Pub* 18, 435-445.

Landau-Ossondo, M., Rabia, N., Jos-Pelage, J., Marquet, L.M., Isidore, Y., Saint-Aime, C., Martin, M., Irigaray, P., Belpomme, D., pesticides, A.i.r.g.o., 2009. Why pesticides could be a common cause of prostate and breast cancers in the French Caribbean Island, Martinique. An overview on key mechanisms of pesticide-induced cancer. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 63, 383-395.

Lawrence, M.S., Stojanov, P., Polak, P., Kryukov, G.V., Cibulskis, K., Sivachenko, A., Carter, S.L., Stewart, C., Mermel, C.H., Roberts, S.A., Kiezun, A., Hammerman,

- 470 P.S., McKenna, A., Drier, Y., Zou, L., Ramos, A.H., Pugh, T.J., Stransky, N.,
471 Helman, E., Kim, J., Sougnez, C., Ambrogio, L., Nickerson, E., Shefler, E., Cortes,
472 M.L., Auclair, D., Saksena, G., Voet, D., Noble, M., DiCara, D., Lin, P.,
473 Lichtenstein, L., Heiman, D.I., Fennell, T., Imielinski, M., Hernandez, B., Hodis, E.,
474 Baca, S., Dulak, A.M., Lohr, J., Landau, D.A., Wu, C.J., Melendez-Zajgla, J.,
475 Hidalgo-Miranda, A., Koren, A., McCarroll, S.A., Mora, J., Lee, R.S., Crompton, B.,
476 Onofrio, R., Parkin, M., Winckler, W., Ardlie, K., Gabriel, S.B., Roberts, C.W.,
477 Biegel, J.A., Stegmaier, K., Bass, A.J., Garraway, L.A., Meyerson, M., Golub, T.R.,
478 Gordenin, D.A., Sunyaev, S., Lander, E.S., Getz, G., 2013. Mutational
479 heterogeneity in cancer and the search for new cancer-associated genes. *Nature*
480 499, 214-218.
- 481 Lee, W.J., Sandler, D.P., Blair, A., Samanic, C., Cross, A.J., Alavanja, M.C., 2007.
482 Pesticide use and colorectal cancer risk in the Agricultural Health Study. *Int J*
483 *Cancer* 121, 339-346.
- 484 Lima, E.E., Queiroz, B.L., 2014. Evolution of the deaths registry system in Brazil:
485 associations with changes in the mortality profile, under-registration of death
486 counts, and ill-defined causes of death. *Cadernos de saude publica* 30, 1721-1730.
- 487 Lodovici, M., Casalini, C., Briani, C., Dolara, P., 1997. Oxidative liver DNA damage
488 in rats treated with pesticide mixtures. *Toxicology* 117, 55-60.
- 489 Maqbool, F., Mostafalou, S., Bahadar, H., Abdollahi, M., 2016. Review of
490 endocrine disorders associated with environmental toxicants and possible involved
491 mechanisms. *Life Sci* 145, 265-273.
- 492 Matuo, Y.K., Lopes, J.N., Casanova, I.C., Matuo, T., Lopes, J.L., 1992.
493 Organochlorine pesticide residues in human milk in the Ribeirao Preto region, state

494 of Sao Paulo, Brazil. Archives of environmental contamination and toxicology 22,
495 167-175.

496 Meyer, A., Chrisman, J., Moreira, J.C., Koifman, S., 2003. Cancer mortality among
497 agricultural workers from Serrana Region, state of Rio de Janeiro, Brazil.
498 Environmental research 93, 264-271.

499 Nagao, M., Sugimura, T., 1993. Carcinogenic factors in food with relevance to
500 colon cancer development. Mutat Res 290, 43-51.

501 Nebert, D.W., Dalton, T.P., 2006. The role of cytochrome P450 enzymes in
502 endogenous signalling pathways and environmental carcinogenesis. Nat Rev
503 Cancer 6, 947-960.

504 Parron, T., Requena, M., Hernandez, A.F., Alarcon, R., 2014. Environmental
505 exposure to pesticides and cancer risk in multiple human organ systems. Toxicol
506 Lett 230, 157-165.

507 Poirier, M.C., 2016. Linking DNA adduct formation and human cancer risk in
508 chemical carcinogenesis. Environ Mol Mutagen 57, 499-507.

509 Sakita, J.Y., Gasparotto, B., Garcia, S.B., Uyemura, S.A., Kannen, V., 2017. A
510 critical discussion on diet, genomic mutations and repair mechanisms in colon
511 carcinogenesis. Toxicol Lett 265, 106-116.

512 Siegel, R., Desantis, C., Jemal, A., 2014. Colorectal cancer statistics, 2014. CA
513 Cancer J Clin 64, 104-117.

514 Soliman, A.S., Smith, M.A., Cooper, S.P., Ismail, K., Khaled, H., Ismail, S.,
515 McPherson, R.S., Seifeldin, I.A., Bondy, M.L., 1997. Serum organochlorine
516 pesticide levels in patients with colorectal cancer in Egypt. Archives of
517 environmental health 52, 409-415.

- 518 Soto, A.M., Sonnenschein, C., 2010. Environmental causes of cancer: endocrine
519 disruptors as carcinogens. *Nature reviews. Endocrinology* 6, 363-370.
- 520 Tellez-Banuelos, M.C., Haramati, J., Franco-Topete, K., Peregrina-Sandoval, J.,
521 Franco-Topete, R., Zaitseva, G.P., 2016. Chronic exposure to endosulfan induces
522 inflammation in murine colon via beta-catenin expression and IL-6 production. *J*
523 *Immunotoxicol* 13, 842-849.
- 524 Tomasetti, C., Vogelstein, B., 2015. Cancer etiology. Variation in cancer risk
525 among tissues can be explained by the number of stem cell divisions. *Science* 347,
526 78-81.
- 527 Torre, L.A., Bray, F., Siegel, R.L., Ferlay, J., Lortet-Tieulent, J., Jemal, A., 2015.
528 Global cancer statistics, 2012. *CA Cancer J Clin* 65, 87-108.
- 529 Ulm, K., 1990. A simple method to calculate the confidence interval of a
530 standardized mortality ratio (SMR). *Am J Epidemiol* 131, 373-375.
- 531 Uyemura, S.A., Stopper, H., Martin, F.L., Kannen, V., 2017. A Perspective
532 Discussion on Rising Pesticide Levels and Colon Cancer Burden in Brazil.
533 *Frontiers in public health* 5, 273.
- 534 Vogelstein, B., Kinzler, K.W., 2015. The Path to Cancer --Three Strikes and You're
535 Out. *The New England journal of medicine* 373, 1895-1898.
- 536 Walker, D.M., Gore, A.C., 2011. Transgenerational neuroendocrine disruption of
537 reproduction. *Nature reviews. Endocrinology* 7, 197-207.
- 538 Wu, S., Powers, S., Zhu, W., Hannun, Y.A., 2016. Substantial contribution of
539 extrinsic risk factors to cancer development. *Nature* 529, 43-47.

540 Yi, S.W., 2013. Cancer incidence in Korean Vietnam veterans during 1992-2003:
541 the Korean veterans health study. Journal of preventive medicine and public health
542 = Yebang Uihakhoe chi 46, 309-318.

543

544

Figure Legends

Fig.1. Smoothed standard mortality rates for CC in the Brazilian male population in each state of the country, as calculated by the Bayesian model.













Fig.2. Smoothed standard mortality rates for CC in the Brazilian female population in each state of the country, as calculated by the Bayesian model.













Fig.3. Heatmaps show the amount of sold pesticide recorded in each Brazilian state by total cultivated area (1000 x tonne /hectare) from 2000 to 2012.















Fig.4. Scatterplots of the relationship in the Brazilian male population between SMR values and the amount of sold pesticide recorded in each state of the country by total cultivated area (1000 x tonne/ hectare) from 2000 to 2012.














Fig.5. Scatterplots of the relationship in the Brazilian female population between SMR values and the amount of sold pesticide recorded in each state of the country by total cultivated area (1000 x tonne/ hectare) from 2000 to 2012.














Fig.6. Credible intervals for the effects β_{st} of the amount of sold pesticide on the SMR values for each year (t) and gender (s), obtained from the Bayesian spatiotemporal regression models.













(a) Male**(b) Female**

Highlights

- Human exposure to xenobiotics occurs worldwide, largely;
- Pesticides may promote cancer risk;
- Brazil is the world major pesticides consumer;
- Colon cancer (CC) mortality is steadily increasing in Brazil;
- We found CC mortality and pesticide levels may be correlated events in Brazil.