

## Central Lancashire Online Knowledge (CLoK)

Title	Significance and potential of marine microbial natural bioactive compounds against biofilms/biofouling: necessity for green chemistry
Type	Article
URL	<a href="https://clock.uclan.ac.uk/23379/">https://clock.uclan.ac.uk/23379/</a>
DOI	<a href="https://doi.org/10.7717/peerj.5049">https://doi.org/10.7717/peerj.5049</a>
Date	2018
Citation	Adnan, Mohd, Alshammari, Eyad, Patel, Mitesh, Amir Ashraf, Syed, Khan, Saif and Hadi, Ss (2018) Significance and potential of marine microbial natural bioactive compounds against biofilms/biofouling: necessity for green chemistry. PeerJ, 6. e5049.
Creators	Adnan, Mohd, Alshammari, Eyad, Patel, Mitesh, Amir Ashraf, Syed, Khan, Saif and Hadi, Ss

It is advisable to refer to the publisher's version if you intend to cite from the work.  
<https://doi.org/10.7717/peerj.5049>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

# Significance and potential of marine microbial natural bioactive compounds against biofilms/biofouling: necessity for green chemistry

Mohd Adnan<sup>1</sup>, Eyad Alshammari<sup>2</sup>, Mitesh Patel<sup>3</sup>, Syed Amir Ashraf<sup>2</sup>, Saif Khan<sup>1</sup> and Sibte Hadi<sup>4</sup>

<sup>1</sup> Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, University of Hail, Hail, Saudi Arabia

<sup>2</sup> Department of Clinical Nutrition, College of Applied Medical Sciences, University of Hail, Hail, Saudi Arabia

<sup>3</sup> Department of Biosciences, Bapalal Vaidhya Botanical Research Centre, Veer Narmad South Gujarat University, Surat, India

<sup>4</sup> School of Forensic and Applied Sciences, University of Central Lancashire, Preston, UK

## ABSTRACT

Natural products from the unique environments of sea water and oceans represent a largely unfamiliar source for isolation of new microbes, which are potent producers of secondary bioactive metabolites. These unique life-forms from the marine ecosphere have served as an important source of drugs since ancient times and still offer a valuable resource for novel findings by providing remedial treatments. Therefore, it can be expected that many naturally bioactive marine microbial compounds with novel structures and bioactivities against those from terrestrial environments may be found among marine metabolites. Biofilms in aquatic environment possess serious problems to naval forces and oceanic industries around the globe. Current anti-biofilm or anti-biofouling technology is based on the use of toxic substances that can be harmful to their surrounding natural locales. Comprehensive research has been done to examine the bioactive potential of marine microbes. Results are remarkably varied and dynamic, but there is an urgent need for bioactive compounds with environmentally friendly or “green” chemical activities. Marine microbes have the potential as upcoming and promising source of non-toxic compounds with sustainable anti-biofouling/anti-biofilm properties as they can produce substances that can inhibit not only the chemical components required for biofilm production but also the attachment, microorganism growth, and/or cell–cell communication.

**Subjects** Ecology, Marine Biology, Microbiology, Ecotoxicology, Green Chemistry

**Keywords** Biofouling, Biofilm, Bioactive compounds, Green chemistry, Marine microbes

## INTRODUCTION

Little is known about the ecology of marine microorganisms, which is probably the reason that attention was not given by chemists and ecologists to these organisms for many years. As they flourish in diverse types of ecological pressures, climates, food supplies, and darkness, these marine organisms develop certain adaptation mechanisms. This can

Submitted 2 March 2018

Accepted 1 June 2018

Published 27 June 2018

Corresponding authors

Mohd Adnan,

drmohdadnan@gmail.com

Sibte Hadi, shadi@uclan.ac.uk

Academic editor

Najat Saliba

Additional Information and  
Declarations can be found on  
page 11

DOI 10.7717/peerj.5049

© Copyright

2018 Adnan et al.

Distributed under

Creative Commons CC-BY 4.0


**OPEN ACCESS**

include the struggle for space, avoidance of predators, ability to effectively reproduce, and many other unknown defense mechanisms. Evolution/production of unique and natural bioactive metabolites is one such outcome resulting from these adaptations, which can be beneficial for human beings in many ways. They can possibly be responsible for the interaction with chemical components of biofilms (Joint, Muhling & Querellou, 2010).

Naturally bioactive chemical compounds produced by bacteria and diatoms can cause disruption in biofilm formation (Ganapiriya, Maharajan & Kumarasamy, 2012); therefore, they can be useful in development of environmentally friendly compounds for protection against marine bio-fouling (Holmström & Kjelleberg, 1999). It is just a matter of finding the correct naturally bioactive compound for a specific application. Coating and application of active ingredients from marine organisms (invertebrates, microorganisms, algae) that prevent the growth and settlement of fouling organisms has been proposed since the 1980s (Maki et al., 1988) but none of the potent, non-toxic, naturally bioactive compounds from diverse marine life-forms are found to be active against biofouling. When compared with the natural anti-biofoulants from macro-organisms (sponges, mollusks, tunicates, bryozoans, polychetes, and many other marine invertebrates), very limited information is available from marine microorganisms (Fusetani, 2004).

Our aim through this literature review was to search for potential biologically active compounds, whose chemical information can be useful in facilitating the development of new anti-biofouling or anti-biofilm agents from natural marine sources (Li et al., 2018). These agents should possess some extent of bioactivity, either against biofilms or any other biofilm producing microorganism and must rely on two actions: (1) synergistic action and (2) continuous production of enough potent molecules/defensive compounds by microbes so that they may also be induced upon attacks on the host. Thus far, after comparing the available information on bioactive metabolites from macro-organisms, only a small number of microorganisms have been investigated for bioactive metabolites (Debbab et al., 2010). This review covers future research with new biologically active natural marine microbial compounds and highlights their mechanism of action with focus on chemical potential. Meticulous research in the field of bioactive compounds from marine microbes may open the gates for many prospect implications in oceanic industries as well as in the field of biomedical sciences.

The ocean is called the “mother of origin of life,” and an enormous proportion of all life on Earth exists within the oceans (Mora et al., 2011). In densely populated marine environments, space is often a limiting factor. When free space is not available, one organism grows on top of another one. This process is defined as epibiosis (Wahl et al., 2012). Adaptation to epibiosis arises via three methods: (1) tolerance; (2) avoidance; and (3) defense (Wahl et al., 2012). Overgrowth is controlled by employing either one or a combination of ecological, physical, and chemical defense mechanisms (Wahl, Jensen & Fenical, 1994), which includes the major factor, production of bioactive metabolites (Wahl & Banaigs, 1991). Studies unraveling the epibiotic chemical defense of marine microorganisms can provide insight into the development of novel ecofriendly antifouling compounds and strategies (Jensen & Fenical, 1996).



**Figure 1** Comparative statistical data about number of publications in PubMed since 1989 till 2017. Comparative general data about the number of publications in PubMed when searching for several keywords/phrases: (1) microbial natural bioactive compounds (blue column); (2) marine microbial natural bioactive compounds (yellow column); (3) microbial anti-biofilm compounds (red column); and (4) marine microbial anti-biofilm compounds (green column). Moving average trend lines show the importance and urgent need for research concerning on marine microbial natural anti-biofouling/anti-biofilm compounds. [Full-size !\[\]\(5f471a71b78d7676bc356df190b88ab4\_img.jpg\) DOI: 10.7717/peerj.5049/fig-1](https://doi.org/10.7717/peerj.5049/fig-1)

Biofouling occurrence in the marine environment is a sequential process, commencing with micro-fouling and concluding with macro-fouling. The adverse effects of biofouling and economic penalties to the marine industries are well known (Bekiari *et al.*, 2015; Dang & Lovell, 2016). In addition, fouling is also known to cause destruction of metallic surfaces by accelerating the corrosion rate (Edyvean, Thomas & Brook, 1988). Numerous efforts are being made to control biofouling with applications of physical, chemical, biological compounds, but it is achieved to the greatest extent only with the use of anti-fouling paint coatings (Cao *et al.*, 2011; Iorhemen, Hamza & Tay, 2016; Lade, Paul & Kweon, 2014).

Use of toxic anti-foulants and its related environmental concerns have caused an increase in development of non-toxic alternatives (Callow & Callow, 2011; Nurioglu, Esteves & de With, 2015). In this perspective, studies concerning marine micro-organisms' epibiotic chemical defense in the control of biofouling show great promise. Efforts in these areas have proven that the wide range of marine organisms such as algae, sponges, gorgonians, bryozoans, and ascidians are a potential source of anti-fouling metabolites (Ganapiriya, Maharajan & Kumarasamy, 2012; Zhang *et al.*, 2014). However, not much detail is known to date regarding anti-biofouling/anti-biofilm natural compounds from marine microorganisms. Figure 1 demonstrates statistics about the number of publications in PubMed from 1989 until 2017 regarding this subject. These data explain the urgent need to explore the enigma of favorable marine microbes' activities against biofouling. Since the beginning of this type of research, studies concerning marine microbial metabolites, have

**Table 1** List of areas commonly known for applications/uses of marine microbial natural bioactive compounds.

<b>Medical significance of marine microbial metabolites</b>	Antibacterial
	Antiviral
	Antiparasitic
	Antifungal
	Anticoagulant
	Antimutagenic
	Antihyperglycaemic
	Antitumoral
	Antiinflammatory
	Anticarcinogenic
	Antioxidant
	Taste astringent
	Photoprotection
	Immunomodulator
	Emulsifier
	Gelling agent
Cosmetics industry	
Textile industry	
Stimulant	

shown significance that is only related to therapeutic and pharmaceutical agents for various commercial and medical applications (Table 1); however, none have been identified to date, which can be viewed as the existence of potent anti-biofouling/anti-biofilm marine microbial derivatives without any synthetic modifications.

Enormous numbers of studies have been done to discover natural product-derived anti-foulants as a substitute for toxic anti-fouling paints (Ciriminna, Bright & Pagliaro, 2015; Omae, 2003; Qian, Xu & Fusetani, 2009; Wang et al., 2017). Epibiotic chemical defense of marine organisms have been evaluated, but some restrictions hindered those investigations. For instance, these studies were carried out in the laboratory against one or two groups of epibionts (Davis, 1991; Laudien & Wahl, 2004), and such unrealistic assay conditions do not exist in nature (Henrikson & Pawlik, 1998). This restricts the scope of studies concerning epibiotic chemical defense in marine organisms, which present a wide range of epibionts in nature. In this respect, evaluation of metabolites obtained from marine microorganisms against broad spectrum of epibionts provides better insight into epibiotic chemical defense (Penesyan, Kjelleberg & Egan, 2010).

In the real world, identification of bioactive metabolites origins is complicated as some marine invertebrates are home for a wide range of bacteria. For example, in some sponge species, bacteria contribute up to 60% of the total biomass (Mohamed et al., 2008) and in

some studies they have been reported to produce bioactive metabolites similar to their host (Graça *et al.*, 2015; Elyakov *et al.*, 1991). Such cases are a subject of scientific debate. Several other studies have reported the isolation of bioactive metabolites from microorganisms associated with their invertebrate hosts (Kelecom, 2002; Bernan, Greenstein & Maiese, 1997; Hay, 2009). As an example, the marine sponge was the known source for the isolation of surfactant-like depsipeptides (Prasad *et al.*, 2011; Anjum *et al.*, 2016). Later, it was found that the bacterium *Bacillus pumilus* produces these compounds which were previously associated with just the sponge (Kalinovskaya *et al.*, 1995).

In view of this, it is important to understand the role of associated bacteria in the production of bioactive metabolites and the importance of their association with the invertebrate host, which is still not clear at this time. However, studies to explore the relationship between anti-microbial activity and anti-fouling activities are lacking. In this context, studies to elucidate the effectiveness/isolation of various types of extracts against/ from microorganisms will give a better understanding of these microorganisms' activities.

## SURVEY METHODOLOGY

All peer reviewed scientific papers were used for this review article. Extensive literature searches have been performed using various literature search engines, including Science Direct and PubMed with several terms: (1) *microbial natural bioactive compounds*; (2) *marine microbial natural bioactive compounds*; (3) *microbial anti-biofilm compounds*; and (4) *marine microbial anti-biofilm compounds*. All relevant studies meeting search criteria were included in this review.

## Need for antibiofilm or antibiofouling bioactive compounds with green chemistry

What is green chemistry and why do we need it? Chemistry is undeniably a very obvious part of our daily lives, and new chemical developments bring new environmental complications with unexpected harmful side effects. Hence, there is continued pressure on various chemical industries to reduce chemical waste and use their creativity to develop novel synthetic methods, reaction conditions, analytical tools, catalysts, and processes under the new paradigm of “green chemistry” (Erickson, Nelson & Winters, 2012).

The term green chemistry, also known as sustainable chemistry, was introduced by Paul Anastas in 1991 (Matthews, 2000). Twelve principles of green chemistry were created that encompass a new attitude toward industrial practices and chemical syntheses (Winterton, 2016). This set of principles describes guidelines for reduction/elimination of the generation of hazardous substances while designing, manufacturing, and applying any chemical products. It is the only science that focuses on the intrinsic hazards of a chemical or chemical process. Thus, to realize more sustainable thinking and implementation of sustainability criteria in research, development, and production, educational strategies have to include green chemistry principles and indicators of sustainability (Burmeister, Rauch & Eilks, 2012; Wilson & Schwarzman, 2009).

Formation of biofilms leads to biofouling, which is a process of gradual accumulation of waterborne organisms on natural and artificial surfaces of marine environment that

leads to corrosion and decline in the efficiency of moving parts (*Adnan et al., 2010; Adnan, Morton & Hadi, 2011; Yebra, Kiil & Dam-Johansen, 2004*). Advances in polymer chemistry in the early 1970s led to biofouling control, but it was based on the use of toxic substances such as tributyl tin (TBT) (*Rouhi, 1998*), copper or organic compounds (such as Sea-Nine, isothiazolone) (*Yebra, Kiil & Dam-Johansen, 2004*), or self-polishing copolymer anti-fouling paints, in which organotin compounds are the biocides. So far, organotin compounds are the most effective methods of biofouling control; however, these chemicals are toxic to aquatic environments (*Yebra, Kiil & Dam-Johansen, 2004*). The use of such toxic biocides presents environmental concerns involving anomalies in non-target organisms that include shell malformation in bivalves, depletion of oyster population, imposex in gastropods, and others (*Park et al., 2016*). TBT has been labeled as the most lethal substance to have been deliberately introduced into the marine environment by man (*Park et al., 2016*). In general, the use of such toxins in marine environments poses a real threat to the marine biota (*Fleming et al., 2006*). Effective from January 2008, the International Maritime Organization (IMO) and Marine Environmental Protection Committee (MEPC) introduced resolutions to ban the practice of using TBT or other substances containing tin as biocides (*Champ, 2003*). Therefore, there is a need for the development of environmental friendly non-toxic and natural anti-biofouling agents.

### **Advancement in methods used to prevent microbial biofilm and biofouling**

Since removal of biofilms is a very difficult, challenging, and demanding process, a complete and cost-effective cleaning procedure should be developed (*Caixeta et al., 2012*) because an inappropriate cleaning strategy generally leads to biofilm formation (*Garrett, Bhakoo & Zhang, 2008*). A number of methods can be used to prevent microbial biofilm formation and/or biofouling. Currently, these measures fall into three economically lucrative categories: (1) physical; (2) chemical; and (3) biological methods (*Cao et al., 2011*). In this review, we will discuss the biological methods that are currently used. Adsorption of bioactive compounds such as bacteriocins and nisin onto food-contact surfaces reduces bacterial adhesion. Enzyme mixtures have also proven to be effective for cleaning and biofilm removal (*Garrett, Bhakoo & Zhang, 2008; Stiefel et al., 2016; Guerra et al., 2005; Van Houdt & Michiels, 2010; Pimentel-Filho et al., 2014*). Unique properties of Endo H (endoglycosidases) for removal of bacteria such as *Staphylococci* and *E. coli* from glass and cloth surfaces makes it useful in buffers and detergent solutions (*Carpenter, Goldstein & Lad, 1995*). Colanic acid-degrading enzymes, derived from a *Streptomyces* isolate, are also useful for removal and prevention of biofilm formation (*Van, Bruggeman & Van, 1996*).

Marine benthic organism surfaces act as potential locations for the settlement of fouling organisms, including bacteria, algae, and invertebrates (*Cao et al., 2011; Pereira et al., 2002*). However, at the same time these organisms present a great variety of potential defense mechanisms against fouling organisms, which include surface sloughing, possession of spines, production of mucus, low surface energy, and secondary metabolite production (*Pereira et al., 2002*). Secondary metabolites demonstrating anti-fouling

properties opened a new perception into inhibition of overgrowth by epibionts and could possibly be used as commercial anti-biofoulants ([Wahl et al., 2012](#)). In fact, a vast range of tested anti-biofoulants have been patented. However, these anti-biofoulants have been tested under laboratory conditions, using the larvae of fouling organisms such as barnacles and bryozoans ([Pereira et al., 2002](#); [Zimmerman et al., 1997](#)). Moreover, there are only few studies in which this test has been performed under ecological settings, in which real concentrations of metabolites found in source organisms were used ([Da Gama et al., 2002](#)). From a molecular perspective, secondary marine metabolites seem to hinder bacterial colonization and can control biofilm formation/biofouling ([Kjelleberg et al., 1997](#)). To justify and prove this statement, further investigations, including ecological and molecular tests, are required.

### Ecological role of marine bioactive compounds

Natural marine products, secondary metabolites, enzymes, lipids, and heteropolysaccharides derived from marine sources are biologically active. Moreover, they are safer, cheaper, and less toxic than existing medicines ([D'Ayala, Malinconico & Laurienzo, 2008](#); [Bergé & Barnathan, 2015](#)). Out of 36 known living phyla, 34 are found in throughout the marine environment with more than 300,000+ known species of flora and fauna ([Baskaran et al., 2017](#)) (and still counting). However, the detailed ecological role of these marine animal-extracted secondary metabolites, which includes prevention of fouling, anti-predation, protection from ultraviolet radiation, mediation of spatial competition, and other functions ([Armstrong, Boyd & Burgess, 2000](#)) is still not clear and present areas for novel and extensive studies.

Many natural marine products are currently being tested in clinical trials, and various others are presently used for treating microbial infections (fungal, malarial, bacterial, viral, and nematode) and pain management in addition to cancer and inflammation control ([Martins et al., 2014](#)). The first marine compound to enter a human cancer clinical trial as a purified natural product was Didemnin-B from the Caribbean tunicate *Trididemnum solidum* ([Ram et al., 2008](#)). Unfortunately, secondary metabolites of macro-organisms are usually present in trace amounts, and that presents a drawback to use of the marine compounds. Due to less stock, the marine compound cannot compete with the development of widely available medicines. The developmental ways such as bioprocess engineering is currently the most important method for obtaining large quantities of beneficial secondary metabolites. This can only be accomplished by extracting and testing less investigated drug sources such as marine fungi and bacteria, which are vast untapped reservoirs of metabolic diversity ([Debbab et al., 2010](#)).

Microorganisms have some advantages such as isolation of the same compound in large-scale cultivation using biotechnological fermentations and advance bioprocessing engineering techniques with different parameters without ecological exploitation. They can also easily be genetically manipulated. There is still opportunity for an advanced scale of research and investigation to discover the potential of marine microorganisms as producers of novel drugs ([Malve, 2016](#)).

## Bioactive compounds from marine bacteria, fungi and cyanobacteria

Over the last several decades, microorganisms have been accepted as significant and unexploited resources for many unique bioactive compounds with clinical significance (Malve, 2016; Zhang *et al.*, 2005). It is very clear that microbial diversity found in the oceans is poorly understood, and <5% of marine bacterial and fungal species are known (de Felício *et al.*, 2015). Microbes thrive not only in the sea's surface waters but also in the lower and immeasurable depths (Das, Lyta & Khan, 2006). For survival, soft-bodied marine organisms rely on chemical defenses from the production of bioactive secondary metabolites either by themselves or in association with microflora (Mehubub *et al.*, 2014). These secondary metabolites may represent a diverse structure of classes that include terpenes, peptides, polypeptides, and compounds of mixed biosynthetic origin. A total of 961 new compounds from marine microorganisms were described in the year 2007 (Blunt *et al.*, 2009). These studies and the sharp rise in numbers of these compounds indicate that the study on secondary metabolites from marine bacteria and fungi as source of new bioactive metabolites has been steadily increasing. So far, only a few anti-larval settlement compounds have been isolated and identified from bacteria (Table 2).

One of the ways of discovering novel bioactive metabolites from marine microorganisms is via isolation of new microorganisms. Research over the years has demonstrated that out of 10% of cultivable microorganisms, only 1% were found to have clinical and industrial significance (Vartoukian, Palmer & Wade, 2010). Due to the complex nature of the oceans, marine bacteria have developed sophisticated physiological and biochemical systems with which they uniquely adapt to extreme habitats and various unfavorable marine environmental conditions. They live in a biologically competitive environment with unique conditions of pH, temperature, pressure, oxygen, light, nutrients, and salinity, which is especially rich in chlorine and bromine elements. Microbes can quickly sense, adapt, and respond to their environments (Adnan *et al.*, 2010; Adnan, Morton & Hadi, 2011) and can compete for defense and survival via the generation of unique secondary metabolites. Although the response to stress initiates the production of these compounds, they have shown value in pharmaceutical and biotechnological applications (Darvishi Harzevili & Chen, 2014).

Among all microorganisms, fungi are found to be a potentially useful source of pharmacologically active metabolites because they have the capability to adapt and survive in marine environments and to produce unique secondary metabolites. Fungi have also been widely distributed in marine habitats such as seawater, sediment, marine animals, and plants. Even though decomposition of dead plants and animal tissues has been suggested to play an important ecological role in recycling nutrients (Zhang *et al.*, 2015), the role of marine fungi is poorly understood. Marine fungi have been isolated from different marine habitats for the investigation of natural products (Imhoff, 2016). They found to form a mutual synergetic relationship with other marine organisms such as algae, sponges, mollusks and corals, but this association and related functions are rarely known (Ding *et al.*, 2011). Culture-based technology is used for the isolation of various fungi from marine habitats. Due to their uniqueness when compared with their terrestrial

**Table 2** List of few marine microbial species reported to produce bioactive metabolites with antibiofilm and antifouling activities (*Abu Sayem et al., 2011; Arai, Niikawa & Kobayashi, 2013; Buseti et al., 2015; Dos Santos et al., 2010; Estrela & Abraham, 2016; Fusetani, 2011; Hong & Cho, 2013; Jiang et al., 2011; Salta et al., 2013; Satheesh, Ba-Akdah & Al-Sofyani, 2016; Scopel et al., 2014; Shao et al., 2015; Shao et al., 2011; Yang et al., 2007*).

Microorganism	Bioactive compound	Biological activity	Effects against	Relevance
<b>Bioactive compounds isolated from marine bacteria</b>				
<i>Bacillus licheniformis</i>	$\alpha$ -D-galactopyranosyl-(1 $\rightarrow$ 2)-glycerol-phosphate	Antibiofilm	<i>Escherichia coli</i> and <i>Pseudomonas fluorescens</i>	Clinical
<i>Alteromonas</i> sp.	Ubiquinone-8	Antifouling	Larval settlement of barnacle <i>Balanus amphitrite</i>	Environmental
<i>Acinetobacter</i> sp.	6-bromoindole-3-carboxaldehyde	Antifouling	Larval settlement of barnacle <i>Balanus amphitrite</i>	Environmental
<i>Pseudomonas</i> sp.	Pyolipic acid	Antifouling	Larval settlement under laboratory and field experiment assays	Environmental
<i>Pseudomonas</i> sp.	Phenazine-1-carboxylic acid	Antifouling	Larval settlement under laboratory and field experiment assays	Environmental
<i>Pseudomonas</i> sp.	2-alkylquinol-4-ones	Antifouling	Larval settlement under laboratory and field experiment assays	Environmental
<i>Streptomyces praecox</i>	Diketopiperazines	Antifouling	Larval settlement under laboratory and field experiment assays	Environmental
<i>Streptomyces violaceoruber</i>	3-octa-1',3'-dienyl-4-methylfuran-2(5H)-one	Antifouling	Zoospores of <i>Ulva pertusa</i> , the diatom <i>Navicula annexa</i> , and the mussel <i>Mytilus edulis</i>	Environmental
<i>Streptomyces violaceoruber</i>	3-octa-1'-enyl-4-methylfuran-2(5H)-one	Antifouling	Zoospores of <i>Ulva pertusa</i> , the diatom <i>Navicula annexa</i> , and the mussel <i>Mytilus edulis</i>	Environmental
<i>Vibrio</i> sp. QY101	Exopolysaccharide A101	Antibiofilm	<i>Pseudomonas aeruginosa</i> FRD1	Clinical
<b>Bioactive compounds isolated from marine fungi</b>				
<i>Letendraea helminthicola</i>	3-methyl-N-(2-phenylethyl) butanamide	Antifouling	Larval settlement of barnacle <i>Balanus amphitrite</i>	Industrial/Environmental
<i>Scopulariopsis</i> sp.	Dihydroquinolin-2(1H)-one	Antifouling	Larval settlement of barnacle <i>Balanus amphitrite</i>	Industrial/Environmental
<i>Letendraea helminthicola</i>	Cyclo (D-Pro-D-Phe)	Antifouling	Larval settlement of barnacle <i>Balanus amphitrite</i>	Industrial/Environmental
<i>Cochliobolus lunatus</i>	Resorcyclic acid lactones	Antifouling	Larval settlement of barnacle <i>Balanus amphitrite</i>	Industrial/Environmental
<i>Emericella varicolor</i>	Ophiobolin K	Antibiofilm	<i>Mycobacterium bovis</i>	Clinical
<i>Emericella varicolor</i>	6-epi-ophiobolin K	Antibiofilm	<i>Mycobacterium smegmatis</i>	Clinical
<i>Emericella varicolor</i>	6-epi-ophiobolin	Antibiofilm	<i>Mycobacterium smegmatis</i>	Clinical
Unidentified marine fungus	Mevalonolactone	Antibiofilm	<i>Staphylococcus epidermidis</i>	Clinical
Marine <i>Penicillium</i> sp.	Cyclo(L-Tyr-L-Leu)	Antibiofilm	<i>Staphylococcus epidermidis</i>	Clinical/Environmental
<i>Penicillium commune</i>	Cyclo(L-Leu-L-Pro)	Antibiofilm	<i>Staphylococcus aureus</i>	Clinical/Environmental
<i>Cladosporium</i> sp.	F14 cyclo-(Phe-Pro)	Antibiofilm	<i>Loktanella hongkongensis</i> , <i>Micrococcus luteus</i> and <i>Ruegeria</i> sp.	Clinical/Environmental
<i>Cladosporium</i> sp.	cyclo-(Val-Pro) 4	Antibiofilm	<i>Loktanella hongkongensis</i>	Clinical/Environmental
<i>Aspergillus flavipes</i>	Flavipesin A 49	Antibiofilm	<i>Staphylococcus aureus</i> and <i>Bacillus subtilis</i>	Clinical

(Continued)

Table 2 (continued).

Microorganism	Bioactive compound	Biological activity	Effects against	Relevance
<b>Bioactive compounds isolated from marine algae</b>				
<i>Chondrus crispus</i>	(+)-Usnic acid	Antibiofilm	<i>Cobetia marina</i> and <i>Marinobacter hydrocarbonoclasticus</i>	Environmental
<i>Chondrus crispus</i>	Juglone	Antibiofilm	<i>Cobetia marina</i> and <i>Marinobacter hydrocarbonoclasticus</i>	Environmental
<i>Laurencia elata</i>	Elatol	Antifouling	<i>Leishmania amazonensis</i>	Clinical
<i>Halidrys siliquosa</i>	Organic extract	Antibiofilm	<i>Staphylococcus</i> ; <i>Streptococcus</i> ; <i>Enterococcus</i> ; <i>Pseudomonas</i> ; <i>Stenotrophomonas</i> ; and <i>Chromobacterium</i>	Clinical

counterparts, they have also shown to be phylogenetically distant (Reich & Labes, 2017; Rédou et al., 2015). They are a promising source for new bioactive natural products with high structural diversity when compared with other microbial sources isolated from the sea. Because of their living conditions, including tolerance to salinity, high pressure, nutrition requirements, temperature variations, and competition with other microbes, they may have developed unique metabolic pathways (Samuel, Prince & Prabakaran, 2011).

The most famous and first reported group of bioactive compounds obtained from marine-derived fungi were cephalosporins (with cephalosporin C) that were isolated by Brotzu in 1945 from a marine strain of *Acremonium chrysogenum*. They had been discovered in a sewage outlet in the Mediterranean Sea close to the island of Sardinia coast, and they exhibited pronounced antibacterial activities (Silber et al., 2016; Demain & Elander, 1999). After the discovery of cephalosporin C, sicayne was isolated from *Halocyphina vilosa* and was identified as the second antibiotic from marine-derived fungi (Kupka et al., 1981). Marine-derived fungi have been widely studied for novel anti-cancer, anti-bacterial, anti-plasmodial, anti-inflammatory, and anti-viral agents (Rajasekar et al., 2012). However, very few anti-biofilm/anti-biofouling bioactive metabolites have been derived from marine fungi.

More than 270 new natural products isolated from marine fungi between 1990 and 2002 followed by 330 new metabolites between 2002 and 2006 have been reported (Imhoff, 2016). In 2009 and 2010, >472 new structures were reported. The total number of new natural products from marine-derived fungi currently exceeds 1,000, and a few of them are in preclinical and clinical trials (Gomes et al., 2015). The ecological role of these multiple compounds include prevention of fouling, competition for space/resources, protection from ultraviolet radiation, and facilitation of reproduction (Nunez-Pons & Avila, 2015). These marine-derived fungal metabolites have been proven to be promising anti-fouling/anti-biofilm agents, but still required additional investigation and implementation in oceanic-related industries.

Marine cyanobacteria (blue-green algae) are another such group of prokaryotic photosynthetic organism and are potent sources of pharmacological and industrial products having diverse structures and habitats that are known to produce various

bioactive compounds (*Mourelle, Gómez & Legido, 2017*). The potential use of cyanobacteria has gained worldwide attention in view of their importance in agriculture, industry, pharmaceutical markets, and other areas (*Singh et al., 2017*). They are the richest sources of known and unique bioactive compounds, including toxins with potential for therapeutic applications (*Singh et al., 2017*). Marine cyanobacteria demonstrate a wide range of biological activities, including its use as biocontrol agents against various bacterial and fungal pathogens in addition to its use as a potential source of novel antibiotics. They are known to produce anti-bacterial and anti-fungal compounds (*Shantanu et al., 2013*), anti-cancerous and anti-neoplastic agents, and compounds that may be useful in the treatment of human immunodeficiency virus (*Vijayakumar & Menakha, 2015; Swain, Padhy & Singh, 2015*). Screening of cyanobacteria for potent anti-biofouling/anti-biofilm natural agents will open a new horizon for marine industries (*Table 2*). More than 50% of marine cyanobacteria are potentially exploitable for extracting bioactive substances, which can be effective anti-biofouling agents. Cyanobacterial blooms are rich sources of secondary metabolites with novel chemical and molecular structures (*Engene et al., 2011*). Exploring the efficiency of cyanobacterial products has an advantage; it can be grown in mass culture, which can be manipulated to attain optimal production of bioactive substances.

## CONCLUSIONS

Ever since the beginning of development of a civilized society, biologically active compounds, which are obtained from diverse range of microbes, have been extensively investigated. Bacterial biofilms are the dominant reason for biofouling in most commercial systems with no permanent solution toward removal of biofilms. Microbial secondary bioactive metabolites possess quite a few pharmaceutical applications. Use of ecofriendly biocides as an alternative to synthetic chemicals has recently emerged because the sole purpose of green chemistry is to either discover or to produce the best and novel chemical products that are safe for use with increased productivity. Unquestionably, our understanding of the field of microbial metabolites has significantly improved over the past several years, but there are still many steps to achieve a better understanding about the potential of marine microbial metabolites. We are in a promising era of science, which proves to be the right time; we can uncover the potential of marine microbes in the field of biofouling prevention and not in biofouling causation. In order to explore the natural anti-fouling/anti-biofilm compounds (green chemistry) and potential of marine microbes, this review will definitely draw attention to the search for bioactive metabolites that can resolve many difficulties and obstacles not only for oceanic industries, but many other industrial systems.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

The authors received no funding for this work.

## Competing Interests

The authors declare that they have no competing interests.

## Author Contributions

- Mohd Adnan conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Eyad Alshammari analyzed the data, authored or reviewed drafts of the paper, approved the final draft.
- Mitesh Patel performed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables.
- Syed Amir Ashraf contributed reagents/materials/analysis tools, prepared figures and/or tables.
- Saif Khan performed the experiments, analyzed the data, approved the final draft.
- Sibte Hadi conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, approved the final draft.

## Data Availability

The following information was supplied regarding data availability:

This is a review article that examined peer reviewed articles available in Science Direct and Pubmed.

## REFERENCES

- Abu Sayem AM, Manzo E, Ciavatta L, Tramice A, Cordone A, Zanfardino A, De Felice M, Varcamonti M. 2011. Anti-biofilm activity of an exopolysaccharide from a sponge-associated strain of *Bacillus licheniformis*. *Microbial Cell Factories* **10**(1):74 DOI 10.1186/1475-2859-10-74.
- Adnan M, Morton G, Hadi S. 2011. Analysis of *rpoS* and *bolA* gene expression under various stress-induced environments in planktonic and biofilm phase using  $2^{-\Delta\Delta CT}$  method. *Molecular and Cellular Biochemistry* **357**(1–2):275–282 DOI 10.1007/s11010-011-0898-y.
- Adnan M, Morton G, Singh J, Hadi S. 2010. Contribution of *rpoS* and *bolA* genes in biofilm formation in *Escherichia coli* K-12 MG1655. *Molecular and Cellular Biochemistry* **342**(1–2):207–213 DOI 10.1007/s11010-010-0485-7.
- Anjum K, Abbas SQ, Shah SAA, Akhter N, Batool S, ul Hassan SS. 2016. Marine sponges as a drug treasure. *Biomolecules & Therapeutics* **24**(4):347–362 DOI 10.4062/biomolther.2016.067.
- Arai M, Niikawa H, Kobayashi M. 2013. Marine-derived fungal sesterterpenes, ophiobolins, inhibit biofilm formation of Mycobacterium species. *Journal of Natural Medicines* **67**(2):271–275 DOI 10.1007/s11418-012-0676-5.
- Armstrong E, Boyd KG, Burgess JG. 2000. Prevention of marine biofouling using natural compounds from marine organisms. *Biotechnology Annual Review*. Vol. 6. Sixth Edition. Elsevier, 221–241.
- Baskaran R, Subramanian T, Zuo W, Qian J, Wu G, Kumar A. 2017. Major source of marine Actinobacteria and its biomedical application. In: Kalia VC, ed. *Microbial Applications Vol.2: Biomedicine, Agriculture and Industry*. Cham: Springer International Publishing, 55–82.
- Bekiari V, Nikolaou K, Koromilas N, Lainioti G, Avramidis P, Hotos G, Kallitsis JK, Bokias G. 2015. Release of polymeric biocides from synthetic matrices for marine biofouling applications. *Agriculture and Agricultural Science Procedia* **4**:445–450 DOI 10.1016/j.aaspro.2015.03.051.

- Bergé JP, Barnathan G. 2015.** Fatty acids from lipids of marine organisms: molecular biodiversity, roles as biomarkers, biologically active compounds, and economical aspects. In: Ulber R, Le Gal Y, eds. *Marine Biotechnology I*. Berlin, Heidelberg: Springer Berlin Heidelberg, 49–125.
- Bernan VS, Greenstein M, Maiese WM. 1997.** Marine microorganisms as a source of new natural products. In: Neidleman SL, Laskin AI, eds. *Advances in Applied Microbiology*. Vol. 43. Cambridge: Academic Press, 57–90.
- Blunt JW, Copp BR, Hu WP, Munro MH, Northcote PT, Prinsep MR. 2009.** Marine natural products. *Natural Product Reports* 26:170–244.
- Burmeister M, Rauch F, Eilks I. 2012.** Education for sustainable development (ESD) and chemistry education. *Chemistry Education Research and Practice* 13:59–68.
- Busetti A, Thompson PT, Tegazzini D, Megaw J, Maggs CA, Gilmore BF. 2015.** Antibiofilm activity of the brown alga *Halidrys siliquosa* against clinically relevant human pathogens. *Marine Drugs* 13(6):3581–3605 DOI 10.3390/md13063581.
- Caixeta DS, Scarpa TH, Brugnera DF, Friere DO, Alves E, De Abreu LR, Piccoli RH. 2012.** Chemical sanitizers to control biofilms formed by two *Pseudomonas* species on stainless steel surface. *Food Science and Technology* 32(1):142–150 DOI 10.1590/s0101-20612012005000008.
- Callow JA, Callow ME. 2011.** Trends in the development of environmentally friendly fouling-resistant marine coatings. *Nature Communications* 2(1):244 DOI 10.1038/ncomms1251.
- Cao S, Wang J, Chen H, Chen D. 2011.** Progress of marine biofouling and antifouling technologies. *Chinese Science Bulletin* 56(7):598–612 DOI 10.1007/s11434-010-4158-4.
- Carpenter RS, Goldstein IJ, Lad PJ. 1995.** Cleaning composition containing a type II endoglycosidase. *Google Patents*.
- Champ MA. 2003.** Economic and environmental impacts on ports and harbors from the convention to ban harmful marine anti-fouling systems. *Marine Pollution Bulletin* 46(8):935–940 DOI 10.1016/s0025-326x(03)00106-1.
- Ciriminna R, Bright FV, Pagliaro M. 2015.** Ecofriendly antifouling marine coatings. *ACS Sustainable Chemistry & Engineering* 3(4):559–565 DOI 10.1021/sc500845n.
- Da Gama BAP, Pereira RC, Carvalho AGV, Coutinho R, Valentin Y. 2002.** The effects of seaweed secondary metabolites on biofouling. *Biofouling* 18(1):13–20 DOI 10.1080/08927010290017680.
- Dang H, Lovell CR. 2016.** Microbial surface colonization and biofilm development in marine environments. *Microbiology and Molecular Biology Reviews* 80:91–138 DOI 10.1128/mnbr.00037-15.
- Darvishi Harzevili F, Chen H. 2014.** *Microbial Biotechnology: Progress and Trends*. Boca Raton: CRC Press.
- Das S, Lyla PS, Khan SA. 2006.** Marine microbial diversity and ecology: importance and future perspectives. *Current Science* 90(10):1325–1335.
- Davis AR. 1991.** Alkaloids and ascidian chemical defense: evidence for the ecological role of natural products from *Eudistoma olivaceum*. *Marine Biology* 111(3):375–379 DOI 10.1007/bf01319409.
- D’Ayala GG, Malinconico M, Laurienzo P. 2008.** Marine derived polysaccharides for biomedical applications: chemical modification approaches. *Molecules* 13(9):2069–2106 DOI 10.3390/molecules13092069.
- de Felício R, Pavão GB, de Oliveira ALL, Erbert C, Conti R, Pupo MT, Furtado NA, Ferreira EG, Costa-Lotufo LV, Young MCM, Yokoya NS. 2015.** Antibacterial, antifungal and cytotoxic activities exhibited by endophytic fungi from the Brazilian marine red alga *Bostrychia tenella*

- (Ceramiales). *Revista Brasileira de Farmacognosia* **25**(6):641–650  
DOI [10.1016/j.bjp.2015.08.003](https://doi.org/10.1016/j.bjp.2015.08.003).
- Debbab A, Aly AH, Lin WH, Proksch P. 2010.** Bioactive compounds from marine bacteria and fungi. *Microbial Biotechnology* **3**(5):544–563 DOI [10.1111/j.1751-7915.2010.00179.x](https://doi.org/10.1111/j.1751-7915.2010.00179.x).
- Demain AL, Elander RP. 1999.** The beta-lactam antibiotics: past, present, and future. *Antonie van Leeuwenhoek* **75**:5–19.
- Ding B, Yin Y, Zhang F, Li Z. 2011.** Recovery and phylogenetic diversity of culturable fungi associated with marine sponges *Clathrina luteoculcitella* and *Holoxea* sp. in the South China Sea. *Marine Biotechnology* **13**:713–721 DOI [10.1007/s10126-010-9333-8](https://doi.org/10.1007/s10126-010-9333-8).
- Dos Santos AO, Veiga-Santos P, Ueda-Nakamura T, Filho BP, Sudatti DB, Bianco EM, Pereira RC, Nakamura CV. 2010.** Effect of elatol, isolated from red seaweed *Laurencia dendroidea*, on *Leishmania amazonensis*. *Marine Drugs* **8**(11):2733–2743 DOI [10.3390/md8112733](https://doi.org/10.3390/md8112733).
- Edyvean RGJ, Thomas CJ, Brook R. 1988.** The effect of marine fouling on fatigue and corrosion-fatigue of offshore structures. In: Houghton DR, Smith RN, Eggins HOW, eds. *Biodeterioration 7*. Dordrecht: Springer Netherlands, 385–390.
- Elyakov GB, Kuznetsova T, Mikhailov VV, Maltsev II, Voinov VG, Fedoreyev SA. 1991.** Brominated diphenyl ethers from a marine bacterium associated with the sponge *Dysidea* sp. *Experientia* **47**(6):632–633 DOI [10.1007/bf01949894](https://doi.org/10.1007/bf01949894).
- Engene N, Choi H, Esquenazi E, Rottacker EC, Ellisman MH, Dorrestein PC, Gerwick WH. 2011.** Underestimated biodiversity as a major explanation for the perceived rich secondary metabolite capacity of the cyanobacterial genus *Lyngbya*. *Environmental Microbiology* **13**(6):1601–1610 DOI [10.1111/j.1462-2920.2011.02472.x](https://doi.org/10.1111/j.1462-2920.2011.02472.x).
- Erickson B, Nelson JE, Winters P. 2012.** Perspective on opportunities in industrial biotechnology in renewable chemicals. *Biotechnology Journal* **7**(2):176–185 DOI [10.1002/biot.201100069](https://doi.org/10.1002/biot.201100069).
- Estrela AB, Abraham WR. 2016.** Fungal metabolites for the control of biofilm infections. *Agriculture* **6**(3):37 DOI [10.3390/agriculture6030037](https://doi.org/10.3390/agriculture6030037).
- Fleming LE, Broad K, Clement A, Dewailly E, Elmir S, Knap A, Pomponi SA, Smith S, Solo Gabriele H, Walsh P. 2006.** Oceans and human health: emerging public health risks in the marine environment. *Marine Pollution Bulletin* **53**(10–12):545–560  
DOI [10.1016/j.marpolbul.2006.08.012](https://doi.org/10.1016/j.marpolbul.2006.08.012).
- Fusetani N. 2004.** Biofouling and antifouling. *Natural Product Reports* **21**(1):94–104  
DOI [10.1039/b302231p](https://doi.org/10.1039/b302231p).
- Fusetani N. 2011.** Antifouling marine natural products. *Natural Product Reports* **28**(2):400–410  
DOI [10.1039/c0np00034e](https://doi.org/10.1039/c0np00034e).
- Ganapiriya V, Maharajan A, Kumarasamy P. 2012.** Antifouling effect of bioactive compounds from marine sponge *Acanthella elongata* and different species of bacterial film on larval attachment of *Balanus amphitrite* (cirripedia, crustacea). *Brazilian Archives of Biology and Technology* **55**(3):395–402 DOI [10.1590/s1516-89132012000300010](https://doi.org/10.1590/s1516-89132012000300010).
- Garrett TR, Bhakoo M, Zhang Z. 2008.** Bacterial adhesion and biofilms on surfaces. *Progress in Natural Science* **18**(9):1049–1056 DOI [10.1016/j.pnsc.2008.04.001](https://doi.org/10.1016/j.pnsc.2008.04.001).
- Gomes NGM, Lefranc F, Kijjoa A, Kiss R. 2015.** Can some marine-derived fungal metabolites become actual anticancer agents? *Marine Drugs* **13**(6):3950–3991  
DOI [10.3390/md13063950](https://doi.org/10.3390/md13063950).
- Graça AP, Viana F, Bondoso J, Correia MI, Gomes L, Humanes M, Reis A, Xavier JR, Gaspar H, Lage OM. 2015.** The antimicrobial activity of heterotrophic bacteria isolated from the marine sponge *Erylus deficiens* (Astrophorida, Geodiidae). *Frontiers in Microbiology* **6**:389  
DOI [10.3389/fmicb.2015.00389](https://doi.org/10.3389/fmicb.2015.00389).

- Guerra NP, Araujo AB, Barrera Am, Agrasar AT, Macías CL, Carballo J, Pastrana L. 2005. Antimicrobial activity of nisin adsorbed to surfaces commonly used in the food industry. *Journal of Food Protection* 68(5):1012–1019 DOI 10.4315/0362-028x-68.5.1012.
- Hay ME. 2009. Marine chemical ecology: chemical signals and cues structure marine populations, communities, and ecosystems. *Annual Review of Marine Science* 1(1):193–212 DOI 10.1146/annurev.marine.010908.163708.
- Henrikson AA, Pawlik JR. 1998. Seasonal variation in biofouling of gels containing extracts of marine organisms. *Biofouling* 12(1–3):245–255 DOI 10.1080/08927019809378357.
- Holmström C, Kjelleberg S. 1999. Marine Pseudoalteromonas species are associated with higher organisms and produce biologically active extracellular agents. *FEMS Microbiology Ecology* 30(4):285–293 DOI 10.1111/j.1574-6941.1999.tb00656.x.
- Hong YK, Cho JY. 2013. Effect of seaweed epibiotic bacterium *Streptomyces violaceoruber* SCH-09 on marine fouling organisms. *Fisheries Science* 79(3):469–475 DOI 10.1007/s12562-013-0604-y.
- Imhoff JF. 2016. Natural products from marine fungi—still an underrepresented resource. *Marine Drugs* 14(1):19 DOI 10.3390/md14010019.
- Iorhemen OT, Hamza RA, Tay JH. 2016. Membrane bioreactor (MBR) technology for wastewater treatment and reclamation: membrane fouling. *Membranes* 6(2):33 DOI 10.3390/membranes6020033.
- Jensen PR, Fenical W. 1996. Marine bacterial diversity as a resource for novel microbial products. *Journal of Industrial Microbiology & Biotechnology* 17(5–6):346–351 DOI 10.1007/bf01574765.
- Jiang P, Li J, Han F, Duan G, Lu X, Gu Y, Yu W. 2011. Antibiofilm activity of an exopolysaccharide from marine bacterium *Vibrio* sp. QY101. *PLOS ONE* 6(4):e18514 DOI 10.1371/journal.pone.0018514.
- Joint I, Muhling M, Querellou JI. 2010. Culturing marine bacteria—an essential prerequisite for biodiscovery. *Microbial Biotechnology* 3(5):564–575 DOI 10.1111/j.1751-7915.2010.00188.x.
- Kalinovskaya NI, Kuznetsova TA, Rashkes YV, Milgrom YM, Milgrom EG, Willis RH, Wood AI, Kurtz HA, Carabedian C, Murphy P, Elyakov GB. 1995. Surfactin-like structures of five cyclic depsipeptides from the marine isolate of *Bacillus pumilus*. *Russian Chemical Bulletin* 44(5):951–955 DOI 10.1007/bf00696935.
- Kelecom A. 2002. Secondary metabolites from marine microorganisms. *Anais da Academia Brasileira de Ciências* 74(1):151–170 DOI 10.1590/s0001-37652002000100012.
- Kjelleberg S, Steinberg P, Givskov M, Gram L, Manefield M, de Nys R. 1997. Do marine natural products interfere with prokaryotic AHL regulatory systems? *Aquatic Microbial Ecology* 13:85–93 DOI 10.3354/ame013085.
- Kupka J, Anke T, Giannetti BM, Steglich W. 1981. Antibiotics from basidiomycetes. *Archives of Microbiology* 130:223–227.
- Lade H, Paul D, Kweon JH. 2014. Quorum quenching mediated approaches for control of membrane biofouling. *International Journal of Biological Sciences* 10(5):550–565 DOI 10.7150/ijbs.9028.
- Laudien J, Wahl M. 2004. Associational resistance of fouled blue mussels (*Mytilus edulis*) against starfish (*Asterias rubens*) predation: relative importance of structural and chemical properties of the epibionts. *Helgoland Marine Research* 58(3):162–167 DOI 10.1007/s10152-004-0181-7.
- Li Y, Xu Y, Fleischer CC, Huang J, Lin R, Yang L, Mao H. 2018. Impact of anti-biofouling surface coatings on the properties of nanomaterials and their biomedical applications. *Journal of Materials Chemistry B* 6(1):9–24 DOI 10.1039/c7tb01695f.

- Maki JS, Rittschof D, Costlow JD, Mitchell R. 1988.** Inhibition of attachment of larval barnacles, *Balanus amphitrite*, by bacterial surface films. *Marine Biology* **97**(2):199–206 DOI [10.1007/bf00391303](https://doi.org/10.1007/bf00391303).
- Malve H. 2016.** Exploring the ocean for new drug developments: Marine pharmacology. *Journal of Pharmacy and Bioallied Sciences* **8**(2):83–91 DOI [10.4103/0975-7406.171700](https://doi.org/10.4103/0975-7406.171700).
- Martins A, Vieira H, Gaspar H, Santos S. 2014.** Marketed marine natural products in the pharmaceutical and cosmeceutical industries: tips for success. *Marine Drugs* **12**(2):1066–1101 DOI [10.3390/md12021066](https://doi.org/10.3390/md12021066).
- Matthews MA. 2000.** *Green Chemistry. Kirk-Othmer Encyclopedia of Chemical Technology*. Hoboken: John Wiley & Sons, Inc.
- Mehub FM, Lei J, Franco C, Zhang W. 2014.** Marine sponge derived natural products between 2001 and 2010: trends and opportunities for discovery of bioactives. *Marine Drugs* **12**(8):4539–4577 DOI [10.3390/md12084539](https://doi.org/10.3390/md12084539).
- Mohamed NM, Rao V, Hamann MT, Kelly M, Hill RT. 2008.** Monitoring bacterial diversity of the marine sponge *Ircinia strobilina* upon transfer into aquaculture. *Applied and Environmental Microbiology* **74**(13):4133–4143 DOI [10.1128/aem.00454-08](https://doi.org/10.1128/aem.00454-08).
- Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. 2011.** How many species are there on earth and in the ocean? *PLOS Biology* **9**(8):e1001127 DOI [10.1371/journal.pbio.1001127](https://doi.org/10.1371/journal.pbio.1001127).
- Mourelle LM, Gómez PC, Legido LJ. 2017.** The potential use of marine microalgae and cyanobacteria in cosmetics and thalassotherapy. *Cosmetics* **4**(4):46 DOI [10.3390/cosmetics4040046](https://doi.org/10.3390/cosmetics4040046).
- Nunez-Pons L, Avila C. 2015.** Natural products mediating ecological interactions in Antarctic benthic communities: a mini-review of the known molecules. *Natural Product Reports* **32**(7):1114–1130 DOI [10.1039/c4np00150h](https://doi.org/10.1039/c4np00150h).
- Nurioglu AG, Esteves AC, de With G. 2015.** Non-toxic, non-biocide-release antifouling coatings based on molecular structure design for marine applications. *Journal of Materials Chemistry B* **3**(32):6547–6570 DOI [10.1039/c5tb00232j](https://doi.org/10.1039/c5tb00232j).
- Omae I. 2003.** Organotin antifouling paints and their alternatives. *Applied Organometallic Chemistry* **17**(2):81–105 DOI [10.1002/aoc.396](https://doi.org/10.1002/aoc.396).
- Park MS, Kim YD, Kim BM, Kim YJ, Kim JK, Rhee JS. 2016.** Effects of antifouling biocides on molecular and biochemical defense system in the gill of the pacific oyster *Crassostrea gigas*. *PLOS ONE* **11**(12):e0168978 DOI [10.1371/journal.pone.0168978](https://doi.org/10.1371/journal.pone.0168978).
- Penesyan A, Kjelleberg S, Egan S. 2010.** Development of novel drugs from marine surface associated microorganisms. *Marine Drugs* **8**(3):438–459 DOI [10.3390/md8030438](https://doi.org/10.3390/md8030438).
- Pereira RC, Carvalho AGV, Gama BAP, Coutinho R. 2002.** Field experimental evaluation of secondary metabolites from marine invertebrates as antifoulants. *Brazilian Journal of Biology* **62**(2):311–320 DOI [10.1590/s1519-69842002000200015](https://doi.org/10.1590/s1519-69842002000200015).
- Pimentel-Filho NdJ, Martins MCdF, Nogueira GB, Mantovani HC, Vanetti MC. 2014.** Bovicin HC5 and nisin reduce *Staphylococcus aureus* adhesion to polystyrene and change the hydrophobicity profile and Gibbs free energy of adhesion. *International Journal of Food Microbiology* **190**:1–8 DOI [10.1016/j.ijfoodmicro.2014.08.004](https://doi.org/10.1016/j.ijfoodmicro.2014.08.004).
- Prasad P, Aalbersberg W, Feussner KD, Van Wagoner RM. 2011.** Papuamides E and F, cytotoxic depsipeptides from the marine sponge *Melophlus* sp. *Tetrahedron* **67**(44):8529–8531 DOI [10.1016/j.tet.2011.08.100](https://doi.org/10.1016/j.tet.2011.08.100).
- Qian PY, Xu Y, Fusetani N. 2009.** Natural products as antifouling compounds: recent progress and future perspectives. *Biofouling* **26**(2):223–234 DOI [10.1080/08927010903470815](https://doi.org/10.1080/08927010903470815).

- Rajasekar T, Balaji S, Kumaran S, Deivasigamani B, Pugzhavendhan SR. 2012. Isolation and characterization of Marine fungal metabolites against clinical pathogens. *Asian Pacific Journal of Tropical Disease* 2:S387–S392 DOI 10.1016/s2222-1808(12)60187-x.
- Ram S, Mukul S, Penny J, Diwan SR. 2008. Clinical status of anti-cancer agents derived from marine sources. *Anti-Cancer Agents in Medicinal Chemistry* 8(6):603–617 DOI 10.2174/1871520610808060603.
- Rédou V, Navarri M, Meslet-Cladière L, Barbier G, Burgaud G. 2015. Species richness and adaptation of marine fungi from deep-subseafloor sediments. *Applied and Environmental Microbiology* 81(10):3571–3583 DOI 10.1128/aem.04064-14.
- Reich M, Labes A. 2017. How to boost marine fungal research: a first step towards a multidisciplinary approach by combining molecular fungal ecology and natural products chemistry. *Marine Genomics* 36:57–75 DOI 10.1016/j.margen.2017.09.007.
- Rouhi AM. 1998. The squeeze on tributyltins. *Chemical & Engineering News* 76(17):41–42 DOI 10.1021/cen-v076n017.p041.
- Salta M, Wharton JA, Dennington SP, Stoodley P, Stoke KR. 2013. Anti-biofilm performance of three natural products against initial bacterial attachment. *International Journal of Molecular Sciences* 14(11):21757–21780 DOI 10.3390/ijms141121757.
- Samuel P, Prince L, Prabakaran P. 2011. Antibacterial activity of marine derived fungi collected from South East Coast of Tamilnadu, India. *Journal of Microbiology and Biotechnology Research* 1(4):86–94.
- Satheesh S, Ba-Akdah MA, Al-Sofyani AA. 2016. Natural antifouling compound production by microbes associated with marine macroorganisms—a review. *Electronic Journal of Biotechnology* 21:26–35 DOI 10.1016/j.ejbt.2016.02.002.
- Scopel M, Abraham WR, Antunes AL, Henriques AT, Macedo AJ. 2014. Mevalonolactone: an inhibitor of staphylococcus epidermidis adherence and biofilm formation. *Medicinal Chemistry* 10(3):246–251 DOI 10.2174/15734064113096660055.
- Shantanu B, Deep PR, Binata N, Monalisa P, Bharati M. 2013. Antimicrobial activity of two diazotropic cyanobacteria against Staphylococcus aureus. ID—20133248222. *International Journal of Medicinal and Aromatic Plants* 3:283–292.
- Shao CL, Xu RF, Wang CY, Qian PY, Wang KL, Wei MY. 2015. Potent antifouling marine dihydroquinolin-2(1H)-one-containing alkaloids from the gorgonian coral-derived fungus Scopulariopsis sp. *Marine Biotechnology* 17(4):408–415 DOI 10.1007/s10126-015-9628-x.
- Shao CL, Wu HX, Wang CY, Liu QA, Xu Y, Wei MY, Qian PY, Gu YC, Zheng CJ, She ZG, Lin YC. 2011. Potent antifouling resorcylic acid lactones from the gorgonian-derived fungus *Cochliobolus lunatus*. *Journal of Natural Products* 74:629–633.
- Silber J, Kramer A, Labes A, Tasdemir D. 2016. From discovery to production: biotechnology of marine fungi for the production of new antibiotics. *Marine Drugs* 14(7):137 DOI 10.3390/md14070137.
- Singh R, Parihar P, Singh M, Bajguz A, Kumar J, Singh S, Singh VP, Prasad SM. 2017. Uncovering potential applications of cyanobacteria and algal metabolites in biology, agriculture and medicine: current status and future prospects. *Frontiers in Microbiology* 8:515 DOI 10.3389/fmicb.2017.00515.
- Stiefel P, Mauerhofer S, Schneider J, Maniura-Weber K, Rosenberg U, Ren Q. 2016. Enzymes enhance biofilm removal efficiency of cleaners. *Antimicrobial Agents and Chemotherapy* 60(6):3647–3652 DOI 10.1128/aac.00400-16.
- Swain SS, Padhy RN, Singh PK. 2015. Anticancer compounds from cyanobacterium *Lyngbya* species: a review. *Antonie van Leeuwenhoek* 108(2):223–265 DOI 10.1007/s10482-015-0487-2.

- Van SMMP, Bruggeman G, Van PJ. 1996.** Exopolysaccharide degrading enzyme and use of the same. *Google Patents*.
- Van Houdt R, Michiels CW. 2010.** Biofilm formation and the food industry, a focus on the bacterial outer surface. *Journal of Applied Microbiology* **109**(4):1117–1131  
DOI [10.1111/j.1365-2672.2010.04756.x](https://doi.org/10.1111/j.1365-2672.2010.04756.x).
- Vartoukian SR, Palmer RM, Wade WG. 2010.** Strategies for culture of ‘unculturable’ bacteria. *FEMS Microbiology Letters* **309**:1–7 DOI [10.1111/j.1574-6968.2010.02000.x](https://doi.org/10.1111/j.1574-6968.2010.02000.x).
- Vijayakumar S, Menakha M. 2015.** Pharmaceutical applications of cyanobacteria—a review. *Journal of Acute Medicine* **5**(1):15–23 DOI [10.1016/j.jacme.2015.02.004](https://doi.org/10.1016/j.jacme.2015.02.004).
- Wahl M, Banaigs B. 1991.** Marine epibiosis. III. Possible antifouling defense adaptations in *Polysyncraton lacazei* (Giard) (Didemnidae, Ascidiacea). *Journal of Experimental Marine Biology and Ecology* **145**(1):49–63 DOI [10.1016/0022-0981\(91\)90005-h](https://doi.org/10.1016/0022-0981(91)90005-h).
- Wahl M, Goecke F, Labes A, Dobretsov S, Weinberger F. 2012.** The second skin: ecological role of epibiotic biofilms on marine organisms. *Frontiers in Microbiology* **3**:292  
DOI [10.3389/fmicb.2012.00292](https://doi.org/10.3389/fmicb.2012.00292).
- Wahl M, Jensen PR, Fenical W. 1994.** Chemical control of bacterial epibiosis on ascidians. *Marine Ecology Progress Series* **110**:45–57 DOI [10.3354/meps110045](https://doi.org/10.3354/meps110045).
- Wang KL, Wu ZH, Wang Y, Hao YY, Wang Y. 2017.** Mini-review: antifouling natural products from marine microorganisms and their synthetic analogs. *Marine Drugs* **15**(9):266  
DOI [10.3390/md15090266](https://doi.org/10.3390/md15090266).
- Wilson MP, Schwarzman MR. 2009.** Toward a new U.S. chemicals policy: rebuilding the foundation to advance new science, green chemistry, and environmental health. *Environmental Health Perspectives* **117**(8):1202–1209 DOI [10.1289/ehp.0800404](https://doi.org/10.1289/ehp.0800404).
- Winterton N. 2016.** Green chemistry: deliverance or distraction? *Clean Technologies and Environmental Policy* **18**(4):991–1001 DOI [10.1007/s10098-016-1118-y](https://doi.org/10.1007/s10098-016-1118-y).
- Yang LH, Miao L, Lee OO, Li X, Xiong H, Pang KL, Vrijmoed L, Qian PY. 2007.** Effect of culture conditions on antifouling compound production of a sponge-associated fungus. *Applied Microbiology and Biotechnology* **74**(6):1221–1231 DOI [10.1007/s00253-006-0780-0](https://doi.org/10.1007/s00253-006-0780-0).
- Yebra DM, Kiil S, Dam-Johansen K. 2004.** Antifouling technology—past, present and future steps towards efficient and environmentally friendly antifouling coatings. *Progress in Organic Coatings* **50**(2):75–104 DOI [10.1016/j.porgcoat.2003.06.001](https://doi.org/10.1016/j.porgcoat.2003.06.001).
- Zhang L, An R, Wang J, Sun N, Zhang S, Hu J, Kuai J. 2005.** Exploring novel bioactive compounds from marine microbes. *Current Opinion in Microbiology* **8**(3):276–281  
DOI [10.1016/j.mib.2005.04.008](https://doi.org/10.1016/j.mib.2005.04.008).
- Zhang T, Wang NF, Zhang YQ, Liu HY, Yu LY. 2015.** Diversity and distribution of fungal communities in the marine sediments of Kongsfjorden, Svalbard (High Arctic). *Scientific Reports* **5**(1):14524 DOI [10.1038/srep14524](https://doi.org/10.1038/srep14524).
- Zhang XY, Xu XY, Peng J, Ma CF, Nong XH, Bao J, Zhang GZ, Qi SH. 2014.** Antifouling potentials of eight deep-sea-derived fungi from the South China Sea. *Journal of Industrial Microbiology & Biotechnology* **41**(4):741–748 DOI [10.1007/s10295-014-1412-9](https://doi.org/10.1007/s10295-014-1412-9).
- Zimmerman RC, Alberte RS, Todd JS, Crews P. 1997.** Phenolic acid sulfate esters for prevention of marine biofouling. *Google Patents*.