

Central Lancashire Online Knowledge (CLOK)

Title	Physiological, kinematic, and electromyographic responses to kinesiology-type patella tape in elite cyclists
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/24887/
DOI	https://doi.org/10.1016/j.jelekin.2018.11.009
Date	2019
Citation	Hébert-Losier, K, Yin, NS, Beavena, CM, Li, CTC and Richards, James (2019) Physiological, kinematic, and electromyographic responses to kinesiology-type patella tape in elite cyclists. Journal of Electromyography and Kinesiology, 44. pp. 36-45. ISSN 1050-6411
Creators	Hébert-Losier, K, Yin, NS, Beavena, CM, Li, CTC and Richards, James

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.jelekin.2018.11.009>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

1 **Physiological, kinematic, and electromyographic responses to kinesiology-type patella**
2 **tape in elite cyclists**

3

4 Kim Hébert-Losier^{a*}, Ngieng Siew Yin^b, C. Martyn Beaven^a, Chris Tee Chow Li^b, and Jim
5 Richards^c

6

7 *^aFaculty of Health, Sport and Human Performance, University of Waikato, Tauranga, New*
8 *Zealand*

9

10 *^bDepartment of Sports Science, National Sports Institute of Malaysia, Kuala Lumpur,*
11 *Malaysia*

12

13 *^bSchool of Health Sciences, University of Central Lancashire, Preston, Lancashire, United*
14 *Kingdom*

15

16 *Corresponding Author: Kim Hébert-Losier,

17 e-mail: kim.hebert-losier@waikato.ac.nz

18

19 **Keywords:** 3D analysis; biomechanics; electromyography; kinesiology; performance;
20 physiology

21

22 **CONFLICT OF INTEREST**

23 None.

24

25 **FINANCIAL DISCLOSURE**

26 None.

27

28 **ABSTRACT**

29

30 Kinesiology-type tape (KTT) has become popular in sports for injury prevention,
31 rehabilitation, and performance enhancement. Many cyclists use patella KTT; however, its
32 benefits remain unclear, especially in uninjured elite cyclists. We used an integrated
33 approach to investigate acute physiological, kinematic, and electromyographic responses to
34 patella KTT in twelve national-level male cyclists. Cyclists completed four, 4-minute
35 submaximal efforts on an ergometer at 100 and 200 W with and without patella KTT.
36 Economy, energy cost, oxygen cost, heart rate, efficiency, 3D kinematics, and lower-body
37 electromyography signals were collected over the last minute of each effort. Comfort levels
38 and perceived change in knee stability and performance with KTT were recorded.
39 The effects of KTT were either unclear, non-significant, or clearly trivial on all collected
40 physiological and kinematic measures. KTT significantly, clearly, and meaningfully
41 enhanced *vastus medialis* peak, mean, and integrated electromyographic signals, and *vastus*
42 *medialis-to-lateralis* activation. Electromyographic measures from *biceps femoris* and
43 *biceps-to-rectus femoris* activation ratio decreased in either a significant or clinically
44 meaningful manner. Despite most cyclists perceiving KTT as comfortable, increasing
45 stability, and improving performance, the intervention exerted no considerable effects on all
46 physiological and kinematic measures. KTT did alter neuromuscular recruitment, which has
47 potential implications for injury prevention.

48 INTRODUCTION

49 Many health professionals, athletes, and coaches use kinesiology-type tape (KTT), with the
50 intent to manage musculoskeletal sport injuries; however, growing evidence suggests no
51 additional benefit from KTT application compared to placebo taping or active control
52 treatment methods when managing musculoskeletal conditions (Ouyang et al., 2018,
53 Williams et al., 2012). On the other hand, several beneficial effects from KTT application
54 have been reported, including enhancement in muscle activation (Gilleard et al., 1998);
55 improved biomechanics, joint, and patella alignment (Lyman et al., 2017, Merino-Marban et
56 al., 2013); and decreased pain (Bockrath et al., 1993, Merino-Marban et al., 2013). It is
57 worth noting, however, that many studies report no effect from KTT on these measures
58 (Halski et al., 2015) or athletic performance (Lins et al., 2013, Reneker et al., 2018).
59 Underlying reasons for such contrasting scientific findings likely include the varied
60 application methods, differences in the mechanical properties of KTT across brands
61 (Matheus et al., 2017), targeted population, and individuals' perceived benefits of KTT with
62 a potential for placebo effect (Mak et al., 2018) .

63

64 Cycling is a popular recreational and competitive sporting activity worldwide, and a
65 common exercise modality used during rehabilitation. At an elite level, athletes and coaches
66 continually seek for ways to improve performance through marginal gains and prevent the
67 occurrence of injuries. KTT is routinely used by coaches and athletes as an ergogenic aid
68 (Reneker et al., 2018), with various forms of taping employed to prevent injury occurrence
69 or recurrence (Zech and Wellmann, 2017). Given that up to 94% of professional cyclists
70 suffer from at least one overuse injury annually (Silberman, 2013), the visible increase in
71 use of KTT amongst elite cyclists for prophylactic purposes is not surprising.

72

73 Taping or bracing are frequently used to alleviate patellofemoral pain (PFP) symptoms and
74 can impact knee motion during cycling (Theobald et al., 2012). Non-specific KTT
75 application has been shown to be as effective as specific application for reducing pain and

inducing significant changes in lower-body cycling biomechanics in a symptomatic population group (Theobald et al., 2014). However, any potential positive effect of taping on energy cost of cycling, neuromuscular recruitment patterns, and performance of asymptomatic high-level cyclists has not been examined, despite the visibly increased prevalence of use in the cycling community and KTT marketing campaigns.

Using an integrated approach, our aim was therefore to investigate the acute physiological, kinematic, and electromyographic outcomes in response to applying KTT to the knee of elite cyclists. As perceptions may influence outcomes of interventions, individual perceptions were also assessed. We hypothesised that taping would be accompanied by changes in muscle recruitment patterns, cycling economy and efficiency, and perceived stability that have the potential to modulate cycling performance.

MATERIALS & METHODS

Participants

All male cyclists of the National Cycling Team training at the National Sports Institute for Malaysia ($n = 12$) were invited and accepted to participate in this study. These 12 national cyclists (mean \pm standard deviation (SD): age, 21.7 ± 2.8 years; body mass, 65.6 ± 5.4 kg; and height, 172.7 ± 3.4 cm) with at least four years of training experience provided written informed consent to participate in this study, which adhered to The Code of Ethics of the World Medical Association (*Declaration of Helsinki*) and was approved by our Institutional Ethical Review Board (ISNRP 29/2015). Inclusion criteria were cyclists training at the National Sports Institute of Malaysia for the National Cycling Team, good self-reported general health, and at least 18 years of age. Cyclists with current or recent (< 1 month) musculoskeletal injuries, joint pathologies, or medical contraindications to physical exertion were excluded.

Design

104 All cyclists attended 3 sessions at the biomechanics laboratory of the National Sports
105 Institute of Malaysia, one week apart. The first two weeks were familiarization sessions and
106 the third week was used to investigate the acute effects of KTT application through a
107 repeated-measures randomized experimental study design.

108

109 Given that ergometer versus outdoor cycling can affect cycling physiological measures
110 (Bertucci et al., 2012) and pedalling biomechanics (Bertucci et al., 2007), cyclists brought
111 their road bikes to the laboratory the first week of testing. Bike setup parameters were
112 recorded and employed to individualize setup on a Lode cycling ergometer (Excalibur Sport,
113 Lode B.V., Groningen, Netherlands), with all cyclists using their habitual cleats. The final
114 ergometer setup was recorded and used across the three weeks. Baseline demographics,
115 including leg dominance (self-perceived stronger cycling side), were recorded. Body mass
116 (kg) was measured weekly and subsequently used to calculate relative physiological
117 variables.

118

119 Cyclists began all sessions with a 2-min cycling warm-up on the ergometer after being set-
120 up with the monitoring equipment. Cyclists then performed a submaximal 4-minute cycling
121 effort at 100 W, followed immediately by a second submaximal 4-minute cycling effort at
122 200 W. The efforts were completed without KTT during the familiarization weeks, and
123 twice on the third week: once with and once without KTT. The no tape (NT) and KTT
124 conditions were completed in a block-randomized order and separated by a 5-minute passive
125 seated rest. Thus, all cyclists completed four 4-minute efforts: NT 100 W, NT 200 W, KTT
126 100 W, and KTT 200 W. The powers of 100 and 200 W were selected to ensure that cycling
127 efforts were below the anaerobic threshold and to compute delta efficiency (Coyle et al.,
128 1992) (see **Physiology**). Furthermore, the application of patella KTT at these powers has
129 been shown to alter lower-body cycling biomechanics in previous studies (Theobald et al.,
130 2014), with these power levels set alongside the National Cycling Team of Malaysia to
131 inform their practice and use of KTT application in longer steady-state riding situations.

132

133 *Taping method*

134 The taping application we used was based on a method previously reported to induce
135 changes in cycling biomechanics within the power ranges here examined (Theobald et al.,
136 2012, Theobald et al., 2014). With cyclists seated on the ergometer with the leg at the
137 bottom of their power stroke , a strip of KTT (RockTape™, RockTape Inc., California) of
138 length equal to 50% of individual knee circumference was applied on to the centre of the
139 patella with light tension (approximately 25% of stretch to the tape). The medial and lateral
140 tape edges were aligned with the medial and lateral knee-joint lines (**Figure 1**). The same
141 experienced physiotherapist applied the tape to all cyclists. This simple KTT method (i.e.,
142 across the patella) was selected given its ease-of-use and findings from previous studies
143 indicating that such a method impacts cycling biomechanics in a manner that is comparable
144 to that of a more intricate KTT application method (Theobald et al., 2014). Given the
145 minimalist KTT method applied, a “placebo” taping method was not implemented.

146 *****Insert Figure 1*****

147

148 *Physiology*

149 Oxygen consumption (VO_2), carbon dioxide output (VCO_2), and heart rate were monitored
150 throughout the 4-minute experimental efforts using a calibrated K5 wearable metabolic
151 technology system (COSMED, Rome, Italy). All physiological measures were averaged over
152 the last minute where steady state was observed. VO_2 was used to determine steady-state
153 relative oxygen cost (mL/kg/km) and absolute cycling economy (W/L/min). From the VO_2
154 and CO_2 data, the relative energy cost of efforts (kcal/kg/km) was estimated using the energy
155 expenditure equations described by Jeukendrup and Wallis (2005). Gross efficiency (%) and
156 delta efficiency (%) were calculated as suggested by Coyle et al. (1992) using the ratio of
157 work accomplished (watts converted to kcal/min) to absolute energy cost (kcal/min) for
158 gross efficiency, and the reciprocal of the slope that describes the relationship between the
159 absolute energy cost and work accomplished for delta efficiency.

160

161 *Kinematics*

162 Lower-body, trunk, and pelvis movements were captured in 3D during the last minute of
163 each 4-minute cycling effort at 300 Hz using 10 Oqus 300 infrared cameras and the Qualisys
164 Track Manager Software version 2.12 (Qualisys AB, Gothenburg, Sweden). Forty-six retro-
165 reflective markers (12 mm in diameter) were affixed to the skin, clothes, and shoes of
166 cyclists based on the Calibrated Anatomical System Technique (Cappozzo et al., 1997) and
167 following established guidelines (Grood and Suntay, 1983). All 46 markers were used for
168 static calibration; whereas 14 markers were removed for the cycling efforts (**Figure 2**).

169

170 *****Insert Figure 2*****

171

172 An 8-segment biomechanical model with 6 degrees of freedom at each joint was constructed
173 in Visual3D Professional™ Software version 5.02.30 (C-Motion Inc., Germantown, MD,
174 USA), with the local coordinates of the trunk, pelvis, thighs, shanks, and feet derived from
175 the static calibration and the pelvis used to define hip-joint centres (Bell et al., 1989). Prior
176 to each session, the measurement volume was calibrated using a 750-mm wand and L-frame
177 that defined the Cartesian origin of the laboratory. Cyclists were then requested to sit on the
178 saddle of the ergometer, with legs hanging to the side, and remain motionless to allow static
179 calibration.

180

181 *Electromyography*

182 The electromyography (EMG) signals from the following four muscles were recorded on
183 both the dominant and non-dominant sides: *vastus medialis* (VM), *vastus lateralis* (VL),
184 *rectus femoris* (RF) and *biceps femoris* (BF). Signals were recorded using Noraxon's Dual
185 EMG surface Ag/AgCl electrodes (17.5 mm inter-electrode distance), wireless EMG
186 sensors, and Desktop DTS data logger (Noraxon USA Inc., Scottsdale, AZ). EMG data were
187 sampled at 1500 Hz, low-pass filtered at 500 Hz, and digitally integrated through the

188 Qualisys Track Manager Software. Skin preparation and electrode positioning followed the
189 Surface EMG for Noninvasive Assessment of Muscle (Hermens et al., 2000), International
190 Society of Electrophysiology and Kinesiology (Merletti and di Torino, 1999), and published
191 protocols (Gilleard et al., 1998). Cyclists completed a few cycling revolutions before
192 experimentation to allow visual inspection of EMG signal quality. Sensors were checked and
193 reapplied if artefacts were observed.

194

195 ***Perception***

196 Perceived change in knee stability and performance with KTT compared to NT was assessed
197 at the end of the experimental session using a 5-point Likert (1932) Scale from negative (1)
198 to positive (5) perception, with the mid-point value representing no change (3). Comfort
199 level of KTT was also assessed using a similar method. Anchor points ranged from very
200 uncomfortable (1), much less stable (1), and much worse (1) to very comfortable (5), much
201 more stable (5), and much better (5) for comfort, knee stability, and performance,
202 respectively.

203

204 ***Data processing***

205 Kinematic and EMG data were exported to the C3D format and processed in Visual 3D.
206 Marker data were filtered using a 4th order zero-lag 15 Hz Butterworth bidirectional filter.
207 Kinematic parameters were then calculated using rigid-body analysis and Euler angles
208 obtained from the static calibration. Hip, knee, and ankle angles in the sagittal (flexion-
209 extension), coronal (adduction-abduction), and transverse (internal-external rotation) planes
210 were calculated using an x-y-z Cardan sequence equivalent to the Joint Coordinate System
211 (Grood and Suntay, 1983), with the pelvis angles in the sagittal (anterior-posterior), coronal
212 (dominant, non-dominant obliquity), and transverse (dominant, non-dominant rotation)
213 planes defined relative to the laboratory. Trunk angles in the sagittal (flexion-extension),
214 coronal (dominant, non-dominant lateral flexion), and transverse (dominant, non-dominant
215 rotation) planes were also defined in relation to the laboratory coordinates. Data were

216 divided into movement cycles and time-normalized based on maximal knee flexion events.
217 Ensemble-average kinematic curves were generated for each participant and cycling effort,
218 and range of motion (ROM) values extracted.
219
220 EMG signal data were zeroed to remove any baseline offset and a 20-Hz high-pass filter
221 applied to remove movement artefacts. Signals were subsequently rectified and linear
222 envelopes generated by smoothing the data using a low-pass, 4th order, zero-lag 15 Hz
223 Butterworth filter. The linear envelope for each muscle was then normalized to the highest
224 observed signal across all four conditions examined (% max). Similar to the kinematic data,
225 ensemble-average EMG signal curves time normalized to maximal knee flexion events were
226 generated from which mean and peak EMG signal values were extracted. An integrated
227 EMG (iEMG) signal was also generated by integrating the linear envelop from the start to
228 the end of each movement cycle, which was then normalized to the maximal observed iEMG
229 across all four efforts (% max).

230

231 *Statistical analysis*

232 Mean and SD values were computed for all parameters for both the 100 and 200 W efforts
233 and dominant and non-dominant sides. Changes in mean (Δ_{mean}) and standardized effect
234 sizes (ES) were computed to quantify the acute effect of KTT; with ES considered small,
235 moderate, large, and very large when reaching thresholds of 0.2, 0.6, 1.2, 2.0, and trivial
236 when < 0.2 (Smith and Hopkins, 2011). An effect was deemed ‘clear’ when its 90%
237 confidence limit did not overlap the thresholds for small positive and small negative effects
238 (i.e., 5%); and ‘likely’ to be clinically meaningful when its probability exceeded 75% (Smith
239 and Hopkins, 2011).

240

241 Paired *t*-tests were used to investigate differences between the tape and no-tape condition for
242 the measures of interest, with the threshold for statistical significance set at $P \leq 0.05$. All

243 data were analysed using customized statistical spreadsheets (Microsoft Excel 2013,
244 Microsoft Corp, Redmond WA, USA).

245

246 **RESULTS**

247 *Physiology*

248 KTT had clear and trivial effects on oxygen cost and energy cost measures at 200 W that did
249 not reach statistical significance. The effect of KTT on all other physiological parameters
250 was unclear or unlikely, and not statistically significant (**Table 1**).

251

252 *****Insert Table 1*****

253

254 *Kinematics*

255 The clear and likely effects of KTT on ROM values at 100 W (**Table 2**) and 200 W (**Table**
256 **3**) were trivial, except for the mean ankle ROM in the transverse plane at 100 W on the
257 dominant side, where a small non-significant increase was noted (ES, 0.35; *P*, 0.097; **Table**
258 **2**). In all other cases, the effect of KTT was unclear or unlikely, and not statistically
259 significant.

260

261 *****Insert Table 2*****

262

263 *****Insert Table 3*****

264

265 *Electromyography*

266 The effect of KTT on certain VM, VM-to-VL ratio, BF, and RF-to-BF ratio measures were
267 clear, likely, and significant at 100 W (**Table 4**) and 200 W (**Table 5**). Changes primarily
268 affected the efforts performed at 100 W.

269

270 At 100 W, the effect of KTT on the non-dominant side was clear, likely, and significant for
271 increasing VM peak (ES, 1.35; P , 0.044), and decreasing the RF-to-BF ratio peak (ES, -0.42;
272 P , 0.021) and mean (ES, -0.62; P , 0.016) measures. There was also clear and likely non-
273 significant increases in VM iEMG (ES, 0.72; P , 0.128); and VM-to-VL ratio peak (ES, 2.20;
274 P , 0.118), iEMG (ES, 1.26; P , 0.097), and mean (ES, 1.21; P , 0.08) measures.

275

276 At 100 W, the effect of KTT on the dominant side was clear, likely, and significant for
277 increasing VM iEMG (ES, 0.98; P , 0.024) and mean (ES, 0.95; P , 0.030); increasing VM-to-
278 VL ratio peak (ES, 2.19; P , 0.009), iEMG (ES, 1.63; P , 0.020), and mean (ES, 1.21; P ,
279 0.029); and decreasing BF mean (ES, -0.36; P , 0.047) measures. There was also a clear and
280 likely non-significant increase in VM peak (ES, 0.87; P , 0.056); and decrease in RF peak
281 (ES, -0.39; P , 0.135) and mean (ES, -0.51; P , 0.137) measures.

282

283 *****Insert Table 4*****

284

285 At 200 W, the effect of KTT on the non-dominant side was clear, likely, and significant for
286 increasing VM iEMG (ES, 1.04; P , 0.014). There was also a clear and likely non-significant
287 increase in VM peak (ES, 0.92; P , 0.122) and mean (ES, 0.92; P , 0.088); increase in VM-to-
288 VL ratio peak (ES, 1.41; P , 0.157), iEMG (ES, 0.88; P , 0.124), and mean (ES, 2.07; P ,
289 0.098); and decrease in BF mean (ES, -0.39; P , 0.194). At 200 W, there was a clear and
290 likely non-significant effect of KTT on decreasing peak BF (ES, -0.69; P , 0.077) measures.

291

292 *****Insert Table 5*****

293

294 ***Perception***

295 Most cyclists perceived KTT as being comfortable, providing additional stability to the
296 knee, and enhancing performance (**Figure 3**). However, three cyclists felt that KTT was
297 uncomfortable, with one cyclist feeling more unstable with KTT.

298

299 *****Insert Figure 3*****

300

301 **DISCUSSION**

302 Despite most cyclists perceiving enhanced performance and knee stability with patella KTT;
303 the effects of the intervention were either unclear, non-significant, or clearly trivial for all
304 physiological and kinematic measures, except for a small non-significant increase in ankle
305 ROM on the dominant side in the transverse plane at 100 W. KTT affected the EMG-
306 determined muscle activation patterns the most, notably increasing VM and VM-to-VL ratio
307 measures at both powers; and decreasing BF and RF-to-BF ratio measures. Overall, our
308 findings indicate a potential for patella KTT to alter the neuromuscular recruitment patterns
309 of elite cyclists with no current musculoskeletal injury at low powers, which could have
310 implications in the prevention of overuse injuries.

311

312 ***Physiology***

313 Cycling biomechanics and neuromuscular function can alter energy cost, oxygen cost, and
314 cycling efficiency. For instance, cycling in a more aerodynamic than upright position can
315 increase oxygen cost by 1.5% (Gnehm et al., 1997). This increase is speculated to result in
316 part from a shift in mean hip-joint angles towards greater flexion, which alters the operating
317 points of the hip- and knee-joint muscles on the force-velocity and force-length curves, as
318 well as an increase in hip adductor activation to prevent out-of-plane motion in extreme hip
319 flexion. The biomechanical and neuromuscular differences associated with changing cycling
320 positions from aerodynamic to upright are inherently much larger than those potentially
321 resulting from KTT application, especially proximally at the trunk, pelvis, and hip (Dorel et
322 al., 2009). It is likely that the neuromuscular changes observed here in VM, VM-to-VL ratio,
323 BF, and RF-to-BF ratio measures with KTT were not sufficient to cause significant or clear
324 alterations in the physiological parameters monitored.

325

326 ***Kinematics and muscle activation***

327 Ideally, the legs should act as pistons during cycling (Sanner and O'Halloran, 2000), with
328 lower-body motion mainly directed upwards and downwards, and cyclists in a saddle
329 position that allows knee extension with minimal valgus angulation. Most studies addressing
330 lower-body kinematics during cycling have focused on sagittal plane motion, with our
331 sagittal ROM values agreeing with those typically reported (Bini et al., 2011). Although a
332 certain amount of 'out-of-plane' motion is anticipated, lower-body misalignment and
333 excessive out-of-plane motion are reported to contribute to musculoskeletal injuries in
334 cyclists (Bini et al., 2011, Gregor and Wheeler, 1994). One of the proposed benefits of KTT
335 is to assist in joint alignment through improvements in proprioception, which in turn can
336 improve movement patterns and cycling efficiency. Hence, we anticipated less out-of-plane
337 motion at the knee with KTT; however, such a reduction was not evident.

338

339 Previous studies have shown that patellar taping can affect movement patterns in both
340 healthy and symptomatic individuals (Theobald et al., 2014), as well as muscle recruitment
341 of VM (Gilleard et al., 1998), VL (Gilleard et al., 1998), and RF (Konishi, 2013). These
342 changes in neuromuscular function are suggested to result from the tactile stimulation of the
343 skin (Konishi, 2013), rather than by the actual tape configuration or alterations in patellar
344 positioning (Bockrath et al., 1993). Conversely, several other studies have observed no
345 effect from therapeutic taping on neuromuscular function (Halski et al., 2015, Lins et al.,
346 2013), with little evidence supporting improved athletic performance or muscle strength
347 (Csapo and Alegre, 2015, Lins et al., 2013). Our results support the hypothesis that applying
348 KTT across the knee stimulates VM activation and increases the VM-to-VL ratio in
349 asymptomatic elite cyclists during submaximal efforts, without inducing significant or clear
350 changes in knee biomechanics.

351

352 The VM muscle is the dynamic medial stabilizer of the patella and functionally important in
353 aligning the patella within the patella-femoral joint trochlea, which cannot be readily

354 examined using skin-markers and 3D motion capture. Our KTT application had no
355 mechanical intent, and the altered muscle activation seen here most likely resulted from the
356 enhanced tactile input that altered the excitability of the central nervous system and
357 modulated proprioceptive afferent feedback loops (Simoneau et al., 1997). The enhanced
358 VM activation seen in our cyclists when wearing KTT could be beneficial for preventing
359 patellofemoral pain given that VM is important for the dynamic alignment of the patella.
360 Studies have shown that individuals with PFP exhibit lower activity levels of all vastus
361 muscles during walking (Powers et al., 1996) and VM-to-VL ratios across a range of
362 functional and isometric contraction tasks (Souza and Gross, 1991). Furthermore, delayed
363 onset of EMG activity of the VM in relation to VL (-0.67 ms) has been identified as a
364 contributing factor to the development of PFP in one prospective study (Van Tiggelen et al.,
365 2009). That said, prospective studies on this topic in elite cyclists are needed to confirm the
366 prophylactic effect of patella KTT on knee injury occurrence in this population group.

367

368 Although the VM and VL muscles play a critical role in power output during cycling, there
369 is also a high activation of the RF and BF muscles (Akima et al., 2005), with proper co-
370 activation of the hamstrings, which has been suggested to reduce stress at the knee during
371 cycling (So et al., 2005). Hence, reducing the RF-to-BF ratio may have meaningful clinical
372 implications for athletes who exhibit imbalances between knee extensor and flexor strength,
373 poor coordination, and non-optimal activation patterns (i.e., athletes who are quadriceps
374 dominant). With KTT application; there was a clear, likely, and significant decrease in mean
375 BF signals on the dominant side, as well as and peak and mean RF-to-BF ratio values at 100
376 W on the non-dominant side; with only a small non-significant decrease in RF-to-BF mean
377 observed at 200 W. Despite our results indicating some potential for alterations in RF-to-BF
378 muscle activation patterns, larger sample sizes would be needed to confirm outcomes and
379 implications of these changes.

380

381 Most of the neuromuscular effects observed at 100 W became unclear and non-significant at
382 200 W, pointing to an interaction effect between power output and neuromuscular responses
383 to taping. It is well established that muscle contraction forces increase primarily due to an
384 increase in the number of motor units active and associated firing rates in a non-linear
385 fashion (Merletti and Parker, 2004), with previous cycling studies showing progressive
386 increase in muscle activation with progressive loads from ~150, 220, 290, and 370 W
387 (Carpes et al., 2010a). It is plausible that the effect of KTT on the neuromuscular control
388 diminished with increased overall muscle recruitment, explaining the attenuated effects of
389 KTT at 200 W; however, the underlying mechanisms are unclear given the paucity of
390 literature investigating the effect of KTT at different contraction levels and loads within a
391 given exercise.

392

393 The non-dominant and dominant holistically demonstrated comparable responses to KTT,
394 although some clear effects and significant findings were only detected on one side. This
395 discrepancy might be linked to preferred movement patterns of our cyclists, previous injuries
396 with residual neuromuscular inhibition or muscle weakness, or our limited sample size that
397 reduced our statistical power. Most studies suggest that bilateral pedalling asymmetries in
398 terms of power, work, or force increase as the workload decreases (Carpes et al., 2010b),
399 which might explain some of the differential responses between legs that were observed.

400 However, given that work was controlled, and power and force not monitored, the
401 mechanistic reasons behind the between-leg differences remain undetermined.

402

403 ***Perception***

404 Applying RockTape™ to the anterior aspects of the arms and legs and posterior aspects of
405 the neck and back has previously been shown to decrease ‘overall’ and ‘chest’ ratings of
406 perceived exertion of trained cyclists, but not alter ‘arm’ and ‘leg’ ratings of perceived
407 exertions or gross efficiency (Miller et al., 2015). The physiological findings from this same
408 investigation were unable to support improved athletic performance with RockTape™ use

409 (Miller et al., 2015). Although most of our cyclists perceived additional knee stability and
410 enhanced performance with KTT; the biomechanical and physiological findings were unable
411 to support that KTT improved knee stability or economy, with KTT application exerting
412 unclear, non-significant, or clearly trivial effects on knee ROM and physiological measures.
413 It is likely that our cyclists' perceptions result from the EMG changes observed or a placebo
414 effect (Mak et al., 2018). Nonetheless, KTT application may still provide some benefits to
415 certain cyclists given the changes observed in the EMG parameters, notably the increased
416 VM activation and alterations in the VM-to-VL and RF-to-BF activation ratios.

417

418 *Individual responses*

419 One cyclist perceived KTT as uncomfortable and decreasing knee stability. Nonetheless, this
420 particular cyclist felt that KTT improved his performance. This cyclist's data indicated a
421 slight worsening in cycling economy measures at 200 W, with a general increase in knee
422 ROM in all planes of motion with KTT. Simultaneously, EMG signals for VM, VL, and RF,
423 and the VM-to-VL and RF-to-BF ratios increased with KTT, and decreased for BF. In this
424 particular case, perceptions matched well with biomechanical findings, but not necessarily
425 with the physiological ones. In contrast, several cyclists who perceived an increased knee
426 stability, an improved performance, and felt comfortable with KTT application showed
427 'negative' responses, with their perceptual ratings disagreeing with their objective measures.
428 Hence, although individual data suggest the presences of 'positive responders', 'negative
429 responders', and 'non-responders', we were unable to clearly define subgroups from the
430 subjective data collected.

431

432 *Limitations*

433 Small sample sizes are an inherent limitation in any high performance sport environment,
434 which reduced our statistical power. All male National Team cyclists available for testing
435 accepted to participate. Our sample size could not be increased further without
436 compromising the external validity of our findings (i.e., testing lower-level cyclists). Future

research should examine the repeatability of the effect of KTT application and the potential for any long-term effect or habituation to KTT more thoroughly. We tested only elite male cyclists since the national-level female cyclists were training overseas at the time of data collection and thus the findings may be specific to this population. Female athletes differ physiologically, morphologically, and with respect to injury risk factors compared to male athletes, therefore specific investigations of how female cyclists respond to KTT are warranted. We also acknowledge that the power settings selected were submaximal for elite cyclists and that responses at higher powers might differ. Using lower powers was a necessity to calculate steady state oxygen consumption, economy, and efficiency, and for practical relevance to the National Cycling Team of Malaysia. It should be noted however, that during tour events cyclists often perform for prolonged periods at relatively low levels of power production. For example, Alexander Kristoff's average power output during the first hour of Stage 4 of the 2017 Tour de France was 118 W and his average power output over the entire 4:53:54 of the stage (in which he finished second), was 189 W (www.trainingpeaks.com). Finally, given the minimalist taping technique applied, it was not possible to implement a "placebo" taping method or different taping configurations to confer differences in proprioceptive input.

CONCLUSIONS

Most cyclists perceived increased performance and knee stability with patella KTT, but the intervention had little impact on physiological measures and mostly trivial non-significant effects on knee ROM values. However, patella KTT decreased ROM at the pelvis and trunk at the higher power and appeared to stabilize the segments proximally, which could be a favourable adaptation in cyclists (McDaniel et al., 2005). KTT application did alter EMG responses, notably increasing VM activation and altering the VM-to-VL activation ratio at 100 and 200 W, and changes indicating an increase in BF recruitment in relation to RF at 100 W. Our findings imply that there is a potential for patella KTT to alter neuromuscular recruitment patterns in elite uninjured cyclists, which could have implications for injury

465 prevention and especially the development of PFP by assisting with patella alignment and
466 alleviating knee-joint stress through neuromuscular pathways as opposed to altering knee
467 biomechanics. As such, the neuromuscular changes we observed indicate that cyclists may
468 benefit acutely from patella KTT, although the longitudinal effects of KTT use have not yet
469 been established.

470

471 **ACKNOWLEDGEMENTS**

472 The authors would like to thank Dr. Yeo Wee Kian and Mohd Izham Bin Mohamad for
473 assistance in coordinating the project, Mee Chee Chong for helping during data collection,
474 and all cyclists for their voluntary participation.

475

476

477 **REFERENCES**

- 478 Akima H, Kinugasa R, Kuno S. Recruitment of the thigh muscles during sprint cycling by
479 muscle functional magnetic resonance imaging. *Int J Sports Med*. 2005;26:245-52.
- 480 Bell AL, Brand RA, Pedersen DR. Prediction of hip joint centre location from external
481 landmarks. *Hum Mov Sci*. 1989;8:3-16.
- 482 Bertucci W, Grappe F, Gros Lambert A. Laboratory versus outdoor cycling conditions:
483 Differences in pedaling biomechanics. *J Appl Biomech*. 2007;23:87-92.
- 484 Bertucci WM, Betik AC, Duc S, Grappe F. Gross efficiency and cycling economy are higher
485 in the field as compared with on an Axiom stationary ergometer. *J Appl Biomech*.
486 2012;28:636-44.
- 487 Bini R, Hume PA, Croft JL. Effects of bicycle saddle height on knee injury risk and cycling
488 performance. *Sports Med*. 2011;41:463-76.
- 489 Bockrath K, Wooden C, Worrell T, Ingersoll CD, Farr J. Effects of patella taping on patella
490 position and perceived pain. *Med Sci Sports Exerc*. 1993;25:989-92.
- 491 Cappozzo A, Cappello A, Croce UD, Pensalfini F. Surface-marker cluster design criteria for
492 3-d bone movement reconstruction. *IEEE Trans Biomed Eng*. 1997;44:1165-74.

493 Carpes FP, Diefenthaler F, Bini RR, Stefanyshyn DJ, Faria IE, Mota CB. Influence of leg
 494 preference on bilateral muscle activation during cycling. *J Sports Sci.* 2010a;29:151-9.

495 Carpes FP, Mota CB, Faria IE. On the bilateral asymmetry during running and cycling - A
 496 review considering leg preference. *Phys Ther Sport.* 2010b;11:136-42.

497 Coyle EF, Sidossis LS, Horowitz JF, Beltz JD. Cycling efficiency is related to the
 498 percentage of type I muscle fibers. *Med Sci Sports Exerc.* 1992;24:782-8.

499 Csapo R, Alegre LM. Effects of Kinesio® taping on skeletal muscle strength—A meta-
 500 analysis of current evidence. *J Sci Med Sport.* 2015;18:450-6.

501 Dorel S, Couturier A, Hug F. Influence of different racing positions on mechanical and
 502 electromyographic patterns during pedalling. *Scand J Med Sci Sports.* 2009;19:44-54.

503 Gilleard W, McConnell J, Parsons D. The effect of patellar taping on the onset of vastus
 504 medialis obliquus and vastus lateralis muscle activity in persons with patellofemoral pain.
 505 *Phys Ther.* 1998;78:25-32.

506 Gnehm P, Reichenbach S, Altpeter E, Widmer H, Hoppeler H. Influence of different racing
 507 positions on metabolic cost in elite cyclists. *Med Sci Sports Exerc.* 1997;29:818-23.

508 Gregor RJ, Wheeler JB. Biomechanical factors associated with shoe/pedal interfaces.
 509 Implications for injury. *Sports Med.* 1994;17:117-31.

510 Grood ES, Suntay WJ. A joint coordinate system for the clinical description of three-
 511 dimensional motions: Application to the knee. *J Biomech Eng.* 1983;105:136-44.

512 Halski T, Dymarek R, Ptazkowski K, Słupska L, Rajfur K, Rajfur J, et al. Kinesiology
 513 Taping does not modify electromyographic activity or muscle flexibility of quadriceps
 514 femoris muscle: A randomized, placebo-controlled pilot study in healthy volleyball players.
 515 *Med Sci Monit.* 2015;21:2232-9.

516 Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G. Development of recommendations for
 517 SEMG sensors and sensor placement procedures. *J Electromyogr Kinesiol.* 2000;10:361-74.

518 Jeukendrup AE, Wallis GA. Measurement of substrate oxidation during exercise by means
 519 of gas exchange measurements. *Int J Sports Med.* 2005;26:S28-37.

520 Konishi Y. Tactile stimulation with Kinesiology tape alleviates muscle weakness attributable
 521 to attenuation of Ia afferents. *J Sci Med Sport*. 2013;16:45-8.
 522 Likert R. A technique for the measurement of attitudes. *Arch Psychol*. 1932;22:5-55.
 523 Lins CADA, Neto FL, de Amorim ABC, Macedo LdB, Brasileiro JS. Kinesio Taping® does
 524 not alter neuromuscular performance of femoral quadriceps or lower limb function in
 525 healthy subjects: Randomized, blind, controlled, clinical trial. *Man Ther*. 2013;18:41-5.
 526 Lyman KJ, Keister K, Gange K, Mellinger CD, Hanson TA. Investigating the effectiveness
 527 of Kinesio(R) taping space correction method in healthy adults on patellofemoral joint and
 528 subcutaneous space. *Int J Sports Phys Ther*. 2017;12:250-7.
 529 Mak DN, Au IP, Chan M, Chan ZY, An WW, Zhang JH, et al. Placebo effect of facilitatory
 530 Kinesio tape on muscle activity and muscle strength. *Physiother Theory Pract*. 2018:1-6. doi:
 531 10.1080/09593985.2018.1441936
 532 Matheus JPC, Zille RR, Gomide Matheus LB, Lemos TV, Carregaro RL, Shimano AC.
 533 Comparison of the mechanical properties of therapeutic elastic tapes used in sports and
 534 clinical practice. *Phys Ther Sport*. 2017;24:74-8.
 535 McDaniel J, Subudhi A, Martin JC. Torso stabilization reduces the metabolic cost of
 536 producing cycling power. *Can J Appl Physiol*. 2005;30:433-41.
 537 Merino-Marban R, Mayorga-Vega D, Fernandez-Rodriguez E. Effect of kinesio tape
 538 application on calf pain and ankle range of motion in duathletes. *J Hum Kinet*. 2013;37:129-
 539 35.
 540 Merletti R, di Torino P. Standards for reporting EMG data. *J Electromyogr Kinesiol*.
 541 1999;9:III–IV.
 542 Merletti R, Parker P. *Electromyography: Physiology, Engineering, and Noninvasive*
 543 *Applications*. Hoboken, NJ: IEEE/John Wiley & Sons; 2004.
 544 Miller MG, Michael TJ, Nicholson KS, Petro RV, Hanson NJ, Prater DR. The effect of
 545 Rocktape on rating of perceived exertion and cycling efficiency. *J Strength Cond Res*.
 546 2015;29:2608-12.

547 Ouyang JH, Chang KH, Hsu WY, Cho YT, Liou TH, Lin YN. Non-elastic taping, but not
 548 elastic taping, provides benefits for patients with knee osteoarthritis: Systemic review and
 549 meta-analysis. *Clin Rehabil.* 2018;32:3-17.
 550 Powers CM, Landel R, Perry J. Timing and Intensity of Vastus Muscle Activity During
 551 Functional Activities in Subjects With and Without Patellofemoral Pain. *Phys Ther.*
 552 1996;76:946-55.
 553 Reneker JC, Latham L, McGlawn R, Reneker MR. Effectiveness of kinesiology tape on
 554 sports performance abilities in athletes: A systematic review. *Phys Ther Sport.* 2018;31:83-
 555 98.
 556 Sanner W, O'Halloran W. The biomechanics, etiology, and treatment of cycling injuries. *J*
 557 *Am Podiatr Med Assoc.* 2000;90:354-76.
 558 Silberman MR. Bicycling injuries. *Curr Sports Med Rep.* 2013;12:337-45.
 559 Simoneau GG, Degner RM, Kramper CA, Kittleson KH. Changes in ankle joint
 560 proprioception resulting from strips of athletic tape applied over the skin. *J Athl Train.*
 561 1997;32:141.
 562 Smith TB, Hopkins WG. Variability and predictability of finals times of elite rowers. *Med*
 563 *Sci Sports Exerc.* 2011;43:2155-60.
 564 So RCH, Ng JKF, Ng GYF. Muscle recruitment pattern in cycling: A review. *Phys Ther*
 565 *Sport.* 2005;6:89-96.
 566 Souza DR, Gross MT. Comparison of vastus medialis obliquus: vastus lateralis muscle
 567 integrated electromyographic ratios between healthy subjects and patients with
 568 patellofemoral pain. *Phys Ther.* 1991;71:310-6.
 569 Theobald G, Parry S, Richards J, Thewlis D, Tootill I, Selfe J. Biomechanical effects of
 570 different treatment modalities used in knee pain during cycling. *Physiother Pract Res.*
 571 2012;33:16-21.
 572 Theobald G, Selfe J, Richards J, Roddam H. An investigation of the biomechanical efficacy
 573 and clinical effectiveness of patello-femoral taping in elite and experienced cyclists. *J Sci*
 574 *Cycling.* 2014;3:57-63.

575 Van Tiggelen D, Cowan S, Coorevits P, Duvigneaud N, Witvrouw E. Delayed vastus
576 medialis obliquus to vastus lateralis onset timing contributes to the development of
577 patellofemoral pain in previously healthy men: A prospective study. *Am J Sports Med.*
578 2009;37:1099-105.

579 Williams S, Whatman C, Hume PA, Sheerin K. Kinesio taping in treatment and prevention
580 of sports injuries. *Sports Med.* 2012;42:153-64.

581 Zech A, Wellmann K. Perceptions of football players regarding injury risk factors and
582 prevention strategies. *PLoS One.* 2017;12:e0176829.

Table 1. Mean \pm SD values for oxygen cost (mL/kg/km), energy cost (kcal/kg/km), cycling economy (W/L/min), heart rate (bpm), and gross efficiency (%) in the No Tape (NT) and Kinesiology-Type Tape (KTT) conditions during the 100 and 200 W cycling efforts. Delta efficiency (%) in NT and KTT conditions is also presented. Differences between conditions are expressed using mean change (Δ_{mean}); standardized effect size (ES); and paired *t*-test statistical significance values (*P*). Thresholds for clear ES are provided (trivial, small, large, and very large) and significant changes ($P \leq 0.05$) are highlighted in grey.

Parameters	NT	KTT	Δ_{mean}	ES (threshold)	<i>P</i>
100 W					
Oxygen cost	175.4 \pm 30.6	175.2 \pm 33.5	-0.2 \pm 14.1	-0.01 \pm 0.42 (unclear)	0.963
Energy cost	0.86 \pm 0.15	0.86 \pm 0.16	0.001 \pm 0.07	0.001 \pm 0.43 (unclear)	0.993
Economy	53.4 \pm 5.6	53.6 \pm 5.5	0.2 \pm 4.5	0.03 \pm 0.78 (unclear)	0.885
Heart rate	115.3 \pm 9.8	115.6 \pm 9.5	0.3 \pm 5.8	0.03 \pm 0.58 (unclear)	0.845
Gross efficiency	15.5 \pm 1.6	15.6 \pm 1.6	0.04 \pm 1.31	0.02 \pm 0.79 (unclear)	0.926
200 W					
Oxygen cost	277.3 \pm 48.0	273.4 \pm 46.1	-3.9 \pm 12.6	-0.08 \pm 0.26 (trivial) [§]	0.311
Energy cost	1.37 \pm 0.23	1.35 \pm 0.22	-0.02 \pm 0.06	-0.09 \pm 0.26 (trivial) [§]	0.263
Economy	67.4 \pm 5.2	68.4 \pm 5.9	1.0 \pm 3.1	0.19 \pm 0.54 (trivial)	0.303
Heart rate	147.6 \pm 9.2	146.8 \pm 8.4	-0.8 \pm 6.4	-0.08 \pm 0.70 (unclear)	0.691
Gross efficiency	19.5 \pm 1.5	19.8 \pm 1.7	0.3 \pm 0.9	0.21 \pm 0.54 (small)	0.260
Delta efficiency	26.5 \pm 3.1	27.3 \pm 2.3	0.8 \pm 3.0	0.25 \pm 1.04 (unclear)	0.380

Note. An effect was deemed ‘unclear’ when its 90% confidence limit overlapped the thresholds for small positive and small negative effects (i.e., 5%). [§]Probability of the effect exceeds 75% and is ‘likely’ to be clinically meaningful.

Table 2. Mean \pm SD values for range of motion values ($^{\circ}$) in the sagittal (X), coronal (Y), and transverse (Z) planes in the No Tape (NT) and Kinesiology-Type Tape (KTT) conditions during the 100 W cycling efforts for the dominant (D) and non-dominant (ND) sides. Differences between conditions are expressed using mean change (Δ_{mean}); standardized effect size (ES); and paired t -test statistical significance values (P). Thresholds for clear ES are provided (trivial, small, large, and very large) and significant changes ($P \leq 0.05$) are highlighted in grey.

Joint	Side	Plane	NT	KTT	Δ_{mean}	ES	P
Ankle	ND	X	19.5 \pm 8.1	19.9 \pm 8.7	0.4 \pm 2.1	0.05 (trivial) [§]	0.517
		Y	5.3 \pm 1.7	5.5 \pm 1.9	0.2 \pm 0.6	0.11 (trivial) [§]	0.289
		Z	5.7 \pm 2.3	5.7 \pm 1.9	0.0 \pm 1.5	0.02 (unclear)	0.920
	D	X	21.4 \pm 4.9	20.3 \pm 6.0	-1.1 \pm 3.9	-0.22 (small)	0.355
		Y	4.3 \pm 1.8	5.0 \pm 1.7	0.6 \pm 1.2	0.35 (small) [§]	0.097
		Z	5.5 \pm 0.7	5.2 \pm 1.0	-0.3 \pm 1.1	-0.44 (unclear)	0.350
Knee	ND	X	76.7 \pm 3.1	76.2 \pm 3.6	-0.5 \pm 1.5	-0.16 (trivial)	0.270
		Y	9.2 \pm 3.6	9.0 \pm 3.3	-0.2 \pm 1.5	-0.04 (trivial) [§]	0.727
		Z	9.2 \pm 4.9	9.3 \pm 3.9	0.1 \pm 2.6	0.03 (unclear)	0.867
	D	X	78.8 \pm 2.3	78.2 \pm 2.2	-0.5 \pm 1.7	-0.23 (small)	0.304
		Y	8.3 \pm 2.6	8.1 \pm 2.3	-0.2 \pm 1.1	-0.08 (trivial) [§]	0.526
		Z	12.3 \pm 5.3	12.4 \pm 4.7	0.0 \pm 1.8	0.003 (trivial) [§]	0.976
Hip	ND	X	48.7 \pm 3.5	48.8 \pm 3.8	0.1 \pm 1.5	0.02 (unclear)	0.901
		Y	6.3 \pm 3.4	6.2 \pm 2.8	0.0 \pm 1.3	-0.01 (trivial) [§]	0.915
		Z	11.5 \pm 3.8	11.4 \pm 4.3	0.0 \pm 1.8	-0.01 (unclear)	0.937
	D	X	47.5 \pm 2.9	47.4 \pm 2.8	-0.1 \pm 1.1	-0.04 (trivial) [§]	0.694
		Y	6.6 \pm 1.8	6.6 \pm 1.9	0.0 \pm 1.4	-0.005 (unclear)	0.983
		Z	10.0 \pm 3.7	9.5 \pm 3.2	-0.5 \pm 1.4	-0.12 (trivial)	0.283
Pelvis		X	8.1 \pm 1.8	8.3 \pm 2.2	0.2 \pm 1.4	0.10 (unclear)	0.647
		Y	3.5 \pm 0.9	3.6 \pm 1.1	0.1 \pm 0.5	0.09 (trivial)	0.561
		Z	3.7 \pm 1.9	3.6 \pm 1.3	-0.1 \pm 2.2	-0.05 (unclear)	0.881
Trunk		X	8.8 \pm 3.9	10.1 \pm 8.2	1.3 \pm 8.9	0.34 (unclear)	0.620
		Y	0.8 \pm 0.2	0.7 \pm 0.2	0.0 \pm 0.1	-0.07 (trivial) [§]	0.643
		Z	8.4 \pm 4.1	9.6 \pm 8.5	1.2 \pm 9.4	0.30 (unclear)	0.663

Notes. Sagittal (X): ankle dorsiflexion and plantar flexion, knee and hip flexion and extension, pelvis and trunk anterior and posterior tilt; Coronal (Y): ankle inversion and eversion, knee valgus and varus, hip abduction and adduction, pelvis and trunk non-dominant side and dominant side tilt; Transverse (Z) ankle, knee, and hip internal and external rotation, pelvis and trunk non-dominant side and dominant side rotation. An effect was deemed ‘unclear’ when its 90% confidence limit overlapped the thresholds for small positive and small negative effects (i.e., 5%). [§]Probability of the effect exceeds 75% and is ‘likely’ to be clinically meaningful.

Table 3. Mean \pm SD values for range of motion values ($^{\circ}$) in the sagittal (X), coronal (Y), and transverse (Z) planes in the No Tape (NT) and Kinesiology-Type Tape (KTT) conditions during the 200 W cycling efforts for the dominant (D) and non-dominant (ND) sides. Differences between conditions are expressed using mean change (Δ_{mean}); standardized effect size (ES); and paired *t*-test statistical significance values (*P*). Thresholds for clear ES are provided (trivial, small, large, and very large) and significant changes ($P \leq 0.05$) are highlighted in grey.

Joint	Side	Plane	NT	KTT	Δ_{mean}	ES	<i>P</i>
Ankle	ND	X	21.7 \pm 7.8	22.7 \pm 8.3	1.0 \pm 2.6	0.12 (trivial) [§]	0.235
		Y	5.9 \pm 2.0	6.2 \pm 2.6	0.2 \pm 1.0	0.11 (trivial)	0.450
		Z	5.9 \pm 1.8	5.7 \pm 1.9	-0.2 \pm 1.0	-0.11 (trivial)	0.529
	D	X	24.0 \pm 6.9	24.7 \pm 5.7	0.7 \pm 3.0	0.10 (trivial) [§]	0.435
		Y	4.7 \pm 1.3	4.9 \pm 1.5	0.2 \pm 0.6	0.14 (trivial)	0.342
		Z	6.0 \pm 1.6	5.8 \pm 1.7	-0.2 \pm 0.4	-0.14 (trivial) [§]	0.123
Knee	ND	X	78.4 \pm 3.7	78.9 \pm 3.1	0.5 \pm 1.7	0.14 (trivial)	0.302
		Y	8.3 \pm 2.6	8.2 \pm 2.6	-0.7 \pm 1.4	-0.03 (unclear)	0.849
		Z	9.4 \pm 4.1	8.8 \pm 3.1	-0.6 \pm 1.8	-0.14 (trivial)	0.277
	D	X	80.1 \pm 2.3	80.9 \pm 2.5	0.8 \pm 2.1	0.34 (small)	0.224
		Y	7.7 \pm 2.1	8.1 \pm 2.1	0.4 \pm 0.9	0.19 (trivial)	0.146
		Z	11.5 \pm 4.5	11.2 \pm 4.5	-0.3 \pm 1.6	-0.07 (trivial) [§]	0.496
Hip	ND	X	49.6 \pm 3.9	49.3 \pm 4.5	-0.9 \pm 2.1	-0.07 (unclear)	0.661
		Y	7.4 \pm 3.7	7.4 \pm 3.4	0.0 \pm 1.0	0.001 (trivial) [§]	0.988
		Z	11.5 \pm 3.4	12.1 \pm 3.4	0.6 \pm 1.7	0.18 (trivial)	0.245
	D	X	47.4 \pm 2.8	47.5 \pm 2.4	0.1 \pm 1.4	0.02 (unclear)	0.874
		Y	7.1 \pm 2.5	6.9 \pm 2.2	-0.3 \pm 1.2	-0.10 (trivial)	0.490
		Z	10.3 \pm 3.1	10.4 \pm 3.2	0.1 \pm 1.5	0.03 (unclear)	0.812
Pelvis		X	8.4 \pm 2.4	7.9 \pm 2.2	-0.5 \pm 1.1	-0.19 (trivial)	0.178
		Y	3.5 \pm 1.3	3.7 \pm 1.5	0.2 \pm 0.5	0.14 (trivial)	0.269
		Z	4.0 \pm 1.8	3.6 \pm 1.1	-0.4 \pm 1.8	-0.22 (unclear)	0.447
Trunk		X	9.5 \pm 4.2	8.6 \pm 2.7	-0.9 \pm 3.4	-0.21 (unclear)	0.391
		Y	0.9 \pm 0.5	0.9 \pm 0.3	0.0 \pm 0.3	-0.04 (unclear)	0.834
		Z	8.6 \pm 3.8	7.8 \pm 2.6	-0.9 \pm 3.6	-0.22 (unclear)	0.430

Notes. Sagittal (X): ankle dorsiflexion and plantar flexion, knee and hip flexion and extension, pelvis and trunk anterior and posterior tilt; Coronal (Y): ankle inversion and eversion, knee valgus and varus, hip abduction and adduction, pelvis and trunk non-dominant side and dominant side tilt; Transverse (Z) ankle, knee, and hip internal and external rotation, pelvis and trunk non-dominant side and dominant side rotation. An effect was deemed ‘unclear’ when its 90% confidence limit overlapped the thresholds for small positive and small negative effects (i.e., 5%). [§]Probability of the effect exceeds 75% and is ‘likely’ to be clinically meaningful.

Table 4. Mean (%_{max}), peak (%_{max}), and integrated EMG (iEMG, %_{max}) signal values (mean ± SD) for the *vastus medialis* (VM), *vastus lateralis* (VL), *rectus femoris* (RF), and *biceps femoris* (BF) muscles in the No Tape (NT) and Kinesiology-Type Tape (KTT) conditions during the 100 W cycling efforts for the dominant (D) and non-dominant (ND) sides. The VM-to-VL and RF-to-BF activation ratios are also presented. Differences between conditions are expressed using mean change (Δ_{mean}); standardized effect size (ES); and paired *t*-test statistical significance values (*P*). Thresholds for clear ES are provided (trivial, small, large, and very large) and significant changes ($P \leq 0.05$) are highlighted in grey.

Muscle	Side	EMG	NT	KTT	Δ_{mean}	ES (threshold)	<i>P</i>
VM	ND	Peak	29.5 ± 5.9	37.4 ± 5.1	8.0 ± 7.4	1.35 (large) [§]	0.047
		iEMG	39.9 ± 10.1	47.3 ± 7.9	7.3 ± 9.9	0.72 (moderate) [§]	0.128
		Mean	9.4 ± 1.7	9.2 ± 2.6	-0.2 ± 2.6	-0.13 (unclear)	0.840
	D	Peak	31.4 ± 9.3	39.5 ± 8.0	8.1 ± 10.1	0.87 (moderate) [§]	0.056
		iEMG	38.1 ± 11.9	49.7 ± 16.0	11.7 ± 8.9	0.98 (moderate) [§]	0.024
		Mean	7.9 ± 2.1	9.9 ± 2.2	2.0 ± 2.1	0.95 (moderate) [§]	0.030
VL	ND	Peak	39.3 ± 11.0	36.8 ± 8.0	-2.5 ± 15.8	-0.23 (unclear)	0.717
		iEMG	47.3 ± 11.0	36.8 ± 8.0	-0.5 ± 18.0	-0.05 (unclear)	0.950
		Mean	9.4 ± 1.7	9.2 ± 2.6	-0.2 ± 2.6	-0.13 (unclear)	0.840
	D	Peak	37.0 ± 5.4	36.9 ± 8.2	-0.2 ± 10.0	-0.03 (unclear)	0.965
		iEMG	44.7 ± 10.1	47.2 ± 12.4	2.5 ± 10.2	0.25 (unclear)	0.545
		Mean	8.7 ± 1.8	9.0 ± 2.4	0.2 ± 1.8	0.13 (unclear)	0.725
RF	ND	Peak	24.7 ± 4.5	22.7 ± 2.5	-2.0 ± 5.7	-0.44 (unclear)	0.482
		iEMG	35.1 ± 6.8	33.0 ± 4.4	-2.1 ± 4.8	-0.30 (unclear)	0.391
		Mean	7.5 ± 1.0	7.0 ± 0.6	-0.6 ± 1.0	-0.54 (unclear)	0.279
	D	Peak	29.6 ± 11.0	25.2 ± 8.7	-4.3 ± 4.3	-0.39 (small) [§]	0.135
		iEMG	39.0 ± 10.0	36.1 ± 5.8	-3.0 ± 4.6	-0.30 (unclear)	0.287
		Mean	7.3 ± 1.3	6.7 ± 0.7	-0.7 ± 0.6	-0.51 (small) [§]	0.137
BF	ND	Peak	37.2 ± 11.9	39.2 ± 8.6	1.9 ± 6.0	0.16 (unclear)	0.514
		iEMG	42.8 ± 11.7	42.8 ± 6.7	-0.9 ± 9.3	-0.08 (unclear)	0.830
		Mean	7.9 ± 2.0	8.4 ± 1.5	0.5 ± 1.1	0.25 (unclear)	0.351
	D	Peak	42.0 ± 5.2	38.5 ± 3.3	-3.5 ± 3.2	-0.67 (moderate) [§]	0.120
		iEMG	42.7 ± 5.1	40.7 ± 4.7	-2.1 ± 3.3	-0.41 (unclear)	0.303
		Mean	7.6 ± 1.3	7.4 ± 1.4	-0.6 ± 0.3	-0.36 (small) [§]	0.044
VM:VL	ND	Peak	81.6 ± 8.0	99.2 ± 12.2	17.6 ± 19.8	2.20 (very large) [§]	0.118
		iEMG	82.2 ± 16.0	102.4 ± 18.3	20.1 ± 24.2	1.26 (large) [§]	0.097
		Mean	76.8 ± 11.0	95.4 ± 14.2	18.6 ± 22.8	0.95 (moderate) [§]	0.088
	D	Peak	75.1 ± 14.7	107.2 ± 29.2	32.1 ± 18.8	2.19 (large) [§]	0.009
		iEMG	84.9 ± 14.0	107.7 ± 26.0	22.8 ± 16.6	1.63 (large) [§]	0.020
		Mean	91.9 ± 19.6	115.6 ± 27.7	23.7 ± 23.4	1.21 (large) [§]	0.029
RF:BF	ND	Peak	68.0 ± 21.8	58.8 ± 18.7	-9.1 ± 6.5	-0.42 (small) [§]	0.021
		iEMG	85.6 ± 19.8	80.7 ± 16.0	-5.0 ± 13.1	-0.25 (unclear)	0.445
		Mean	99.5 ± 22.8	85.5 ± 17.2	-14.0 ± 7.8	-0.62 (moderate) [§]	0.016
	D	Peak	69.5 ± 18.8	65.4 ± 20.2	-4.1 ± 8.1	-0.22 (unclear)	0.388
		iEMG	91.6 ± 23.6	89.5 ± 18.2	-2.1 ± 5.7	-0.09 (trivial)	0.514
		Mean	93.9 ± 23.6	91.6 ± 18.2	-2.3 ± 5.5	-0.10 (trivial)	0.456

Note. An effect was deemed ‘unclear’ when its 90% confidence limit overlapped the thresholds for small positive and small negative effects (i.e., 5%). An effect was deemed ‘unclear’ when its 90% confidence limit overlapped the thresholds for small positive and

635 small negative effects (i.e., 5%). §Probability of the effect exceeds 75% and is ‘likely’ to be
636 clinically meaningful.

Table 5. Mean (%_{max}), peak (%_{max}), and integrated EMG (iEMG, %_{max}) signal values (mean ± SD) for the *vastus medialis* (VM), *vastus lateralis* (VL), *rectus femoris* (RF), and *biceps femoris* (BF) muscles in the No Tape (NT) and Kinesiology-Type Tape (KTT) conditions during the 200 W cycling efforts for the dominant (D) and non-dominant (ND) sides. The VM-to-VL and RF-to-BF activation ratios are also presented. Differences between conditions are expressed using mean change (Δ_{mean}); standardized effect size (ES); and paired *t*-test statistical significance values (*P*). Thresholds for clear ES are provided (trivial, small, large, and very large) and significant changes ($P \leq 0.05$) are highlighted in grey.

Muscle	Side	EMG	NT	KTT	Δ_{mean}	ES (threshold)	<i>P</i>
VM	ND	Peak	39.4 ± 9.4	48.0 ± 4.4	8.6 ± 12.7	0.92 (moderate) [§]	0.122
		iEMG	58.2 ± 9.6	68.1 ± 5.7	9.9 ± 7.7	1.04 (moderate) [§]	0.014
		Mean	10.7 ± 1.5	12.1 ± 1.8	1.4 ± 1.8	0.92 (moderate) [§]	0.088
	D	Peak	47.0 ± 9.4	48.2 ± 5.3	1.2 ± 10.9	0.13 (unclear)	0.741
		iEMG	69.1 ± 10.7	70.1 ± 6.2	1.0 ± 11.4	0.09 (unclear)	0.830
		Mean	13.0 ± 2.4	14.1 ± 1.7	1.1 ± 3.0	0.46 (unclear)	0.380
VL	ND	Peak	46.4 ± 9.5	47.7 ± 6.2	1.3 ± 12.4	0.13 (unclear)	0.797
		iEMG	66.4 ± 6.9	68.9 ± 11.5	2.5 ± 13.7	0.37 (unclear)	0.643
		Mean	12.8 ± 1.8	12.8 ± 3.2	0.1 ± 2.2	0.05 (unclear)	0.924
	D	Peak	50.2 ± 7.0	47.7 ± 9.0	-2.6 ± 9.3	-0.37 (unclear)	0.433
		iEMG	65.8 ± 8.1	69.4 ± 9.0	3.6 ± 9.4	0.45 (unclear)	0.284
		Mean	13.0 ± 1.6	13.1 ± 2.1	0.1 ± 1.7	0.09 (unclear)	0.831
RF	ND	Peak	34.4 ± 12.6	37.1 ± 2.5	2.8 ± 14.5	0.22 (unclear)	0.657
		iEMG	54.4 ± 15.9	53.7 ± 6.7	-0.7 ± 15.1	-0.04 (unclear)	0.906
		Mean	11.9 ± 3.0	11.9 ± 1.7	-0.1 ± 3.4	-0.02 (unclear)	0.971
	D	Peak	39.5 ± 10.1	40.8 ± 6.6	1.3 ± 7.2	0.13 (unclear)	0.672
		iEMG	62.1 ± 11.9	63.7 ± 7.8	1.6 ± 13.6	0.14 (unclear)	0.761
		Mean	11.6 ± 1.7	11.4 ± 1.8	-0.2 ± 2.2	-0.11 (unclear)	0.844
BF	ND	Peak	42.6 ± 11.6	45.1 ± 6.4	2.6 ± 16.0	0.22 (unclear)	0.711
		iEMG	57.8 ± 12.9	58.8 ± 8.7	1.0 ± 12.9	0.08 (unclear)	0.846
		Mean	12.7 ± 4.8	12.2 ± 3.0	-0.5 ± 3.7	-0.11 (unclear)	0.735
	D	Peak	44.1 ± 8.1	38.5 ± 5.9	-5.6 ± 6.2	-0.69 (moderate) [§]	0.077
		iEMG	52.8 ± 13.1	52.8 ± 6.8	0.0 ± 13.4	-0.01 (unclear)	0.999
		Mean	10.7 ± 3.1	9.6 ± 2.1	-1.0 ± 2.4	-0.34 (unclear)	0.338
VM:VL	ND	Peak	85.6 ± 8.5	97.6 ± 15.5	11.9 ± 17.6	1.41 (large) [§]	0.157
		iEMG	87.8 ± 12.8	101.5 ± 20.9	13.7 ± 20.3	0.88 (moderate) [§]	0.124
		Mean	84.6 ± 6.8	98.7 ± 21.5	14.2 ± 19.2	2.07 (very large) [§]	0.098
	D	Peak	94.8 ± 19.3	98.5 ± 19.8	3.8 ± 27.2	0.20 (unclear)	0.707
		iEMG	96.6 ± 20.9	104.4 ± 18.3	7.8 ± 23.1	0.37 (unclear)	0.340
		Mean	100.2 ± 13.1	110.8 ± 27.9	10.6 ± 25.2	0.80 (unclear)	0.311
RF:BF	ND	Peak	82.6 ± 24.3	83.3 ± 9.7	0.7 ± 30.6	0.03 (unclear)	0.955
		iEMG	98.3 ± 34.4	92.9 ± 18.2	-5.3 ± 27.1	-0.16 (unclear)	0.620
		Mean	120.1 ± 34.2	106.7 ± 16.2	-13.4 ± 19.2	-0.39 (small) [§]	0.194
	D	Peak	92.5 ± 27.2	106.8 ± 17.0	14.3 ± 24.7	0.53 (unclear)	0.214
		iEMG	124.1 ± 36.9	121.5 ± 15.1	-2.7 ± 40.3	-0.07 (unclear)	0.867
		Mean	118.3 ± 43.9	121.8 ± 27.7	3.6 ± 37.6	0.09 (unclear)	0.825

Note. An effect was deemed ‘unclear’ when its 90% confidence limit overlapped the thresholds for small positive and small negative effects (i.e., 5%). [§]Probability of the effect exceeds 75% and is ‘likely’ to be clinically meaningful.

648 **Figure captions**

649

650 **Figure 1.** Cyclist set-up for data collection with the patella kinesiology-type tape (KTT)
651 applied.

652

653 **Figure 2.** Marker placement for 3D motion capture from anterior (left), posterior (middle),
654 and lateral (right) views. Anatomical reference markers were placed bilaterally on the
655 acromial processes, anterior superior iliac spines, posterior superior iliac spines, greater
656 trochanters, medial and lateral femoral epicondyles, medial and lateral malleoli, and 1st and
657 5th metatarsal heads. Tracking markers were placed bilaterally on the heel, mid-foot, and
658 forefoot, and 4-marker rigid clusters were placed on the lateral aspect of the pelvis and
659 bilaterally on the lateral aspects of the thighs and shanks. Anterior superior iliac spine,
660 greater trochanter, femoral epicondyle, malleolus, and 1st metatarsal head markers were
661 removed before the dynamic cycling efforts (red circles).

662

663 **Figure 3.** Ratings of comfort levels and perceived change in knee stability and cycling
664 performance with the application of kinesiology-type tape (KTT) compared to no tape (NT)
665 on a 5-point Likert scale. Data presented are the number of cyclists (*n*) that provided a given
666 rating. Comfort level: 1, very uncomfortable; 2, uncomfortable; 3, no change; 4,
667 comfortable; 5, very comfortable. Knee stability: 1, much less stable; 2, less stable; 3, no
668 change; 4, more stable; 5, much more stable. Performance: 1, much worse; 2, worse; 3, no
669 change; 4, better; 5, much better.