



## Article

# Need for early, minimally invasive cancer diagnosis

Paraskevaidi, Maria, Martin-Hirsch, Pierre L and Martin, Francis L

Available at <http://clock.uclan.ac.uk/26416/>

*Paraskevaidi, Maria, Martin-Hirsch, Pierre L and Martin, Francis L ORCID: 0000-0001-8562-4944 (2019) Need for early, minimally invasive cancer diagnosis. Proceedings of the National Academy of Sciences of the United States of America . ISSN 0027-8424*

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1073/pnas.1817551116>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

## **Need for early, minimally-invasive cancer diagnosis**

Maria Paraskevasidi<sup>a,1</sup>, Pierre L. Martin-Hirsch<sup>b</sup> and Francis L. Martin<sup>a,c</sup>

*<sup>a</sup>School of Pharmacy and Biomedical Sciences, University of Central Lancashire, Preston PR1 2HE, UK; <sup>b</sup>Department of Obstetrics and Gynaecology, Lancashire Teaching Hospitals NHS Foundation, Preston PR2 9HT, UK; <sup>c</sup>Biocel Ltd, Hull, Yorkshire, UK*

<sup>1</sup>To whom correspondence should be addressed. Email: [mparaskevasidi@uclan.ac.uk](mailto:mparaskevasidi@uclan.ac.uk)

Dear Editor,

We have read the article by Mittal et al. (1) and we would like to congratulate the authors for an excellent piece of work demonstrating spectroscopy's capability for accurately subtyping epithelial cells and the tumor-associated microenvironment. Herein we would like to add some insights and comment on how this work could take the field forward.

In this recent paper, a high-definition Fourier-transform infrared imaging method using a quantum cascade laser (QCL) source has been employed for the characterization of 100-case breast tissue samples. This advancement has the potential to facilitate label-free molecular histopathology and bring spectroscopy one step closer to the clinic: undoubtedly, this study would be of great benefit with regards to expediting the diagnostic workflow as well as in terms of saving the pathologist's and / or surgeon's precious time and caseloads - not to mention providing high-quality data. Having said that, the most critical point with regards to patient benefit still remains unanswered:

“Could this technique be used for improving cancer patients' prognosis and survival?” “Could this method allow for an earlier, non-invasive diagnosis while there is still room for timely intervention and favorable outcomes (2, 3)?”

For instance, the technological development of hand-held spectroscopic devices and fiber optics has previously shown tremendous promise when it comes to characterization of cancer and could be used for *in-vivo* diagnostics without the need for tissue resection (4-6). Recent studies have also shown the promise of biological fluids for a minimally-invasive and pre-symptomatic cancer detection (7-9). Different types of machine learning algorithms have been developed to interpret spectroscopic data (10) with the aim of detecting cancer. A major field will be the development of predictive algorithms in order to identify pre-symptomatic cases that are destined to develop the disease; this would allow earlier intervention.

The results presented in this carefully-conducted work are undeniably a progressive step towards an accurate and differential diagnosis; however, we also propose that further spectroscopic / clinical research should focus on the development of screening tools for an earlier and minimally invasive detection of cancer.

## References

1. Mittal S, *et al.* (2018) Simultaneous cancer and tumor microenvironment subtyping using confocal infrared microscopy for all-digital molecular histopathology. *Proc Natl Acad Sci USA*:201719551.
2. Richards M (2009) The national awareness and early diagnosis initiative in England: assembling the evidence. *Br J Cancer* 101(S2):S1.
3. Hiom S (2015) Diagnosing cancer earlier: reviewing the evidence for improving cancer survival. (Nature Publishing Group).
4. Mohs AM, *et al.* (2010) Hand-held spectroscopic device for in vivo and intraoperative tumor detection: contrast enhancement, detection sensitivity, and tissue penetration. *Anal Chem* 82(21):9058-9065.
5. Wang W, Zhao J, Short M, & Zeng H (2015) Real-time in vivo cancer diagnosis using raman spectroscopy. *J Biophotonics* 8(7):527-545.
6. Mackanos MA & Contag CH (2010) Fiber-optic probes enable cancer detection with FTIR spectroscopy. *Trends Biotechnol* 28(6):317-323.
7. Baker MJ, *et al.* (2016) Developing and understanding biofluid vibrational spectroscopy: a critical review. *Chem Soc Rev* 45(7):1803-1818.
8. Shaw RA & Mantsch HH (2000) Infrared spectroscopy in clinical and diagnostic analysis. *Encyclopedia Anal Chem* 1:83-102.
9. Gajjar K, *et al.* (2013) Fourier-transform infrared spectroscopy coupled with a classification machine for the analysis of blood plasma or serum: a novel diagnostic approach for ovarian cancer. *Analyst* 138(14):3917-3926.
10. Goodacre R. (2003) Explanatory analysis of spectroscopic data using machine learning of simple, interpretable rules. *Vibr Spectrosc* 32(1):33-45.