



Article

Implementation in rehabilitation: a roadmap for practitioners and researchers

Morris, Jacqui H., Bernhardsson, Susanne, Bird, Marie-Louise, Connell, Louise, Lynch, Elizabeth, Jarvis, Kathryn, Kayes, Nicola M., Miller, Kim, Mudge, Suzie and Fisher, Rebecca

Available at <http://clock.uclan.ac.uk/28316/>

Morris, Jacqui H., Bernhardsson, Susanne, Bird, Marie-Louise, Connell, Louise ORCID: 0000-0002-0629-2919, Lynch, Elizabeth, Jarvis, Kathryn ORCID: 0000-0001-5963-7346, Kayes, Nicola M., Miller, Kim, Mudge, Suzie et al (2020) Implementation in rehabilitation: a roadmap for practitioners and researchers. Disability and Rehabilitation, 42 (22). pp. 3265-3274. ISSN 0963-8288

It is advisable to refer to the publisher's version if you intend to cite from the work.
<http://dx.doi.org/10.1080/09638288.2019.1587013>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.



Implementation in rehabilitation: A roadmap for practitioners and researchers

Journal:	<i>Disability and Rehabilitation</i>
Manuscript ID	TIDS-11-2018-108.R1
Manuscript Type:	Perspectives in Rehabilitation
Keywords:	Knowledge Translation, Rehabilitation, Implementation science, Implementation Context

SCHOLARONE™
Manuscripts

1
2
3 **Implications for Disability and Rehabilitation**
4

- 5 • Effective implementation of research evidence into rehabilitation practice has many
6 interconnected steps and a roadmap analogy is helpful in defining them
7
- 8 • Understanding context for implementation is critically important and using theory can facilitate
9 development of understanding
10
- 11 • Research methods for implementation in rehabilitation should be carefully selected and
12 outcomes should evaluate implementation success as well as clinical change
13
- 14 • Sustainability requires regular revisiting of the interconnected steps
15

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

ABSTRACT

Purpose

Despite growth in rehabilitation research, implementing research findings into rehabilitation practice has been slow. This creates inequities for patients and is an ethical issue. However, methods to investigate and facilitate evidence implementation are being developed. This paper aims to make these methods relevant and accessible for rehabilitation researchers and practitioners.

Methods

Rehabilitation practice is varied and complex and occurs within multilevel healthcare systems. Using a “road map” analogy, we describe how implementation concepts and theories can inform implementation strategies in rehabilitation. The roadmap involves a staged journey that considers: the nature of evidence; context for implementation; navigation tools for implementation; strategies to facilitate implementation; evaluation of implementation outcomes; and sustainability of implementation. We have developed a model to illustrate the journey, and four case studies exemplify implementation stages in rehabilitation settings.

Results and Conclusions

Effective implementation strategies for the **complex** world of rehabilitation are urgently required. The journey we describe unpacks that complexity to provide a template for effective implementation, to facilitate translation of the growing evidence base in rehabilitation into improved patient outcomes. It emphasises the importance of understanding context and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

application of relevant theory, and highlights areas which should be targeted in new implementation research in rehabilitation.

KEY WORDS: Knowledge Translation, Rehabilitation, Implementation Science, Implementation Context

For Peer Review

Introduction

Rehabilitation science is a rapidly growing field, leading to an explosion in research to inform rehabilitation practice [1]. Just in the past four years, the number of published randomized controlled trials in rehabilitation (RCTs) have increased by approximately 2,200 per year [1]. Rigorous trials of rehabilitation interventions, some of which have been conducted internationally [2], innovative evidence synthesis [3], and thousands of evidence based guidelines are all contributing high quality evidence to inform rehabilitation practice [1].

Despite these developments, translating research findings into rehabilitation practice is slow. Use of research evidence among rehabilitation practitioners varies considerably [4-12] and evidence-based practice guidelines often are not fully implemented [13-15]. Typically patients receive a mix of evidence-based and non-evidence-based interventions [9, 16, 17], indicating that research does not automatically influence practice [18, 19]. However, where evidence-informed rehabilitation recommendations have been systematically implemented, benefits for practitioners *and* patients occur. [20]

Failure to translate research into practice creates ethical issues. Moral and actual harm occurs when strategies known to improve peoples' lives are received only by a few, or when ineffective interventions continue to be delivered [19].

However, translating research knowledge into rehabilitation practice is not straightforward, and publication of research evidence is only one step in the process. Rehabilitation is a complex multi-disciplinary process with many moving parts. It is conducted within complex multi-level healthcare

1
2
3 systems, involving interactions between healthcare professionals, teams and managers, patients,
4 families and carers. Organisational context and culture, clinical and professional context and team
5 working and leadership all determine if and how research knowledge is implemented [21]
6
7
8 Transitions from acute and inpatient rehabilitation settings to outpatient primary care and
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

systems, involving interactions between healthcare professionals, teams and managers, patients, families and carers. Organisational context and culture, clinical and professional context and team working and leadership all determine if and how research knowledge is implemented [21] Transitions from acute and inpatient rehabilitation settings to outpatient primary care and community rehabilitation settings make translating knowledge into rehabilitation practice even more complex. Understanding these factors and their interactions will inform strategy development and testing to facilitate knowledge translation for practice, increasing the likelihood for rehabilitation research efforts to make a meaningful difference to rehabilitation practice and outcomes.

Implementation science is a rapidly evolving field concerned with learning how to translate and embed research evidence into practice to improve health outcomes [21-24]. Although the field of implementation research has grown in the last decade, the emergence of implementation research in rehabilitation is relatively recent. Although many guides to implementation science exist, the evolving nature of rehabilitation research means an introduction for rehabilitation researchers and clinicians interested in implementation science, and its potential impact for rehabilitation practice, is timely. This is particularly so, given the complex and highly specific context of rehabilitation, which requires unique strategies for translating research knowledge into practice.

Methods

Using the analogy of a journey towards effective implementation, we have developed a “Roadmap” for implementation research in rehabilitation. This is not an exhaustive review, but by sharing lessons learned from our international group, we intend to offer insight, and create momentum for implementation research in rehabilitation. The roadmap and its stages are described in Figure 1. It

1
2
3 illustrates considerations for each stage of the journey, and shows how these must be revisited to
4
5 ensure sustainability of implementation over time through a central circle denoting how
6
7
8 sustainability can be facilitated.
9

10
11
12
13 We developed the roadmap from four case studies describing research conducted in international
14
15 rehabilitation settings by some of the authors of this paper to provide examples of key
16
17 implementation concepts, theories and methods, and to demonstrate their application in a
18
19 rehabilitation research context (Table 1). The roadmap was developed by examining the strategies
20
21 used in these studies, and by exploration of the processes researchers adopted for implementation
22
23 within the case studies. At the roadmap destination, travellers will have a broad sense of the current
24
25 landscape in terms of what is currently known about factors which may help or hinder
26
27 implementation efforts and related strategies, as they apply in a rehabilitation context. We hope
28
29 this will provide a solid platform on which to further advance knowledge and practice for more
30
31 timely and efficient knowledge translation in rehabilitation.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

40 Insert table 1 about here.

45 Insert Figure 1 about here

50 **The Roadmap**

54 **A: The journey begins. What do we mean by evidence?**

56
57 As rehabilitation researchers, our primary focus is on evidence derived from research. However,
58
59 evidence takes multiple forms. Local data, clinical reasoning, evaluation, and even tacit knowledge
60

1
2
3 [25], patient experience, and organisational information all provide evidence to inform
4
5 implementation efforts. Reconciling tension between clinical trial evidence cited in clinical
6
7 guidelines, derived from the average patient within a highly standardised sample, against what is
8
9 best for individual patients presenting in rehabilitation, is critical. Slavish devotion to evidence based
10
11 guidelines irrespective of factors relevant to the individual, is amoral, placing patients at risk of
12
13 inappropriate treatment [26]. Thus, Bowen and Graham [27] (page S4) suggest evidence-informed
14
15 (rather than evidence-based) practice because “(1) research to inform complex problems facing
16
17 health systems is often incomplete, (2) much research is contradictory, and (3) non-research forms
18
19 of evidence can be legitimately used in decision-making”. These arguments reflect our pragmatic
20
21 approach to real-world rehabilitation research, and we fully support the use of multiple forms of
22
23 evidence drawn from different sources.
24
25
26
27
28
29
30
31
32

33 **Roadmap B: Planning the route to avoid roadblocks: The importance of context**

34
35 Why do some therapists adopt evidence-informed interventions, when others do not [15, 17, 28]
36
37 and why is it that improvement initiatives work in some settings but not in others? [29, 30]. In
38
39 implementation research, the term context has been used to describe factors that influence
40
41 intervention adoption, effectiveness or sustainability in real world settings. Appreciation and
42
43 planned investigation of context aids identification of possible routes to take in this type of research
44
45 and enhances clarity about potential obstacles on the journey [31].
46
47
48
49
50
51

52
53 Investigating context facilitates understanding of “what happens when a particular intervention is
54
55 joined together with an individual, team, organisation or healthcare system” [32] (page 109).
56
57 Considering context at each of these levels is important for informing targeted strategy
58
59 development. For example, investigating effects of strategies designed to influence therapy practice
60

1
2
3 requires exploration of therapist knowledge, behaviour, and actual treatment of patients (i.e.
4 informing strategies targeted at the individual level) [15] [28]. Conversely, investigating adoption of
5 evidence-based services, such as hospital-based stroke rehabilitation or early supported discharge
6 (ESD), requires investigation of team working and organisational influences (i.e. strategies targeted
7 at the team and organisational level) [33, 34].
8
9
10
11
12
13
14
15
16
17

18 Context is dynamic and changes over time, therefore it should be considered as a process. Often
19 context is reduced to environment or place of intervention delivery, to identify discrete contextual
20 factors [35]. However context can also include individual factors and human agency, making
21 understanding staff and patient perspectives crucial. As Bate points out “how we think about
22 context will determine how we go about researching and ultimately trying to manage it” [36] (page
23 20).
24
25
26
27
28
29
30
31
32
33
34

35 In rehabilitation, implementation research is particularly challenging, because intervention and
36 context overlap. For example in group therapy, it may not be clear which intervention components
37 are essential for the therapeutic effect, and which can be deemed context. Yet, this is critical,
38 because adaptations are often necessary in order for interventions to be adopted, or sustained over
39 time. Damschroder’s emphasis on identifying core intervention components (active ingredients)
40 and clear hypotheses about intervention mechanisms of action are crucial [21]. If the active
41 ingredients of an intervention are known, other aspects of the intervention can be adapted to
42 facilitate adoption without fundamentally changing the intervention. This approach should maintain
43 the effectiveness of the intervention [37].
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Implementation research requires a shift in thinking from the gold standard randomised controlled
4 trial model [36, 38, 39]. This design was developed for testing drugs, and requires identical
5
6 interventions to be delivered across sites under controlled circumstances to ensure any observed
7
8 effect can be attributed to the drug. However, in a complex rehabilitation intervention, intervention
9
10 effect cannot be removed from the context in which it is delivered and indeed, observed effect may
11
12 arise through a combination of intervention content, delivery and context. As such, investigations
13
14 into how interventions should be adapted for complex rehabilitation delivery contexts whilst
15
16 maintaining their active ingredients are necessary, but lacking. [37]. Such investigation is crucial to
17
18 optimise the likelihood for rehabilitation interventions to be adopted and sustained in real world
19
20 settings, and is why implementation research is so important.
21
22
23
24
25
26
27
28
29

30 **Roadmap C: What tools can help us navigate? Implementation theory**

31
32 Important tools for implementation research are theories, models and frameworks developed to
33
34 understand implementation success and failure, and to guide and evaluate implementation efforts
35
36 [40]. Theories exist to facilitate generalisation, replicability and effectiveness of implementation
37
38 strategies [41]. Using theory in implementation research is recommended by many and its use has
39
40 increased considerably in recent years [42] [43]. However, a scoping review published in 2010,
41
42 examining use of three theories in implementation research in rehabilitation, medicine and nursing,
43
44 [44] demonstrated infrequent use of these theories to inform implementation research in
45
46 rehabilitation, compared to studies within medicine and nursing. The review suggests the use of
47
48 theory in rehabilitation is in its infancy, compared to its more prevalent use in other fields of health
49
50 research.
51
52
53
54
55
56
57
58
59
60

1
2
3 Selecting appropriate theoretical approaches is challenging as there are more than 100 relevant
4 theories [45]. To aid theory selection, theories can be categorised according to three general aims,
5
6 as proposed by Nilsen [40]:
7
8

- 9
10 1) Determinant frameworks, classic or implementation theories: to optimise understanding
11 and/or explain what influences implementation outcomes
12
- 13 2) Process models: to describe or guide the translation of research into practice
14
- 15 3) Evaluation frameworks: to evaluate implementation.
16
17
18
19

20
21 Here we exemplify how these aims influenced selection of theoretical approaches in our case study
22 examples (see Table 1).
23
24
25

26 27 28 ***Theories used to facilitate understanding of factors influencing implementation:***

29
30 *The Consolidated Framework for Implementation Research (CFIR)* integrates multiple existing
31 implementation theories. It provides a comprehensive framework for understanding multifactorial
32 influences on implementation [21]. These influences span intervention characteristics, individuals
33 and teams involved in implementation and their interactions, internal organisational characteristics,
34 and wider external drivers for change. In Case study 1, Fisher and colleagues used CFIR to analyse
35 and identify core components of an Early Supported Discharge intervention, and to understand how
36 components should be adapted to real-world contextual factors. In Case study 2, Connell and
37 colleagues used CFIR to explore how an upper limb rehabilitation intervention for stroke
38 rehabilitation was implemented in practice. It was also used for survey design to elicit characteristics
39 of rehabilitation therapists and settings where implementation occurred. In both case studies, the
40 CFIR guided identification of how networks, team communication, leadership and preconceptions
41 of the intervention influenced implementation, facilitating strategy development for improving
42 implementation.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Normalisation Process Theory (NPT)* is a conceptual framework for understanding how new
4 technologies and complex interventions are operationalized in everyday work [35]. It examines how
5 people make sense of the intervention, how they commit to implementing it, how they work
6 together to make implementation happen, and how they individually and collectively appraise
7 implementation. In Case study 4, Mudge and Kayes used NPT to analyse data from patients and staff
8 participating in a pilot trial. Experiences of implementation in practice were explored and findings
9 informed revisions to the Toolkit intervention as well as the planned implementation processes. In
10 Case Study 2, NPT was used to explore processes involved in identifying, integrating and embedding
11 an evidence-based upper limb training program for stroke in practice. Although useful in identifying
12 individual and collective experiences, it did not capture wider contextual factors that influenced
13 implementation, confirming that one theory in isolation is often insufficient for identification of the
14 diverse influences on implementation.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 ***Theories used to guide the process of translating research into practice:***

36
37 The *Grol and Wensing Implementation model* [46] is a process model, developed to support and
38 guide implementation. It was used to guide development of a tailored implementation strategy for
39 clinical practice guidelines in primary care physiotherapy (Case study 3). The model was selected
40 because it integrates theories from several research fields, including behavioural change theories,
41 and provides a structured, pragmatic, approach for implementation. It proved to be a
42 comprehensive tool for addressing factors requiring consideration in a rehabilitation
43 implementation project.
44
45
46
47
48
49
50
51
52
53
54
55
56

57 *The Behaviour Change Wheel (BCW)* [47] was developed by systematic synthesis of behaviour
58 change frameworks to support behaviour change intervention development. It seeks to identify
59
60

1
2
3 and understand aspects of capability, opportunity and motivation that need to be addressed to
4
5 facilitate the desired behaviours, and to match them to relevant behaviour change techniques.
6
7
8 Whilst the BCW focuses on changing an individual's behaviour, it also includes policy categories to
9
10 address system-level influences. In Case study 2, the BCW was used to guide development of a
11
12 behaviour change strategy to enable therapists to implement strategies to increase intensity of
13
14 upper limb exercise in stroke rehabilitation. In this context, the taxonomy of behaviour change
15
16 techniques was useful to inform implementation strategies development.
17
18
19
20
21
22

Theories used to guide evaluation of implementation

23
24 Evaluation of implementation is often an iterative process, with findings facilitating tailoring of
25
26 implementation strategies, for subsequent re-evaluation. In Case study 4, NPT informed our analysis
27
28 of a pilot study of implementing the Living Well Toolkit in practice in four rehabilitation settings. We
29
30 were able to respond to the findings by making some changes to the implementation of the toolkit,
31
32 including the development of introductory videos to simplify the explanation of the toolkit,
33
34 development of an additional version of the toolkit and development of a tiered training package.
35
36 The first tier of training involved everyone (including administrators) to ensure a shared philosophy
37
38 for the toolkit. Stakeholders were trained to embed systems and to reflect on practice, organising
39
40 opportunities to share reflections of successes, challenges and tensions in embedding the toolkit
41
42 into practice. This in turn facilitated development of strategies for managing the implementation
43
44 process [48].
45
46
47
48
49
50
51
52
53

54 From our case studies examples, it is clear that targeted selection of frameworks for specific
55
56 implementation purpose is **critical**. One framework or theory is unlikely to address all facets of
57
58
59
60

1
2
3 implementation and selection of the appropriate framework that is fit for purpose requires careful
4
5 consideration.
6
7
8
9

10 **Roadmap D: Making sure the journey happens: Implementation strategies**

11
12 Despite the theories and frameworks available to guide implementation processes, a 2015 review
13
14 [49] highlights that implementation strategies in rehabilitation have relied on educational strategies
15
16 as the primary intervention tool. This is despite some evidence suggesting active, multi-component
17
18 implementation strategies might be more effective [20, 50, 51]. Educational strategies alone may
19
20 improve clinician knowledge and skill, however they do not necessarily result in implementation,
21
22 particularly when didactic and passive strategies are used [52, 53] are examples, but there are
23
24 others. Other implementation strategies that have been shown useful in rehabilitation settings
25
26 include multifaceted training packages, journal clubs, and contextualised guidelines [54].
27
28
29
30
31
32
33
34

35 As we have alluded to above, differentiating mandatory 'active ingredients' and effective elements
36
37 of desired practice, from those that are optional and can be adapted, helps to address tensions
38
39 between the need for local contextualisation and fidelity [28, 55]. Furthermore, tools and strategies
40
41 informed by theories discussed above, are necessary for uptake and sustained implementation. A
42
43 plethora of different implementation strategies have been described and used, and, as with
44
45 theories, choosing the appropriate ones can be challenging. A recent overview of implementation
46
47 strategies by Leeman et al [56] provides a useful classification of strategies and suggested
48
49 terminology. The purpose was to increase consistency in reporting and facilitate comparisons and
50
51 synthesis of findings, as well as aligning strategies with relevant theories, and finding strategies that
52
53 best match specific needs [56].
54
55
56
57
58
59
60

1
2
3 In considering what tools are appropriate and when, we define three stages of the implementation
4 process (planning, implementation and monitoring), provide examples of tools to support each
5 stage, and associated considerations for research and practice. These are presented in Table 2. A
6 number of the tools referred to in Table 2 are derived from the theories and frameworks described
7 above. For example, CFIR offers a practical framework for exploring in-depth the factors which may
8 help or hinder uptake within a specific context during the planning stage. Similarly, the NOMAD tool
9 was developed drawing on NPT to use with staff involved in implementation activities and can
10 inform the development of more targeted implementation strategies. As well as drawing on tools
11 derived from relevant implementation theories, Table 2 also provides examples of a range of other
12 strategies discussed in the literature which have been argued to support the implementation
13 process, such as structural supports, communities of practice, knowledge brokerage; and review
14 and monitoring processes which may contribute to more sustained change (discussed in more detail
15 below). The practical considerations presented alongside each stage are not exhaustive, but rather
16 are some of the authors' collective learning from the case studies in Table 1.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 **Roadmap E: How do we know we have followed the right road? Implementation Research Designs**

41
42 The complexity of rehabilitation interventions coupled with the multi-factorial systems-orientated
43 nature of implementation means multiple methodological approaches and research designs are
44 required to investigate implementation. Peters et al. (2013) summarise implementation research
45 objectives, which, together with our case studies, we use to illustrate some implementation
46 research designs [57].
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Explore

Here, the objective is to explore a phenomenon, develop understanding and make generalisations for subsequent testing. Case study 1 aimed to explore implementation contexts for ESD services after stroke, and to identify factors that facilitate or impede implementation. The purpose was to identify adaptable elements of context that could be modified to enable embedding of ESD in practice. The design was an exploratory qualitative design using interviews with key stakeholders. The novel exploratory work informed a subsequent implementation study [58]. In Case study 3, determinants of guideline use by musculoskeletal physiotherapists were investigated in a survey, as a first step in a model-supported implementation project, and the implementation strategy was tailored to address the identified determinants.

Describe

A starting point in implementation research is often description of a phenomenon or context, with identification of correlates and possible causes as another potential objective. Within Case study 2, a qualitative study and a survey, informed by CFIR, together described rehabilitation contexts. Contextual information informed development of specific behaviour change strategies to support implementation of increased dose of upper limb rehabilitation [28].

Influence

Influencing change in specific outcomes is of course the overall aim of implementation research. Research designs for evaluating implementation strategies fall along a continuum between uncertainty and certainty of effectiveness. Design selection is informed by the desired degree of probability that effects are due to the intervention [57]. In most implementation research, the rehabilitation interventions being implemented are likely to have proven efficacy (demonstrated

1
2
3 through an RCT conducted in ideal circumstances). What is of most interest is the effectiveness of
4
5 the rehabilitation intervention in practice (effect in a real world setting). The real-world nature of
6
7 implementation research therefore challenges researchers to see beyond the RCT design [39]. It
8
9 thus may be appropriate to utilise uncontrolled before and after studies (Case study 1) or time series
10
11 designs to examine whether implementation is occurring over time.
12
13
14
15
16
17

18 In participatory action research, researchers and participants work together to collect data, take
19
20 action where this is supported and reflect on the process [59]. It is another rigorous method by
21
22 which change can be iteratively implemented and evaluated in a given context, and that tolerates
23
24 uncertainty, but is not expected to generalise beyond that setting.
25
26
27
28
29

30 Realist methodologies may also be useful to gain an understanding of what works, for whom and in
31
32 what circumstances [60]. Using this approach in implementation research could lead to more
33
34 targeted implementation activities which build on an understanding of how the intervention
35
36 operates in specific contexts and then leads to specific outcomes [29].
37
38
39
40
41

42 Design selection should therefore stem from evaluation of both implementation aims, tolerance of
43
44 uncertainty and an appreciation of context.
45
46
47
48

49 **Roadmap F: How do we know if we've got to the destination? Implementation outcomes**

50
51
52 The primary outcomes within rehabilitation are clinical and patient-recorded outcomes, which are
53
54 relevant when examining how well rehabilitation interventions operate in real world settings [58].
55

56
57 However, clinical outcomes are distal when establishing effects of implementation or facilitation
58
59 strategies, as they depend on successful implementation. To understand, develop and evaluate how
60

1
2
3 well implementation strategies are working it is salient to examine conceptually distinct
4
5 implementation outcomes. Proctor et al. define specific implementation outcomes, which include
6
7 acceptability, adoption, appropriateness, costs, feasibility, fidelity, penetration and sustainability of
8
9 the implementation strategy [61]. These are implementation outcomes thought necessary to
10
11 mediate and sustain change in clinical outcomes [62]. Implementation outcomes can be examined
12
13 using quantitative or qualitative methods.
14
15
16
17
18
19

20 In both Case study 2 and 3, implementation outcomes were examined. In Case study 2, the effects
21
22 of an implementation strategy aiming to change therapists' behaviour in delivery of dose of upper
23
24 limb rehabilitation after stroke were measured. Therapists' behaviours, as well as their perceptions
25
26 of acceptability, appropriateness and compatibility with practice, were examined using qualitative
27
28 methods informed by NPT. Feasibility, or practicability of implementation within the context of the
29
30 study sites was indicated through audit, which recorded performance of the desired staff
31
32 behaviours, and discrepancies between sites were examined through evaluation within the
33
34 interviews. Accounts of difficulties experienced in maintaining momentum for implementation in a
35
36 planned way were also elicited in Case study 2 by capturing implementation rather than clinical
37
38 outcomes. In this way, knowledge to inform future adaptation of the implementation strategy and
39
40 processes was generated. This comprehensive case study illustrates how critical implementation
41
42 outcomes can be examined, and how they reflect the success, or not, of implementation efforts in
43
44 different settings. The study identified differences in implementation success across sites,
45
46 developing generalizable understanding about how implementation success can ultimately
47
48 influence clinical outcomes.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 In Case study 3, implementation outcomes were also examined. Effects of the multi-component
4 implementation strategy to increase use of clinical practice guidelines were measured in terms of
5
6 therapists' attitudes (acceptability), access (feasibility) and behaviours, particularly those related to
7
8 guideline use (adoption). Sustainability of the implementation is currently being measured in a long-
9
10 term follow-up study.
11
12
13
14
15
16
17

18 **Roadmap G: How do we remain roadworthy over time? Sustaining implementation**

19
20 The behavioural and evolving nature of rehabilitation science and practice means sustainability
21 requires action to preserve fidelity and intended effects of the implemented intervention. However,
22 as CFIR suggests, sustainability is challenging, requiring continuous adaptation of the implemented
23 intervention *and* the implementation strategy (Table 2). Clearly planning and systems for
24 sustainability **should be adopted**. Identifying, anticipating and responding to factors that influence
25 long-term sustainability should be considered as integral to implementation strategies [62].
26
27
28
29
30
31
32
33
34
35
36

37 Explicitly adopting a cyclical approach to review of implementation should ensure new evidence is
38 implemented **whilst** accounting for internally and externally generated system changes [62, 63].
39 Research evidence, once established as usual care, can become outdated. Considering how
40 monitoring and review for sustainability can be **conducted** at the individual and organisational levels
41 is **critical**. Tools such as the NOMAD mentioned above and models such as the Plan, Do, Study Act
42 model (PDSA) [64] (page 135) within quality improvement can structure the process of
43 sustainability. **These methods are highly applicable and accepted in rehabilitation conducted within**
44 **health service settings, and can facilitate** continuous review and updating of rehabilitation practice,
45 given its rapidly expanding knowledge base [65]. At any point in the sustainability cycle, return to
46 other stages on the roadmap is possible, as highlighted in Figure 1. Building capacity for adaptation
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 as the evidence base expands is also vital and requires continued learning and problem solving, and
4
5 continuous evaluation of fit between the rehabilitation intervention and multi-layered context
6
7 within the system [66].
8
9

10 11 12 13 **Conclusions and implications** 14

15 This paper provides a snapshot of the large and growing landscape of implementation research and
16
17 was not intended as a comprehensive overview. Instead, our intention was to provide an
18
19 introduction and guidance to implementation science for rehabilitation researchers and
20
21 practitioners. In it we share empirical experiences of implementation research in rehabilitation
22
23 contexts and map them to theoretical perspectives derived from the current implementation
24
25 literature.
26
27
28
29
30
31

32 We have highlighted the importance of understanding the context in which implementation is to
33
34 occur, and how implementation strategies should emerge from that understanding. We recognise,
35
36 however, that future research needs to investigate more fully how specific contexts in rehabilitation
37
38 influence implementation. We acknowledge too that engaging practitioners to understand what
39
40 their knowledge needs are, will support better implementation in future. Developing creative ways
41
42 to generate and provide knowledge to support implementation **should be a focus** for future
43
44 research.
45
46
47
48
49
50

51 We have highlighted just a few of the many theoretical approaches and strategies that can be used
52
53 to guide knowledge generation about context and to guide implementation. Our arguments are
54
55 based on our own experiences, but we want to emphasize the importance to other researchers of
56
57 choosing one or several of those theories and strategies so knowledge gained by research can be
58
59
60

1
2
3 generalised and theories developed to match rehabilitation concepts. There is an urgent need for
4
5 more theory-informed research to generate comparable and generalizable knowledge that will
6
7 contribute to expanding the modest evidence base for successful implementation in rehabilitation.
8
9
10 Research on implementation strategies which accounts for the unique and specific complexities of
11
12 the rehabilitation context, will allow for a better understanding of what works, and what does not
13
14 work, ultimately increasing our likelihood to meaningfully impact rehabilitation practice and
15
16 outcomes.
17
18
19

20
21
22 We used a roadmap analogy to make the stages of conducting implementation research in
23
24 rehabilitation explicit. These stages were exemplified by mapping of implementation components
25
26 to the different parts of the road journey through rehabilitation studies conducted by some of the
27
28 study authors. It is our hope that this roadmap, and the sharing of our own implementation research
29
30 experiences, can lead to more rehabilitation researchers embarking on this important journey. We
31
32 hope it will support their endeavours in exploring, implementing, and sustaining evidence-informed
33
34 practices within their context.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Kamper SJ, Moseley AM, Herbert RD, et al 15 years of tracking physiotherapy evidence on PEDro, where are we now? Br J Sports Med 2015; 49: 907-909.
2. AVERT Trial Collaboration Group. Efficacy and safety of very early mobilisation within 24 hours of stroke onset (AVERT): a randomised controlled trial. Lancet. 2015; 386: 46-55.
3. Pollock A, Farmer SE, Brady MC et al. Interventions for Improving Upper Limb Function After Stroke. Cochrane Database Syst Rev. 2014; 11: CD010820. doi: 10.1002/14651858.CD010820.pub2.
4. Jette DU, [Bacon K](#), [Batty C](#). Evidence-based practice: beliefs, attitudes, knowledge, and behaviors of physical therapists. Phys Ther. 2003; 83: 786-805.
5. Iles R, Davidson M. Evidence based practice: a survey of physiotherapists' current practice. Phys Res Int. 2006; 11: 93-103.
6. Hodder RK, Wolfenden L, Kamper SJ. Developing implementation science to improve the translation of research to address low back pain: A critical review. Best Pract Res Clin Rheumatol. 2017; 30: 1050-1073.
7. Holden MA, Nicholls EE, Hay EM et al. Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations? Phys Ther. 2008; 88; 1109-1121.
8. Salbach NM, Guilcher SJT, Jaglal SB. Determinants of research use in clinical decision making among physical therapists providing services post-stroke: a cross-sectional study. Implement Sci. 2010; doi: [10.1186/1748-5908-5-77](#)
9. Liddle, S.D., Physiotherapists' use of advice and exercise for the management of chronic low back pain: A national survey. Man Ther. 2009; 14: 189-196.

- 1
2
3 10. Hodder, R.K., et al., Developing implementation science to improve the translation of
4
5 research to address low back pain: A critical review. *Best Pract Res Clin Rheumatol.* 2016;
6
7 30: 1050-1073.
8
9
- 10 11. Bernhardsson, S., et al., Determinants of guideline use in primary care physical therapy: a
12
13 cross-sectional survey of attitudes, knowledge, and behavior. *Phys Ther.* 2014; 94: 343-354
14
- 15 12. Lindström AC and B. S., Evidence-based practice in primary care occupational therapy:
16
17 a survey in Western Sweden. *Occup Ther Int.* 2018; doi.org/10.1155/2018/5376764
18
- 19 20 13. Royal College of Physicians. National sentinel stroke audit 2010. Clinical Effectiveness
21
22 and Evaluation Unit Royal College of Physicians, London
23
- 24 25 14. Bernhardsson S, Larsson ME, Eggertsen R, Olsén MF, Johansson K, Nilsen P, et al.
26
27 Evaluation of a tailored, multi-component intervention for implementation of evidence-
28
29 based clinical practice guidelines in primary care physical therapy: a non-randomized
30
31 controlled trial. *BMC Health Serv Res.* 2014; 14: 105.
32
33
- 34 35 15. Korner-Bitensky NJ, Desrosiers J, Rochette A. A national survey of occupational therapists'
36
37 practices related to participation post-stroke. *J Rehabil Med.* 2008; 40: 291-297.
38
- 39 40 16. Mikhail C. Physical Therapists' Use of Interventions With High Evidence of Effectiveness in
41
42 the Management of a Hypothetical Typical Patient With Acute Low Back Pain. *Phys Ther.*
43
44 2005; 85:11.
45
- 46 47 17. Bernhardsson S, Öberg B, Johansson K, Nilsen P, Larsson ME. Clinical practice in line
48
49 with evidence? A survey among primary care physiotherapists in western Sweden. *J Eval*
50
51 *Clin Pract.* 2015; 21: 1169-1177.
52
53
- 54 55 18. McGlynn EA¹, Asch SM, Adams J, Keeseey J, Hicks J, DeCristofaro A, Kerr EA. The quality of
56
57 health care delivered to adults in the United States. *New Engl J Med.* 2003; 348: 2635-
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2645.

19. Solomon, MZ. The ethical urgency of advancing implementation science. *Am J Bioethics*. 2010; 10: 31-32.
20. Salbach NM, Wood-Dauphinee S, Desrosiers J, Eng JJ, Graham ID, Jaglal SB et al. Facilitated interprofessional implementation of a physical rehabilitation guideline for stroke in inpatient settings: process evaluation of a cluster randomized trial. *Implement Sci*. 2017: doi: 10.1186/s13012-017-0631-7
21. Damschroder LJ, Aaron DC, Keith RE, Kirsh S, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci*. 2009; doi.org/10.1186/1748-5908-4-50
22. Madon T, Hofman KJ, Kupfer L, Glass RI. Public Health. *Implementation Science*. Science. 2007; 318: 1728-1729.
23. Eccles MP, Mittman BS. Welcome to Implementation Science. 2006. *Implementation Science*; doi.org/10.1186/1748-5908-1-1
24. Lang ES, Wyer PC, Haynes RB. Knowledge Translation: Closing the Evidence-to-Practice Gap. *Ann Emerg Med*. 2007; 49: 355-363.
25. Dijkers, MP, SL Murphy, J Krellman. Evidence-based practice for rehabilitation professionals: concepts and controversies. *Arch Phys Med Rehab*. 2012; 93: S164-S176.
26. Greenhalgh T. Of lamp posts, keys, and fabled drunkards: A perspectival tale of 4 guidelines. *J Eval Clin Pract*. 2018; 24:1132-1138.

- 1
2
3 27. Bowen SJ and ID Graham. From knowledge translation to engaged scholarship: promoting
4 research relevance and utilization. *Arch Phys Med Rehab.* 2013; 94: S3-S8.
5
6
7
8 28. Connell, L.A, McMahon NE, Watkins CL, Eng JJ. Therapists' use of the Graded Repetitive
9 Arm Supplementary Program (GRASP) intervention: a practice implementation survey
10 study. *Phys Ther.* 2014; 94: 632-643.
11
12
13
14
15 29. Cadilhac DA, Andrew NE, Stroil Salama E On behalf of the Australian Stroke Clinical
16 Registry Consortium, *et al* Improving discharge care: the potential of a new
17 organisational intervention to improve discharge after hospitalisation for acute stroke, a
18 controlled before–after pilot study *BMJ Open* 2017; **7**: e016010. doi: 10.1136/bmjopen-
19 2017-016010
20
21
22
23
24
25
26
27 30. Power M, Tyrell P, Rudd AG, Tully M, Dalton D, Marshall M et al. Did a quality
28 improvement collaborative make stroke care better? A cluster randomized trial.
29
30
31
32 Implementation science. 2014; 9: 40. doi.org/10.1186/1748-5908-9-40
33
34
35 31. Papoutsi C, Boden R, Foy R, Grimshaw J, Rycroft-Malone J. Challenges for
36 implementation science. *In* Raine R, Fitzpatrick R, Barratt H, Bevan G, Black N, Boaden,
37 R et al. Challenges, solutions and future directions in the evaluation of service
38 innovations in health care and public health. *Health Services and Delivery Research*
39 2016; 16: 1-139. DOI: 10.3310/hsdr04160-105
40
41
42
43
44
45
46
47 32. Howarth E, Denvers K, Moore G, O'Cathain A, Dixon-Woods M. Contextual issues and
48 qualitative research. *In* Raine R, Fitzpatrick R, Barratt H, Bevan G, Black N, Boaden,
49 R et al. Challenges, solutions and future directions in the evaluation of service innovations in
50 health care and public health. *Health Services and Delivery Research* 2016. 16; 4: 1-139. DOI:
51 10.3310/hsdr04160-105
52
53
54
55
56
57
58
59 33. Bayley MT, Hurdowar A, Richards CL, Korner-Bitensky N, Wood-Dauphinee S, Eng JJ.
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Barriers to implementation of stroke rehabilitation evidence: findings from a multi-site pilot project. *Disabil Rehab.* 2012; 34: 1633-1638.

34. Chouliara, N, Fisher RJ, Kerr M, Walker MF. Implementing evidence-based stroke Early Supported Discharge services: a qualitative study of challenges, facilitators and impact. *Clin Rehab.* 2014; 28: 370-377.
35. May CR, Mair F, Finch T, MacFarlane A, Dowrick C, Treweek S, Rapley T. Development of a theory of implementation and integration: Normalization Process Theory. *Implement Sci.* 2009; 4: 29. doi: 10.1186/1748-5908-4-29.
36. Bate P, Robert G, Fulop N, Ovretveit J, Dixon-Woods M. Perspectives on Context: A Collection of Essays Considering the Role of Context in Successful Quality Improvement. 2014. The Health Foundation. London.
37. Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W. Process evaluation of complex interventions. *Complex interventions in health: an overview of research methods.* 2015; 350: 1258. doi.org/10.1136/bmj.h1258
38. May CR, Johnson M, Finch T. Implementation, context and complexity. *Implement Sci.* 2016; 11: 141. doi.org/10.1186/s13012-016-0506-3
39. Kersten, P, Ellis-Hill C, McPherson KM, Harrington R. Beyond the RCT—understanding the relationship between interventions, individuals and outcome—the example of neurological rehabilitation. *Disabil Rehab.* 2010; 32: 1028-1034.
40. Nilsen P. Making sense of implementation theories, models and frameworks. *Implement Sci.* 2015; 10: 53. doi.org/10.1186/s13012-015-0242-0

- 1
2
3 41. Rycroft-Malone J, Bucknall T (Eds.). Evidence based nursing series, Volume 2. Models and
4 frameworks for implementing evidence-based practice: linking evidence to action.
5
6 Singapore, John Wiley & Sons; 2011.
7
8
9
10 42. Hudon A, Gervais MJ, and Hunt M. The contribution of conceptual frameworks to knowledge
11 translation interventions in physical therapy. *Phys Ther.* 2015; 95: 630-639.
12
13
14 43. Liang L, Bernhardsson S, Vernooij RW, Armstrong MJ, Bussièrès A, Brouwers
15 MC, Gagliardi AR. Members of the Guidelines International Network Implementation
16 Working Group. Use of theory to plan or evaluate guideline implementation among
17 physicians: a scoping review. *Implement Sci.* 2017;12:26. doi: 10.1186/s13012-017-
18 0557-0
19
20
21
22
23
24
25
26
27 44. Colquhoun, H.L Letts LJ, Law MC, MacDermid JC, Missiuna CA. A scoping review of the
28 use of theory in studies of knowledge translation. *Can J Occup Ther.* 2010; 77: 270-279.
29
30
31
32 45. Birken SA, Powell BJ, Shea CM, Haines ER, Kirk AM, Leeman J, et al. Criteria for selecting
33 implementation science theories and frameworks: results from an international survey.
34
35 *Implement Sci.* 2017; 12: 124. doi.org/10.1186/s13012-017-0656-y.
36
37
38
39
40
41 46. Grol R, Wensing M, Eccles M, Davis D (Eds.). *Improving patient care: the implementation of*
42 *change in health care, Second Edition.* John Wiley & Sons, Oxford, 2013:
43
44
45
46 47. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, Eccles
47 MP, Cane J, Wood CE. The behavior change technique taxonomy (v1) of 93
48 hierarchically clustered techniques: building an international consensus for the reporting
49 of behavior change interventions. *Ann Behav Med.* 2013; 46: 81-95.
50
51
52
53
54
55 48. Mudge S , Sezier A, Payne D, Kersten P, Kayes N. Developing and putting into practice a
56 Living Well Toolkit: a pilot study drawing on Normalisation Process Theory. Presented at
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

New Zealand Rehabilitation Association Conference 2017: Christchurch, New Zealand.

49. Jones CA, Roop SC, Pohar SL, Albrecht L, Scott SD. Translating knowledge in rehabilitation: systematic review. *Phys Ther.* 2015; 95: 663-677.
50. Bekkering, G., Hendriks HJ, van Tulder MW, Knol DL, Hoeijenbos M, Oostendorp RA, Bouter LM. Effect on the process of care of an active strategy to implement clinical guidelines on physiotherapy for low back pain: a cluster randomised controlled trial. *BMJ Qual Safe.* 2005; 14: 107-112.
51. Menon, A Korner-Bitensky N, Kastner M, McKibbin KA, Straus S. Strategies for rehabilitation professionals to move evidence-based knowledge into practice: a systematic review. *J Rehabil Med.* 2009; 41: 1024-1032.
52. Glegg SM, Holsti L, Stanton S, Hanna S, Velikonja D, Ansley B, Sartor D, Brum C. Evaluating change in virtual reality adoption for brain injury rehabilitation following knowledge translation. *Disabil Rehab: Assist Tech.* 2017; 12: 217-226.
53. Levac D, Glegg SM, Camden C, Rivard LM, Missiuna C. Best practice recommendations for the development, implementation, and evaluation of online knowledge translation resources in rehabilitation. *Phys Ther.* 2015; 95: 648-662.
54. Bernhardsson S, Lynch E, Dizon JM, Fernandes J, Gonzalez-Suarez C, Lizarondo L, Luker J, Wiles L, Grimmer K. Advancing evidence-based practice in physical therapy settings: multinational perspectives on implementation strategies and interventions. *Phys Ther.* 2016; 97: 51-60.
55. Clinton-McHarg T, Young SL, Tzelepis F, Regan T, Fielding A, Skelton E et al. Psychometric properties of implementation measures for public health and community settings and mapping of constructs against the Consolidated Framework for Implementation Research: a systematic review. *Implement Sci.* 2016; 11:148. DOI 10.1186/s13012-016-0512-5

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
56. Leeman J, Birkin SA, Powell BJ, Rohweder C, Shea C. Beyond “implementation strategies”: classifying the full range of strategies used in implementation science and practice. *Implement Sci.* 2017; 12:125. DOI 10.1186/s13012-017-0657-x
57. Peters DH. Implementation research: what it is and how to do it. *BMJ.* 2013;347:f6753.
58. Fisher RJ, Cobley CS, Potgeiter I, Moody A, Nouri F, Gaynor C et al. Is stroke early supported discharge still effective in practice? A prospective comparative study. *Clin Rehab.* 2016; 30: 268-276.
59. Kindon S, R. Pain, Kesby M (Eds). *Participatory action research: approaches and methods.* 2007; Routledge, London.
60. Pawson R, Tilley N. *Realistic evaluation.* 1997: Sage Publications, London.
61. Proctor E, Silmere H, Raghaven R, Hovmand P, Aarons G, Bunger A et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Administration and Policy in Mental Health and Mental Health Services Research.* 2011; 38: 65-76.
62. Haji F, Morin MP, and Parker K. Rethinking programme evaluation in health professions education: beyond ‘did it work?’. *Med Ed.* 2013; 47: 342-351.
63. Bar-Yam, Y., *Complexity rising: From human beings to human civilization, a complexity profile.* 2002; NESCO Encyclopedia of Life Support Systems, Cambridge MA.
64. Deming, W. E. 1993. *The New Economics For Industry, Government & Education.*
Cambridge: Massachusetts Institute of Technology Center for Advanced Engineering Study
65. Sanders J, Fitzpatrick JM. Improving service user self-management: development and implementation of a strategy for the Richmond Response and Rehabilitation Team. *BMJ Open Qual.* 2017; 6: e000126.

- 1
2
3 66. Chambers DA, Glasgow, Stange KC. The dynamic sustainability framework: addressing the
4 paradox of sustainment amid ongoing change. *Implement Sci.* 2013; 8: 117.
5
6
7 doi.org/10.1186/1748-5908-8-117
8
9
- 10 67. Langhorne P, Widen-Holmqvist L. Early supported discharge trialists. Early supported
11 discharge after stroke. *J Rehabil Med.* 2007; 39: 269-269.
12
13
14
- 15 68. Graham ID, Logan J, Harrison MB, Straus SE, Tetroe J, Caswell W, Robinson N. Lost in
16 knowledge translation: time for a map? *Journal of continuing education in the health*
17
18
19
20
21
22 professions. 2006; 26: 13-24.
- 23 69. Fisher RJ, Gaynor C, Kerr M, Langhorne P, Anderson C, Bautz-Holter E et al. A consensus on
24 stroke: early supported discharge. *Stroke.* 2011; 42: 1392-1397.
25
26
27
- 28 70. Cobley CS, Fisher RJ, Chouliara N, Kerr M, Walker M. A qualitative study exploring patients'
29 and carers' experiences of Early Supported Discharge services after stroke. *Clin Rehab.* 2013;
30
31
32
33
34 27: 750-757.
- 35 71. Connell LA, McMahon NE, Harris JE, Watkins CL, Eng JJ. A formative evaluation of the
36 implementation of an upper limb stroke rehabilitation intervention in clinical practice: a
37
38
39
40
41
42 qualitative interview study. *Implement Sci.* 2014;9:1:90. doi: 10.1186/s13012-014-0090-3.
- 43 72. Connell LA, McMahon NE, Tyson SF, Watkins CL, Eng JJ. Case Series of a Knowledge
44 Translation Intervention to Increase Upper Limb Exercise in Stroke Rehabilitation. *Phys.*
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
73. Connell LA, McMahon NE, Tyson SF, Watkins CL, Eng JJ. Development of a behaviour change
intervention to increase upper limb exercise in stroke rehabilitation. *Implement Sci.* 2015;
10: 34. doi.org/10.1186/s13012-015-0223-3

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
74. Connell LA, McMahon NE, Tyson SF, Watkins CL, Eng JJ. Mechanisms of action of an implementation intervention in stroke rehabilitation: a qualitative interview study. *BMC Health Serv Res.* 2016; 1: 534.
75. Bernhardsson S, Larsson ME. Does a tailored guideline implementation strategy have an impact on clinical physiotherapy practice? A nonrandomized controlled study. *J Eval Clin Pract.* 2018; e-publication ahead of print: doi: 10.1111/jep.12958
76. Sezier A, Mudge S, Kayes S, Kersten P, Payne D, Harwood M, et al. Development of a toolkit to enhance care processes for people with a long-term neurological condition: a qualitative descriptive study. *BMJ Open.* 2018; 8: e022038. Doi: 10.1136/bmjopen-2018-022038.
77. Harvey, G. and A. Kitson, PARIHS revisited: from heuristic to integrated framework for the successful implementation of knowledge into practice. *Implement Sci.* 2015;11:1:33.
78. Michie S, Van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci.* 2011; 6: 42. <https://doi.org/10.1186/1748-5908-6-42>
79. Barwick M. Checklist to Assess Organizational Readiness (CARI) for EIP Implementation. 2011; Hospital for Sick Children, Toronto.
80. Rapley T, Girling M, Mair F, Murray E, Treweek S, McColl E et al. Improving the normalization of complex interventions: part 1 - development of the NoMAD instrument for assessing implementation work based on normalization process theory (NPT). *BMC Medical Research Methodology.* 2018; 18: 133. doi.org/10.1186/s12874-018-0590-y
81. Huijg JM, Gebhardt WA, Dusseldorp E, Verheijdeb MW, van der Zouwe, Middelkoop BJC et al. Measuring determinants of implementation behavior: psychometric properties of a questionnaire based on the theoretical domains framework. *Implement Sci.* 2014; 9: 33. doi.org/10.1186/1748-5908-9-33.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

82. Alcalde-Rabanal JE, Becerril-Montekio VM, and Langlois EV. Evaluation of Communities of Practice performance developing implementation research to enhance maternal health decision-making in Mexico and Nicaragua. *Implement Sci.* 2018; 13: 41. doi.org/10.1186/s13012-018-0735-8.
83. Kinsella E and Caty M. Reflective practice for allied health: theory and applications *in* English LM (Ed), Adult education and health. 2012; University of Toronto Press, London.
84. Jones F and Bailey N, How can we train stroke practitioners about patient self-management? Description and evaluation of a pathway wide training programme. *European Journal for Person Centered Healthcare.* 2013; 1: 246-254.
85. Ward VL, House A, Hamer S. Knowledge brokering: the missing link in the evidence to action chain? *Evidence & policy: a journal of research, debate and practice.* 2009; 5: 267-279.
86. Ward VL, House A, and Hamer S. Knowledge brokering: exploring the process of transferring knowledge into action. *BMC Health Serv Res.* 2009; 9: 12. doi.org/10.1186/1472-6963-9-1

Table 1. Case studies illustrating implementation approaches in rehabilitation

Case study 1: Evaluating the benefits of Early Supported Discharge after stroke in real world settings

Aims: Although national stroke guidelines in many countries recommend the provision of ESD, based on randomised controlled trial (RCT) evidence [67], it is unclear if the benefits seen in RCT conditions are still apparent when ESD is implemented in practice. This series of studies aimed to examine if Early Supported Discharge (ESD) is still beneficial when operating in the unpredictable and complex context of frontline healthcare provision.

Methods: A qualitative evaluation of effectiveness of two ESD services, that had adopted defined evidence based core components, and had made context-dependent modifications. Damschroder's Consolidated Framework for Implementation Research (CFIR) and Graham's Knowledge to Action model offered theoretical frameworks with which to guide the research [21] [68]. First, the aim was to identify essential characteristics ('core components') that theory suggests need to be implemented for ESD to be effective in practice (knowledge creation phase) [69]. Then the aim was to investigate the context in which ESD services were operating. This enabled investigation of the 'adaptable periphery' – adaptable elements, structures or systems that may need to be modified for the intervention to be embedded successfully in practice [34] [70]. Improvement activities were designed and delivered to support and sustain evidence based ESD delivery, using prospective comparative study methods.

Findings: Studies demonstrated that ESD services reduced length of hospital stay and accelerated stroke patient recovery, showing that benefits were achievable in a real world setting [58].

Case study 2: improving upper limb exercise provided to people with stroke in inpatient rehabilitation

Aims: To develop and implement a clinically feasible upper limb exercise programme after stroke. This study evaluated how RCT-informed upper limb intervention was used in real world clinical practice [71] [28] using implementation theories.

Methods: Interviews with therapists already using the intervention in practice:

- Interview guide (how therapists found out about intervention, processes used to implement intervention, how therapists used intervention) informed by Normalisation Process Theory (NPT)
- CFIR used to guide data analysis of factors influencing implementation and use of intervention
- Survey design (exploring therapists' opinions of using intervention) informed by CFIR to elicit information about intervention characteristics and characteristics of therapists

Findings: The studies identified factors that influenced implementation, and also highlighted challenges with fidelity when the intervention was used in the real world. We then developed and evaluated a behaviour change intervention to increase upper limb exercise in 3 stroke rehabilitation units:

- Behaviour change wheel guided selection of behaviour change strategies [72, 73]
- Theoretical Domains Framework guided analysis of mechanisms of action to explain how behaviour change intervention produced change [74]

This case series is an example of how a rigorous research-informed knowledge translation process resulted in practice change.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Case study 3: Implementation of clinical practice guidelines in primary care physiotherapy

Aims: To develop regional physiotherapy clinical practice guidelines for low back pain, neck pain and subacromial pain, and to develop a theory- and evidence-based strategy for their implementation, and evaluated the implementation in a prospective, non-randomised, controlled trial [14].

Methods: The strategy was based on the Grol and Wensing implementation of change model [46], which is a synthesis of many different behaviour change theories that are relevant for implementation research. The theories on which the model is based relate to individual professionals, as well as to the influence of social interaction and organisational and economic context. The model, comprising five steps, was adapted to the clinical and organisational context of primary care physiotherapy in the region of Västra Götaland in Sweden. As part of this model, determinants of guideline use were investigated in a baseline survey [11], and the implementation strategy was tailored to address these determinants. The strategy comprised several components, the core component being an educational seminar in which the guidelines were presented and discussed. This seminar was conducted in physiotherapy clinics across half the region of Västra Götaland, while the other half served as control group. Effects of guideline implementation were evaluated in a post-implementation survey, which showed that the tailored, multi-component guideline implementation significantly increased awareness of, knowledge of, and access to guidelines [14].

Findings: Use of guidelines was significantly affected among those who attended the implementation seminar. However, the impact of the guideline implementation on clinical practice was very limited, due to practice being in line with evidence already before the guidelines were implemented [75].

Case study 4: Implementation of a Living Well Toolkit into diverse rehabilitation settings

Aims: We aimed to develop, and put into practice, a Living Well Toolkit to support key processes for people with long-term neurological conditions (person-centred communication, building self-management skills, and coordination of long-term care).

Methods: We employed a participatory informed design approach, premised on the notion that understanding and valuing healthcare users' experiences in addition to engaging key stakeholders in the development of the toolkit would be fundamental to its success [48]. We explored perspectives of people with a long-term neurological condition, family, clinicians and other key stakeholders on three key processes. We developed themes from this data, which we used to develop the content of the toolkit [76]. The toolkit was then introduced in four rehabilitation settings. Individual and focus group interviews were carried out with patients (n=11) and clinicians (n=14). Data were analysed using NPT as a framework to inform revisions to the toolkit and future implementation processes.

Findings: There was widespread support for the principles underpinning the toolkit. However, it was less clear how it was envisaged the toolkit could support those practices (NPT: coherence) impacting buy-in from clinicians (NPT: cognitive participation). The built-in flexibility regarding who, how and to what extent the toolkit can be used made it difficult to operationalise in practice (NPT: collective action). Use of the Toolkit appeared more likely when it was perceived to augment existing processes (e.g. goal setting) rather than detract from 'doing' therapy (NPT: collective action, reflexive monitoring).

Table 2. Examples of tools related to implementation processes

Stage of Implementation	Key processes	Examples of tools that could support these processes	Pragmatic considerations for research and practice
Planning	Understanding context (with intended users)	Promoting Action on Research in Health Services (PARIHS) [77] Consolidated Framework for Implementation Research (CFIR) [21]	Partnering with intended users to: <ul style="list-style-type: none"> • identify clinical priorities • optimally tailor to context • identify and minimise implementation barriers • revise implementation plan and/or intervention as needed to optimise implementation • ensure you work <i>with</i> the team/ staff who will be delivering the intervention • consider what works for them in terms of timing and location of meetings • ensure 'buy-in' from all levels
	Understanding the intervention that is to be implemented and what behaviours need to change	Behaviour Change Wheel (BCW) [78] Checklist to Assess <i>Organisational</i> Readiness for evidence informed practice Intervention (CARI) [79] NOMAD: survey instrument for assessing implementation processes from the perspective of staff involved in implementation work. [80]	
	Identifying possible barriers and facilitators to change	Determinants of Implementation Behaviour Questionnaire [81]	
Influencing practice	Critical reflection on practice	Communities of practice [82-84]	<ul style="list-style-type: none"> • Reflective practices can include peer support, peer review, case reflections, supervision, facilitated group discussion, self-reflection • Clinicians need opportunity to practice new skills; role play with peers can be a first step, however recognise that practice with clients is also necessary; needs to be integrated into reflection and refinement and more practice • Knowledge broker needs to be respected by
	Support and encouragement	Knowledge brokerage or similar - boundary spanners, implementation champions, etc. [85, 86]	
	Structural supports	Structures and processes to support new way of working	

	Tailoring the intervention to context	Behaviour Change Techniques (selection of most appropriate tools will be context dependent, hence the planning phase being so important)	<p>clinicians</p> <ul style="list-style-type: none"> • If clinicians perceive an immediate benefit (to clients or themselves) from intervention, then more likely to implement • If clinicians perceive intervention fits with practice, then more likely to implement • Clinicians more likely to make change if perceive work environment (including colleagues) supportive of change
Monitoring and review	<p>Ongoing review and professional development</p> <p>Intervention refinement</p> <p>Sustainability</p>	<p>Recognise that implementation is a dynamic and ongoing process.</p> <p>Review needs to happen at:</p> <p>a) an individual level: individual clinicians reflecting in an ongoing way over knowledge use, challenges and tensions, and recognising need for ongoing professional development</p> <p>b) at broader implementation strategy level: ongoing evaluation and monitoring of knowledge use and behaviour change to determine if implementation intervention is being effective or whether refinements need to be made to optimise uptake</p> <p>NOMAD, linked to NPT, is an evaluative tool to focus on implementation, findings can be used to modify implementation process constructively.</p>	<ul style="list-style-type: none"> • Recognise change can be gradual, so build on change through cycle that includes monitoring and review • Acknowledge and celebrate any positive change • Ensure you have a plan for training/ involving new staff • If implementation not happening, it may be due to organisational or external influences- consider if now is the right time, or if you need to pause?

Figure 1. The Implementation Roadmap for Rehabilitation

