



Article

Nutritional quality, cost and availability of gluten free food in England

Soon, Jan Mei and Hopkins, Sarah

Available at <http://clock.uclan.ac.uk/29650/>

Soon, Jan Mei ORCID: 0000-0003-0488-1434 and Hopkins, Sarah (2019) Nutritional quality, cost and availability of gluten free food in England. British Food Journal, 121 (11). pp. 2867-2882. ISSN 0007-070X

It is advisable to refer to the publisher's version if you intend to cite from the work.
<http://dx.doi.org/10.1108/BFJ-09-2018-0607>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

Nutritional quality, cost and availability of gluten free food in England

Abstract

Purpose: Coeliac disease is a life-long condition requiring strict adherence to a gluten free diet. Due to wide claims of availability and lower costs of gluten free food (GFF) and Clinical Commissioning Groups (CCGs) in England needing to save costs, access to prescriptions for patients with coeliac disease (CD) is being limited in England. The purpose of this study is to investigate the availability and cost of GFF in an area where patients with coeliac disease have restricted access to prescriptions and to assess the nutritional composition of GFFs available in comparison with foods containing gluten.

Methodology: Eight food categories that were representative of a range of commonly purchased GFFs were selected. Availability and cost of cheapest and most expensive branded and non-branded GFFs and gluten containing equivalents were surveyed at physical stores (n=19) and online stores (n=8). The nutritional composition of some of the widely available GFFs identified (n=190) and comparable foods containing gluten (n=218) were calculated using MyFitnessPal.

Findings: None of the budget stores or corner shops surveyed stocked any of the surveyed cereal-based GFFs. Online stores had more availability than physical stores, however there was no significant difference in cost. Gluten free foods cost on average 2.18 times more than food containing gluten. When making nutritional comparisons with gluten-containing food, protein content was lower across 55% of GFF categories. There was significantly less sugar in gluten free (GF) brown bread, crackers, and wholegrain pasta compared with those containing gluten. Another main finding was GF ready-meals contained significantly less salt than ready-meals containing gluten.

Originality: Limited resources and perceived wide availability of gluten-free products resulted in reduced GF prescriptions to patients in England. The findings in this study revealed that there is no availability of cereal-based GFFs in budget stores, high cost and limited access to prescriptions can influence adherence to a gluten free diet and is most likely to affect patients from deprived groups. This study recommends that the prescription of gluten free food to patients with CD should be continued.

Keywords: availability; coeliac disease; cost; gluten-free; nutrition

36

37 **Introduction**

38 Coeliac disease (CD) is a systemic immune-mediated disorder activated by gluten and
39 related prolamines ingestion in genetically susceptible individuals. It is characterised by the
40 presence of CD specific antibodies and human leukocyte antigen (HLA)-DQ2 and HLA-DQ8
41 (Husby et al., 2012; Nelsen, 2002). The prevalence of CD in the Western World is estimated
42 at approximately 1-2% of the population and is considered an autoimmune disease affecting
43 both adults and children (Simón et al., 2017). In the UK, 0.24% of the population
44 (approximately 150,000 people) are clinically diagnosed with CD, however seroprevalence is
45 1 – 1.6%, and prevalence is increasing (Crooks, 2016). Gluten is an insoluble storage
46 protein found in wheat, rye, barley and triticale, and is made up of peptides gliadin and
47 glutenin (Wieser, 2007). These peptides are used in many foods to give them structure and
48 improve palatability (Gobbeti and Gänzle, 2013), however, when a patient with coeliac
49 disease ingests gluten, an immune reaction promotes secretion of high volumes of
50 proinflammatory cytokines and inflammatory mediators which attack the intestinal villi,
51 causing it to become inflamed (Mazzarella, 2015). Inflammation reduces the surface area of
52 the villi making it difficult to effectively absorb nutrients (Buttriss, 2002). Left untreated, CD
53 can result in villous atrophy, where the intestinal villi have completely eroded causing
54 malabsorption (Theethira et al., 2014). Study found that 43% of CD patients exhibited
55 persistent villous atrophy on a follow up biopsy, which was caused by non-adherence to a
56 gluten-free (GF) diet, thought to be a consequence of social differences due to access/or
57 education (Lebwohl et al., 2014). The only treatment available for individuals suffering with
58 CD or associated complications is to follow a strict life-long, GF diet. This consists of foods
59 that are naturally GF, and manufactured foods such as bread and pasta where substitutes
60 replace wheat (e.g. corn, rice or potato starch) (Ciacci et al., 2015). Products containing
61 less than 20ppm are considered GF by the Codex Alimentarius (FAO/WHO, 2015). Initiation
62 of the GF diet has been found to restore integrity of the intestinal wall, and improve
63 symptoms and deficiencies of CD, however, certain nutritional limitations exist (Barone et
64 al., 2016; Theethira et al., 2014).

65

66 Adhering to a GF diet is difficult and takes a lot of self-control. Other factors influencing
67 adherence include availability, cost, nutritional quality and access to gluten-free food (GFF)
68 on prescription (MacCulloch and Rashid, 2014; Hall et al., 2013). It has previously been
69 demonstrated that GFF cost on average 4.1 times more than food containing gluten (FCG)
70 (Burden et al., 2015). Missbach et al. (2015) observed that GF foods were significantly

71 higher in cost than similar foods containing gluten (range= 205% - 267%) in Austria.
72 Previous studies have also demonstrated limited availability of GF foods, with no availability
73 of cereal-based GF staples in budget stores and corner shops. A persistent finding is the
74 higher cost of GFFs when making comparisons with FCG (Singh and Whelan, 2011; do
75 Nascimento et al., 2014; Fry et al., 2018). The cost of food is an important factor which
76 influences food choice (Lennernäs et al., 1997). The socioeconomic status of patients may
77 determine which stores they shop in (Ellaway and Macintyre, 2000), and so the lack of
78 availability in budget stores and corner shops may affect adherence to a GF diet. This may
79 particularly burden patients who have no or limited access to GF foods on prescription
80 (Muhammad et al., 2017).

81

82 Studies focussing on analysing the nutritional composition of GFFs have found they contain
83 low levels of essential micronutrients such as iron, folic acid, calcium, potassium, zinc and
84 magnesium (Thompson, 2000; Estévez et al., 2016; Missbach et al., 2015). High fat content
85 has been another observation. Studies have demonstrated that GF breads have significantly
86 more fat than breads containing gluten (Kulai and Rashid, 2014; Fry et al., 2018). Studies
87 have also reported lower levels of protein and fibre content when compared with similar
88 gluten containing foods (do Nascimento et al., 2014; Miranda et al., 2014; Missbach et al.,
89 2015. Meanwhile Fry et al. (2018) found fibre content to be significantly higher in GF items
90 such as white and brown bread whilst protein was found to be lower in GF products
91 compared to gluten containing foods.

92

93 Gluten-free foods have been available on prescription to patients with CD since the late
94 1960's (Department of Health, 2017), however, due to wider availability and reduced cost,
95 as of 5th December 2016, the National Health Service (NHS) made budget cuts, and the
96 Clinical Commissioning Groups (CCGs) in England changed their policies on GF prescriptions
97 (Coeliac UK, 2019a). CCGs are groups of general practices (GPs) that come together in an
98 area to commission the best services for patients and the population. CCGs buy services for
99 the local community from any local service provider that meets the NHS standards and cost
100 and aims to provide better care for patients (NHS, 2019). This resulted in 40% of 207 local
101 CCG's restricting or completely stopping GF prescriptions (Coeliac UK, 2019a). The rationale
102 of wider availability and lower costs of GF foods conflicts with evidence found in previous
103 research (Singh and Whelan, 2011; do Nascimento et al., 2014; Burden et al., 2015; Fry et
104 al., 2018; Hanci and Jeanes, 2018).

105

106 Before CCGs made policy changes, 90% of patients with CD relied on prescriptions for GFF
107 (Robins et al., 2008), this offered fundamental support to those on a low income, benefits,
108 and the elderly. Furthermore, Kinsey et al. (2008), found that prescription GFFs were higher
109 in non-starch polysaccharides (NSP), calcium and iron compared with commercially available
110 GF foods. However, since the studies conducted by Robins et al. and Kinsey et al. in 2008,
111 more recent studies have found a lack of availability of GF foods and higher costs associated
112 with such items. Therefore, the removal of GFF prescriptions is likely to affect the nutritional
113 adequacy of a patient with coeliac disease GF diet, which could lead to health complications.
114 A combination of restricting prescriptions, high cost of GFF and limited availability could
115 potentially influence adherence to a gluten free diet (Coeliac UK, 2019.b; Estévez et al.,
116 2016). Health risks associated with non-adherence include osteoporosis, osteopenia,
117 infertility, anaemia, vitamin D deficiency, increased risk of fractures and cancer (BDA, 2015;
118 Kinsey et al., 2008).

119

120 Stopping access to GFFs for primary care patients has recently been rejected by the
121 government, and the Department of Health and Social Care (2018) has decided to restrict
122 gluten-free prescribing to breads and flour mixes only (NHSCC, 2018). However, CCGs have
123 the ability to control their own policies and can ultimately decide how much food can be
124 prescribed, and who is entitled to it. Currently, patients with coeliac disease living in
125 Greater Preston still have no access to GF prescriptions and until regulations have been
126 developed and formally agreed by Parliament, the CCG for Greater Preston do not have to
127 make changes to current policy. With increasing prevalence of CD and restrictions to
128 prescriptions for patients with coeliac disease, the purpose of this study was to assess the
129 availability of GFF and compare cost and nutritional quality with FCG in an area where GF
130 prescribing is currently on hold and may be limited once new prescription regulations have
131 been implemented.

132

133 **Methodology**

134 *Store selection*

135 The area chosen for assessment was Greater Preston situated in North West England, as
136 this was one of the areas where prescriptions for GFF were withdrawn on 5th December
137 2016 (Coeliac UK, 2019a). Stores such as regular stores, quality stores, budget stores,
138 health food shops, convenience stores and corner shops, as previously categorised by Singh
139 and Whelan (2011) were visited. The availability and cost of branded and non-branded GFFs

140 were surveyed as were similar FCG at different stores after receiving consent from the store
141 managers.

142

143 *Food categories*

144 An itinerary of eight food items were chosen and were representative of a range of
145 commonly purchased GFFs (do Nascimento et al., 2014; Fry et al., 2018). Food items were
146 categorised as biscuits, breads, cakes, cereals (breakfast), crackers, flour, pasta and ready-
147 meals were analysed for availability and cost. Additionally, similar categories of FCG were
148 also analysed for cost, so a direct comparison could be made.

149

150 *Survey*

151 Stores were surveyed between November 2017 and January 2018. Availability was
152 measured by counting how many different GF brands were stocked in each of the eight
153 food categories in all stores visited, and a photograph of the same stock was captured on
154 camera by smart phone. The cost of the cheapest and most expensive item was recorded,
155 and for items with wide availability, the cost of the cheapest and most expensive branded
156 and none branded foods was recorded. The same method was adopted for similar FCG and
157 all costs were captured by camera using a smart phone. Package weight was not used as
158 covariate as food comparisons may be weighted differently hence costs were calculated as
159 pence (£) per 100g.

160

161 *Online stores*

162 Appropriate keywords were entered onto a google search engine to find stores with
163 availability of GFF that delivered to homes in Greater Preston. Nine stores were selected
164 and represented online regular stores, quality stores, health food stores and stores
165 specialising in GF foods. This was to achieve a comparable analysis of online store
166 availability and cost. The stores selected were those that landed on the first google result
167 page, as this is where 95% of websites receive most visits (Brafton, 2013). The eight GFFs
168 categories were separately entered onto the search bar of each online store and listed foods
169 available were noted. The cost in pence per 100g of the cheapest and most expensive
170 foods was recorded, and for items with wide availability, the cost of the cheapest and most
171 expensive branded and none branded foods were recorded (Hanci and Jeanes, 2018).

172

173 *Nutritional content*

174 Nutritional quality of GFFs and FCG were calculated using MyFitnessPal (MyFitnessPal.com,
175 2015). This involved foods from the eight food categories, however bread was separated
176 into white and brown, flour into white and wholemeal, and pasta into white and wholegrain
177 so that comparisons could be made with previous research (Miranda et al., 2014; Fry et al.,
178 2018). Food items were entered into MyFitnessPal search bar and products were selected if
179 listed. This app was selected as the nutritional data available from MyFitnessPal contained
180 information obtained from product labels which should be declared according to European
181 Union legislation 1169/2011 (EUFIC, 2011). These mandatory nutrients (per 100 grams)
182 included energy, protein, carbohydrates, sugars, total fat, saturated fat and salt, which were
183 systematically collected for GFFs and similar FCG. Additionally, fibre content was obtained
184 where data was available and any food items, which could not be found, were excluded
185 from analysis.

186

187 *Statistical Analysis*

188 Statistical analyses were performed using SPSS version 24 (IBM Corp, 2016). Normal Q-Q
189 plots and Shapiro-Wilk test of normality ($p > 0.05$) were used to assess if data deviate from
190 a normal distribution. Independent samples t-tests were used to compare the availability
191 and cost of GFFs at online stores and stores located around Greater Preston, and to
192 compare the average nutritional content of GFF and similar FCG across all product
193 categories. Equal variances were assumed if the Levene's test for homogeneity of variance
194 was non-significant ($p > .05$). A one-way analysis of variance (ANOVA) was performed to
195 observe differences in the cost and availability of GFF compared with FCG across all store
196 categories and independent samples *t*-tests were applied to make cost comparisons
197 between two store categories. $p \leq 0.05$ was considered statistically significant.

198

199 **Results**

200 A total of 27 stores were investigated, of which 19 were located around Greater Preston and
201 eight online stores that delivered within the Greater Preston area. Out of the 27 stores
202 visited, seven (25.9%) stocked no GFF, with the remaining 20 (74.1%) stocking at least four
203 categories of GFF. Across the seven different store types, 2,780 GF foods were identified
204 (crossovers occurred), 1,800 of which were available online, and 980 stocked in physical
205 stores around Greater Preston). The number of GFFs available from each food category per
206 store type is shown in Table 1. A one-way ANOVA revealed food category had a significant
207 effect on availability except for flour ($p = 0.156$). Store types stocking one or more GFFs

208 from every food category included online, regular, quality and health food shops (100%),
209 with convenience stores stocking on average 87.5%. In general, budget stores and corner
210 shops stocked no cereal-based GFFs, whilst online stores were found to stock the widest
211 range. Of the physical stores, regular and quality stores had the highest availability and all
212 stores stocked at least one GFF item from each food category with the exception of
213 convenience stores, that carried no stock of GF ready meals.

214

215 Insert Table 1 here

216

217 A one-way ANOVA revealed that store type had a significant effect on cost ($f(4,1008) =$
218 $12.40, p < .001, \eta^2 = .047$). To explore the location of significance, a Tukey's HSD *post hoc*
219 was conducted, which revealed some GFFs significantly differed in cost depending on type
220 of store (Table 2). For example, the average cost of cereals was significantly more
221 expensive in health food shops than regular stores (1.10 and 0.67 pence per 100g
222 respectively, $p < .001$). The cost of 1,047 GFFs within the eight food categories (cheapest
223 and most expensive items) were compared with 750 similar foods containing gluten. Only
224 items with A one-way ANOVA revealed cost had a significant effect on whether or not food
225 contained gluten ($f(1,1781) = 870.299, p < .001, \eta^2 = .328$). Independent *t*-tests showed
226 that all GFFs were significantly more expensive than FCG (Table 2), costing on average 2.18
227 times more, with bread representing the highest price difference (62.2%) and ready meals
228 the lowest price difference (38.89%). GF bread costs an average 2.65 times more than GC
229 bread products.

230

231 Insert Tale 2 here

232

233 A comparison of energy and macronutrients of widely available forms of GFFs (n=190) and
234 a gluten containing equivalent (n=218) from each food category was analysed using
235 independent t-tests and results are displayed in Table 3. Across all food categories, the
236 nutritional content of GFF compared with a gluten free equivalent did not significantly differ
237 between biscuits, cakes and cereals ($p > 0.05$). The only significant difference in energy was
238 observed in flour, where GF white flour contained significantly more than white flour
239 containing gluten ($p = 0.05$). In the breads, crackers, flour and white pasta categories, the
240 average protein content in GF foods was significantly lower than their gluten containing
241 equivalents ($p < 0.05$). When compared with FCG, gluten free flours and white pasta

242 contained significantly more carbohydrates ($p < 0.05$), although less carbohydrates were
243 observed in GF white bread ($p = 0.001$).

244

245 There was significantly less sugar in GF brown bread, crackers, and wholegrain pasta
246 compared with those containing gluten ($p < 0.05$). Another finding was GF ready-meals
247 contained significantly less salt than ready-meals containing gluten ($p = 0.002$). When
248 comparing the gluten containing bread categories, more total fat was found in both GF
249 brown and white bread ($p = .02$ and $p = .007$ respectively). Interestingly, GF white bread was
250 found to contain significantly more fibre than white bread containing gluten (7.15 ± 2.13 and
251 2.72 ± 1.46 g/100g respectively, $p < .001$), however, white GF pasta contained less fibre
252 ($p < .001$). In general, protein content was found to be significantly lower in 55% of GFF
253 compared with FCG.

254

255 Insert Table 3 here

256

257 **Discussion**

258

259 *Availability of gluten-free food*

260 The present study identified 2,780 GFFs within eight food categories, across seven store
261 types, demonstrating availability of a wide range of products, however availability varied
262 significantly between store type ($p < .001$). This could be attributable to the difference in
263 store sizes which was not factored into the analysis. Overall, online stores offered the
264 widest range of GFF compared to physical stores, which was also an observation made by
265 Burden et al. (2015) and could be due to several reasons. Online stores run their operations
266 from warehouses which have the capacity to carry large volumes of stock, their profitability
267 margins are likely to be high due to lower running costs, their logistics are very efficient, for
268 example fast turnaround between order time and delivery time; and marketing campaigns
269 reach out to new customers (Fernie et al., 2010). These may all contribute to increased
270 availability to meet consumer demand. The wider range of GFF available online offers
271 consumers more choice and is a service which may help CD patients who have limited
272 mobility. However, access is only available to those who are computer literate, have access
273 to the internet and can afford delivery costs (Webb, 2013).

274

275 Of all physical stores surveyed, regular stores carried the most stock of cereal-based GFF
276 followed by quality stores, health food shops and convenience stores, with budget stores

277 and corner shops stocking none. This is similar with previous findings, however the present
278 study observed a wider choice, for example, Singh and Whelan. (2011), reported less
279 availability of GF breads in regular supermarkets than the present study. Furthermore,
280 Burden et al. (2015), observed the highest availability of GFF in quality supermarkets, and
281 although there was more availability in quality stores in the present study, the highest
282 availability was found in regular stores. Similarly, Hanci and Jeanes (2018) reported that GF
283 food availability has increased in premium and online stores. This could be due to
284 manufacturers responding to increasing demand of GFF due to the increasing prevalence of
285 CD and associated complications. There has also been a growth in supermarkets own-label
286 GFFs (Smith, 2010), which may offer further explanation as to why there is a wider range of
287 GFF available than previously reported. With CD on the increase, future studies may see an
288 increase in availability, particularly in stores with own-brand labels.

289

290 Although the present study supports a wider range of GF items available than in recent
291 years, it also demonstrates availability is poor in budget stores and corner shops, a
292 consistent finding in previous studies (do Nascimento et al., 2014; Hanci and Jeanes, 2018;
293 Singh and Whelan, 2011; Burden et al., 2015). Budget stores may choose not to stock
294 cereal-based GFF because offering them for sale at prices their shoppers are accustomed to
295 may not be profitable. Gluten-free ingredients are expensive, and there are additional costs
296 incurred during manufacturing such as dedicated GF factories, dedicated areas within
297 factories, and investment in new technologies (Coeliac UK, 2016). Therefore, suppliers of
298 GF foods may not have the flexibility to negotiate low prices and until ingredient and
299 production costs of GFFs reduce, it is unlikely that budget stores will stock them.

300

301 A wider range of GFFs offer patients with coeliac disease more options, allowing them to
302 choose food based on personal preferences which can contribute to adhering to a GF diet
303 (Wright, 2017). However, this only applies to patients with higher disposable expenditures.
304 Gluten-free food options remain limited for patients shopping on a budget, or are unable to
305 access larger stores, or have no internet access. This study has demonstrated there are
306 more cereal-based GFF items available than previously reported, which supports the NHS's
307 principle of wider availability of GFF, however, availability remains poor in budget stores and
308 corner shops, and until all stores stock 100% of common foods that are GF then availability
309 should be considered limited.

310

311 *Cost of gluten-free food*

312 The present study found GFF within eight food categories presented significantly higher
313 prices than FCG, costing on average 2.18 times more (118% price difference). This could
314 be attributable to the use of more expensive grains in replacement of wheat (Jnawali et al.,
315 2016), or the usage of innovative processing techniques to improve the nutritional quality
316 and organoleptic properties of GFF (Padalino et al., 2016). Therefore, it is understandable
317 that they are costlier than FCG.

318

319 A difference in cost of GFF and FCG has been a consistent finding, however the price
320 difference in the present study is lower than previously reported (Singh and Whelan, 2011;
321 Lee et al., 2007; Stevens and Rashid, 2008). In the UK, Burden et al. (2015) found GFF cost
322 on average 4.1 times more than FCG. More recently, Fry et al. (2018) surveyed websites of
323 four leading UK supermarkets as well as a leading online food retailer and found that 10
324 common GFFs were on average 159% more expensive than FCG. The high cost of GFF is a
325 key influential factor for adherence to a GF diet (Muhammad et al., 2017). This can be
326 supported by responses from a patient survey which revealed that 82% of 1,000 primary
327 care patients revealed it was harder to manage their diet due to GF foods costing more than
328 regular foods (BSNA, 2013). Therefore, due to the CCG ceasing prescriptions for GFF in
329 Greater Preston, local patients with coeliac disease who have heavily relied on GF
330 prescriptions may find it difficult to adhere to a gluten free diet due to affordability.
331 Furthermore, patients have been found to be more likely to adhere to a GF diet when
332 obtaining their food on prescription (Hall et al., 2013).

333

334 *Nutritional content of gluten-free food*

335 The present data shows that 75% of GFF across all categories displayed differences in
336 nutritional composition compared with FCG. A key finding was a significantly lower protein
337 content in GF bread, crackers, flour and pasta compared with gluten-containing equivalents,
338 with bread and flours having more than two times less protein. Gluten is an important
339 constituent for food manufacturers, as its major storage proteins gliadin and glutenin gives
340 food elasticity and structure, producing an end-product which is visually appealing and
341 highly palatable (Gobbetti and Gänzle, 2013). Manufacturers use a wheat substitute, such
342 as corn and rice flour and add proteins, gums and emulsifiers to substitute gluten to
343 formulate GFF to represent FCG (Padalino et al., 2016). Rice flour is an inexpensive widely
344 available ingredient which is commonly used to formulate GF breads (Sandri et al., 2017),

345 however the protein content in rice flour is almost 50% less than wheat flour (Lásztity,
346 1983).

347

348 Lower protein content in GFF compared with FCG has been a consistent finding in previous
349 research (Miranda et al., 2014; Fry et al., 2018; Wu et al., 2015; do Nascimento et al.,
350 2014; Missbach et al., 2015). This has resulted in food technologists investigating the re-
351 formulation of GFF with pseudo-cereals, which naturally contain high amounts of essential
352 amino acids (Drzewieki et al., 2003). Gambus et al. (2002) found that replacing corn starch
353 with amaranth flour in the re-formulation of GF bread increased its protein content by 32%
354 with no effect on its sensory quality. A more recent study by Alvarez-Jubete et al. (2009),
355 found that formulating GF bread with 50% rice flour or 50% potato starch with amaranth,
356 buckwheat or quinoa flour increased protein, fibre, calcium, iron, vitamin E and polyphenol
357 contents. Therefore, the use of pseudo-cereals in the formulation of GF foods will enhance
358 its protein content and overall quality, providing an improved product for the consumer.
359 Although some manufacturers have made advances in improving quality of GFFs (Wang et
360 al., 2017), their availability on the market may be limited. This could be due to the extra
361 cost of ingredients and the advanced technology needed in their re-formulation. Also,
362 certain food manufacturers may choose least expensive ingredients to keep production costs
363 at a minimum to keep costs down for consumers.

364

365 Carbohydrates are an important source of energy, and over a third of the diet should consist
366 of starchy foods (NHS Choices, 2017). The present study found that flours and white pasta
367 had significantly more carbohydrates than their gluten-containing equivalents. This could be
368 due to the poor protein content in GF flours that are also used to produce pasta. When
369 making comparisons with breads however, a significantly lower carbohydrate content was
370 found in GF white bread compared with gluten-containing equivalents. This is an interesting
371 finding, considering starch ingredients such as corn, cassava, potato and rice are commonly
372 used as main ingredients or partial substitutes in GF breads. Additionally, gel forming
373 starches such as pregelatinized starches and gums act as stabilising agents to achieve
374 structure, texture and crumb in the absence of gliadin and glutenin (Horstmann et al.,
375 2017). Previous studies have observed no significant differences in GF breads when
376 comparing them with breads containing gluten (CG) (Kulai and Rashid, 2014; do Nascimento
377 et al., 2013). This further supports the notion of application of better-quality ingredients
378 during reformulation of GF white bread.

379

380 The present study found that ready-meals containing gluten had over 50% more salt than
381 GF ready-meals. This could be due to GFF manufacturers attempting to meet salt targets
382 set by the Department of Health (DH, 2014). In 2004, the World Health Organisation
383 (WHO) reported that people were consuming too much salt (9-12g/day) and set a global
384 goal to reduce salt intake to <5g/day by 2025 (WHO, 2016). This prompted governments to
385 design salt-reduction initiatives, and the food industry was asked to voluntarily reformulate
386 salt content in foods as an attempt to improve public health (DH, 2014; FSA, 2008; WHO,
387 2004). As the production of GF ready meals is relatively recent, manufacturers can
388 formulate foods with less salt without the consumer being affected. This may offer further
389 explanation as to why GF ready-meals contained less salt compared to GC ready-meals.

390

391 The present study found that GF white pasta contained significantly less fibre than white
392 pasta CG. This could be due to the removal of cereals, which can leave food with a high
393 starch and low fibre content (Saturni et al., 2010). However, a key finding in the current
394 study was significantly more fibre in GF white bread than white breads CG. This could be an
395 outcome of the re-formulation of GF breads with pseudo-cereals to improve overall quality.
396 For example, when Gambus et al. (2002), replaced corn starch with amaranth flour in GF
397 bread, a 152% increase in fibre was observed. Furthermore, buckwheat flour has been
398 found to improve nutritional quality of bread as well as improve its viscosity due to its high
399 fibre content (Mariotti et al., 2013). Similar findings were reported by Fry et al. (2018),
400 however, do Nascimento et al. (2014), found significantly lower fibre content in GF bread
401 compared with bread CG. Therefore, based on results in the present study and by Fry et al.
402 (2018), higher fibre content in UK GF white breads may be attributable to several factors;
403 more manufacturers are re-formulating their GF breads to achieve a better-quality product;
404 and GFF manufacturers may be responding to new government dietary recommendations
405 published in 2016 regarding daily increase of fibre from 18g a day to 30g a day (PHE, 2016).
406 Total fat content of GF breads in the present study was significantly higher than bread CG,
407 which supports previous findings (Fry et al., 2018; Kulai and Rashid, 2014). This could be
408 attributable to the addition of fats during formulation to create consistency, increase volume
409 and reduce staling (Mancebo et al., 2017).

410

411 **Limitations**

412

413 There were several limitations for the present study. Firstly, nutritional composition of food
414 was not obtained via direct chemical analysis. Instead, data was obtained from

415 'MyFitnessPal', which is an application that obtains nutritional information from
416 manufacturers' labels. Only selected GFF (n=190) and FCG (n=218) products were
417 analysed. Secondly, a comparison of micronutrients in GF foods and foods containing gluten
418 could not be investigated using the method adopted due to micronutrient contents on labels
419 not being a mandatory requirement under current (EU) legislation 1169/2011 (FSA, n.d.).
420 Direct chemical analysis is the only viable way to acquire micronutrient composition of foods.
421 Thirdly, the price and nutritional content of all food analysed was based on g/100g and
422 serving sizes have not been factored in to the analysis. Fourth, surveillance of foods in
423 physical stores was based on observations of items which were displayed on the shelves,
424 and any foods which had not been replenished may have been missed. The authors also
425 recommend that the full range of GF breads and the price difference between types of GF
426 bread in comparison with regular equivalents should be further studied. Finally, the sample
427 size of stores and surveyed GFFs were limited to 27 stores within the Greater Preston area.

428

429 **Conclusion**

430

431 The present study has shown wider availability of cereal-based GFF, however access
432 remains poor in budget stores and corner shops which are reported to be frequented by
433 people on lower incomes. The price of GFF remains significantly high compared with FCG,
434 which has been shown to affect some patients' adherence to a GF diet. Nutritional quality
435 has improved based on previous research, and advantages were observed in fibre content,
436 however, protein quality was significantly lower in 55% of GFFs analysed, and a higher fat
437 content was found in breads. With increasing prevalence of CD, a demand for more GFFs
438 may see manufacturers reduce their prices further as the market becomes more
439 competitive. This is quite promising, however, ingredients equal in quality to FCG may be
440 compromised to keep production costs down which could result in larger disparities in
441 nutritional quality. In view of restrictions to GFF prescriptions, further longitudinal studies
442 are required to evaluate the true impact that limiting access to GFF will have on health and
443 adherence across all population groups. Therefore, full prescriptions for GF food should still
444 be available for primary care patients with CD and should remain until availability,
445 accessibility, price and nutritional content of GF food is improved.

446

447 **References**

448 Alvarez-Jubete, L.A., Arendt, E.K. and Gallagher, E. (2010), "Nutritive value of pseudocereals
449 and their increasing use as functional gluten-free ingredients", *Trends in Food Science &*
450 *Technology*, Vol. 21, pp. 106-113.

451
452 Barone, M., Della Valle, N., Rosania, R., Facciorusso, A., Trotta, A., Cantatore, F.P., Falco,
453 S., Pignatiello, S., Viggiani, M.T., Amorusso, A., De Filippis, R., Di Leo, A. and Francavilla, R.
454 (2016), "A comparison of the nutritional status between adult celiac patients on a long-term,
455 strictly gluten-free diet and healthy subjects", *European Journal of Clinical Nutrition*, Vol. 70,
456 No. 1, pp. 23-27.

457
458 Brafton (2013). *95 Percent of web traffic goes to sites on page 1 of Google SERPS [Study]*.
459 Available at: [https://www.brafton.com/news/95-percent-of-web-traffic-goes-to-sites-on-](https://www.brafton.com/news/95-percent-of-web-traffic-goes-to-sites-on-page-1-of-google-serps-study/)
460 [page-1-of-google-serps-study/](https://www.brafton.com/news/95-percent-of-web-traffic-goes-to-sites-on-page-1-of-google-serps-study/) [Accessed 11 April 2018].

461
462 British Dietetics Association (BDA) (2015), *Policy Statement: Gluten Free Food on*
463 *Prescription*. Available at:
464 [https://www.bda.uk.com/improvinghealth/healthprofessionals/policy_statement_gluten_free](https://www.bda.uk.com/improvinghealth/healthprofessionals/policy_statement_gluten_free_food_on_prescription)
465 [_food_on_prescription](https://www.bda.uk.com/improvinghealth/healthprofessionals/policy_statement_gluten_free_food_on_prescription) [Accessed 17 April 2018].

466
467 BSNA (2013). *The impact of the restrictions to gluten-free prescription foods on people with*
468 *Coeliac Disease: Report of Findings*. British Specialist Nutrition Association. Available at:
469 [https://bsna.co.uk/uploads/files/Report-on-Impact-of-Restrictions-to-Gluten-Free-](https://bsna.co.uk/uploads/files/Report-on-Impact-of-Restrictions-to-Gluten-Free-Prescription-Foods.pdf)
470 [Prescription-Foods.pdf](https://bsna.co.uk/uploads/files/Report-on-Impact-of-Restrictions-to-Gluten-Free-Prescription-Foods.pdf) [Accessed 15 April 2018].

471
472 Burden, M., Mooney, P. D., Blanshard, R. J., White, W. L., Cambray-Deakin, D. R and
473 Sanders, D. S. (2015), "Cost and availability of gluten-free food in the UK: in store and
474 online", *Postgraduate Medical Journal*, Vol. 91, No. 1081, pp. 622-626.

475
476 Buttriss, J. (2002). *Adverse Reactions to Food*. Oxford: Blackwell Science.

477
478 Ciacci, C., Ciclitira, P., Hadjivassiliou, M., Kaukinen, K., Ludvigsson, J.F., McGough, N.,
479 Sanders, D. S., Woodward, J., Leonard, J. N. and Swift, G. L. (2015), "The gluten-free diet
480 and its current application in coeliac disease and dermatitis herpetiformis. *United European*
481 *Gastroenterology Journal*, Vol. 3, No. 2, pp. 121-135.

482
483 Coeliac UK, 2016. *Campaigning on the cost of gluten-free*. [online] Available at:
484 <https://www.coeliac.org.uk/about-us/news/campaigning-on-the-cost-of-gluten-free/> >
485 [Accessed 14 April 2018].

486
487 Coeliac UK. (2019a). *Prescription policies*. Available at: [https://www.coeliac.org.uk/gluten-](https://www.coeliac.org.uk/gluten-free-diet-and-lifestyle/prescriptions/prescription-policies/)
488 [free-diet-and-lifestyle/prescriptions/prescription-policies/](https://www.coeliac.org.uk/gluten-free-diet-and-lifestyle/prescriptions/prescription-policies/) [Accessed 16 July 2019].

489
490 Coeliac UK. (2019b). *Gluten-free product certification*. Available at:
491 <https://www.coeliac.org.uk/food-industry-professionals/the-crossed-grain-symbol/> [Accessed
492 16 July 2019].

493
494 Crooks, C. (2016). *Epidemiology of coeliac disease*. Available at: <https://www.coeliac.org.uk/>
495 [Accessed 28 March 2018].

496
497 Department of Health (DH), 2014. *Responsibility Deal Food Network – New salt targets: F9*
498 *Salt Reduction 2017 pledge & F10 Out of Home Salt reduction pledge*. Available at:
499 [http://webarchive.nationalarchives.gov.uk/20180201175712/https://responsibilitydeal.dh.go](http://webarchive.nationalarchives.gov.uk/20180201175712/https://responsibilitydeal.dh.gov.uk/responsibility-deal-food-network-new-salt-targets-f9-salt-reduction-2017-pledge-f10-out-of-home-salt-reduction-pledge/)
500 [v.uk/responsibility-deal-food-network-new-salt-targets-f9-salt-reduction-2017-pledge-f10-](http://webarchive.nationalarchives.gov.uk/20180201175712/https://responsibilitydeal.dh.gov.uk/responsibility-deal-food-network-new-salt-targets-f9-salt-reduction-2017-pledge-f10-out-of-home-salt-reduction-pledge/)
501 [out-of-home-salt-reduction-pledge/](http://webarchive.nationalarchives.gov.uk/20180201175712/https://responsibilitydeal.dh.gov.uk/responsibility-deal-food-network-new-salt-targets-f9-salt-reduction-2017-pledge-f10-out-of-home-salt-reduction-pledge/) [Accessed 16 April 2018].

502

503 Department of Health (2017). The availability of gluten free foods on prescription in primary
504 care. Available at:
505 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_d](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/604842/Gluten_free_foods_cons.pdf)
506 [ata/file/604842/Gluten_free_foods_cons.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/604842/Gluten_free_foods_cons.pdf) [Accessed 18 Sept 2018]
507
508 Department of Health and Social Care (2018). Report of responses following the public
509 consultation on gluten free prescribing. Availability of gluten free food on prescription in
510 primary care. Available at:
511 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_d](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/678181/report_of_responses_-_gluten_free_food_prescribing_consultation.pdf)
512 [ata/file/678181/report_of_responses_-_gluten_free_food_prescribing_consultation.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/678181/report_of_responses_-_gluten_free_food_prescribing_consultation.pdf)
513 [Accessed 16 July 2019].
514
515 do Nascimento, A. B., Fiates, G. M. R., dos Anjos, A. and Teixeira, E. (2013), "Analysis of
516 ingredient lists of commercially available gluten-free and gluten-containing food products
517 using the text mining technique", *International Journal of Food Sciences and Nutrition*, Vol.
518 64, No. 2, pp. 217-222.
519
520 do Nascimento, A., Medeiros Rataichesk Fiates, G., dos Anjos, A. and Teixeira, E. (2014),
521 "Availability, cost and nutritional composition of GF products", *British Food Journal*, Vol. 116
522 No. 12, pp.1842-1852.
523
524 Drzewiecki, J., Delgado-Licon, E., Haruenkit, R., Pawelzik, E., Martin-Belloso, O., Park, Y.S.,
525 Jung, S.T., Trakhtenberg, S. and Gorinstein, S. (2003), "Identification and differences of
526 total proteins and their soluble fractions in some pseudocereals based on electrophoretic
527 patterns", *Journal of Agricultural and Food Chemistry*, Vol. 51, No. 26, pp. 7798-7804.
528
529 Ellaway, A. and Macintyre, S. (2000), "Shopping for food in socially contrasting localities",
530 *British Food Journal*, Vol. 102, No. 1, pp. 52-59.
531
532 Estévez, V., Ayala, J., Vespa, C. and Araya, M. (2016), "The gluten-free basic food basket: a
533 problem of availability, cost and nutritional composition", *European Journal of Clinical*
534 *Nutrition*, Vol. 70, No. 10, pp. 1215-1217.
535
536 EUFIC (2011). Regulation (EU) no 1169/2011 of the European Parliament and of the council.
537 European Union Food Information Council. Available at: [http://eur-lex.europa.eu/legal-](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32011R1169&from=en)
538 [content/EN/TXT/PDF/?uri=CELEX:32011R1169&from=en](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32011R1169&from=en) [Accessed 11 April 2018].
539
540 Fernie, J., Sparks, L. and McKinnon, A. C. (2010), "Retail logistics in the UK: past, present
541 and future", *International Journal of Retail & Distribution Management*, Vol. 38, No. 11/12,
542 pp. 894-914.
543
544 FAO/WHO (2015). Standard for Foods for Special Dietary Use for Persons Intolerant to
545 Gluten: Codex stan 118-1979. Available at:
546 http://www.fao.org/input/download/standards/291/CXS_118e_2015.pdf [Accessed 19 April
547 2018].
548
549 FSA (n.d.) *European Food Information to Consumers Regulation No 1169/2011 (FIC)*. Food
550 Standards Agency. Available at:
551 <https://www.food.gov.uk/enforcement/regulation/fir/labelling> [Accessed 18 April 2018].
552

553 FSA (2008). *UK Salt Reduction Initiative*. Food Standards Agency. Available at:
554 <https://www.food.gov.uk/sites/default/files/multimedia/pdfs/saltreductioninitiatives.pdf>
555 [Accessed 18 April 2018].
556
557 Fry, L., Madden, A. M. and Fallaize, R. (2018), "An investigation into the nutritional
558 composition and cost of gluten-free versus regular food products in the UK", *Journal of*
559 *Human Nutrition and Dietetics*, Vol. 31, No. 1, pp.108-120.
560
561 Gambus, H., Gambus, F. and Sabat, R. (2002), "The research on quality improvement of
562 gluten-free bread by amaranthus flour addition", *Żywnosc*, Vol. 2, No. 31, pp. 99-112.
563
564 Gobbeti, M. and Gänzle, M., 2013. *Handbook on Sourdough Biotechnology*. New York:
565 Springer.
566
567 Hanci, O. and Jeanes, Y. M. (2018), "Are gluten-free food staples accessible to all patients
568 with coeliac disease?" *Frontline Gastroenterology*, Vol. 10, No. 3, pp. 222-228.
569
570 Hall, N. J., Rubin, G. P. and Charnock, A. (2013), "International and inadvertent non-
571 adherence in adult coeliac disease: A cross-sectional survey", *Appetite*, Vol. 68, pp. 56-62.
572
573 Horstmann, S. W., Lynch, K. M. and Arendt, E. K. (2017), "Starch characteristics linked to
574 gluten-free products", *Foods*, Vol. 6, No. 4, pp. 29.
575
576 IBM Corp (2016). IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp.
577
578 Husby, S., Koletzko, S., Korponay-Szabo, I. R., Mearin, M. L., Phillips, A., Shamir, R. et al.
579 (2012), "European Society for Pediatric Gastroenterology, Hepatology, and Nutrition
580 Guidelines for the diagnosis of coeliac disease", *Journal of Pediatric Gastroenterology and*
581 *Nutrition*, Vol. 54, No. 1, pp. 572-573.
582 Jnawali, P., Kumar, V. and Tanwar, B. (2016), "Celiac disease: Overview and considerations
583 for development of gluten-free foods", *Food Science and Human Wellness*, Vol. 5, No. 4, pp.
584 169-176.
585
586 Kinsey, L., Burden, S. T. and Bannerman, E. (2008), "A dietary survey to determine if
587 patients with coeliac disease are meeting current healthy eating guidelines and how their
588 diet compares to that of the British general population", *European Journal of Clinical*
589 *Nutrition*, Vol. 62, No. 11, pp. 1333-1342.
590
591 Kulai, T. and Rashid, M. (2014), "Assessment of nutritional adequacy of packaged gluten-
592 free food products", *Canadian Journal of Dietetic Practice and Research*, Vol. 75, No. 4, pp.
593 186-190.
594
595 Lacey, C., Clark, B., Frewer, L. and Kuznesof, S. (2016), "Reaching its limits": industry
596 perspectives on salt reduction. *British Food Journal*, Vol. 118, No. 7, pp. 1610-1624.
597
598 Lasztity, R. (1983). *Amino Acid Composition and Biological Value of Cereal Proteins*.
599 Budapest: Kluwer Academic Publishers.
600
601 Lebwohl, B., Murray, J., Rubio-Tapia, A., Green, P. and Ludvigsson, J. (2014), "Predictors of
602 persistent villous atrophy in coeliac disease: a population-based study", *Alimentary*
603 *Pharmacology & Therapeutics*, Vol. 39 No. 5, pp. 488-495.
604

605 Lee, A. R., Ng, D. L., Zivin, J. and Green, P. H. R. (2007), "Economic burden of a gluten-free
606 diet," *Journal of Human Nutrition and Dietetics*, Vol. 20, No. 5, pp. 423-430.
607

608 Lennernäs, M., Fjellstrom, C., Becker, W., Giachetti, I., Schmitt, A., Remaut de Winter, A.
609 and Kearney, M. (1997), "Influences on food choice perceived to be important by nationally-
610 representative samples of adults in the European Union", *European Journal of Clinical*
611 *Nutrition*, Vol. 51, No. 2, pp. 8-15.
612

613 MacCulloch, K. and Rashid, M. (2014), "Factors affecting adherence to a gluten-free diet in
614 children with celiac disease", *Paediatrics & Child Health*, Vol. 19, No. 6, pp.305-309.
615

616 Mancebo, C.M., Martínez, M.M., Merino, C., de la Hera, E. and Gómez, M. (2017), "Effect of
617 oil and shortening in rice bread quality: Relationship between rheology and quality
618 characteristics", *Journal of Texture Studies*, Vol. 48, No. 6, pp. 597-606.
619

620 Mariotti, M., Pagani, M.A. and Lucisano, M. (2013), "The role of buckwheat and HPMC on
621 the breadmaking properties of some commercial gluten-free bread mixtures", *Food*
622 *Hydrocolloids*, Vol. 30, No. 1, pp. 393-400.
623

624 Mazzarella, G. (2015), "Effector and suppressor T cells in celiac disease", *World Journal of*
625 *Gastroenterology*, Vol. 21 No. 24, pp. 7349.
626

627 Miranda, J., Lasa, A., Bustamante, A., Churrua, I. and Simon, E. (2014), "Nutritional
628 differences between a gluten-free diet and a diet containing equivalent products with gluten.
629 *Plant Foods for Human Nutrition*, Vol. 69, No. 2, pp. 182-187.
630

631 Missbach, B., Schwingshackl, L., Billmann, A., Mystek, A., Hickelsberger, M., Bauer, G. and
632 König, J. (2015), "Gluten-free food database: the nutritional quality and cost of packaged
633 gluten-free foods", *PeerJ*, Vol. 3, e1337.
634

635 Muhammad, H., Reeves, S., Ishaq, S., Mayberry, J. and Jeanes, Y. M. (2017), "Adherence to
636 a gluten free diet is associated with receiving gluten free foods on prescription and
637 understanding food labelling", *Nutrients*, Vol. 9, No. 7, pp. 705.
638

639 MyFitnessPal (2015). *Myfitnesspal: Food Search*. Available at:
640 <http://www.myfitnesspal.com/food/calorie-chart-nutrition-facts> [Accessed 10 December
641 2017].
642

643 Nelsen, D.A. (2002), "Gluten-sensitive enteropathy (celiac disease): more common than you
644 think", *American Family Physician*, Vol. 66 No. 12, pp. 2259-2266.
645

646 NHS (2019). Clinical commissioning groups (CCGs). Available at:
647 <https://www.england.nhs.uk/commissioning/who-commissions-nhs-services/ccgs/> [Accessed
648 17 July 2019].
649

650 NHSCC (2018). *NHSCC statement on the result of the Department of Health and Social Care*
651 *consultation on the availability of gluten-free foods on NHS prescription*. Available at:
652 [https://www.nhscc.org/latest-news/nhscc-statement-on-the-result-of-the-department-of-](https://www.nhscc.org/latest-news/nhscc-statement-on-the-result-of-the-department-of-health-and-social-care-consultation-on-the-availability-of-gluten-free-foods-on-nhs-prescription/)
653 [health-and-social-care-consultation-on-the-availability-of-gluten-free-foods-on-nhs-](https://www.nhscc.org/latest-news/nhscc-statement-on-the-result-of-the-department-of-health-and-social-care-consultation-on-the-availability-of-gluten-free-foods-on-nhs-prescription/)
654 [prescription/](https://www.nhscc.org/latest-news/nhscc-statement-on-the-result-of-the-department-of-health-and-social-care-consultation-on-the-availability-of-gluten-free-foods-on-nhs-prescription/) [Accessed 17 April 2018].
655

656 NHS Choices (2017). *Starchy foods and carbohydrates*. Available at:
657 <https://www.nhs.uk/Livewell/Goodfood/Pages/starchy-foods.aspx> [Accessed 13 April 2018].
658

659 Padalino, L., Conte, A. and Del Nobile, M. A. (2016), "Overview of the general approaches to
660 improve gluten-free pasta and bread," *Foods*, Vol. 5, No. 4, pp. 87.
661

662 Public Health England (PHE), 2016. *Government Dietary Recommendations: Government*
663 *recommendations for energy and nutrients for males and females aged 1-18 years and 19+*
664 *years*. Available at:
665 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/618167/gov](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/618167/government_dietary_recommendations.pdf)
666 [ernment dietary recommendations.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/618167/government_dietary_recommendations.pdf) [Accessed 28 March 2018].
667

668 Robins, G., Akobeng, A., McGough, N., Merrikin, E. and Kirk, E. (2008). *A systematic*
669 *literature review on the nutritional adequacy of a typical gluten-free diet with particular*
670 *reference to iron, calcium, folate and B vitamins*. Available at:
671 <https://www.food.gov.uk/sites/default/files/research-report-gluten-free.pdf> [Accessed 10
672 April 2018].
673

674 Sandri, L. T. B., Santos, F. G., Fratelli, C. and Capriles, V. D. (2017), "Development of
675 gluten-free bread formulations containing whole chia flour with acceptable sensory
676 properties", *Food Science & Nutrition*, Vol. 5, No. 5, pp. 1021-1028.
677

678 Saturni, L., Ferretti, G. and Bacchetti, T. (2010), "The gluten-free diet: Safety and nutritional
679 quality." *Nutrients*, Vol. 2, No. 1, pp. 16-34.
680

681 Simón, E., Larretxi, I., Churruga, I., Lasa, A., Bustamante, M.A., Navarro, V., Fernández-Gil,
682 and Miranda, J. (2017). *Nutritional and Analytical Approaches of Gluten-Free Diet in Celiac*
683 *Disease*. Springer International Publishing, pp. 1 – 94.
684

685 Singh, J. and Whelan, K. (2011), "Limited availability and higher cost of gluten-free foods",
686 *Journal of Human Nutrition and Dietetics*, Vol. 24, No. 5, pp. 479-486.
687

688 Smith, P. (2010). Gluten-free food supplier faces competition from the supermarkets. *The*
689 *Telegraph online*, 03 May. Available at:
690 [https://www.telegraph.co.uk/finance/businessclub/7674019/Gluten-free-food-supplier-faces-](https://www.telegraph.co.uk/finance/businessclub/7674019/Gluten-free-food-supplier-faces-competition-from-the-supermarkets.html)
691 [competition-from-the-supermarkets.html](https://www.telegraph.co.uk/finance/businessclub/7674019/Gluten-free-food-supplier-faces-competition-from-the-supermarkets.html) [Accessed 14 April 2018].
692

693 Stevens, L. and Rashid, M. (2008), "Gluten-free and regular foods: A cost comparison",
694 *Canadian Journal of Dietetic Practice and Research*, Vol. 69, No. 3, pp. 147-150.
695

696 Theethira, T., Dennis, M. and Leffler, D. (2014), "Nutritional consequences of celiac disease
697 and the gluten-free diet", *Expert Review of Gastroenterology & Hepatology*, Vol. 8 No. 2, pp.
698 123-129.
699

700 Thompson, T. (2000), "Folate, iron and dietary fiber contents of the gluten-free diet",
701 *Journal of the Academy of Nutrition and Dietetics*, Vol. 100, No. 11, pp. 1389-1396.
702

703 Wang, K., Lu, F., Li, Z., Zhao, L. and Han, C. (2017), "Recent developments in gluten-free
704 bread baking approaches: a review", *Food Science and Technology*, Vol. 37, No. 1, pp. 1-9.
705

706 Webb, G. P. (2013). *Nutrition: Maintaining and improving health*. 4th ed. Boca Raton: CRC
707 Press.

708
709 WHO (2004). *Global Strategy on Diet, Physical Activity and Health*. Geneva: World Health
710 Organisation. Available at:
711 http://www.who.int/dietphysicalactivity/strategy/eb11344/strategy_english_web.pdf
712 [Accessed 17 April 2018].
713
714 WHO (2016). *Salt reduction: fact sheet*. Geneva: World Health Organisation. Available at:
715 <http://www.who.int/mediacentre/factsheets/fs393/en/> [Accessed 19 April 2018].
716
717 Wieser, H. (2007), "Chemistry of gluten proteins", *Food Microbiology*, Vol. 24, No. 2, pp.
718 115-119.
719
720 Wright, H., 2017. *Availability of gluten-free products on NHS prescription: Consultation*
721 *response form*. Available at:
722 <https://www.rpharms.com/Portals/0/RPS%20document%20library/Open%20access/Consultations/June%202017/consdoc3991.pdf> [Accessed 14 April 2018].
723
724
725 Wu, J. H. Y., Neal, B., Trevena, H. and Dunford, E. (2015), "Are gluten free foods healthier
726 than non-gluten free foods? An evaluation of supermarket products in Australia", *British*
727 *Journal of Nutrition*, Vol. 114, No. 3, pp. 448-454.
728