



Article

Profile of repeat victimisation within multi-agency referrals

Shorrocks, Sarah, Mcmanus, Michelle Ann and Kirby, Stuart

Available at <http://clock.uclan.ac.uk/31564/>

Shorrocks, Sarah, Mcmanus, Michelle Ann ORCID: 0000-0002-0095-1071 and Kirby, Stuart ORCID: 0000-0002-3049-1248 (2020) Profile of repeat victimisation within multi-agency referrals. International Review of Victimology, 26 (3). pp. 332-343. ISSN 0269-7580

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1177/0269758020902890>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

1 **A neuromuscular training programme enhances dynamic neuromuscular control and physical**
2 **performance in court-sport athletes**

3 Thomas I. Gee^a, Ryan A. Morrow^a, Mark R. Stone^b, Daniel C. Bishop^a

4
5 ^aSchool of Sport and Exercise Science,

6 University of Lincoln,

7 Lincoln,

8 United Kingdom

9
10 ^bSchool of Sport and Wellbeing

11 University of Central Lancashire,

12 Preston,

13 Lancashire,

14 United Kingdom

15
16
17 **Acknowledgments**

18 Funding for this research was provided by the School of Sport and Exercise Science at the University of
19 Lincoln. There was no conflict of interest associated with the study. The authors would like to thank the
20 athletes who volunteered to participate in the research.

26 **Abstract**

27 This study aimed to investigate the effects of an 8 week neuromuscular training protocol on dynamic
28 neuromuscular control and physical performance. Twenty female court-sport athletes were randomly
29 allocated to two groups, a control (n = 10) and an intervention group (n = 10). All participants performed
30 pre-testing inclusive of; a 20-m sprint, countermovement jump (CMJ), Illinois agility test and the qualitative
31 analysis of a single-leg squat (QASLS), which is an assessment of dynamic neuromuscular control.
32 Subsequently the intervention group engaged in a modified version of the FIFA 11+ neuromuscular training
33 programme twice weekly for eight weeks. At post-testing there were between-group differences in CMJ
34 (Intervention; +3.96-cm vs Control; -1.36-cm, $p = 0.003$) and QASLS for both legs ($p < 0.001$) (Right-Leg,
35 Intervention; Pre: 3.75 Post: 1.91, Control; Pre: 4.11 Post: 4.08) (Left-Leg, Intervention; Pre: 3.72 Post:
36 0.98, Control; Pre: 4.05 Post: 4.23). There was a within-group improvement in 20-m sprint for the
37 intervention group only post-training (Pre: 3.69-s Post: 3.60-s, $p = 0.043$), whilst no differences occurred
38 in Illinois agility test for either group. A modified FIFA 11+ protocol can be considered an effective
39 neuromuscular training programme for enhancing dynamic neuromuscular control and sport-specific
40 physical performance in female court-sport athletes.

41

42

43

44

45

46

47

48

49

50

51

52 Introduction

53 Court-sport athletes, such as those involved in badminton, squash, netball, basketball and volleyball are
54 required to repetitively perform movements patterns such as jumping, landing and changing direction
55 quickly whilst competing.^{1,2} Irrespective of sex, the movement patterns performed by court-sport athletes
56 inherently present a risk of injury if performed with compromised technique and body mechanics.³
57 Repetitive quick changes in direction can cause anterior translation of the tibia relative to femur, which
58 places a significant amount of force on the anterior cruciate ligament (ACL).⁴ This excessive force exerted
59 on the ACL can cause ligament rupture, with females reported to be up to eight times more likely to suffer
60 an ACL injury in comparison to males.⁵ Subsequently, ACL annual injury risk in female court athletes (up
61 to 7.3% of surveyed population suffering a current ACL injury) are some of the highest reported when
62 compared to variety of sporting populations.⁶ The speculated reasons why females are at an increased risk
63 of injuring their ACL include hormonal and anatomical factors which affect the rigidity of the ACL.⁷

64 Abnormal dynamic alignment and neuromuscular-control of the lower limb during athletic tasks
65 such as jumping, landing and changing direction has been associated with numerous lower limb pathologies
66 including ACL injury.³ Kinematic analysis of the single-leg squat has been shown to predict and mirror
67 movement dysfunction in a netball specific leap-landing task, and subsequently is recognized as an effective
68 field-based screening method for identifying netball players who display biomechanical deficits during
69 landing.¹ In addition, mixed sport elite collegiate athletes displaying poor movement performance on a
70 single-leg squat had significantly greater subsequent incidence of lower extremity injury, when compared
71 to athletes displaying superior movement competency.⁸ The qualitative analysis of single-leg squat
72 (QASLS) scoring system was developed to quantify neuromuscular control at various body regions during
73 a single-leg squat. This method has demonstrated high validity in relation to 3D motion capture (percentage
74 of agreement; 98%, range 97%-100%).⁹

75 Females often demonstrate neuromuscular deficiencies in comparison to males in actions such as
76 eccentric muscular landing control and trunk to lower-limb alignment during landing and changing
77 direction.³ Electromyography (EMG) research has demonstrated females appear to have a lower ability to

78 recruit the muscle groups needed to protect the integrity of the ACL, notably the hamstrings group and local
79 trunk stabilisers such as the transversus abdominis and multifidus.^{3,10} However, it has been demonstrated,
80 that if females engage in an effective neuromuscular training programme it can increase their ability to
81 recruit such protective muscles groups.¹¹ Neuromuscular training programmes include plyometric, balance,
82 perturbation and strength exercises. These forms of training induce specific neural adaptations within the
83 descending corticospinal tracts, motor units, neuromuscular junctions as well as improving the reflex
84 potential of the muscle spindles.^{12,13} One neuromuscular training programme that has demonstrated
85 significant promise for decreasing the risk of injury and improving performance variables is the FIFA
86 11+.^{14,15} Soligard et al.¹⁵ reported significant reductions in minor and severe injury within female soccer
87 players performing the FIFA 11+ protocol over an eight-month period compared to a control group.
88 Subsequent authors have also reported the FIFA 11+ programme to have beneficial effects in reducing
89 injuries.^{11,14,16} The FIFA 11+ protocol has shown to improve physical performance variables such jump
90 height, and sprint ability when implemented both acutely and longitudinally as a training intervention.^{17,18}
91 The FIFA 11+ protocol has demonstrated to reduce injury risk and increase performance indicators in
92 mostly soccer players.^{14,15} However there is noticeable lack of research on how the FIFA 11+ protocol
93 benefits those in other sporting contexts. Considering a female's increased risk of an ACL injury and the
94 additional risk of injuring this ligament through participating in a court-sport,³ it seems appropriate to
95 investigate the effects of the FIFA 11+ programme on physical performance and lower limb neuromuscular
96 control, which has been emphasised as a predictive marker of ACL injury.^{1,8}

97 The purpose of this study was to investigate the effects of an 8 week neuromuscular training
98 protocol on dynamic neuromuscular control of the lower limb and physical performance in female court-
99 sport athletes. Subsequently, it was hypothesised that the intervention group performing neuromuscular
100 training would experience significant improvements in 20-m sprint, agility performance, countermovement
101 jump height and neuromuscular control of the lower-limb as assessed via single-leg squat, when compared
102 to the control group.

103

104 **Materials and Methods**

105 *Participants*

106 Twenty female court-sport athletes participated (stature: 169 ± 7 -cm, mass: 61.3 ± 8.3 -kg, age: 22.3 ± 2.0 -
107 years). The participants came from a variety of different sporting backgrounds which included badminton,
108 squash, futsal, netball, basketball and volleyball. Participants were randomly assigned to either an
109 intervention group ($n = 10$) or a control group ($n = 10$). Participants had no history of knee, thigh, hip or
110 lower back injuries within the past year and had not previously suffered an ACL tear. Participants provided
111 written informed consent to participate in the study, which was approved by the local ethics committee in-
112 line with the Helsinki Declarations for research with human volunteers. Prior to data collection, the
113 reproducibility of the featured assessments over three separate trials spaced by a 48-h period, was
114 established using 10 female university netball players who were part of the study cohort. The data was
115 analysed using procedures published by Hopkins¹⁹ to establish typical error as a percentage (TE %). Mean
116 typical error as a percentage observed for each assessment is provided below.

117

118 *Experimental assessment protocol*

119 Before the assessments were conducted, all participants took part in a standardised warm-up consisting of
120 full-body static stretching followed by 5-min of jogging. All participants performed pre-testing; 20-m sprint
121 (TE = 1.3%) recorded with Smart Speed electronic timing gates (Fusion Sport, Queensland, Australia). This
122 involved three maximal 20-m sprint attempts with 60-s rest in-between each sprint. Each sprint begin from
123 a standing start with the toe of the leading leg was placed behind start line, participants were instructed to
124 sprint as fast as possible on each sprint. Three individual countermovement jumps (CMJ) (TE = 1.1%) with
125 60-s rest in-between. The maximal vertical displacement of the participant's jump was assessed with the
126 use of a Just Jump mat (Just Jump, Probotics, Huntsville, AL, USA). Participants were instructed to place
127 their hands on hips and when prompted by the assessor squat down to a self-selected depth, and jump
128 vertically as high as possible. The Illinois agility test (TE = 2.7%) recorded with Smart Speed electronic
129 timing gates (Fusion Sport, Queensland, Australia), which involved three maximal test attempts with 180-

130 s rest in-between each attempt. The test began from a standing start with the toe of the leading leg placed
131 behind start line, participants were instructed to complete the test as quickly as possible. For 20-m sprint,
132 CMJ and Illinois agility the mean value of the three maximal assessment attempts was used for subsequent
133 data analysis.

134 The QASLS was performed on both the right-leg (TE = 2.0%) and left-leg (TE = 2.3%) to assess
135 the dynamic neuromuscular control of the lower limb. All assessments were conducted by the same member
136 of the experimental team, who was experienced in using the assessment with athletes in an applied setting.
137 The protocol from Herrington and Munro⁹ was adopted, which requires the single-leg squat movement to
138 be recorded and reviewed, scoring the movement at different regions of the body (arm, trunk, pelvis, thigh,
139 knee and foot). The associated scoring scale runs from 0-10, a higher score indicates poor dynamic
140 neuromuscular control.⁹ This required each participant to position themselves 5-m away from the video
141 camera (Sony Handycam CX250). Participants' were then required to stand on one leg and squat down to
142 a depth at which their knee angle was between 45°-60°, they were asked to perform this movement over a
143 5-s period. The participants were given the opportunity to familiarise themselves with the depth of single-
144 leg squat with the use of a goniometer and three trials on each leg with a timer in front of them so they
145 stayed within the 5-s period. Participants were then required to perform three recorded single-leg squats on
146 each leg, and trials were only accepted if the squat was within the desired depth via goniometer monitoring.
147 Once the scores for each leg were collected, the three scores for each leg were averaged and this gave the
148 overall result for the left and right-leg. Both groups then completed post-testing, featuring all the described
149 assessments, following the training intervention (described below). Testing was performed two days after
150 training or competition to allow for sufficient recovery. All tests were carried out on basketball court
151 surface.

152

153 *Neuromuscular training intervention*

154 After pre-testing the intervention group engaged in a modified FIFA 11+ neuromuscular training
155 programme, twice weekly for eight weeks (Table 1). The original FIFA 11+ protocol included football

156 specific drills, subsequently these were removed from the current protocol and replaced with alternative
157 exercises specific to court-based sports to increase the opportunity for neuromuscular improvements
158 specific to the sports involved.²⁰ All participants began each exercise of the training protocol at level one,
159 progressions for an individual were not considered until their participation of the third session. Each session
160 took ~35-min to complete. During the intervention period both groups continued with their habitual training
161 regimes as delivered by their sport coaches.

162 **INSERT TABLE 1 HERE**

163

164 *Statistical analysis*

165 Results are presented as mean (\pm standard deviation) unless stated. A 2x2 mixed ANOVA was performed
166 using the IBM SPSS version 22 to determine any significant differences ($p < 0.05$) between the two groups
167 performance measurements (20-m sprint, CMJ, Illinois agility test and QASLS). Assumptions of sphericity
168 were assessed using Mauchly's test of sphericity, post-hoc pairwise-comparisons were conducted using the
169 Bonferroni correction. Within effect size for intervention and control groups pre to post change on assessed
170 measures was calculated using Cohen's d , with interpretation of observed effect sizes are as follows; trivial
171 < 0.2 , small 0.2-0.6, moderate 0.6-1.2, large 1.2-2.0, very large > 2.0 .²¹

172

173 **Results**

174 There was a significant effect over time for CMJ height ($F = 9.998$, $p = 0.005$) and a significant trial x group
175 interaction ($F_{1,9} = 36.571$, $p < 0.001$). At post-testing CMJ significantly differed between the intervention
176 and control group (6.58-cm [95% CI: 2.48 – 10.69-cm], $p = 0.003$; $d = 1.51$). The intervention group's CMJ
177 height was significantly increased from pre to post-testing (3.96-cm [95% CI: 2.68 – 5.24-cm], $p < 0.001$;
178 $d = 0.99$), whilst the control group experienced no significant change in from pre to post-testing CMJ height
179 (-1.36-cm [95% CI: -2.52 – -0.37-cm], $p = 0.056$; $d = -0.27$).

180 There were no between-group differences for 20-m sprint ($F_{1,9} = 1.704$, $p = 0.208$). A significant
181 within-group effect for 20-m performance was evident for the intervention group as performance

182 significantly improved following the training intervention (-0.09-s [95% CI: -0.003 - -0.18-s], $p = 0.043$, d
 183 = -0.40). No significant within-group effect was evident for the control group (-0.02-s [95% CI: -0.10 -
 184 0.08-s], $p = 0.747$, $d = 0.07$). There was no effect over time for the Illinois agility test ($F = 0.915$, $p = 0.351$),
 185 and no between-group differences ($F_{1,9} = 2.299$, $p = 0.147$). There were no significant changes for either
 186 group pre-to-post (Intervention group; -0.22-s [95% CI: -0.47 - 0.04-s], $p = 0.097$, $d = -0.32$), (Control
 187 group; 0.05-s [95% CI: -0.21 - 0.30-s], $p = 0.697$, $d = 0.04$).

188 There was a significant effect over time for QASLS right-leg scores ($F = 14.814$, $p = 0.001$) and
 189 left-leg scores ($F = 38.400$, $p < 0.001$). A significant trial x group interaction was present for QASLS right-
 190 leg scores ($F_{1,9} = 13.88$, $p = 0.002$) and left-leg scores ($F_{1,9} = 50.00$, $p < 0.001$). At post-testing QASLS
 191 scores for both the right-leg (2.22 [95% CI: 1.09 - 3.25], $p = 0.001$; $d = -1.30$) and left-leg (3.25 [95% CI:
 192 2.00 - 4.50], $p < 0.001$; $d = -1.98$) were significantly different between the two groups. The intervention
 193 group reduced their right-leg (-1.84 [95% CI: -1.12 - -2.56], $p = 0.001$; $d = 1.41$) and left-leg (-2.74 [-2.13
 194 - -3.35], $p < 0.001$; $d = 2.36$) QASLS scores significantly from pre to post, the control group scores were
 195 unchanged (Right-leg; -0.03 [95% CI: -0.75 - 0.69], $p = 0.931$; $d = 0.02$, Left-leg; 0.18 [95% CI: -0.43 -
 196 0.79], $p = 0.545$; $d = 0.11$).

197 **INSERT TABLE 2 HERE**

198

199 Discussion

200 The results demonstrate the FIFA 11+ protocol significantly increased CMJ height and reduced QASLS
 201 scores when compared to the control group outcomes, indicating improved lower limb neuromuscular
 202 control and stability which represents an improvement in a screening measure of ACL injury risk.¹
 203 Significant within-group improvements for the intervention group's CMJ, QASLS and 20-m sprint times
 204 were evidenced. Previous research has demonstrated neuromuscular training programmes can improve
 205 performance indicators such as sprint time, lower body power, agility, and proprioception as well as reduce
 206 the risk of ACL injury in female soccer and basketball populations.^{15,20,22}

207 At post-testing the intervention group significantly increased their CMJ in comparison to the

208 control group. There are several potential reasons why the CMJ height increased significantly within the
209 intervention group. The featured neuromuscular programme included exercises such as box jumps and
210 depth jumps which are designed to elicit increases in lower body power and stretch shortening cycle
211 function.²³ In addition, exercises such as planks, Nordic curls, unilateral balancing and bounding runs also
212 featured. Previous studies have highlighted the positive relationship between increased neuromuscular
213 activity within the core, hip and leg muscles all of which contribute towards the efficiency of a CMJ.^{12,13}
214 Previous cohorts of athletes have recorded increased CMJ performance following prolonged interventions
215 utilizing the FIFA 11+ programme. Bonato et al²⁰ speculated an increase in postural control due to
216 performing the neuromuscular training programme may have attributed toward a significant increase in
217 CMJ height within female basketball athletes. Furthermore youth soccer populations performing FIFA 11+
218 intervention have experienced significant increases in CMJ height in comparison to control following
219 interventions lasting 8-10 weeks.^{12,24}

220 Following the neuromuscular programme there was a significant improvement in neuromuscular
221 control observed during the QASLS test for both right and left-leg. These findings correspond with previous
222 research regarding the benefit of neuromuscular training programmes for improving neuromuscular control
223 and stability and reducing indicators of ACL injury risk in females. Bonato et al²⁰ previously showed that a
224 season-long, neuromuscular-training programme significantly improved postural control and reduced
225 injury occurrence within female basketball players. While Benis et al¹³ found that twice weekly for
226 neuromuscular training for 8 weeks, led to significant improvements in balance when compared to a control
227 group within female basketball players. In addition, researchers have reported a lower injury occurrence
228 within female youth soccer players following a season long, neuromuscular training programme.^{15,22} The
229 training protocol used within this study included exercises designed to elicit neural adaptations between the
230 motor cortex and the targeted muscle.^{11,12,13} The revised FIFA 11+ protocol was designed to condition the
231 following muscles: piriformis; superior and inferior gemelli; obturator internus / externus; quadratus
232 femoris; and the hamstrings and quadriceps muscle groups. Female athletes have been shown to have lower
233 recruitment of the muscles targeted by the intervention programme, which in turn cause misalignments in

234 the lower body.^{3,10} Subsequently this places a significant amount of pressure on the ACL and its ability to
235 prevent anterior tibial subluxation, a common cause of ACL rupture.^{4,7} This study included exercises
236 specially chosen by Santa Monica Sports Medicine Foundation and the Oslo Trauma and Research Centre
237 (i.e. Nordic curls, unilateral balancing, box jumps, inline lunges and depth jumps) with the intention of
238 recruiting the aforementioned muscle groups responsible for protecting the integrity of the knee.¹⁴

239 A significant within-group reduction in 20-m sprint time pre to post for the intervention group was
240 apparent (2.4% decrease). Similar to this study previous researchers have found the FIFA 11+ protocol or
241 similar training programmes to be effective for improving sprint performance; Kilding et al²⁵ recorded a
242 2% significant decrease in male soccer players 20-m sprint time after performing the FIFA 11+ training
243 programme five-times weekly for six weeks. In addition, Reis et al¹⁶ found significant improvements in 30-
244 m (3.3% reduction) sprint for an intervention group of male futsal players performing the FIFA 11+ training
245 programme twice-weekly for 12 weeks. The intervention protocol within the present study included a
246 combination of lower body strengthening exercises (inline lunges, Nordic curls) and plyometric exercises
247 (box jumps, depth jumps). Previous literature has found such exercises to invoke substantial muscular
248 recruitment and activation within hamstring, quadriceps and gluteal group musculature.^{26,27,28,29} These such
249 muscle groups are strongly recruited during sprinting.³⁰ Subsequently the observed enhancements in sprint
250 performance may have been due to an enhanced recruitment and firing rate of muscle fibres needed to
251 improve sprint mechanics and power output.^{11,12,13} However shorter (4 week) FIFA 11+ training
252 interventions have not shown such improvements in 20-m sprint time.³¹ Taking this study and other pieces
253 of evidence into consideration the FIFA 11+ or a slightly amended version of the programme can be
254 considered a useful tool to maintain or reduce 20-m sprint times in court-sport athletes.^{13,15,20}

255 No significant differences between the intervention and control groups Illinois agility test time post
256 eight weeks were discovered. The findings are in accordance with Kilding et al²⁵ who found no significant
257 differences in Illinois agility test time following a FIFA 11+ training programme of 6 weeks duration when
258 compared to a control group within male soccer players. Since court-based athletes frequently engage in
259 change of direction cutting actions within their sport, they may have been less responsive to positively adapt

260 to such movements within the training intervention.¹ It is also important to note, a meta-analysis conducted
261 by Gomes Neto et al¹⁸ who analyzed eleven FIFA 11+ studies, including 4700 participants. The analysis
262 indicated an overall significant improvement in agility performance following engagement in the FIFA 11+
263 programme.¹⁸ This may be indicative of the benefit of longitudinal implementation of the FIFA 11+ on
264 agility.

265 The high adherence rate of participants within this study likely had a meaningful influence on the
266 improvement in numerous post-intervention measures. In total 90% of the intervention group participants
267 completed all 16 sessions. Participant adherence for neuromuscular training programmes which are
268 relatively short in duration such as the one implemented within this study (approximately 35-min to
269 complete), is essential for achieving improvements in physical performance and reducing the risk of sport
270 related injuries. A systematic review supported this claim and concluded that neuromuscular training
271 programmes approximately 25-min duration with compliance rates of 75% significantly reduced the risk of
272 lower extremity injuries.³² Another study created three different groups to assess how adherence rates of
273 the FIFA 11+ programme affect the risk of injury.³³ Participants were either assigned to unsupervised
274 group (control), a group with a coach and physiotherapist (comprehensive) or a group with a coach and no
275 physiotherapist (regular). The comprehensive, regular and control group achieved 86%, 81% and 73%
276 completion of total possible sessions, respectively. The authors discovered that high player adherence to
277 the intervention resulted in significant improvements in functional balance and a reduced injury risk.³³

278 In conclusion, the results demonstrate a modified FIFA 11+ protocol can improve dynamic lower
279 limb neuromuscular control and stability which represents an improvement in a screening measure of ACL
280 injury risk. Considering that females are at greater risk of ACL injury in comparison to males and that the
281 movements performed by court-sport athletes during performance can contribute further to the risk of an
282 ACL injury, implementation of preventative regimes such as the FIFA 11+ programme are justified. The
283 neuromuscular training programme also proved to significantly increase CMJ height in comparison to the
284 control group and lead to a within-group improvement in 20-m sprint for the intervention group. This
285 demonstrated the FIFA 11+ training programme can be used for improving desirable physical attributes for

286 court-sport athletes

287

288 **Perspectives**

289 An amended FIFA 11+ programme performed twice weekly can be considered an appropriate
290 neuromuscular training regime to induce improvements in certain sport-specific performance measures and
291 neuromuscular control within a single-leg squat. This is in support of previous neuromuscular training
292 interventions which have shown to improve physical performance, proprioception and reduce the risk of
293 ACL injury in female soccer and basketball populations.^{15,20,22} Strength and conditioning practitioners
294 should explore the use of neuromuscular training programmes in court-based sports which have a high
295 prevalence of ACL injury especially in female populations.^{1,2,3,8} The QASLS scoring scale is considered a
296 cost and time effective indirect screening method of assessing ACL injury risk, making it a practical and
297 realistic tool for strength and conditioning practitioners.^{1,8,9} Considering that neuromuscular training
298 programmes, such as the FIFA 11+ routine, require minimal equipment to conduct, they can be performed
299 effectively as a strength and conditioning session and do not have to be performed as a warm-up routine
300 alone.

301

302 **References**

303 1. Fox AS, Bonacci J, Saunders N. The relationship between performance of a single-leg squat
304 and leap landing task: moving towards a netball-specific anterior cruciate ligament (ACL)
305 injury risk screening method. *Sports Biomech* 2018;28:1-17.

306

307 2. James LP, Kelly VG, Beckman EM. Injury risk management plan for volleyball athletes: *Sports*
308 *Med* 2014;44:1185-1195.

309

310 3. Hewett TE, Ford KR, Hoogenboom BJ, Myer GD. Understanding and preventing ACL injuries:
311 current biomechanical and epidemiologic considerations-update 2010. *N Am J Sports Phys*

312 Ther 2010;5:234-251.

313

314 4. Hirschmann M, Müller W, Hirschmann MT, Müller W. Complex function of the knee joint:
315 the current understanding of the knee. *Knee Surg Sports Traumatol Arthrosc* 2015;23:2780-
316 2788.

317

318 5. Elliot DL, Goldberg L, Kuehl KS. Young women's anterior cruciate ligament injuries: an
319 expanded model and prevention paradigm. *Sports Med* 2010;40:367-376.

320

321 6. Vauhnik R, Morrissey MC, Rutherford OM, Turk Z, Piliš IA, Perme MP. Rate and risk of
322 anterior cruciate ligament injury among sportswomen in Slovenia. *J Athl Train* 2011;46:92-98.

323

324 7. Shultz SJ. ACL Injury Risk in the Physically Active: Why are Females More Susceptible?
325 *Kinesiol Rev* 2015;4:52-62.

326

327 8. Eckard T, Padua D, Mauntel T, Frank B, Pietrosimone L, Begalle R, Goto S, Clark M, Kucera
328 K. Association between double-leg squat and single-leg squat performance and injury
329 incidence among incoming NCAA Division I athletes: A prospective cohort study. *Phys Ther*
330 *Sport* 2018;34:192-200.

331

332 9. Herrington L, Munro A. A preliminary investigation to establish the criterion validity of a
333 qualitative scoring system of limb alignment during single-leg squat and landing. *JESO*
334 2014;1:1-6.

335

336 10. Hughes G, Dally N. Gender difference in lower limb muscle activity during landing and rapid
337 change of direction. *Sci Sport* 2015;30:163-168.

338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363

11. Zebis MK, Bencke J, Andersen LL, Døssing S, Alkjær T, Magnusson SP, Kjær M, Aagaard, P. The effects of neuromuscular training on knee joint motor control during side cutting in female elite soccer and handball players. *Clin J Sport Med* 2008;18:329-337.

12. Akbari H, Sahebozamani M, Daneshjoo A, Amiri-Khorasani M. Effect of the FIFA 11+ programme on vertical jump performance in elite male youth soccer players. *MJSSM* 2018;7:17-22.

13. Benis R, Bonato M, La Torre A. Elite Female Basketball Players' Body-Weight Neuromuscular Training and Performance on the Y-Balance Test. *J Athl Train* 2016;51:688-695.

14. Silvers-Granelli H, Mandelbaum B, Adeniji O, Insler S, Bizzini M, Pohlig R, Junge A, Snyder-Mackler L, Dvorak J. Efficacy of the FIFA 11+ injury prevention program in the collegiate male soccer player. *Am J Sports Med* 2015;43:2628-2637.

15. Soligard T, Myklebust G, Steffen K, Holme I, Silvers H, Bizzini M, Junge A, Dvorak J, Bahr R, Andersen TE. Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomised controlled trial. *Br Med J* 2008;337:a2469.

16. Reis I, Rebelo A, Krstrup P, Brito J. Performance enhancement effects of Federation Internationale de Football Association's "The 11+" injury prevention training program in youth futsal players. *Clin J Sport Med* 2013;23:318-320.

17. Bizzini M, Impellizzeri FM, Dvorak J, Bortolan L, Schena F, Modena R, Junge A. Physiological and performance responses to the "FIFA 11+"(part 1): is it an appropriate warm-

364 up? J Sports Sci 2013;31:1481-1490.

365

366 18. Gomes Neto M, Conceição CS, de Lima Brasileiro AJA, de Sousa CS, Carvalho VO, de Jesus
367 FLA. Effects of the FIFA 11 training program on injury prevention and performance in football
368 players: a systematic review and meta-analysis. Clin Rehabil 2017;31:651-659.

369

370 19. Hopkins WG. Analysis of reliability with a spreadsheet. [Online]. 2007; Available at:
371 <http://sportssci.org/resource/stats/xrely.xls>.

372

373 20. Bonato M, Benis R, La Torre A. Neuromuscular training reduces lower limb injuries in elite
374 female basketball players. A cluster randomized controlled trial. Scand J Med Sci Sports
375 2018;28:1451-1460.

376

377 21. Hopkins WG. A scale of magnitudes for effect statistics. In: A New View of Statistics 2002;
378 newstats.org/effectmag.html

379

380 22. Steffen K, Myklebust G, Olsen OE, Holme I, Bahr R. Preventing injuries in female youth
381 football—a cluster-randomized controlled trial. Scand J Med Sci Sports 2008;18:605-614.

382

383 23. Hernández S, Ramirez-Campillo R, Álvarez C, Sanchez-Sanchez J, Moran J, Pereira LA,
384 Loturco I. Effects of plyometric training on neuromuscular performance in youth basketball
385 players: a pilot study on the influence of drill randomization. J Sport Sci Med 2018;17:372-
386 378.

387

388 24. Rossler R, Donath L, Bizzini M, Faude O. A new injury prevention programme for children's
389 football—FIFA 11+ Kids—can improve motor performance: a cluster-randomised controlled

- 390 trial. *J Sports Sci* 2016;34:549-556.
- 391
- 392 25. Kilding AE, Tunstall H, Kuzmic D. Suitability of FIFA's "The 11" training programme for
- 393 young football players—impact on physical performance. *J Sport Sci Med* 2008;7:320-326.
- 394
- 395 26. Asadi A. The effects of a 6-week of plyometric training on electromyography changes and
- 396 performance. *Sport Sci* 2011;4:38-42.
- 397
- 398 27. Boudreau SN, Dwyer MK, Mattacola CG, Lattermann C, Uhl TL, McKeon JM. Hip-muscle
- 399 activation during the lunge, single-leg squat, and step-up-and-over exercises. *J Sport Rehabil*
- 400 2009;18:91-103.
- 401
- 402 28. Ditroilo M, De Vito G, Delahunt E. Kinematic and electromyographic analysis of the Nordic
- 403 Hamstring Exercise. *J Electromyogr Kinesiol* 2013;23:1111-1118.
- 404
- 405 29. Pincivero DM, Aldworth C, Dickerson T, Petry C, Shultz T. Quadriceps-hamstring EMG
- 406 activity during functional, closed kinetic chain exercise to fatigue. *Eur J Appl Physiol*
- 407 2000;81:504-509.
- 408
- 409 30. Howard RM, Conway R, Harrison AJ. Muscle activity in sprinting: a review. *Sports Biomech*
- 410 2018;17:1-17.
- 411
- 412 31. Palazón FJR, Noguera CP, Rodríguez FA, Sánchez SH, Romero MTM, de Baranda MDPS,
- 413 Wesolek I. Acute and chronic effects of the FIFA 11+ on several physical performance
- 414 measures in adolescent football players. *Eur J Hum Mov* 2016;36:116-136.
- 415

416 32. Ter Stege MH, Dallinga JM, Benjaminse A, Lemmink KA. Effect of interventions on potential,
417 modifiable risk factors for knee injury in team ball sports: a systematic review. *Sports Med*
418 2014;44:1403-1426.

419
420 33. Steffen K, Emery CA, Romiti M, Kang J, Bizzini M, Dvorak J, Finch CF, Meeuwisse WH.
421 High adherence to a neuromuscular injury prevention programme (FIFA 11+) improves
422 functional balance and reduces injury risk in Canadian youth female football players: a cluster
423 randomised trial. *Br J Sports Med* 2013;47:794-802.

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441 **Tables**

442 Table 1. Adapted version of FIFA 11+ protocol performed by the intervention group. Training session
 443 exercises listed as sets x repetitions.

Exercise and section	Sets and repetitions
<p style="text-align: center;"><i>Warm-up (cones 10 m apart)</i></p> <ul style="list-style-type: none"> • Straight ahead • Hip out • Hip in • Circle partner • Single-leg bounding (added from main work out section) • Quick forwards and backwards 	<p>All exercises are to be performed two times in the allocated area</p>
<p style="text-align: center;"><i>Main training session</i></p> <ul style="list-style-type: none"> • Plank (3 levels of difficulty) • Side plank (3 levels of difficulty) • Nordic curls (3 levels of difficulty) • Test your partners balance (single-leg) <i>(football removed)</i> • Box jumps (3 different heights) <i>(replaced for jumps)</i> • Inline lunges (3 levels of difficulty) • Depth jumps (3 different heights) <i>(added)</i> • Running and cutting (3 levels of difficulty) 	<p>1 min rest between each set</p> <ul style="list-style-type: none"> • 2 x 60 s • 2 x 60 s (each side) • 3 x 6, 8, 10 • 3 x 60 s (each leg) • 2 x 30 s • 2 x 16 (each leg) • 2 x 6 • 2 sets within 10 m space
<p style="text-align: center;"><i>Cool-down (added)</i></p>	<p>Full body static-stretch</p>

444

445

446

447

448

449 Table 2. Mean and (standard deviation) of all assessments pre and post-testing for both groups

Assessment	Control group		Intervention group	
	Pre	Post	Pre	Post
20-m (s)	3.86 (0.24)	3.84 (0.30)	3.69 (0.27)	3.60 (0.17)*
Illinois Agility (s)	18.64 (1.24)	18.69 (1.51)	17.67 (0.70)	17.45 (0.67)
CMJ (cm)	38.58 (4.69)	37.34 (4.55)	39.97 (3.84)	43.93 (4.18)*#
QASLS RL (1-10 scale)	4.11 (1.21)	4.08 (1.24)	3.75 (1.51)	1.91 (1.05)*#
QASLS LL (1-10 scale)	4.05 (1.54)	4.23 (1.62)	3.72 (1.34)	0.98 (0.94)*#

450

451 # = Significant interaction (group x time) pre to post. * = Significant within-group difference pre to post.

452

453