

Central Lancashire Online Knowledge (CLoK)

Title	On-site hygiene and biosecurity assessment: A new tool to assess live bird stalls in wet markets
Type	Article
URL	https://clock.uclan.ac.uk/37312/
DOI	https://doi.org/10.1016/j.foodcont.2021.108108
Date	2021
Citation	Soon, Jan Mei and Wahab, Ikarastika Rahayu Abdul (2021) On-site hygiene and biosecurity assessment: A new tool to assess live bird stalls in wet markets. Food Control. ISSN 0956-7135
Creators	Soon, Jan Mei and Wahab, Ikarastika Rahayu Abdul

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.foodcont.2021.108108>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

1 **On-site hygiene and biosecurity assessment: A new tool to assess live bird stalls in wet** 2 **markets**

3
4 Jan Mei Soon¹, Ikarastika Rahayu Abdul Wahab²

5
6 ¹Faculty of Health and Wellbeing, University of Central Lancashire, Preston, PR1 2HE UK

7 ²Faculty of Agro-Based Industry, Universiti Malaysia Kelantan, 17600 Jeli, Kelantan, Malaysia
8

9 **Abstract**

10 Wet markets play an important role in food security and consumers often view the produce as fresher
11 and cheaper. It is highly prevalent in Asia and a source of livelihood for many small and medium
12 businesses. Studies have revealed that highly unsanitary markets, especially those with live bird stalls
13 operating within the wet market could pose a threat to consumer food safety and public health. This
14 study proposed a new, rapid assessment tool for monitoring hygiene and biosecurity measures of live
15 bird stalls. The design of Hygiene and Biosecurity Assessment Tool (HBAT) was supported by the
16 identification of critical hygiene and biosecurity practices based on empirical evidence that suggests
17 such control measures can prevent or reduce the cross contamination or transmission of zoonoses.
18 An observational, cross sectional study of wet markets selling live birds and/or slaughtered birds was
19 conducted to test the tool. Most wet market stalls slaughter and/or sell chicken, followed by quail,
20 duck and amphibians. 50% of the wet market stalls were rated as moderate, 43.2% as poor and
21 require major improvement, 2.3% as good and 4.5% as excellent. Stalls are in general kept in clean
22 condition and no mixing of species or presence of pests or strays were observed. The cleaning and
23 disinfection practices of slaughter area (after each slaughter) and tools require urgent improvement
24 as majority of stalls cleaned the surfaces with water only. Customers have direct access to live bird
25 stalls and should be reminded (with visible signs) to wash their hands before entering other zones.
26 Toilet and handwashing facilities are highly inadequate and improved physical infrastructure and the
27 provision of sufficient hygiene and handwashing facilities are required to facilitate hand hygiene. This
28 study is highly relevant to countries where wet markets and live bird stalls play a crucial food security
29 role to the local communities. The tool could be used to aid policymakers design evidence-based

¹ Corresponding author: jmsoon@uclan.ac.uk

30 assessments to monitor on-site hygiene and biosecurity measures of live bird / animal stalls in wet
31 markets. To our knowledge, this is the first empirical study to propose an on-site hygiene and
32 biosecurity assessment tool to monitor live bird stalls in wet market.

33

34 **Keywords:** biosecurity; cross contamination; handwashing; hygiene; wet markets; zoning

35

36 **Highlights**

- 37 • Procedure for assessing on-site hygiene and biosecurity measures of wet market stalls is
38 proposed;
- 39 • Cleaning and disinfection practices of slaughter area and tools require urgent improvement to
40 prevent cross contamination;
- 41 • Visible notices to prompt workers and public about importance of hand hygiene and procedures
42 for handwashing is required;
- 43 • Adequate hygiene and handwashing facilities should be provided to workers and consumers.

44

45 **Introduction**

46 As of 26 December 2020, a total of 78,383,527 confirmed cases of COVID-19 have been reported
47 globally including 1,740,390 deaths (WHO, 2020). Although the source of 2019-nCoV is yet to be
48 confirmed, early findings suggest a high possibility of a bat origin (Lu et al., 2020; Zhou et al., 2020)
49 and possibly also involving an intermediary animal species (Junejo et al., 2020; Lam et al. 2020).
50 Cohen (2020) and Li et al. (2020) suggested that marketplace played an early role in the spread of
51 COVID-19. Mizumoto et al. (2020) estimated the reproduction number (R) for market-to-human
52 transmission was 0.24 and 2.37 for human-to-human transmission. Moreover, the reporting rate for
53 market-to-human transmission was estimated to be 2 to 34-fold higher than for cases stemming from
54 human-to-human transmission. This strongly suggests that contact history with wet market played a
55 crucial role in identifying COVID-19 cases. Following the spike case of COVID-19 cases, the Huanan
56 Seafood Wholesale Market in Wuhan, China, for example, created a wild animal section where the
57 animals were slaughtered on-site prior to sale (Maron 2020).

58

59 Wet market is defined as a place that sell fresh produce, including meat, seafood, fruits and
60 vegetables and sometimes live animals that are slaughtered and sold in open-air environments
61 (Nadimpalli & Pickering, 2020). Domestic and wild animal species including poultry, mammalian,
62 reptiles, amphibians and fish are held in cages / tanks, are stressed and located in close proximity to
63 each other, and that makes for ideal conditions for diseases to multiply. Animals are often
64 slaughtered on-site and hung or placed in the open air without ice or refrigeration (Poland, 2020). In
65 Asia, wet markets are prevalent because consumers view the produce from such environments as
66 fresher, cheaper and highly accessible to low-income communities. It is also an important source of
67 livelihood for many small and medium businesses (Petrikova et al., 2020; Zhong et al., 2020).

68

69 Even though viruses causing human foodborne illness cannot grow in foods as reported by Caldwell
70 (2020), various studies have reported high prevalence of foodborne pathogens and viruses found in
71 animal-based and seafood products sold in wet markets. This could be as a result of faecal
72 contamination or handling by infected persons (Seymour & Appleton 2001). *Escherichia coli* and
73 *Salmonella enterica* were isolated from wooden cutting boards used to process raw meat in Hong
74 Kong (Sekoai et al., 2020), avian influenza A virus were found in environmental and animal samples
75 from live poultry markets in China (Yuan et al., 2014), multidrug-resistant *Salmonella* and *Listeria*
76 *monocytogenes* were isolated from chicken, pork and shrimp sold in Thailand (Minami et al., 2010),
77 influenza virus (H5N1) were detected in live bird markets and food markets in Thailand (Amonsin et
78 al., 2008) and Indonesia (Henning et al., 2019) and *Salmonella* were identified in meat from
79 Philippines (Santos et al., 2020). In Malaysia, multidrug-resistant *Salmonella* were isolated from
80 poultry and processing environments (Chuah et al., 2018; Nidaullah et al., 2017), *Listeria*
81 *monocytogenes* were detected in chicken offal sold in wet markets (Kuan et al., 2013) and *Vibrio*
82 *parahaemolyticus* were found in seafood samples in wet markets (Tan et al., 2020). Abatcha et al.
83 (2018) found up to 48% (n=35) of chicken carcasses and 41% (n=202) of environmental samples
84 from wet markers were positive for *Salmonella* spp. *Salmonella* is carried asymptotically in the
85 gastrointestinal tracts of live birds and contamination can occur whilst slaughtering and handling in
86 the wet markets (Abatcha et al., 2018; Trongjit et al., 2017). A higher prevalence of *Salmonella* spp.,
87 including *S. Enteritidis* and *S. Typhimurium* were also reported in wet markets compared to

88 supermarkets (Thung et al., 2016). It is likely that the higher prevalence of foodborne pathogens in
89 wet markets were due to lack of personal hygiene and sanitary conditions in wet markets and high
90 humidity and higher storage temperature in wet markets (Oscar, 2004; Thung et al., 2016). There is
91 an inherent risk of transmitting zoonotic pathogens from live animals to humans and as reported by
92 Poland (2020), it acts as the perfect 'petri dish' environment for a variety of zoonoses to incubate and
93 emerge.

94

95 There was a ban on slaughtering of live birds in wet markets by the National Council of Malaysian
96 government to prevent the spread of infectious diseases (The Star, 2013) but slaughtering of live
97 birds in such markets are still prevalent today (Chuah et al., 2018). However, one of the local
98 authorities in Malaysia i.e. the Penang Island City Council had issued a notice to ban the slaughtering
99 and processing of live birds in all stalls (including wet markets) from October 2021 to prevent and
100 control the spread of infectious diseases from slaughtering live birds, prevent cross contamination of
101 water sources as a result of waste from live bird stalls and to control the spread of pests and wastes
102 (Pers. Communication, Penang Island City Council, 2020). A blanket ban on wet markets selling live
103 animals could potentially drive traders to underground markets where monitoring would be impossible
104 (Lynteris & Fearnley, 2020). Instead of banning wet markets, it would be more effective to improve
105 market biosecurity and hygiene of wet markets and use regulated wet markets to enforce the ban of
106 sale of illegal wildlife (Aguirre et al., 2020; Daszak et al., 2020; Petrikova et al., 2020). There exists a
107 number of tools to measure biosecurity measures at farm level. For example, Biocheck.UGent
108 (<https://biocheck.ugent.be/en>) had been developed to measure biosecurity at broiler farms (Gelaude
109 et al., 2014; Van Limbergen et al., 2018) while Lewerin et al. (2015) developed a risk assessment tool
110 to assess biosecurity measures in cattle and pig farms. BioAsseT (Biosecurity Assessment Tool) was
111 used to measure external, internal biosecurity and diagnostic monitoring of pig farms (Sasaki et al.,
112 2020). To our knowledge, there is a lack of hygiene and biosecurity assessment of live birds delivered
113 post-farm gate including the sale of live birds in wet markets or live poultry markets. This study aims
114 to develop an assessment tool to investigate on-site hygiene and biosecurity measures of live bird
115 stalls in wet markets.

116

117 **Methodology**

118 **Design principles of Hygiene and Biosecurity Assessment Tool (HBAT)**

119 The design of Hygiene and Biosecurity Assessment Tool (HBAT) was guided by previous diagnostic
120 tools utilised in processing environment, slaughterhouses and open-air food markets (Gelaude et al.,
121 2020; Ledo et al., 2016). HBAT was developed in English and supported by the identification of
122 critical hygiene and biosecurity practices based on empirical evidence that suggests such control
123 measures can prevent or reduce the cross contamination or transmission of zoonoses i.e. adequate
124 wet market infrastructure (Chowdhury et al., 2020; Nadimpalli & Pickering, 2020), cleaning and
125 disinfection (Chowdhury et al., 2020; Samaan et al., 2011; Webster, 2004; Yuan et al., 2014; Yuan et
126 al., 2015), zoning (Chowdhury et al., 2020; Indriani et al., 2010; Samaan et al., 2011), waste removal
127 (Indriani et al., 2010), availability of toilets and handwashing facilities (Nadimpalli & Pickering, 2020)
128 and personal hygiene (Ledo et al., 2020). Both authors who are experts in food hygiene, safety and
129 food security reviewed the tool to ensure the content measures the on-site hygiene and biosecurity
130 parameters. The tool is divided into five main sections: Part I: Premises; Part II: Preparation and
131 slaughter area; Part III: Zoning and cross contamination; Part IV: Cleaning and disinfection; Part V:
132 Personal hygiene (Table 1). On-site hygiene and biosecurity scores were quantified by converting the
133 answers in Table 1 into scores where correct application of certain measures = 1 point or 2 points or
134 zero for no application. Food hygiene and biosecurity items that were deemed more likely to result in
135 cross contamination of food and increased public health risks were awarded 2 points. The maximum
136 score of "75" equals full application of hygiene and biosecurity measures on site while "0" represents
137 total absence of any hygienic or biosecurity measures in the live bird / slaughtered bird stall.

138

139 Insert Table 1

140

141 A score of 2, 1 or 0.5 was applied in certain measures where multiple possible scenarios exist. The
142 points were 'graded' to distinguish different level of good hygiene and biosecurity practices. For
143 example, under Handwashing facilities (in Premises Section of Table 1), if handwashing facilities such
144 as handwashing basin with soap and running water was observed, the stall was awarded 2 points. If
145 only a handwashing basin with running water (but no soap or handwashing liquid) was provided, the

146 stall was awarded 1 point. If only buckets of stagnant water were available, the stall was awarded 0.5
147 point and if no handwashing facilities were available at all – no marks were awarded. Table 1 also
148 shows that certain criteria were awarded 2 points in Yes, No or Not Applicable sections. A score of 2
149 is given when the stall demonstrated examples of good hygiene and biosecurity measures. This may
150 include practices where stalls should not be placing animals in overcrowded cages and not
151 slaughtering animals directly near to other live animals.

152

153 **Design of pilot study**

154 Items in HBAT (Table 1) were adapted into an electronic checklist tool using Online Survey so
155 observation could be carried out using a smartphone (Figure 1). HBAT was pilot tested at two wet
156 markets selling live poultry. Three graduate research assistants well versed in participatory and non-
157 participatory observation skills, food safety and hygiene control measures were trained virtually to use
158 HBAT. Before conducting the actual assessment, all users participated in the observation of wet
159 market stalls selling live or slaughtered animals in the pilot study. Results were reviewed and
160 discordant notifications were discussed.

161 Insert Figure 1 here

162 **On-site observation**

163 An observational, cross sectional study of hygienic and biosecurity operations at wet markets selling
164 live birds and/or slaughtered birds was conducted. Wet markets were selected using a convenience
165 sampling approach in cities and sub-urban areas of both East and West Coast of Peninsular Malaysia
166 including one town in East Coast of Malaysia. The selection of wet markets was limited by voluntary
167 participation from the stall owners, markets that remained open for business and national restrictions
168 on inter-state travel during the pandemic. The wet markets in the following states were observed:
169 Selangor (n=10), Perak (n=18), Sarawak (n=2) and Kelantan (n=14). Prior to on-site observation,
170 verbal consents were sought from the owners or sellers at the live birds or slaughtered-birds' stalls
171 for the observers to conduct non-participant observation. The study received institutional ethical
172 approval and abide by the Global Code of Conduct for research in low resource settings requirements
173 (TRUST, 2018). The on-site observation was conducted during the period of June – November 2020.
174 This coincides with the Recovery Movement Control Order (RMCO) period where inter-states travel

175 was allowed. The type of market, animals sold, and number of employees per stall were also
176 observed. A total of 2,822 minutes of observations were carried out at 44 live birds and/or
177 slaughtered birds stalls, with each stall averaging 64 minutes of surveillance.

178

179 **Compliance with Hygiene and Biosecurity Requirements**

180 Hygiene and biosecurity compliance among wet market stall owners selling live and/or slaughtered
181 animals were calculated using the modified formula (Santana et al., 2009; Soon, 2019) below.

182

$$183 \quad P = \left(\frac{TS}{\sum 1 - \sum 2} \right) \times K$$

184

185 P = Part I to Part V (Part I: Premise; Part II: Preparation and slaughter area; Part III: Zoning and
186 cross contamination; Part IV: Cleaning and disinfection; Part V: Personal hygiene);

187 TS = Total score;

188 $\sum 1$ = Total possible points;

189 $\sum 2$ = Total non-applicable points;

190 K = constant value (K values for Part I=16; Part II=24; Part III=14.66; Part IV=16; Part V=29.33)

191

192 The total score represents the points achieved by a specific section e.g. Premises. $\sum 1$ represents the
193 total possible points that could be achieved. In this case, the total possible points for Premises = 12;
194 Preparation and slaughter area = 18; Zoning and cross contamination = 11; Cleaning and disinfection
195 = 12; and Personal hygiene = 22. Total possible points for all parts = 75. The non-applicable points
196 ($\sum 2$) are points that should be deducted from the equation if the requirement is irrelevant to the wet
197 market stall. This is to avoid potentially distorting the final hygiene and biosecurity score. For
198 instance, if animals are not slaughtered at the stall, then the Preparation and Slaughter section will be
199 noted as non-applicable. The average score for all sections (Part I – V) was calculated as:

200

$$201 \quad \left(\frac{PI + PII + PIII + PIV + PV}{10} \right)$$

202

203 The hygiene and biosecurity scores are classified according to the following scale:

204

205 0 – 1.9 = very poor and urgent improvement necessary

206 2.0 – 4.9 = poor and major improvement necessary

207 5.0 – 6.9 = moderate and some improvement required

208 7.0 – 9.0 = Good

209 9.0 – 10.0 = Excellent

210

211 **Statistical analysis**

212 Intraclass coefficient correlation (ICC) was calculated to determine inter-rater reliability using SPSS
213 version 27.0. Values scoring < 0.5 = poor reliability, 0.5-0.75 = moderate reliability, 0.75 – 0.90 =
214 good reliability and > 0.90 = excellent reliability (Koo & Li, 2016). Exploratory factor analysis was
215 conducted to determine construct validity. Principal component analysis (PCA) was performed using
216 varimax rotation.

217

218 **Results and Discussion**

219 More than half of the wet market stalls were situated indoors, and up to 13.6% located partially
220 indoor. However, only 18.2% of the available ventilations systems were working. A number of the
221 wet market stalls were located indoors or partially indoors with limited working ventilation systems in
222 place. All the wet markets were not housed in air-conditioned buildings unlike supermarkets. As the
223 name 'wet market' suggests, floors are continually washed down, certain fresh produce are kept
224 moist to ensure freshness and to keep animals alive, hence wet markets posed an extremely humid
225 environment (Ho, 2014). Wet market stalls located indoors require high ventilation rate to remove
226 moisture, heat and contaminant (Lee & Lee, 2013). Failure to ventilate the damp and warm
227 environment pose a risk for foodborne pathogens and zoonoses to emerge and thrive (Chuah et al.,
228 2018; Rahman et al., 2018). Moreover, high ambient temperatures could lead to heat stress among
229 broiler chickens. Heat stress were found to affect chicken immune functions, increasing the risk of
230 infectious disease outbreaks (Hirakawa et al., 2020). Stalls were mostly operated by one or two staff

231 (including the owner). Majority of the wet market stalls slaughter and/or sell chicken (88.6%)
232 followed by quail, duck and other category (i.e. frogs and toads) (Table 2).

233

234 Insert Table 2

235

236 **On-site Hygiene and Biosecurity Assessment Scores**

237 The Intra-Class Correlation Coefficient (ICC) between the users in the pilot tests measured 0.84, $F(1, 3) = 110.08, p < 0.05$) indicating high inter-rater reliability. The Kaiser-Meyer-Olkin (KMO) measures
238 of sampling value was 0.75. According to Hair, Black, Babin, Anderson and Tatham (2010), the
239 criterion of validity should be > 0.60 and the KMO fulfils the requirement. All factor loadings were $>$
240 0.40 and explained 79.22% of the variance. Based on the observation, market stalls selling live birds
241 and or slaughtered birds were scored using Table 1 and Formulas 1 and 2. 50% of the wet market
242 stalls were rated as moderate, 43.2% as poor and require major improvement, 2.3% as good and
243 4.5% as excellent (Figure 2).

244

245
246 Insert Figure 2

247

248 Twenty-one stalls were found to place live birds in overcrowded cages although none of the birds
249 were mixed with other poultry. Of these 21 stalls with overcrowded cages, 8 stalls were not
250 maintained in a sanitary condition. Overcrowding of animals lead to highly stressed animals and
251 coupled with highly unsanitary conditions, this would serve as breeding grounds for zoonoses
252 (Wiebers & Feigin, 2020). Most stalls were maintained in clean condition (70.5%), cages were kept
253 clean (43.2%) and the stall area free from pests or strays (e.g. rodents, stray cats or dogs and wild
254 birds) (59.1%). One good practice observed amongst all stalls was the absence of mixing different
255 bird species in the same cage. As reported by Chan et al., (2013) other bird species e.g. ducks, geese
256 and quails were segregated from chickens to prevent the spread of avian influenza viruses from
257 asymptomatic birds to chickens. Majority of stalls had some form of handwashing facilities available.
258 However, there is limited number of public handwashing facilities for customers, especially when
259 crossing into zones selling ready-to-eat food. Although 84.1% of the wet market stalls were in a

260 different zone, up to 30% of the live bird stalls were situated less than 3m away from other food
261 stalls (Table 3). In most wet markets, customers were observed to have direct access to the live bird
262 stalls (90.9%). This is a common practice as customers prefer to select their own bird of choice, other
263 than want to see themselves on how the farms/markets handle the birds. However, this increases the
264 opportunity for human transfer of pathogens and zoonoses (Cui et al., 2019) as evidenced by the
265 spill-over of avian influenza virus from infected poultry to humans (Wang et al., 2017; Wang et al.,
266 2020). As the case of avian influenza H5N1 outbreak occurring in 2003 to 2006, it has been reported
267 that most patients who have been infected had recent direct contacts with poultry (Woo et al., 2006).
268 Although customers could access the handwashing facilities at 43.2% of the stalls, this study did not
269 carry out any observation of the customers (i.e. whether they washed their hands after selecting the
270 birds / touching the surfaces at the live bird stall).

271

272 Insert Table 3

273

274 There is also a lack of public handwashing facilities for customers when entering different zones (e.g.
275 ready-to-eat food stalls). Only nine wet markets provided public handwashing facilities for consumers
276 and staff before entering zones selling ready-to-eat foods. Fourteen wet markets placed visible signs
277 and notices to remind customers and staff to wash their hands. This is concerning given that the
278 observations were carried out during the COVID-19 pandemic. Notices to remind or prompt workers
279 and public about importance of hand hygiene and procedures for handwashing is one of the key
280 strategies proposed by the WHO multimodal hand hygiene improvement strategy (WHO, 2009). The
281 live bird stalls were located in a live animal zone and although most were segregated, up to 30%
282 were within 3 meters of other food stalls. Previous studies have shown that aerosol transmission is an
283 important mode of viral transmission in wet market environment (Wei et al., 2018). In closed
284 environments with minimal ventilation, virus in aerosols may persist in air for longer and at higher
285 concentrations, thus increasing rate of transmission (Wu et al., 2020). Formation of aerosols are
286 further aided with the use of hosepipes (81.8%) and brooms (68.2%) for cleaning which is prevalent
287 in this study. Furthermore, the washing down of stalls resulted in waste run-offs contaminating other
288 non-live bird areas and food stalls.

289

290 Table 4 shows the observation of hygiene practices of wet market stalls selling live birds and/or
291 slaughtered birds. Most stalls that sell live birds also slaughtered the birds on-site. Six stalls were
292 observed to slaughter birds on the ground although most staff carried out the process on bench tops
293 that are easy to clean and smooth (e.g. stainless steel, ceramic stone, worktable with aluminium top).
294 Slaughtered birds were often cleaned to remove visible dirt, soil and blood stains before selling them
295 to customers. Only 38.6% of the stalls cleaned the work surface after each slaughter and most stalls
296 only used water to clean the surface area. Similarly, knives and tools were mostly washed with water
297 only. Wastes were collected into dedicated bins with lids (18.2%), without lids (36.4%) but some
298 wastes were also washed into nearby drains (29.5%). Although most stalls used easy-to-clean and
299 smooth surfaces as their work tops (e.g. stainless steel, ceramic stone, worktable with aluminium
300 top). cleaning of work surfaces was not often carried out after each slaughter. Slaughter area and
301 tools were cleaned with water only (> 70%). This posed a risk of transmission of foodborn pathogens
302 and zoonoses if surfaces were not cleaned adequately. *Escherichia coli* and *Salmonella* (Sekoai et al.,
303 2020), *Kelbsiella pneumoniae* (Lo et al., 2020) and H7N9 virus (Wang et al., 2015) were detected in
304 samples collected from surfaces of chopping boards from wet markets. Traditional cleaning method of
305 wooden cutting boards often used by Asian vendors include scraping the surface of the wooden
306 cutting board with a chopping knife until a white layered film has been removed, followed by rinsing
307 with hot water (Lo et al., 2020). In terms of cleaning practices, most stalls used hosepipes and
308 brooms to clean the stall area. It is concerning to note that liquid wastes such as blood were washed
309 into other nearby areas including food stalls (18.2%).

310

311 Workers mostly wear apron and boots, but only slightly more than half of the staff wore masks or
312 gloves. In certain wet markets, the use of masks could be due to the enforcement during the
313 Movement Control Order (MCO). A third of the workers were observed to touch their mouth, nose or
314 eyes whilst handling or after handling live animals. A number of staff also tend to use their mobile
315 devices during and after handling live birds (15.9%). Contamination with faecal residues upon
316 handling live birds can occur on such mobile devices. It has been reported in Olsen et al. (2020) that
317 mobile communication devices can serve as possible breeding grounds for microbial organisms.

318 Handwashing with soap were only observed in small number of stalls (20.5%) in this study. More
319 than half of the staff (56.8%) would wash hands with water only after handling live animals. This
320 finding reflects the study by Alam et al. (2019) where market sellers were found to rarely wash their
321 hands with soap but tend to wipe their hands with a cloth. Although majority wore aprons and boots,
322 slightly less than half of the workers wore face masks and most do not use gloves when handling live
323 animals. One of the main reasons documented by Alam et al. (2019) was that the high temperature
324 and humidity in wet markets discourage workers from wearing protective equipment. Toilet facilities
325 were lacking in half of the wet market stalls and the remaining stalls with access to such facilities
326 were found to be inadequate (dirty, no running water, no soap or hand drying facilities). Although
327 half of the wet market stalls have access to toilet facilities, the provision of clean and adequate
328 potable water supply is seriously lacking and is important to facilitate handwashing practices.

329

330 Insert Table 4

331

332

333

334

335

336

337

338 The above findings reflect the hygiene and biosecurity measures rating for most stalls. Majority were
339 rated as moderate with some improvements required or poor with major improvements necessary.
340 This study reiterates the recommendations of Nadimpalli and Pickering (2020) and WHO (2006) that
341 called for standardised global monitoring of water, sanitation and hygiene (WASH) and to improve the
342 physical infrastructure of wet markets including the provision of sufficient toilets, handwashing
343 facilities, potable water supply and proper drainage. In April 2020, WHO called for stricter food safety
344 and hygiene standards for wet markets and is developing guidance for the safe operation of wet
345 markets (Briggs, 2020). Our findings can aid the design of evidence-based assessments to monitor
346 on-site safety, hygiene and biosecurity measures of live bird stalls in wet markets. Shi et al. (2020)

347 conducted a meta-analysis of 19 studies and found that implementation of interventions in live bird
348 market environment significantly reduce zoonotic infections. This is also the first assessment tool to
349 assess level of on-site hygiene and biosecurity measures of live bird stalls in wet markets, providing a
350 rapid indication of the hygiene and biosecurity scores of live bird stalls.

351

352 **Limitations, practical implications and recommendation for future studies**

353 There are several limitations associated with the study. First, the study was conducted during the
354 Recovery Movement Control Order and Conditional Movement Control Order (stricter measures with
355 no inter-states travel in November 2020). The data collection was restricted to several states, hence
356 limiting the number of sites. The data collected during the on-site observation only provided a
357 snapshot of current hygiene and biosecurity measures. The presence of observers may have
358 introduced the Hawthorne effect among the participants and potentially increases handwashing
359 behaviours. The researchers did not observe other biosecurity measures such as weekly rest days,
360 weekly and monthly disinfection practices, transportation and receipt of live birds at wet markets,
361 whether poultry were kept overnight / days and treatment and transportation of collected wastes.
362 The tool could be easily adapted to suit the local food and requirements of wet markets in different
363 regions. For example, instead of live bird stalls in Malaysia, the tool could potentially be modified and
364 applied to other stalls selling live and/or slaughtered meat and seafood products. If future
365 researchers were to modify the tool, the content and construct validity and inter-reliability must be
366 tested to ensure multiple users could assess the same hygiene and biosecurity measures with no
367 significant differences. One of the key strengths of HBAT is the convenience, ease of use and enables
368 rapid assessments of on-site hygiene and biosecurity measures. Since it could be used in
369 smartphones, it offers covert observation of stalls by veterinary, public health and food safety
370 inspectors. It would be highly valuable to conduct a live poultry supply chain study to assess the
371 hygiene and biosecurity measures from farm to market. The on-site Hygiene and Biosecurity
372 Assessment tool could be further adapted and/or modified to suit other countries and local wet
373 market practices selling live birds.

374

375 **Conclusion**

376 Wet markets play a crucial food security role for local communities and is a source of livelihood for
377 many small-scale food businesses as well as provide essential social interaction for elderly residents in
378 the area. However, highly unsanitary wet markets with minimal or no biosecurity measures is the
379 perfect 'petri dish' environment for a variety of zoonoses to thrive. Instead of banning or trying to
380 outlaw wet markets with live animals, it would be more effective to ensure that live animal stalls in
381 wet markets are practising good hygiene and biosecurity measures. Physical infrastructures including
382 designated or segregated area for live bird stalls, provision of toilet and adequate handwashing
383 facilities for workers and staff and monitoring of wet markets to ensure hygiene and biosecurity
384 measures are met are crucial interventions needed to ensure the safety and welfare of animals and
385 that public health are not at risk. By practising this, the transmission of the viruses to humans can
386 be controlled and reduced. To our knowledge, this is the first study to propose an on-site hygiene
387 and biosecurity assessment tool to monitor live bird stalls in wet market. The study could be used to
388 aid policymakers in developing guidance and training of staff operating live bird stalls and to design
389 evidence-based assessments to monitor hygiene and biosecurity measures.

390

391 **Acknowledgement**

392 The researchers are grateful to all participating wet market stalls and research assistance conducted
393 during the COVID-19 pandemic.

394

395 **References**

396 Abatcha, M. G., Effarizah, M. E., & Rusul, G. (2018). Prevalence, antimicrobial resistance, resistance
397 genes and class 1 integrons of Salmonella serovars in leafy vegetables, chicken carcasses and related
398 processing environments in Malaysian fresh food markets. *Food Control*, 91, 170-180.

399

400 Aguirre, A. A., Catherina, R., Frye, H., & Shelley, L. (2020). Illicit wildlife trade, wet markets, and
401 COVID-19: Preventing future pandemics. *World Medicine & Health Policy*, 12(3), 256-265.

402

403 Alam, M.-U., Rahman, M., Al-Masud, A., Islam, M. A., Asaduzzaman, M., Sarker, S. et al. (2019).
404 Human exposure to antimicrobial resistance from poultry production: Assessing hygiene and waste-
405 disposal practices in Bangladesh. *International Journal of Hygiene and Environmental Health*, 222,
406 1068-1076.

407

408 Amonsin, A., Choatrakol, C., Lapkuntod, J., Tantilertcharoen, R., Thanawongnuwech, R., Suradhat, S.
409 et al. (2008). Influenza virus (H5N1) in live bird markets and food markets, Thailand. *Emerging
410 Infectious Diseases*, 14(11), 1739-1742.

411

412 Briggs, H. (2020). Coronavirus: WHO developing guidance on wet markets. BBC, 21 April. Available
413 at: <https://www.bbc.co.uk/news/science-environment-52369878>

414
415 Caldwell, J. M. (2020). COVID-19, Wet Markets, and Food Safety. *Food Technology Magazine*, 74(7).
416
417 Cohen, J. (2020). Mining coronavirus genomes for clues to the outbreak's origins. *Science*. Available
418 at: <https://www.sciencemag.org/news/2020/01/mining-coronavirus-genomes-clues-outbreak-s-origins>
419
420 Chan, J. F.-W., To, K. K.-W., Tse, H., Yin, D.-Y., & Yuen, K.-Y. (2013). Interspecies transmission and
421 emergence of novel viruses: lessons from bats and birds. *Trends in Microbiology*, 21(10), 544-555.
422
423 Chowdhury, S., Azziz-Baumgartner, E., Kile, J. C., Hogue, M. A., Rahman, R. Z., Hossain, M. E. et al.
424 (2020). Association of biosecurity and hygiene practices with environmental contamination with
425 influenza A viruses in live bird markets, Bangladesh. *Emerging Infectious Diseases*, 26(9), 2087-2096.
426
427 Chuah, L. O., Syuhada, A.K.S., Suhaimi, I. M., Hanim, T. F., & Rusul, G. (2018). Genetic relatedness,
428 antimicrobial resistance and biofilm formation of *Salmonella* isolated from naturally contaminated
429 poultry and their processing environment in northern Malaysia. *Food Research International*, 105,
430 743-751.
431
432 Cui, J., Li, F., Shi, Z.-L. (2019). Origin and evolution of pathogenic coronaviruses. *Nature Review*
433 *Microbiology*, 17(3), 181-192.
434
435 Daszak, P., Olival, K. J., & Li, H. (2020). A strategy to prevent future epidemics similar to the 2019-
436 nCoV outbreak. *Biosafety and Health*, 2(1), 6-8.
437
438 Gelaude, P., Schlepers, M., Verlinden, M., Laanen, M., Dewulf, J. (2014). Biocheck.UGent: A
439 quantitative tool to measure biosecurity at broiler farms and the relationship with technical
440 performances and antimicrobial use. *Poultry Science*, 93(11), 2740-2751.
441
442 Hair, J. F., Black, B., Babin, B., Anderson, R. E., & Tatham, R. I. (2010). *Multivariate analysis: A*
443 *global perspective*. New Jersey: Person Education Inc, Upper Saddle River.
444
445 Henning, J., Hesterberg, U. W., Zenal, F., Schoonman, L., Brum, E., & McGrane, J. (2019). Risk
446 factors for H5 avian influenza virus prevalence on urban live bird markets in Jakarta, Indonesia –
447 Evaluation of long-term environmental surveillance data. *PLoS ONE*, 14(5), e0216984.
448
449 Hirakawa, R., Nurjanah, S., Furukawa, K., Murai, A., Kikusato, M., Nochi, Y. et al. (2020). Heat stress
450 causes immune abnormalities via massive damage to effect proliferation and differentiation in broiler
451 chickens. *Frontiers in Veterinary Science*, 7, 46.
452
453 Ho, H. F. (2014). Meat marketing – wet markets. M. Dikeman and C. Devine (Eds.). In, *Encyclopaedia*
454 *of Meat Sciences*. Academic Press, pp. 244-247.
455
456 Indriani, R. Samaan, G., Gultom, A., Loth, L., Indryani, S., Adjid, R. et al. (2010). Environmental
457 sampling for avian influenza virus A (H5N1) in live-bird markets, Indonesia. *Emerging Infectious*
458 *Diseases*, 16(12), 1889-1895.
459
460 Junejo, Y., Ozaslan, M., Safdar, M., Khailany, R. A., Rehman, A., Yousaf, W. et al. (2020). Novel
461 SARS-CoV-2/COVID-19: Origin, pathogenesis, genes and genetic variations, immune responses and
462 phylogenetic analysis. *Gene Reports*, 20, 100752.
463
464 Koo, T. K., Li, M. Y. (2016). A guideline of selecting and reporting intraclass correlation coefficients
465 for reliability research. *Journal of Chiropractic Medicine*, 15, 155-163.
466
467 Kuan, C. H., Goh, S. G., Loo, Y. Y., Chang, W. S., Lye, Y. L., Puspanadan, S., Tang, J. Y. H.,
468 Nakaguchi, Y., Nishibuchi, M., Mahyudin, N. A., & Radu, S. (2013). Prevalence and quantification of
469 *Listeria monocytogenes* in chicken offal at the retail level in Malaysia. *Poultry Science*, 92(6), 1664-
470 1669.

471
472 Lam, T. T.-Y., Jia, N., Zhang, Y.-W., Shum, M. H.-H., Jiang, J.-F., Zhu, H.-C. et al. (2020). Identifying
473 SARS-CoV-2-related coronaviruses in Malayan pangolins. *Nature*, 583, 282-285.
474
475 Ledo, J., Hettinga, K. A., & Luning, P. A. (2020). A customized assessment tool to differentiate safety
476 and hygiene control practices in emerging dairy chains. *Food Control*, 111,107072.
477
478 Lee, S. H., & Lee, W. L. (2013). Site verification and modelling of desiccant-based system as an
479 alternative to conventional air-conditioning systems for wet markets. *Energy*, 55, 1076-1083.
480
481 Lewerin, S. S., Osterberg, J., Alenius, S., Elvander, M., Fellstrom, C., Traven, M., Wallgren, P., Waller,
482 K. P., & Jacobson, M. (2015). Risk assessment as a tool for improving external biosecurity at farm
483 level. *BMC Veterinary Research*, 11, 171.
484
485 Li, Q., Guan, X., Wu, P., Wang, X., Zhou, L., Tong, Y. et al. (2020). Early transmission dynamics in
486 Wuhan, China, of novel coronavirus – infected pneumonia. *New England Journal of Medicine*, 382,
487 1199-1207.
488
489 Lo, M. Y., Ngan, W. Y., Tsun, S. M., Hsing, H.-L., Lau, K. T., Hung, H. P. et al. (2020). A field study
490 into Hong Kong's wet markets: raised questions into the hygienic maintenance of meat contact
491 surfaces and the dissemination of microorganisms associated with nosocomial infections. *Frontiers in*
492 *Microbiology*, 10, 2618.
493
494 Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H. (2020). Genomic characterisation and epidemiology
495 of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet*, 395 (10224),
496 565-574.
497
498 Lynteris, C., & Fearnley, L. (2020). Why shutting down Chinese 'wet markets' could be a terrible
499 mistake. *The Conversation*. Available at: [https://theconversation.com/why-shutting-down-chinese-](https://theconversation.com/why-shutting-down-chinese-wet-markets-could-be-a-terrible-mistake-130625)
500 [wet-markets-could-be-a-terrible-mistake-130625](https://theconversation.com/why-shutting-down-chinese-wet-markets-could-be-a-terrible-mistake-130625)
501
502 Maron, D. F. (2020). Wet Markets' Likely Launched the Coronavirus. Here's What you Need to
503 Know. *National Geographic*, April 15.
504 <https://www.nationalgeographic.com/animals/2020/04/coronavirus-linked-to-chinese-wet-markets/>
505
506 Minami A., Chaicumpa, W., Chongsa-Nguan, M., Samosornsuk, S., Monden, S., Takeshi, K. et al.
507 (2010). Prevalence of foodborne pathogens in open markets and supermarkets in Thailand. *Food*
508 *Control*, 21(3), 221-226.
509
510 Mizumoto, K., Kagaya, K., & Chowell, G. (2020). Effect of a wet market on coronavirus disease
511 (COVID-19) transmission dynamics in China, 2019-2020. *International Journal of Infectious Diseases*,
512 97, 96-101.
513
514 Nadimpalli, M. L., & Pickering, A. J. (2020). A call for global monitoring of WASH in wet markets. *The*
515 *Lancet Planetary Health*, 4(10), e439-e440.
516
517 Nastasijevic, I., Tomasevic, I., Smigic, N., Milicevic, D., Petrovic, Z. & Djekic, I. (2016). Hygiene
518 assessment of Serbian meat establishments using different scoring systems. *Food Control*, 62, 193-
519 200.
520
521 Nidaullah, H., Abirami, N., Shamila-Syuhada, A. K., Chuan, L.O., Nurul, H., Tan, T. P., Zainal Abidin,
522 F. W., & Rusul, G. (2017). Prevalence of Salmonella in poultry processing environments in wet
523 markets in Penang and Perlis, Malaysia. *Veterinary World*, 10(3). 286-292.
524
525 Olsen, M., Campos, M., Lohning, A., Jones, P., Legget, J., Bannach-Brown, A., McKirdy, S., Alghafri,
526 R., & Tajouri, L. (2020). Mobile phones represent a pathway for microbial transmission: A scoping
527 review. *Travel Medicine and Infectious Disease*, 35, 101704.

528
529 Oscar, T. P. (2004). A quantitative risk assessment model for Salmonella and whole chickens.
530 *International Journal of Food Microbiology*, 93(2), 231-247.
531
532 Petrikova, I., Cole, J., & Farlow, A. (2020). COVID-19, wet markets, and planetary health. *The Lancet*
533 *Planetary Health*, 4(6), e213-e214.
534
535 Poland, G. A. (2020). Another coronavirus, another epidemic, another warning. *Vaccine*, 38(10), v-vi.
536
537 Rahman, M. H. A., Mohd Hairon, S., Hamat, R. A., Tengku Jamaluddin, T. Z. M., Shafei, M. N., Idris,
538 N. et al. (2018). Seroprevalence and distribution of leptospirosis serovars among wet market workers
539 in northeastern Malaysia: a cross sectional study. *BMC Infectious Diseases*, 18, 569.
540
541 Samaan, G., Gultom, A., Indriani, R., Lokuge, K., & Kelly, P. M. (2011). Critical control points for avian
542 influenza A H5N1 in live bird markets in low resource settings. *Preventive Veterinary Medicine*,
543 100(1), 71-78.
544
545 Santos, P. D. M., Widmer, K. W., & Rivera, W. L. (2020). PCR-based detection and serovar
546 identification of Salmonella in retail meat collected from wet markets in Metro Manila, Philippines.
547 *PLoS ONE*, 15(9), e-239457.
548
549 Sasaki, Y., Furutani, A., Furuichi, T., Hayakawa, Y., Ishizeki, S., Kano, R. et al. (2020). Development
550 of a biosecurity assessment tool and the assessment of biosecurity levels by this tool on Japanese
551 commercial swine farms. *Preventive Veterinary Medicine*, 175, 104848.
552
553 Sekoai, P. T., Feng, S., Zhou, W., Ngan, W. Y., Pu, Y., Pan, J., & Habimana, O. (2020). Insights into
554 the microbiological safety of wooden cutting boards used for meat processing in Hong Kong's wet
555 markets: A focus on food-contact surfaces, cross contamination and the efficacy of traditional hygiene
556 practices. *Microorganisms*, 8(4), 579.
557
558 Seymour, I. F. and H. Appleton. (2001). A review: foodborne viruses and fresh produce. *Journal of*
559 *Applied Microbiology*, 91, 759-773.
560
561 Shi, N., Huang, J., Zhang, X., Bao, C., Yue, N., Wang, Q. et al. (2020). Interventions in live poultry
562 markets for the control of avian influenza: A systematic review and meta-analysis. *Journal of*
563 *Infectious Diseases*, 221(4), 553-560.
564
565 Soon, J. M. (2019). Rapid Food Hygiene Inspection Tool (RFHiT) to assess hygiene conformance
566 index (CI) of street food vendors. *LWT*, 113, 108304.
567
568 Tan, C. W., Rukayadi, Y., Hasan, H., Thung, T. Y., Lee, E., Rollon, W. D. et al. (2020). Prevalence and
569 antibiotic resistance patterns of *Vibrio parahaemolyticus* isolated from different types of seafood in
570 Selangor, Malaysia. *Saudi Journal of Biological Sciences*, 27(6), 1602-1608.
571
572 The Star (2013). Poultry slaughtering banned in wet markets to curb river pollution, spread of
573 infectious diseases. 3 September. Available at:
574 <https://www.thestar.com.my/news/nation/2013/09/03/poultry-slaughter-ban-markets/>
575
576 Thung, T. Y., Mahyudin, N. A., Basri, D. F., Wan Mohamed Radzi, C. W. J., Nakaguchi, Y., Nishibuchi,
577 M., & Radu, S. (2016). Prevalence and antibiotic resistance of *Salmonella* Enteritidis and *Salmonella*
578 *Typhimurium* in raw chicken meat at retail markets in Malaysia. *Poultry Science*, 95(8), 1888-1893.
579
580 Trongjit, S., Angkitrakul, S., Tuttle, R. E., Pongseeree, J., Padungtod, P., & Chuanchuen, R. (2017).
581 Prevalence and antimicrobial resistance in *Salmonella enterica* isolated from broiler chickens, pigs and
582 meat products in Thailand-Cambodia border provinces. *Microbiology and Immunology*, 61(1), 23-33.
583

584 TRUST (2018). Global Code of Conduct for Research in Resource-Poor Settings. Available at:
585 <https://www.globalcodeofconduct.org/>
586
587 Van Limbergen, T., Dewulf, J., Klinkenberg, M., Ducatelle, R., Gelaude, P., Mendez, J., Heinola, K.,
588 Papasolomontos, S., Szeleszczuk, P., & Maes, D. (2018). Scoring biosecurity in European conventional
589 broiler production. *Poultry Science*, 97(1), 74-83.
590
591 Wang, X., Jiang, H., Wu, P., Uyeki, T. M., Feng, L., Lai, S. et al. (2017). Epidemiology of avian
592 influenza A H7N9 virus in human beings across five epidemics in mainland China, 2013-2017: an
593 epidemiological study of laboratory-confirmed case series. *Lancet Infectious Disease*, 17(8), 822-832.
594
595 Wang, X., Liu, S., Mao, H., Yu, Z., Chen, E., & Chai, C. (2015). Surveillance of avian H7N9 virus in
596 various environments of Zhejiang Province, China before and after live poultry markets were closed in
597 2013-2014. *PLoS ONE*, 10(8), e0135718.
598
599 Wang, W., Artois, J., Wang, X., Kucharski, A. J., Pei, Y., Tong, X. et al. (2020). Effectiveness of live
600 poultry market interventions on human infection with avian influenza A (H7N9) virus, China. *Emerging
601 Infectious Diseases*, 26(5), 891-901.
602
603 Webster, R. G. (2004). Wet markets – a continuing source of severe acute respiratory syndrome and
604 influenza? *Lancet*, 363 (9404), 234-236.
605
606 Wei, J., Zhou, J., Cheng, K., Wu, J., Zhong, Z., Song, Y. et al. (2018). Assessing the risk of downwind
607 spread of avian influenza virus via airborne particles from an urban wholesale poultry market. *Building
608 and Environment*, 127, 120-126.
609
610 WHO (2006). A guide to health food markets. Available at:
611 <https://www.who.int/foodsafety/publications/healthy-food-market/en/>
612
613 WHO (2009). A guide to the implementation of the WHO multimodal hand hygiene improvement
614 strategy. WHO/IER/PSP/2009.02. Available at:
615 https://www.who.int/gpsc/5may/Guide_to_Implementation.pdf?ua=1
616
617 WHO (2020). WHO Coronavirus disease (COVID-19) dashboard. World Health Organization. Available
618 at: <https://covid19.who.int/>
619
620 Woo, P. C., Lau, S. K., & Yuen, K-Y. (2006). Infectious diseases emerging from Chinese wet-markets:
621 Zoonotic origins of severe respiratory viral infections. *Current Opinion in Infectious Diseases* 19(5),
622 401-7
623
624 Wu, D., Wu, T., Liu, Q., & Yang, Z. (2020). The SARS-CoV-2 outbreak: What we know. *International
625 Journal of Infectious Diseases*, 94, 44-48.
626
627 Yuan, J., Lau, E. H. Y., Li, K., Leung, Y. H. C., Yang, Z., Xie, C. et al. (2015). Effect of live poultry
628 market closure on avian influenza A (H7N9) virus activity in Guangzhou, China, 2014. *Emerging
629 Infectious Diseases*, 21(10), 1784-1793.
630
631 Yuan, J., Tang, X., Yang, Z., Wang, M., & Zheng, B. (2014). Enhanced disinfection and regular
632 closure of wet markets reduced the risk of avian influenza A virus transmission. *Clinical Infectious
633 Diseases*, 58(7), 1037-1038.
634
635 Zhong, S., Cragg, M., & Zeng, G. (2020). Constructing freshness: the vitality of wet markets in China.
636 *Agriculture and Human Values*, 37, 175-185.
637
638 Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W. et al. (2020). A pneumonia outbreak
639 associated with a new coronavirus of probable bat origin. *Nature*, 579, 270-273.
640