

Central Lancashire Online Knowledge (CLOK)

Title	A comparison of the pharmacodynamic effects of intravenous ketamine-xylazine with alfaxalone in mute swans (Cygnus olor) presenting at a wildlife veterinary hospital
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/39037/
DOI	https://doi.org/10.1016/j.vaa.2021.03.014
Date	2021
Citation	Baldrey, Vicki, Stanford, Michael and Bacon, Heather (2021) A comparison of the pharmacodynamic effects of intravenous ketamine-xylazine with alfaxalone in mute swans (Cygnus olor) presenting at a wildlife veterinary hospital. <i>Veterinary Anaesthesia and Analgesia</i> . ISSN 1467-2987
Creators	Baldrey, Vicki, Stanford, Michael and Bacon, Heather

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.vaa.2021.03.014>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Table 2. Distribution of age categories, sex, bodyweight (Kg) and body condition score (BCS 1-5/5) of 58 mute swans given either intravenous (IV) alfaxalone 10 mg kg⁻¹ (Group A, *n* = 27/58) or IV ketamine 12.5 mg kg⁻¹ and xylazine 0.28 mg kg⁻¹ (Group KX, *n* = 31/58) for the induction of anaesthesia before maintenance with isoflurane in oxygen.

	<i>n</i>	Adult	Juvenile	Male	Female	Sex unknown	Weight (kg)*	BCS*
Group A	27	18	9	16	7	4	6.73 ±1.73 (3.65-9.8)	1.5 ±0.63 (0.5-3)
Group KX	31	21	10	13	9	9	7.7 ±1.42 (4.75-10.0)	2.0 ±0.55 (0.5-3)

*Values shown as mean ± standard deviation (range)