

Central Lancashire Online Knowledge (CLoK)

Title	New horizons for the methodology and physiology of training periodization: Block Periodization: New horizon or a false dawn?
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/5139/
DOI	https://doi.org/10.2165/11535130-000000000-00000
Date	2010
Citation	Kiely, J. (2010) New horizons for the methodology and physiology of training periodization: Block Periodization: New horizon or a false dawn? Sports Medicine, 40 (9). pp. 803-805. ISSN 0112-1642
Creators	Kiely, J.

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.2165/11535130-000000000-00000>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Editorial Manager(tm) for Adis, a Wolters Kluwer business
Manuscript Draft

Manuscript Number:

Title: Letter: Block Periodization: New horizon, or a false dawn?

Short Title: Block Periodization: New horizon, or a false dawn?

Article Type: Letter to the Editor

Section/Category: Sports Medicine

Keywords:

Corresponding Author: mr john kiely, HDip, BSc(Hons), MSc

Corresponding Author's Institution: UK Athletics

First Author: john kiely, HDip, BSc(Hons), MSc

Order of Authors: john kiely, HDip, BSc(Hons), MSc

Manuscript Region of Origin: UNITED KINGDOM

Abstract:

Block Periodization: New horizon, or a false dawn?

Professor Issurin's paper is to be commended on its insightful overview of the historical evolution of periodization planning theory, and the interesting general discussion. However, the paper's central contention, i.e., that block periodization represents a 'new horizon' in training planning is, I suggest, both premature and unsupported.

To substantiate this position; consider the rationale and evidence presented within the "New Horizons" paper promoting the superiority of block periodization in elite training contexts.

Essentially the presented argument consists of two layers of rationale. The first layer is anecdotal, and consists solely of exemplar cases of athletes and coaches who have achieved high levels of success employing block training designs. However, within the elite sports environment it would seem readily apparent that high honours are commonly achieved using a variety of training approaches, reflecting distinct coaching philosophies, and differing planning models. Hence, while the offered examples are undoubtedly interesting and deserve consideration, they remain unconvincing as evidence, as they lack both contextual detail and critical comparisons. Selecting tailored examples to substantiate a certain stance is not a particularly persuasive, clinical, or impartial argument.

The second layer of supporting evidence refers to "*two contemporary scientific concepts*" that have been instrumental in the establishment of the block periodized model: namely; the cumulative training effect and the residual training effect. However, within the review the key citations for these concepts are not evidence-led scientific discussions but rather, self-referenced opinion pieces by the author and by well-known block periodization advocate Dr Anatoly Bondarchuk. In reality, acknowledging that the benefits of physical training gradually accumulate over time (the cumulative

effect), and that these benefits persist for some period after training is terminated (the residual effect) are perhaps best described as self-evident truths, as opposed to scientific constructs. In fact, Matveyev (1981), the foremost formulator of the traditional periodization model, discusses the cumulative training effect, and concepts corresponding to the residual training effect (although using a different terminology) in his influential *Fundamentals of Sports Training* (1981)⁽¹⁾. What is not clear is how an awareness of such poorly understood concepts provides scientific support for block periodization principles. In order to discriminate between either traditional or block planning methods on the basis of these very broad concepts, specific knowledge would be required relating to; the projected time-frames for retention or decay of specific fitness attributes, an understanding of how on-going training interacts with previously conducted training to either accelerate or delay the erosion of previously developed fitness components, and an understanding of how these factors interact with a spectrum of individual-specific considerations, such as training histories and genetic predispositions. A knowledge base which clearly does not exist.

Consequently, while the proffered anecdotal examples and accompanying logic may be alluring, block periodization cannot be rightly framed as a scientifically-validated planning construct, any more than could Matveyev's seminal model, or the raft of subsequently proposed periodization derivations^(2,3,4,5). In essence, the presented argument is notional, rather than factual. Here, I hasten to add; experienced coach/scientist opinion is certainly not to be underestimated, devalued or dismissed. Likewise, a lack of evidence does not necessarily invalidate the model. However, before block periodization can rightly claim to be scientifically supported, an evidence-led, conceptually-valid chain of reasoning surely needs to be more coherently outlined.

As an additional concern; while there is an apparent dearth of evidence supporting the block periodization concept, there is existing evidence that would appear to strongly challenge its central premise, i.e. that *“each of these (fitness) targets requires specific physiological, morphological and psychological adaptation, and many of these workloads are not compatible, causing conflicting responses”*, and that hence *“high performance athletes enhance their preparedness and performance through large amounts of training stimuli that can hardly be obtained using multi-targeted mixed training”* (P 194).

Unravelling the interactions between multi-targeted mixed training modes is obviously a complex task to address empirically. However, it has been tangentially explored in studies investigating the effects of concurrent strength and endurance training. The training modes required to simultaneously develop enhanced strength or endurance appear diametrically opposed, and these attributes would appear prime candidates for exhibiting compromised training effects. As a brief recap of the literature; Hickson (1980) classically demonstrated an ‘interference effect’ between concurrent strength and endurance training resulting in compromised strength development in previously untrained subjects⁽⁶⁾, with similar findings subsequently reported by several authors⁽⁷⁻¹⁰⁾. More recently, however, studies have demonstrated that concurrent training can be as effective in developing both strength and endurance as single attribute-focused interventions^(11,12). More pertinently, studies in a wide variety of sports, variously using well-trained, elite, and World class athletes, have established that simultaneously training both strength and endurance can bestow synergistic benefits to a variety of athletic performance measures, above and beyond the benefits realised by single modality training, and without inhibition of strength development⁽¹³⁻²⁶⁾. Without doubt there is still much to be learned in relation to the intricacies of concurrent training, and key questions remain. However, it also appears clear from the spectrum of evidence that multi-modal training can be effective in enhancing specific performance attributes in already well-trained

athletes, and that the potential exists for various training modes to interact synergistically and additively.

An apparently logical interpretation of the available evidence suggests that whether or not concurrent multi-mode training has an antagonistic, neutral, or synergistic effect is dependent on the interaction between training design considerations such as how training modes are blended, timed and sequenced, and athlete-specific variables, such as training histories, genetic predispositions, and transient biological states. As a relevant additional consideration; the potential benefits of regular training variation have been previously elucidated ^(27,28), as have the potential negative effects of monotonous, unremitting, uni-directional training ⁽²⁹⁻³²⁾.

This is certainly not to suggest that multi-modal training is always advisable and, in the interests of balance, it should be noted that a recent study has demonstrated an improvement in outcomes following an 11 day high intensity endurance training intervention in alpine skiers ⁽³³⁾. Although the design does not necessarily conform to the description of block periodization as outlined in the ‘new horizon’ paper, the study authors do suggest that this finding illustrates the potential superiority of block periodization. However, this may be an overly elaborated conclusion, and perhaps a more parsimonious perspective is that such a finding demonstrates the value of periodically interjecting novelty into habituated training patterns, hence potentially offsetting diminishing training returns, and facilitating a heightened adaptive response.

Reflecting on the evidence discussed it would appear pre-mature to herald block periodization as a “new horizon” in training planning; partly because of a fundamental lack of supporting evidence and clearly delineated rationale, and partly as contradictory evidence exists questioning its universal

1 efficacy in elite contexts. What block periodization does positively contribute to current planning
2 methodologies is a more formal description of a particular planning tactic that may be
3
4 advantageously added to the elite coaches menu of potential planning options.
5
6
7
8
9

10 With reference to potential new horizons in training planning, it is interesting to note that all
11 previous periodization incarnations have been based upon a common set of unexamined
12 mechanistic assumptions that have become deeply engrained in training planning culture. Namely,
13 that optimal future training patterns can be adequately predetermined, that the training process is
14 best designed around a pre-formed template of discrete sequential training units (blocks, phases, or
15 periods), and that there exist relatively stable, predictable time-frames for the realisation and decay
16 of the various fitness attributes. In other words, the assumption that future elite training can be
17 adequately pre-planned.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 However, substantial evidence emanating from across the spectrum of biological sciences serves to
35 illustrate that the human adaptive response to any set of imposed stressors vary widely on both
36 inter-, and intra-, individual dimensions⁽³⁴⁻³⁷⁾. Accordingly, individuals are likely to respond uniquely
37 to any given training session, and will similarly respond in an individually-specific manner to any
38 given training organisational scheme^(38,39). So perhaps a universal limitation, shared by both
39 traditional and block periodized models, is the paradoxical assumption that the future training of an
40 inherently unpredictable and complex biological system, can be effectively pre-planned using a logic
41 rooted in mechanistic assumptions and generalised rules.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57 Hence, perhaps the true new horizon in elite training planning lies not in devising additional
58 idealised, rule-based, pre-planned training templates (as per the various periodization conceptions).
59
60
61
62
63
64
65

Perhaps, instead, the way forward lies in the design of sensitive and responsive self-organising training processes that guide the evolution of context-specific training solutions. Such training systems need not necessarily be based upon any single periodization ideology, but would facilitate the emergence of appropriate training systems as guided by the on-going triangulation of such factors as; training objectives, training readiness, and training responses. Trend analysis of this data could hence facilitate the consistent re-calibration and modulation of training to offset diminishing returns consequent to overly habituated training. The implementation of such organic, evolving training systems has historically been inhibited by the lack of sufficiently sensitive monitoring tools. However, this circumstance would appear to be rapidly changing.

Such a radical departure from traditional deterministic periodized planning paradigms would indeed herald a new horizon in sports training planning.

References

1. Matveyev L. Fundamentals of Sports Training. Moscow: Fizkultura i Sport. 1981
2. Brown LE. Nonlinear Versus Linear Periodization Models. National Strength & Conditioning Association 2001: 23:1, 42–44
3. Brown LE, Greenwood M. Periodization Essentials and Innovations in Resistance Training Protocols. Strength and Conditioning Journal 2005: Vol. 27, No. 4, 80–85
4. Rhea MR, Ball SD, Phillips WT, Burkett L. A Comparison of Linear and Daily Undulating Periodized Programs with Equated Volume and Intensity. The Journal of Strength and Conditioning Research 2002: May;16(2):250-5

5. Verkhoshansky YV. Programming and Organization of Training. 1988. Livonia, MI: Sportivny Press. (Original work published in 1985, Moscow, Russia: Fizkultura i Sport)
6. Hickson RC. Interference of strength development by simultaneously training for strength and endurance. European Journal of Applied Physiology and Occupational Physiology 1980: 45: 2-3
7. Hennessy LC, Watson WS. The Interference Effects of Training for Strength and Endurance simultaneously. The Journal of Strength and Conditioning Research 1994: 8:1
8. Dudley GA, Djamil R. Incompatibility of endurance- and strength-training modes of exercise. J Appl Physiol 1985: 59: 1446-1451
9. Hunter G, Demment R, Miller D. Development of strength and maximum oxygen uptake during simultaneous training for strength and endurance. Journal of sports medicine and physical fitness 1987: vol. 27, 3, 269-275
10. Nelson AG, Arnall DA, Loy SF, Silvester LJ, Conlee RK. Consequences of combining strength and endurance training regimens. Phys Ther 1990: May;70(5):287- 94
11. McCarthy JP, Agre JC, Graf BK, Pozniak MA, Vailas AC. Compatibility of adaptive responses with combining strength and endurance training. Med Sci Sports Exerc. 1995: Mar;27(3):429-36.
12. Shaw BS, Shaw I, Brown GA. Comparison of resistance and concurrent resistance and endurance training regimes in the development of strength. Journal of Strength & Conditioning Research. 2009: Dec;23(9):2507-14.
13. Yamamoto LM, Klau JF, Casa DJ, Kraemer WJ, Armstrong LE, Maresh CM. The effects of resistance training on road cycling performance among highly trained cyclists: a systematic review. J Strength Cond Res. 2010 Feb;24(2):560-6.
14. Izquierdo-Gabarren M, González de Txabarri Expósito R, García-Pallarés J, Sánchez-Medina L, Sáez de Villarreal E S, Izquierdo M. Concurrent Endurance and Strength Training Not To

Failure Optimizes Performance Gains. Med Sci Sports Exerc. 2009 Dec 9. [Epub ahead of print]

15. Balabinis CP, Psarakis CH, Moukas M, Vassiliou MP, Behrakis PK. Early phase changes by concurrent endurance and strength training. Journal of Strength & Conditioning Research 2003: May;17(2):393-401.
16. Davis WJ, Wood DT, Andrews RG, Elkind LM, Davis WB. Concurrent training enhances athletes' strength, muscle endurance, and other measures. Journal of Strength & Conditioning Research 2008: Sep;22(5):1487-502.
17. Hickson RC, Dvorak BA, Gorostiaga EM, Kurowski TT, Foster C. Potential for strength and endurance training to amplify endurance performance. J Appl Physiol. 1988: Nov;65(5):2285-90.
18. Mikkola JS, Rusko HK, Nummela AT, Paavolainen LM, Häkkinen K. Concurrent endurance and explosive type strength training increases activation and fast force production of leg extensor muscles in endurance athletes. Journal of Strength & Conditioning Research 2007: May;21(2):613-20.
19. Mikkola J, Rusko H, Nummela A, Pollari T, Häkkinen K. Concurrent endurance and explosive type strength training improves neuromuscular and anaerobic characteristics in young distance runners. Int J Sports Med 2007: Jul;28(7):602-11
20. Paavolainen L, Häkkinen K, Hämläinen I, Nummela A, Rusko H. Explosive-strength training improves 5-km running time by improving running economy and muscle power. J Appl Physiol 1999: May;86(5):1527-33.
21. Millet GP, Jaouen B, Borrani F, Candau R. Effects of concurrent endurance and strength training on running economy and VO₂ kinetics. Med Sci Sports Exerc 2002: 34: 1351–1359.
22. Hickson RC, Dvorak BA, Gorostiaga EM, Kurowski TT, Foster C. Potential for strength and endurance training to amplify endurance performance. J Appl Physiol 1988: 65: 2285–2290.

23. B. R. Rønnestad, E. A. Hansen, T. Raastad Strength training improves 5-min all-out performance following 185 min of cycling. *Scand j sports sci* 2009: [Epub ahead of print]
24. Hoff J, Gran A, Helgerud J. Maximal strength training improves aerobic endurance performance. *Scand J Med Sci Sports* 2002; 12: 288–295.
25. Hoff J, Helgerud J, Wisloff U. Maximal strength training improves work economy in trained female cross country skiers. *Med Sci Sports Exerc* 1999; 31: 870–877
26. Støren O, Helgerud J, Støa EM, Hoff J. Maximal strength training improves running economy in distance runners. *Med Sci Sports Exerc* 2008; 40: 1087–1092.
27. Koutedakis Y, Jamurtas A. The Dancer as a Performing Athlete: Physiological Considerations. *Sports Medicine* 2004; 34(10):651-661
28. Koutedakis Y, Myszkewycz L, Soulas D, Papapostolou V, Sharp NCC. The effects of rest and subsequent training on selected physiological parameters in professional female classical dancers. *Int. J. Sports Med* 1999; 20(6): 379-383
29. Foster C Monitoring training in athletes with reference to overtraining syndrome. *Med. Sci. Sports Exerc.* 1998; 30:1164–1168
30. Foster C, Lehmann M. Overtraining syndrome. In: Guten GN (ed). *Running injuries*. pp 173-188. Philadelphia: WB Saunders. 1997
31. Suzuki S, Sato T, Maeda A, Takahasi Y Program Design Based on a Mathematical model using Rating of Perceived Exertion for an Elite Japanese Sprinter: A case study. *Journal of Strength and Conditioning Research* 2006;20(1), 36–42
32. Suzuki S, Sato T, Takahasi Y. Diagnosis of training program for a Japanese rower by using the index of monotony. *Can. J. Appl. Physiol* 2003;28(Suppl.):105–106
33. Breil FA, Weber SN, Koller S, Hoppeler H, Vogt M. Block training periodization in alpine skiing: effects of 11-day HIT on VO₂max and performance. *Eur J Appl Physiol* 2010: Apr 4. [Epub ahead of print]

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
34. Kudielka BM, Hellhammer DH, Wust S. Why do we respond so differently? Reviewing determinants of human salivary cortisol responses to challenge. *Psychoneuroendocrinology* 2009;34, 2-18
35. Bouchard C, Rankinen T, Chagnon YC, Rice T, Périusse L, Gagnon J, Borecki IB, An P, Leon AS, Skinner JS, Wilmore JH, Province M, Rao DC. Genomic scan for maximal oxygen uptake and its response to training in the HERITAGE Family Study. *J Appl Physiol* 2000;88(2): 551-559
36. Skinner JS, Jaskólski A, Jaskólska A, Krasnoff J, Gagnon J, Leon AS, Rao DC, Wilmore JH, Bouchard C. Age, sex, race, initial fitness, and response to training: the HERITAGE Family Study. *J Appl Physiol*. 2001;May;90(5):1770-6
37. Foster RG, Kreitzman L Rhythms of life: The biological clocks that control the daily lives of every living thing. Yale University Press. 2004
38. Beavan CM, Gill ND, Cook CJ. Salivary testosterone and cortisol responses in professional rugby players after four resistance exercise protocols. *J Strength Cond Res*. 2008 :Mar;22(2):426-431
39. Beavan CM, Cook CJ, Gill ND. Significant strength gains observed in rugby players after specific resistance exercise protocols based on individual salivary testosterone responses. *J Strength Cond Res*. 2008 :Mar;22(2):419-25