

Central Lancashire Online Knowledge (CLOK)

Title	Direct and indirect orthotic management of medial compartment osteoarthritis of the knee
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/5636/
DOI	https://doi.org/10.1016/j.gaitpost.2006.11.098
Date	2006
Citation	Jones, R.K., Nester, C.J., Kim, W.Y., Tyson, S., Laxton, P., Jari, S., Johnson, D.S. and Richards, Jim (2006) Direct and indirect orthotic management of medial compartment osteoarthritis of the knee. <i>Gait & Posture</i> , 24 (Supp2). S141-S142. ISSN 09666362
Creators	Jones, R.K., Nester, C.J., Kim, W.Y., Tyson, S., Laxton, P., Jari, S., Johnson, D.S. and Richards, Jim

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.gaitpost.2006.11.098>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Table 1

Comparison of mean pre- vs. post-operative data for the STFU (short term) and LTFU (long term) follow-up groups ($p < 0.05$ significance level)

	Pre	Short-term	<i>p</i> -value	Pre	Long-term	<i>p</i> -value
Knee extension at initial contact (°) ^a	25.8	18.7	0.000	25.7	18.5	0.000
Peak knee extension in stance (°) ^a	11.5	5.3	0.000	9.3	5.4	0.062
Pelvic tilt—anterior (°)	19.5	22.0	0.048	17.6	19.2	0.114
Ankle plantar flexion at initial contact (°)	−10.2	−4.0	0.002	−8.1	−4.7	0.191
Velocity (m/s)	0.99	1.07	0.008	0.90	1.02	0.006
Stride length (m)	0.85	0.95	0.000	0.84	0.99	0.000
Cadence (steps/min)	143	137	0.129	124	123	0.696
Normalcy Index	308	136	0.000	295	142	0.001
Popliteal angle (°)	49.3	35.0	0.000	48.6	38.2	0.006
	<i>n</i> = 45 limbs			<i>n</i> = 26 limbs		

^a Positive value = knee flexion.

6. Discussion

The results of this study showed that percutaneous hamstring lengthenings were effective at improving knee extension at initial contact for both STFU and LTFU groups. Velocity and stride length were also significantly increased and these effects were maintained at long-term follow up. Increased leg length due to growth was not accounted for in the calculations and therefore caution should be taken with this interpretation. Popliteal angle demonstrated a significant improvement in the STFU and LTFU groups which is contrary to a previous finding [4]. The Normalcy Index demonstrated significant improvement in both the STFU and LTFU suggesting that normalization of gait (overall improvement) was accomplished when PMHL were done in conjunction with other appropriate soft tissue and bony surgeries. This technique minimizes the morbidity (reduces the scarring internally and externally) while maintaining comparable gait results to a group of patients that received open medial hamstring lengthenings [1]. Percutaneous medial hamstring lengthenings could be considered a viable option for lengthening hamstrings in appropriate patients, not only due to its effectiveness in improving gait outcomes, but also because of its simplicity.

References

- [1] DeLuca PA, et al. *J Pediatr Orthop* 1998;18:712–8.
- [2] Damron T, et al. *J Pediatr Orthop* 1991;11:514–9.
- [3] Schutte LM, et al. *Gait Posture* 2000;11:25–31.
- [4] Chang W-N, et al. *Gait Posture* 2004;19:298–304.

doi:10.1016/j.gaitpost.2006.11.097

PP-030

Direct and indirect orthotic management of medial compartment osteoarthritis of the knee

R.K. Jones^{a,*}, C.J. Nester^a, W.Y. Kim^c, S. Tyson^a, P. Laxton^b, S. Jari^c, D.S. Johnson^d, J.D. Richards^e

^a Centre for Rehabilitation and Human Performance Research, University of Salford, Salford, UK

^b Directorate of Podiatry, University of Salford, Salford, UK

^c Hope Hospital, Salford, UK

^d Stockport NHS Foundation Trust, Stockport, UK

^e Allied Health Professions, University of Central Lancashire, Lancashire, UK

1. Introduction

Osteoarthritis (OA) is a painful condition and affects approximately 80% of individuals by the age of 55 [1], with knee OA occurring two times more frequently than OA of the hand or hip [2]. The condition is more prevalent in the medial compartment and restricts the daily lives of individuals due to pain and a lack of functional independence. Patients with medial compartment osteoarthritis often have a varus alignment, with the mechanical axis and load bearing passing through this compartment with a greater adduction moment leading to greater pain and progression of osteoarthritis [3]. Surgery for the condition is possible although in some cases, particularly younger patients or those not yet requiring surgery, clinical management remains a challenge. Before surgery is considered, however, conservative management is advocated, though no one treatment has been shown to be most effective, and there are few quality biomechanical or clinical studies. Of the conservative approaches the principal orthotic treatments are valgus knee braces and laterally wedged foot inlays. Studies of knee valgus bracing have consistently demonstrated an associated decreased pain and improved function [4], and greater confidence [5]. A laterally wedged foot inlay has a thicker lateral border and applies a valgus moment to the heel. It is theorised that by changing the position of the ankle and subtalar joints during weight-bearing [6] the lateral wedges may apply a val-

gus moment across the knee as well as the rearfoot, with the assumed reduction on load in the medial knee compartment [7]. However, there has been no study to directly compare these orthotic treatments in the same study. The aim of this research is to investigate the efficacy of valgus knee braces and laterally wedged foot inlays in reducing the varus knee moment.

2. Statement of clinical significance

The study suggests that direct and indirect orthotic management of osteoarthritis can alter the biomechanical function of the knee and thereby reduce the loading on the medial compartment.

3. Methods

Eight subjects (four male and four female) (ages 64.8 years (S.D. 6.4) mass 80.8 kg (9.8), were recruited for the study. Each individual was radiologically diagnosed as having medial compartment osteoarthritis of the knee joint. The study was approved by the Salford Local Ethical Committee with consent given by the participants. The study took place at the University of Salford Gait laboratory. An eight-camera motion analysis system (Qualisys, Proreflex) capturing at 100 Hz and two bilateral force plates (Kistler) capturing at 200 Hz were used for the data collection. Marker placement was based on the Calibrated Anatomical System Technique [8] with cluster plates over the shank, thigh, pelvis and foot. Anatomical reference frames were modelled to represent the lower limb as segments in the local coordinate system. Standard footwear was worn for each testing session to ensure no confounding results from differing footwear and the participants performed 10 walking replicates, at a self-selected speed, in two conditions. Each participant wore a laterally wedged foot inlay (Algeos, Liverpool, UK) and a knee valgus brace (OAdjuster, Donjoy Orthopaedics, USA), for a period of 2 weeks in a randomised crossover design. To ensure reliability the foot inlays were manufactured by the same podiatrist, and the same individual fitted all braces. All data were exported to Visual 3D (C-Motion, Inc., USA) where interpolation and low pass filtering (6 Hz Kinematic, 15 Hz Kinetic) was undertaken. The peak knee adduction moment during loading was the primary outcome measure used to compare the two treatments. A repeated measures ANOVA test was performed with post hoc pair-wise comparison with a significance level of 5%, with a bonferroni adjustment to reduce the chance of type-1 errors.

4. Results

The peak knee adduction moment during loading in the laterally wedged foot inlays and valgus knee brace significantly

Table 1

Mean knee adduction moment (1S.D.) during the three conditions ($N=8$)

Condition	No orthotic	Knee brace	Foot inlay
Add moment (Nm/kg)	0.58 (0.08)	0.52 (0.08)	0.47 (0.07)

decreased (Table 1) compared to the no orthotic condition ($P<0.05$). There was a larger decrease in the laterally wedged inlays than the valgus knee brace. There was no significant difference in walking speed between all three conditions ($P>0.05$).

5. Discussion

The results suggest that both direct (knee brace) and indirect (laterally wedged inlay) orthotic interventions can reduce the adduction moment of the osteoarthritic knee. The results of this study will inform the sample size calculation for a phase II clinical trial to assess the biomechanical effects and clinical consequences of these interventions in people with medial compartment osteoarthritis of the knee.

It is hoped that the clinical efficacy of these treatments can be further defined by evaluating any correlation between the biomechanical, clinical and patient perceived outcomes. Providing biomechanical, clinical and patient based evidence to orthopaedic clinicians may increase the use of orthoses in those patients who are either not suitable for surgery or waiting for surgery. The next stage of the research is to the efficacy of combining both types of conservative treatment (direct and indirect) in the hope this will reduce the knee adduction moment even further.

Acknowledgements

The authors would like to thanks Algeo's and Donjoy Orthopaedics for their kind support of materials and braces for the study.

References

- [1] Altman R. *Am J Med* 1987;83:65–9.
- [2] Oliveria S, et al. *Arthritis Rheum* 1995;38:1134–41.
- [3] Kaufman KR, et al. *J Biomech* 2001;34:907–15.
- [4] Lindenfeld TN, et al. *Clin Orthop* 1997;344:240–7.
- [5] Richards JD, et al. *J Bone Joint Surg [Br]* 2005;87B:937–9.
- [6] Pollo FE. *Am J Knee Surg* 1998;11:47–50.
- [7] Jones RK, et al. ISB 2005. Cleveland, OH.
- [8] Cappozzo A, et al. *Clin Biomech* 1995;10:171–8.

doi:10.1016/j.gaitpost.2006.11.098