

Central Lancashire Online Knowledge (CLoK)

Title	EURRECA – Estimating zinc requirements for deriving dietary reference values
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/6843/
DOI	https://doi.org/10.1080/10408398.2012.742863
Date	2013
Citation	Lowe, Nicola M, Dykes, Fiona Clare, Skinner, Anna-louise, Patel, Sujata, Warthon-medina, Marisol, Decsi, Tamas, Fekete, Katalin, Souverein, Olga.W, Dullemeijer, Carla et al (2013) EURRECA – Estimating zinc requirements for deriving dietary reference values. Critical Reviews in Food Science & Nutrition, 53 (10). pp. 1110-1123. ISSN 1040-8398
Creators	Lowe, Nicola M, Dykes, Fiona Clare, Skinner, Anna-louise, Patel, Sujata, Warthon-medina, Marisol, Decsi, Tamas, Fekete, Katalin, Souverein, Olga.W, Dullemeijer, Carla, Cavelaars, Adriënne E, Serra-Majem, Lluís, Nissensohn,, Mariela, Bel, Silvia, Moreno, Luis A, Hermoso, Maria, Vollhardt, Christiane, Berti, Cristiana, Cetin, Irene, Gurinovic, Mirjana, Novakovic, Romana, Harvey, Linda J, Collings, Rachel and Moran, Victoria Louise

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1080/10408398.2012.742863>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

EURRECA – Estimating zinc requirements for deriving dietary reference values

Authors: Nicola M Lowe¹, Fiona C Dykes², Anna-Louise Skinner¹, Sujata Patel¹, Marisol Warthon-Medina¹, Tamás Decsi³, Katalin Fekete³, Olga W. Souverein⁴, Carla Dullemeijer⁴, Adriënne E. Cavelaars⁴, Lluís Serra-Majem⁵, Mariela Nissensohn⁵, Silvia Bel⁶, Luis A. Moreno⁶, Maria Hermoso⁷, Christiane Vollhardt⁷, Cristiana Berti⁸, Irene Cetin⁸, Mirjana Gurinovic⁹, Romana Novakovic⁹, Linda J. Harvey¹⁰, Rachel Collings¹⁰, Victoria Hall-Moran².

¹International Institute of Nutrition and Applied Food Safety Studies, University of Central Lancashire, Preston, UK

²Maternal & Infant Nutrition & Nurture Unit, University of Central Lancashire, Preston, UK

³Department of Paediatrics, University of Pécs, Hungary

⁴Division of Human Nutrition, Wageningen University, The Netherlands.

⁵Department of Clinical Sciences, University of Las Palmas de Gran Canaria, Spain.

⁶Escuela Universitaria de Ciencias de la Salud, Universidad de Zaragoza C/Domingo Miral s/n, 50009 Zaragoza, Spain.

⁷Ludwig-Maximilians-University of Munich, Dr. von Hauner Childrens Hospital, Div. Metabolic Diseases and Nutritional Medicine, Lindwurmstr. 4, D-80337, Munich, Germany.

⁸Department of Obstetrics and Gynecology, Ospedale San Paolo, Milan, Italy.

⁹Centre of Research Excellence in Nutrition and Metabolism, Institute for Medical Research, University of Belgrade, Serbia, Tadeusa Kosciuska 1, PO BOX 102, 11000 Belgrade, Serbia

¹⁰Norwich Medical School, Norwich Research Park, University of East Anglia, Norwich, UK

Corresponding author:

Nicola M Lowe

International Institute of Nutritional Sciences and Applied Food Safety Studies

Darwin Building, Room 206

University of Central Lancashire

Preston

Lancashire

PR1 2HE

UNITED KINGDOM

NMLowe@uclan.ac.uk

Tel :0044 (0) 1772 893599

Fax: 0044 (0) 1772 892925

Abbreviations

ACE Angiotensin-converting enzyme

ApoE Apolipoprotein E

EAR Estimated Average Requirement

EURRECA European Micronutrient Recommendations Alligned

DNA Deoxyribonucleic acid

DRV Dietary Reference Value

FFQ Food frequency questionnaire

GDS Geriatric depression scale

GST Glutathione S transferase

HbH Haemoglobin H

LBW Low birth weight

MMSE Mini mental state examination

PSS Perceived stress scale

RA Research activity

RCT Randomised controlled trials

RNA Ribonucleic acid

SES Socioeconomic status

SNP Single nucleotide polymorphism

WHO World Health Organisation

Abstract

Zinc was selected as a priority micronutrient for EURRECA, because there is significant heterogeneity in the Dietary Reference Values (DRVs) across Europe. In addition, the prevalence of inadequate zinc intakes was thought to be high among all population groups worldwide, and the public health concern is considerable. In accordance with the EURRECA consortium principles and protocols, a series of literature reviews were undertaken in order to develop best practice guidelines for assessing dietary zinc intake and zinc status. These were incorporated into subsequent literature search strategies and protocols for studies investigating the relationships between zinc intake, status and health, as well as studies relating to the factorial approach (including bioavailability) for setting dietary recommendations. EMBASE (Ovid), Cochrane Library CENTRAL and MEDLINE (Ovid) databases were searched for studies published up to February 2010 and collated into a series of Endnote databases that are available for the use of future DRV panels. Meta-analyses of data extracted from these publications were performed where possible in order to address specific questions relating to factors affecting dietary recommendations. This review has highlighted the need for more high quality studies to address gaps in current knowledge, in particular the continued search for a reliable biomarker of zinc status and the influence of genetic polymorphisms on individual dietary requirements. In addition, there is a need to further develop models of the effect of dietary inhibitors of zinc absorption and their impact on population dietary zinc requirements.

Key words: Zinc, Dietary recommendations, Zinc intake, systematic review, zinc status, zinc bioavailability, zinc requirements.

Introduction

Zinc is well established as an essential micronutrient for human health, having numerous structural and biochemical functions at the cellular and sub-cellular level, including enzyme function, DNA and RNA metabolism, protein synthesis, gene expression, cell growth and differentiation, and cell mediated immunity. Inadequate zinc intake has profound consequences at all points of the human lifecycle from the point of conception through to old age. Zinc was selected as a priority micronutrient for EURRECA, because the prevalence of inadequate zinc intakes was thought to be high among all population groups, and the public health concern is considerable. In addition, new scientific evidence has recently become available that demonstrates a large heterogeneity among current recommendations on zinc intake across Europe (Cavelaars et al., 2010; Doets et al., 2008) (See Activity 1 in (Dhonukshe-Rutten et al., 2013)).

The total amount of zinc present in the adult human body ranges from 1.5 to 2.5mg, most of which is intracellular, within skeletal muscle tissue (57%), bone (29%) and other tissues including skin and organs (Jackson, 1989). The zinc located within these tissues has a relatively slow turnover rate and is not readily responsive to changes in dietary zinc intake. Kinetic studies suggest that only a small proportion of total body zinc (approximately 10%) represents the “functional pool” of zinc, which is comprised of zinc, located within the liver and other tissues, that exchanges rapidly with the plasma, and when this functional pool is depleted zinc deficiency ensues (King, 1990). Zinc deficiency in adults can lead to dermatitis, hair loss, diarrhoea, loss of appetite, reproductive failure, hypogeusia, loss of cognitive function, susceptibility to infections and depressed immune function (Shankar and Prasad, 1998), delayed wound healing and depression (Andrews and Gallagher-Allred, 1999). Zinc deficiency has also been associated with three major health diseases prevalent in Europe: Diabetes, cancer and coronary heart disease (Singh et al., 1998). Zinc deficiency may be a

serious public health problem that compromises the development of millions of children (Sandstead and Smith, 1996). The recent Lancet series on maternal and child under nutrition concluded that zinc deficiency is responsible for about 4% of child mortality and disability-adjusted life-years (Black et al., 2008). The consequences and manifestations of severe zinc deficiency in infants and children and adolescents can be retardation of linear growth and development, poor appetite, delayed sexual maturation and hypogonadism, frequent infections (Maret and Sandstead, 2008), alopecia, dermatitis, delayed wound healing, diarrhoea, pneumonia, malaria (Fischer Walker et al., 2009), limitation on the senses of taste and smell, night blindness (Christian et al., 2001; Seiler et al., 2002). Zn deficiency may be associated with deficits in activity, attention, and motor development (Bhatnagar and Taneja, 2001). Results of several studies indicated that supplementation with zinc can significantly reduce the rates of diarrhoea and pneumonia in young children and increase the growth rate of stunted children (Brooks et al., 2005a; Brooks et al., 2005b; Brown et al., 2002). During the acute diarrhoea, zinc supplementation reduces the duration and severity of the disease, so that now the WHO recommends zinc supplementation as an adjunct to rehydration therapy, to replace the excessive losses of zinc during periods of diarrhoea (WHO). The involvement of maternal zinc status in pregnancy outcome is still unclear, animal models have shown that severe maternal zinc deficiency results in impaired implantation, abortions and foetal malformations (Keen et al., 2003). The consideration that zinc deficiency is a teratogenic risk in humans may be supported by the correlation of low plasma zinc concentrations in the first and third trimesters of pregnancy with an increased risk for malformations and low birth weight (LBW), respectively. Zinc deficiency is thought to influence embryonic and foetal development through reduced cell proliferation, or reduced protein synthesis or reductions in rates of tubulin polymerization rather than increased rates of cellular oxidative damage or increased rates of apoptosis and reduced binding of hormones and transcription factors

dependent on zinc-finger regions (Jankowski-Hennig et al., 2000; Mackenzie et al., 2002; WHO).

Current dietary recommendations

Dietary recommendations for zinc intake have been mainly based on balance studies focusing on the prevention of deficiency and use the factorial approach which assumes that the zinc requirement is the lowest intake which replaces obligatory zinc endogenous loss. The method computes the dietary zinc requirement by dividing the endogenous zinc loss by the fractional zinc absorption (King, 1986). In Europe and in other non-European countries, zinc recommendations for infants are generally set either based on zinc concentration of breast milk, using a factorial approach or extrapolating values from those given for adults.

There is significant heterogeneity among current recommendations on zinc intake across Europe and worldwide (Doets et al., 2008). Table 1 illustrates the range of dietary zinc recommendations for various countries worldwide. This heterogeneity is due to a number of factors including the data used to derive the value, and differences in expert opinion between panels convened to review the data.

Current European intakes

As part of the EURRECA programme of work, Roman-Vinas *et al* undertook an analysis of population dietary surveys from across Europe in order to determine the prevalence of inadequate nutrient intake in Europe using the Nordic Nutritional Recommendations as the standard (Roman-Vinas et al., 2010). This study revealed that the failure to meet the EAR (estimated average requirement) of 6.4mg/day or 5.7mg/d for adult males and female respectively was greatest in Ireland with dietary zinc intakes falling below this cut off value

in 11.9% men and 28.8% women. A similar picture emerged for elderly people (aged >64 years), with those living in Ireland having the highest percentage failing to meet the EAR, 13.6% and 13.1% of elderly men and women respectively (Vinas et al., 2011)

A review of available micronutrient intake and status data in Europe (Novakovic R, Submitted 2011) showed that data on intake of zinc were very limited for all life stages, so no cross country comparison could have been made. However, available data for zinc status (based on serum/plasma zinc concentrations) in children, adolescents and adults showed no regional differences when Central and Eastern Europe, Scandinavia, Western and Mediterranean countries were compared. All levels were within the optimal range indicating adequacy in zinc status (Novakovic R, Submitted 2011).

A systematic review of the relationship between micronutrient intake and socioeconomic determinants in Europe revealed that there were almost no differences in zinc intake between different socioeconomic status (SES) groups. On the other hand, status data in adults showed 5% higher serum zinc level in the low SES group. In comparison to reference values (Nordic nutrient recommendations for intake and the WHO for status (de Benoist et al., 2007)), all observed intake and status levels were within the optimal range, with the exception of levels of the low SES group in UK children (Novakovic R, Submitted 2012)

On an individual basis, an inadequate dietary intake of zinc could be the result of a strict vegan diet, a diet which is primarily based on grain products (Solomons and Slavin, 2001) or through a restrictive diet due, for example, to anorexia or alcohol or drug addiction. Certain disease states such as acrodermatitis enteropathica, Celiac disease, Crohn's disease and ulcerative colitis may disrupt the absorption of zinc (Solomons and Slavin, 2001). Other

health states may increase zinc losses primarily through diarrhoea, or increase requirements, such as the post-operative state (Solomons and Slavin, 2001).

The purpose of this review is to provide a summary of the methods used, and the results obtained from the systematic literature searches and subsequent meta-analysis of the data retrieved that were performed by partners in the EURRECA network of excellence. These activities were designed to answer specific questions regarding zinc-intake-status relationships. In addition, a comprehensive review of the factorial approach to setting zinc recommendations was undertaken. The overall aim of these activities was to generate new data and approaches that could assist future panels to derive dietary zinc recommendations using robust and transparent methodology.

Methods

Assessing dietary zinc intake:

One of the initial activities in the EURRECA process was to establish the most robust methodology of assessing zinc intake and status ((Matthys et al., 2010) and activity 3 in (Dhonukshe-Rutten et al., 2013)). The accurate determination of dietary micronutrient intake is notoriously problematic. Following a series of reviews of the methods used to assess micronutrient intake in Europe (Serra-Majem L, 2009) best practice guidelines were developed and adopted by the EURRECA network for all subsequent nutrient review activities. These are described in detail in “RA1.1 Best practice guidelines” in www.eurreca.org. In summary, only studies that used the following methodologies were included in the systematic reviews:

- 1) validated FFQ/Dietary History

- 2) validated 24h recall / food records / diary measurements for at least 3 days
- 3) validated 24h recall / food records / diary measurements < 3 days with adjustment for intra-individual variability

Since interventions commonly involve supplements, these were considered, taking into account the possible differences in bioavailability.

Assessing zinc status:

The assessment of zinc status is also problematic and it is generally accepted that there is currently no specific, reliable biomarker of zinc status. A systematic review and meta-analysis of the literature examining the efficacy of potential biomarkers of zinc status was undertaken (Lowe et al., 2009). This review presented an analysis of data from over 32 potential biomarkers however for many there was insufficient evidence to assess their reliability (Table 2)

Table 2. Biomarkers identified in systematic review (Lowe et al., 2009)

Plasma/serum zinc concentration was the most commonly used marker of zinc status and therefore the biomarker for which there were most data. It was found to respond to both increases and decreases in zinc intake, and was identified as being a useful biomarker however there are considerable reservations due to the effect of multiple confounders, such as infection, inflammatory status and time of last meal. Urine and hair were also considered useful biomarkers.

Health outcomes associated with inadequate zinc intake:

Important health problems related to zinc intake in adults and elderly people were identified by a literature search. These include: compromised immunity, dermatitis, hypogeusia, impaired cognitive functioning (dementia), depression, diabetes (reduced glucose tolerance), ischemic heart disease, carcinogenesis and anorexia. These were discussed and prioritized by the experts in zinc research within the RA2 team (Matthys C, 2011). Prioritisation was based on the strength of evidence of the deficiency and role of zinc on the health outcome, the relevance of the health outcome to the European population groups and the amount of evidence based research literature available based on a pilot literature search. The health outcomes that were identified for each population group are shown in Table 3, and are listed in order of priority.

Table 3. Priority health outcomes associated with inadequate zinc intake for each population group.

The best practice guidelines were then used to design the search protocols for the subsequent systematic reviews of the zinc intake- status- health relationships, the factorial approach for assessing dietary zinc requirements, zinc bioavailability and the influence of polymorphisms on zinc requirements. [Details of the search protocols can be found at: www.eurreca.org]

Results

Factorial approach and bioavailability

A technique commonly used when setting dietary zinc recommendations is the factorial approach which combines zinc required to replace obligatory losses with additional needs for zinc during different stages in the life cycle and makes adjustments for the bioavailability of

zinc in the diet (See Table 3 in (Dhonukshe-Rutten et al., 2013). Additional needs for zinc during different stages of life include that required for foetal growth during pregnancy, lactation and growth through infancy to adulthood. Literature searches were therefore designed to answer the following research questions: What are the key factors that affect zinc losses in all population groups? What are the key factors that affect zinc gains in all population groups? What are the additional needs for zinc during pregnancy, lactation and for growth? How well is zinc absorbed from meals and whole diets?

All titles and abstracts were screened for potential relevance and sorted into the population groups; infants, children and adolescents, pregnant and lactating women, adults, and elderly people as defined by the EURRECA consortium (See Activity 1 in (Dhonukshe-Rutten et al., 2013). An EndNote library was created compiling all the papers that met the inclusion criteria. The data regarding zinc losses and gains were extracted and collated using Excel (Microsoft Office Excel 2003). The quality and the risk of bias were assessed as indicators of validity. The studies included in this review were checked for a minimum quality score system developed by the EURRECA consortium which was adapted from The Cochrane Handbook for Systematic Reviews (Higgins and Green, 2008).

Factors affecting zinc losses and gains

From a total of 491 abstracts retrieved from electronic and hand searches, 105 appeared potentially relevant studies and were assessed for inclusion once the full paper had been obtained. Seventy-two papers were finally considered relevant across all population groups (adults and elderly, infants, children and adolescents, and pregnant and lactating women, adults and elderly). Despite the relatively stringent inclusion criteria, the included studies displayed a broad variety of methodological approaches, and were therefore unsuitable for

meta-analysis. Therefore the data extracted from the papers were tabulated and summarised narratively (Silvia Bel-Serrat, In Progress). Overall, balance studies have shown that zinc losses and gains are a function of the initial zinc status of an individual, the amount of bioavailable zinc in the diet and are modulated by homeostatic mechanisms. That means that dietary zinc recommendations should be estimated on the basis of the target population diet making difficult the possibility of establishing a value valid for the entire population. In addition, age, physical activity level, malabsorption syndromes, disease status can all affect zinc losses and gains. Moreover, interactions among nutrients should be also taken into account as they may also play an important role by means of affecting zinc utilization. As suggested by Taylor et al. (1991), the interaction between these homeostatic changes and zinc availability from different dietary sources should be better characterized to improve the accuracy of dietary zinc recommendations.

Factors affecting zinc bioavailability

The systematic review identified 120 studies as relevant to the research question of which 87 studies were conducted in adults and elderly, 2 in pregnancy and lactating women, 14 in children and adolescents and 17 in infants. Potential modifiers of zinc bioavailability were identified as illustrated in **Figure 1**. Phytate was the most frequently investigated modifier of zinc absorption. Twenty four estimates from seventeen studies that investigated the effect of dietary phytate level on zinc absorption were combined in a random effects meta-analysis. A forest plot showing the overall effect size of high versus low dietary phytate intake on zinc absorption is shown in **Figure 2**. The mean difference in fractional zinc absorption between low and high phytate diets was 0.11 (95% CI: 0.07, 0.16) however there was a high degree of between study heterogeneity (I^2 94% , $P < 0.0001$). Further analysis of this data set is underway to examine the factors that contribute to this heterogeneity and the overall effect of phytate:zinc molar ratio on zinc absorption.

The influence of gene polymorphisms zinc metabolism

The primary aim of this activity was to generate a database containing relevant information related to the impact of functional gene polymorphisms on zinc metabolism. Specifically, this involved identifying data assessing the impact of functional polymorphisms (e.g. single nucleotide polymorphisms, or SNPs) on micronutrient status biomarkers and associated health outcomes. The research questions used to develop the search protocol were: How do genetic polymorphisms affect zinc status? Are there any interactions between functional polymorphisms which affect zinc status and various health outcomes? Information was collated from studies of individuals who are either homozygous, heterozygous or wild type for specific polymorphisms. Where data exists for zinc status, functional polymorphisms and linked health outcome, this information was also recorded. Data were collated from all population groups including infants, children, adolescents and adults including the elderly into a database that is available at www.eurreca.org.

Of the 167 papers identified by the systematic search of the literature databases, 12 papers met the inclusion criteria and reported statistically significant results for altered zinc biomarker status in groups of people with differing gene variants. The gene interleukin 6 which regulates the amount of circulating proteins involved in inflammatory responses associated with hyperglycaemia and non-insulin dependent diabetes mellitus and coronary artery disease is thought to be influenced by zinc status (Giacconi et al., 2006; Giacconi et al., 2005). Two papers reported a relationship between SNP's of the Interleukin 6 gene and plasma zinc concentration and health outcomes, including perceived stress scale (PSS), geriatric depression scale (GDS) and mini mental state examination (MMSE) (Mariani et al., 2008; Mocchegiani et al., 2008)..

Another gene angiotensin-converting enzyme (ACE) was also reported. An impaired zinc status can alter enzyme activity, with adverse effects on angiotensin- conversion from I to II affecting vasoconstriction and hypertension and subjects with the DD genotype polymorphism in the ACE gene have been shown to have the highest enzyme activity increasing the risk of hypertension (Tamura et al., 1996). Tamura *et al* found a significant correlation between plasma zinc concentration and ACE activity in pregnant women at 33 weeks of gestation. However as this was the only significant correlation found in this study, it was stated that the significant result may have occurred by chance.

The metallothioneine gene has been included in the zinc database. Some polymorphisms of the metallothionein gene have been correlated to chronic inflammation and may affect zinc release (Mocchegiani et al., 2006; Richards et al., 2002). Another gene reported in the zinc database was apolipoprotein E (ApoE). Gonzalez *et al* (1999), reported that serum zinc concentrations in epsilon 4 ApoE carriers were significantly higher in patients with Alzheimer's disease than in healthy control patients.

The TP53 mutation in exon 5 through to 8, found in esophageal squamous cell carcinoma tumours was reported in a paper by Dar *et al* (2008). There is a notion that an imbalance of copper and zinc levels may lead to a higher prevalence of TP53 tumour mutations. Dar *et al* found that cancer patients with the TP53 tumour mutation had lower plasma zinc levels than those with no mutation.

The glutathione S transferase (GST) gene was reported by 3 papers (Jin et al., 2011; Reszka et al., 2007; Reszka et al., 2005). The detoxifying enzyme GST metabolises tobacco smoke derived compounds; zinc deficiency therefore can increase the risk of mutations occurring and can decrease the activity of the antioxidant GST enzyme increasing the risk of some cancers including lung cancer. Some polymorphisms in the GST gene may be associated with

an elevated risk of lung cancer and therefore the effect of zinc status on each variant allele needs to be investigated.

The gene CYP1A1 was also investigated by Jin *et al* (2011). The CYP1A1 gene has a polymorphism at exon 7 where a new MspI restriction site is introduced. The CYP1A1 gene is thought to influence metabolic activation and detoxification of some toxins and therefore can increase susceptibility to increasing risk of lung cancer. Jin *et al* reported that the risk of lung cancer decreased with a zinc level >1200ng/ml for both CYP1A1 variants and CYP1A1 carriers suggesting that a higher concentration of serum zinc may protect against lung cancer.

Haemoglobin H disease (HbH) was also included in the database and was thought to influence zinc status. Zinc deficiency is thought to be involved with impaired growth and hypogonadism traits observed in patients with polymorphic diseases such as the thalassemic diseases (Ajayi, 1997; Kajanachumpol *et al.*, 1997) and cystic fibrosis (Van Biervliet *et al.*, 2007).

The final gene reported as having a significant association with blood biomarker zinc status is SLC30A4 gene with a polymorphism on exon 5 915 T-C. The SLC30A4 gene encodes one of the zinc transport proteins and therefore it is thought that a polymorphism in this gene may affect zinc absorption and foetal development. Akar *et al* (2006) studied this gene and zinc status association and found that three hours after a zinc tolerance test there was a significant difference in plasma zinc level for TT and CC carriers, indicating a functional property of this polymorphism.

Table 5. Results of the systematic search for studies of the effect of gene polymorphisms on zinc metabolism.

Intake- status-health relationships

EURRECA is developing a method for the quantitative integration of evidence for deriving nutrient intake recommendations using bivariate dose-response relationships for intake-health (I-H) as well as intake-status (I-S) and status-health (S-H) relationships. These data will be combined in a new integrated trivariate intake-status-health (I-S-H) dose-response model with data from classical nutrition studies and bioavailability factors (See Activity 6 in (Dhonukshe-Rutten et al., 2013). Search protocols were therefore designed to answer the following questions: What is the effect of zinc intake on functional or clinical outcomes (intake-health) and what factors affect this relationship? What is the effect of zinc intake on indicators of exposure or body stores/biomarkers (intake-status) and what factors affect this relationship? What is the effect of indicators of exposure or body stores (i.e. biomarkers) on functional or clinical outcome (status-health) and what factors affect this relationship?

The result of the systematic search for studies addressing zinc intake status health relationships yielded over 1000 articles that were obtained in full text for eligibility evaluation. Due to the heterogeneity of the methodological approaches and outcome measures used in these studies, it is unlikely that meta-analysis of the data will be possible.

Intake- status relationships

Sufficient high quality RCT studies were identified to enable a meta-analysis of data describing the relationship between plasma zinc intake and plasma zinc concentration in each of the population categories. Units of measurement were converted to a standard form to facilitate comparison across studies. Intake-status regression coefficients ($\hat{\beta}$) were estimated for each individual study as described in detail elsewhere (Souverein et al., 2012). An overall pooled $\hat{\beta}$ and $SE(\hat{\beta})$ was calculated using random effects meta-analysis The statistical

transformations to obtain $\hat{\beta}$ s and $SE(\hat{\beta})$'s were performed using GenStat version 13-SP2 (VSN International Ltd., <http://www.vsnl.co.uk/>) and the meta-analysis was performed using STATA version 11.0 (College Station, TX), with statistical significance defined as $P < 0.05$.

For all population groups, with the exception of lactating women, the intake-status analyses revealed a positive and significant relationship between zinc intake and plasma zinc concentration, however a high degree of heterogeneity between the studies was observed (Table 4).

Table 4: Results of the Meta-analysis of intake-status in all populations groups.

Discussion

The determination of dietary zinc recommendations has relied primarily on the factorial approach, with extrapolation to population groups for which data are limited or missing. A complementary approach involves examining the associations between dietary intake, status and health, to arrive at intakes that result in optimal status levels and are sufficient to prevent disease due to deficiency at one end of the spectrum, or toxic effects due to excess dietary zinc at the other end of the spectrum. The difficulty of this approach for zinc is the lack of a reliable and sensitive marker of zinc status, and the non-specific nature of the diseases symptoms associated with sub-optimal zinc intake. However, one of the overarching aims guiding the work described in this review was to gather the best quality data using the most robust methodology to provide a database for future panels to use when setting recommendations. This included both a comprehensive review of the data available using both the classical factorial approach and a more novel intake-status-health association approach for zinc.

Regarding the factorial approach, a key factor that has been highlighted in this review and in discussion with experts is the need to consider bioavailability more closely as a modifier of the amount of dietary zinc required to meet requirements (Hambidge, 2010). This review identified a broad range of dietary components that may impact on the amount of dietary zinc that is absorbed and utilised, the majority of which had a deleterious effect on zinc bioavailability (figure 1). Many of these food components require further studies to generate sufficient high quality data to enable conclusive evaluation of their effect at different levels of intake, however the most widely studied modifier of zinc absorption is dietary phytate. This systematic review and meta-analysis confirms that phytate is a potent modifier of zinc absorption and should be taken into consideration when using the factorial approach to setting dietary zinc recommendations for any given population. Mathematical models that combine the effects of varying levels of phytate and zinc intake on true zinc absorption are potentially valuable tools in this process. A trivariate model (zinc intake-absorption-phytate intake), published by Hambidge *et al* in 2010 has helped to explain much of the variability of zinc absorption from human diets (Hambidge et al., 2010). This mathematical model is based on the accepted view that zinc absorption is a carrier-mediated process, phytate inhibits absorption by binding with zinc in the gut to form an insoluble complex, and that dietary zinc and phytate are the primary dietary factors determining zinc absorption. The model predicts that the quantity of zinc absorbed from 40 mg dietary zinc at zero phytate intake is 6.4 mg Zn/d and that the dietary zinc intake required to meet the requirements for zinc doubles with every 1000 mg phytate consumed in the diet per day (Hambidge et al., 2010).

Another potential modifier of the amount of dietary zinc needed to meet the requirement is genotype (Hambidge, 2010). This has been shown to have profound effects on the bioavailability of some micronutrients such as folate and iron (Casgrain et al., 2010). Most ZnT and Zip families show evidence of polymorphisms, which could produce structurally

different proteins and hence, transporter activity and /or specificity for zinc. Such polymorphisms could influence the amount of dietary zinc needed to meet the requirements and alter zinc metabolism (Cousins et al., 2006; Liuzzi and Cousins, 2004). The systematic search for studies on micronutrient metabolism yielded a very small number of relevant studies. This is clearly an important area for future research development.

Investigation of the zinc intake-status relationships in some population groups yielded some potentially useful new data. A dose-response curve was constructed from the extracted data, where the slope was based on the pooled $\hat{\beta}$ from the meta-analysis of the RCTs expressed on a \log_e -scale. Reported means and standard deviations of zinc intake and zinc status were extracted from the observational studies and was used to estimate the intercept of this curve (See Activity 6 in (Dhonukshe-Rutten et al., 2013)). The dose response curves for the Adult and Elderly, and the Pregnant and Lactating women population groups are shown in **Figure 3** and **Figure 4** respectively. These data can be used as complementary evidence for underpinning zinc reference values, however, the limitations of serum/plasma zinc concentration as a biomarker for zinc status should be acknowledged. Serum/plasma zinc is recognised as being a relatively insensitive index of zinc nutritional status due to efficient homeostatic regulation which responds to alterations in zinc intake, up-regulating absorption and conserving losses via the gastrointestinal tract and kidneys when intakes fall. In addition, whilst all studies included in the analysis were undertaken in apparently healthy individuals, factors such as stress, infection and inflammation, which are known to affect plasma zinc concentrations, may have gone unreported. Unfortunately, more sensitive indexes of zinc status have yet to be identified and plasma serum zinc remains by far the most commonly used biomarker of zinc status (Lowe et al., 2009). It is anticipated that this approach may be used to model the relationships between zinc intake or status with the health outcomes for zinc, and that these can be combined with the intake status relationships described above to

form a trivariate model of intake status and health (See Activity 6 in (Dhonukshe-Rutten et al., 2013). It is unclear at the moment whether or not our systematic searches have yielded sufficient data to enable this but it is likely that further studies are required.

This process has highlighted the need for more high quality studies to address gaps in current knowledge. Some of the key issues that came out of EURRECA workshops through discussion with experts external to the EURRECA Network focussed around zinc bioavailability and the need to model the effect of inhibitors of zinc absorption such as phytate, calcium and iron (Casgrain et al., 2010). Most current knowledge of zinc homeostasis is based on research in healthy adult males. In order to avoid scaling efforts should be made to obtain data on both genders at all ages, including pregnancy and lactation. In particular there is a paucity of data from studies in young children which necessitates the use of scaling to arrive at dietary recommendations, where this is the only option, there needs to be consensus regarding which growth/weight data to use.

In summary, a series of systematic reviews and meta-analyses were conducted in accordance with the protocols and procedures developed by the EURRECA consortium. This process has gathered together information relating to the setting dietary zinc recommendations which will be available as a valuable resource for future DRV panels. It has also generated new intake-status-health association data that may be used in combination with the classical factorial approach to model dietary zincs necessary to meet physiological requirements. This process has also highlighted the key areas for further research, in particular the urgent need for a reliable biomarker of zinc status, the further development of models of the impact of dietary factors on zinc bioavailability and the influence of genetic polymorphisms on individual dietary requirements.

Acknowledgements

The Authors would like to acknowledge the assistance of Joseph Saavedra, Sarah Richardson-Owen, Nick Kenworthy and Christine Cockburn with the data extraction and Lee Hooper for helpful advice. Our sincere thanks to Professor K Michael Hambidge for expert opinion and guidance throughout the EURRECA process. The preparation of this manuscript was coordinated by Rachel Collings from the University of East Anglia and the copy-editing by EUFIC.

This review was carried out with partial financial support from the Commission of the European Communities, specific RTD Programme "Quality of Life and Management of Living Resources", within the 6th Framework Programme (Contract No. FP6-036196-2 EURRECA: EUropean micronutrient RECommendations Aligned). This review does not necessarily reflect the views of the Commission and in no way anticipates the future policy in this area.

References

- Adams, C. L., Hambidge, M., Raboy, V., Dorsch, J. A., Sian, L., Westcott, J. L., and Krebs, N. F. (2002). Zinc absorption from a low-phytic acid maize. *Am J Clin Nutr.* **76**: 556-559.
- Ajayi, G. (1997). Zinc, magnesium, and copper concentrations in serum, erythrocytes, and urine in sickle cell homozygotes and heterozygote Nigerian women. *Trace Elements and Electrocytes.* **14** (2): 69-71.
- Akar, N., Akar, E., Arac, B., Ulu, A., and Ekim, M. (2006). Does SLC30A4 (ZNT4) gene exon 5 915 T-C alteration play a role in neural tube defects? *Trace Elements and Electrolytes.* **23** (4): 266-269.
- Andrews, M., and Gallagher-Allred, C. (1999). The role of zinc in wound healing. *Advances in wound care : the journal for prevention and healing.* **12**: 137-138.
- Australian Government Department of Health and Aging New Zealand Ministry of Health and National Health and Medical Research Council (2005). Nutrient reference values for Australia and New Zealand. pp. 221-227. Commonwealth of Australia.
- de Benoist B, Darnton-Hill I, Davidsson L, Fontaine O, Hotz C. Conclusions of the Joint WHO/UNICEF/IZiNCG Interagency meeting on zinc status indicators. *Food and Nutrition Bulletin* 2007; 28 (3) suppl. S480-4
- Bhatnagar, S., and Taneja, S. (2001). Zinc and cognitive development. *British Journal of Nutrition.* **85**: S139-S145.
- Black, R. E., Allen, L. H., Bhutta, Z. A., Caulfield, L. E., de, O. M., Ezzati, M., Mathers, C., Rivera, J., Maternal, and Child Undernutrition Study, G. (2008). Maternal and child undernutrition: global and regional exposures and health consequences. [Review] [147 refs]. *Lancet.* **371**: 243-260.

- Brooks, W. A., Santosham, M., Naheed, A., Goswami, D., Wahed, M. A., Diener-West, M., Faruque, A. S., and Black, R. E. (2005a). Effect of weekly zinc supplements on incidence of pneumonia and diarrhoea in children younger than 2 years in an urban, low-income population in Bangladesh: randomised controlled trial. *Lancet*. **366**: 999-1004.
- Brooks, W. A., Santosham, M., Roy, S. K., Faruque, A. S., Wahed, M. A., Nahar, K., Khan, A. I., Khan, A. F., Fuchs, G. J., and Black, R. E. (2005b). Efficacy of zinc in young infants with acute watery diarrhea. *Am J Clin Nutr*. **82**: 605-610.
- Brown, K. H., Peerson, J. M., Rivera, J., and Allen, L. H. (2002). Effect of supplemental zinc on the growth and serum zinc concentrations of prepubertal children: a meta-analysis of randomized controlled trials. *American Journal of Clinical Nutrition*. **75**: 1062-1071.
- Casgrain, A., Collings, R., Harvey, L. J., Boza, J. J., and Fairweather-Tait, S. J. (2010). Micronutrient bioavailability research priorities. *Am J Clin Nutr*. **91**: 1423S-1429S.
- Cavelaars, A. E., Kadvan, A., Doets, E. L., Tepsic, J., Novakovic, R., Dhonukshe-Rutten, R., Renkema, M., Glibetic, M., Bucchini, L., Matthys, C., Smith, R., van't Veer, P., de Groot, C. P., and Gurinovic, M. (2010). Nutri-RecQuest: a web-based search engine on current micronutrient recommendations. *European Journal of Clinical Nutrition*. **64**: Suppl-7.
- Christian, P., Khatry, S. K., Yamini, S., Stallings, R., LeClerq, S. C., Shrestha, S. R., Pradhan, E. K., and West, K. P. (2001). Zinc supplementation might potentiate the effect of vitamin A in restoring night vision in pregnant Nepalese women. *The American journal of clinical nutrition*. **73**: 1045-1051.
- Cousins, R. J., Liuzzi, J. P., and Lichten, L. A. (2006). Mammalian zinc transport, trafficking, and signals. [Review] [65 refs]. *Journal of Biological Chemistry*. **281**: 24085-24089.

- Couzy, F., Kastenmayer, P., Mansourian, R., Guinchard, S., Munoz-Box, R., and Dirren, H. (1993). Zinc absorption in healthy elderly humans and the effect of diet. *American Journal of Clinical Nutrition*. **58**: 690-694.
- Couzy, F., Mansourian, R., Labate, A., Guinchard, S., Montagne, D. H., and Dirren, H. (1998). Effect of dietary phytic acid on zinc absorption in the healthy elderly, as assessed by serum concentration curve tests. **In:** The British journal of nutrition, pp. 177-182.
- Dar, N. A., Mir, M. M., Salam, I., Malik, M. A., Gulzar, G. M., Yattoo, G. N., Ahmad, A., and Shah, A. (2008). Association between copper excess, zinc deficiency, and TP53 mutations in esophageal squamous cell carcinoma from Kashmir Valley, India--a high risk area. *Nutrition & Cancer*. **60**: 585-591.
- Department of Health (1991). Dietary Reference Values for Food Energy and Nutrients for the United Kingdom. **In:** Report on Health and Social Subjects 41, pp. 174-177. Panel on Dietary Reference Values for Food Energy and Nutrients for the United Kingdom, London.
- Dhonukshe-Rutten, R., Bouwman, J., Brown, K. A., Cavelaars, A. E., Collings, R., de Groot, L., Grammatikaki, E., Gurinovic, M., Harvey, L., Hermoso, M., Hurst, R., Kremer, B., Ngo, J., Novakovic, R., Raats, M. M., Rollin, F., Serra-Majem, L., Souverein, O. W., Timotijevic, L., and van 't Veer, P. (2013). EURRECA – Evidence-based methodology for deriving micronutrient recommendations. *Critical Reviews in Food Science and Nutrition*. **XX**: XX-XX.
- Doets, E. L., de Wit, L. S., Dhonukshe-Rutten, R. A., Cavelaars, A. E., Raats, M. M., Timotijevic, L., Brzozowska, A., Wijnhoven, T. M., Pavlovic, M., Totland, T. H., Andersen, L. F., Ruprich, J., Pijls, L. T., Ashwell, M., Lambert, J. P., van, t. V., and de Groot, L. C. (2008). Current micronutrient recommendations in Europe: towards

- understanding their differences and similarities. *European Journal of Nutrition*. **47**: Suppl-40.
- Egli, I., Davidsson, L., Zeder, C., Walczyk, T., and Hurrell, R. (2004). Dephytinization of a complementary food based on wheat and soy increases zinc, but not copper, apparent absorption in adults. *J Nutr*. **134**: 1077-1080.
- Fairweather-Tait, S. J., Fox, T. E., Wharf, S. G., Eagles, J., and Kennedy, H. (1992). Zinc absorption in adult men from a chicken sandwich made with white or wholemeal bread, measured by a double-label stable-isotope technique. *Br J Nutr*. **67**: 411-419.
- Farah, D. A., Hall, M. J., Mills, P. R., and Russell, R. I. (1984). Effect of wheat bran on zinc absorption. *Hum Nutr Clin Nutr*. **38**: 433-441.
- Fischer Walker, C. L., Ezzati, M., and Black, R. E. (2009). Global and regional child mortality and burden of disease attributable to zinc deficiency. *European Journal of Clinical Nutrition*. **63**: 591-597.
- Fredlund, K., Bergman, E. L., Rossander-Hulthén, L., Isaksson, M., Almgren, A., and Sandberg, A. S. (2003). Hydrothermal treatment and malting of barley improved zinc absorption but not calcium absorption in humans. *In: European Journal of Clinical Nutrition*, pp. 1507-1513.
- Fredlund, K., Isaksson, M., Rossander-Hulthen, L., Almgren, A., and Sandberg, A.-S. (2006). Absorption of zinc and retention of calcium: dose-dependent inhibition by phytate. *Journal of Trace Elements in Medicine & Biology*. **20**: 49-57.
- Giacconi, R., Cipriano, C., Muti, E., Costarelli, L., Malavolta, M., Caruso, C., Lio, D., and Mocchegiani, E. (2006). Involvement of -308 TNF-alpha and 1267 Hsp70-2 polymorphisms and zinc status in the susceptibility of coronary artery disease (CAD) in old patients. *Biogerontology*. **7**: 347-356.

- Giacconi, R., Cipriano, C., Muti, E., Costarelli, L., Maurizio, C., Saba, V., Gasparini, N., Malavolta, M., and Mocchegiani, E. (2005). Novel -209A/G MT2A polymorphism in old patients with type 2 diabetes and atherosclerosis: relationship with inflammation (IL-6) and zinc. *Biogerontology*. **6**: 407-413.
- Gonzalez, C., Martin, T., Cacho, J., Brenas, M. T., Arroyo, T., Garcia-Berrocal, B., Navajo, J. A., and Gonzalez-Buitrago, J. M. (1999). Serum zinc, copper, insulin and lipids in Alzheimer's disease epsilon 4 apolipoprotein E allele carriers. *European Journal of Clinical Investigation*. **29**: 637-642.
- Hambidge, K. M. (2010). Micronutrient bioavailability: Dietary Reference Intakes and a future perspective. *Am J Clin Nutr*. **91**: 1430S-1432S.
- Hambidge, K. M., Huffer, J. W., Raboy, V., Grunwald, G. K., Westcott, J. L., Sian, L., Miller, L. V., Dorsch, J. A., and Krebs, N. F. (2004). Zinc absorption from low-phytate hybrids of maize and their wild-type isohybrids. *Am J Clin Nutr*. **79**: 1053-1059.
- Hambidge, K. M., Miller, L. V., Westcott, J. E., Sheng, X., and Krebs, N. F. (2010). Zinc bioavailability and homeostasis. *Am J Clin Nutr*. **91**: 1478S-1483S.
- Higgins, J. P., and Green, S. (2008). Cochrane Handbook for Systematic Reviews Version 5.0.0. The Cochrane Collaboration.
- Hunt, J. R., and Beiseigel, J. M. (2009). Dietary calcium does not exacerbate phytate inhibition of zinc absorption by women from conventional diets. **In:** The American journal of clinical nutrition, pp. 839-843.
- Hunt, J. R., Matthys, L. A., and Johnson, L. K. (1998). Zinc absorption, mineral balance, and blood lipids in women consuming controlled lactoovovegetarian and omnivorous diets for 8 wk. *American Journal of Clinical Nutrition*. **67**: 421-430.

- Institute of Medicine (2000). Dietary Reference Intakes: vitamin C, vitamin E, selenium, and carotenoids. Washington, DC: National Academies Press.
- Jackson, M. J. (1989). Physiology of zinc: general aspects. In: Mills CF, ed. Zinc in Human Biology. London: Springer-Verlag. pp. 1-14.
- Jankowski-Hennig, M. A., Clegg, M. S., Daston, G. P., Rogers, J. M., and Keen, C. L. (2000). Zinc-deficient rat embryos have increased caspase 3-like activity and apoptosis. *Biochem Biophys Res Commun.* **271**: 250-256.
- Jin, Y., Zhang, C., Xu, H., Xue, S., Wang, Y., Hou, Y., Kong, Y., and Xu, Y. (2011). Combined effects of serum trace metals and polymorphisms of CYP1A1 or GSTM1 on non-small cell lung cancer: A hospital based case-control study in China. *Cancer Epidemiology.* **35 (2)**: 182-187.
- Kajanachumpol, S., Tatu, T., Sasanakul, W., Chuansumrit, A., and Hathirat, P. (1997). Zinc and copper status of thalassemic children. *Southeast Asian Journal of Tropical Medicine & Public Health.* **28**: 877-880.
- Keen, C. L., Clegg, M. S., Hanna, L. A., Lanoue, L., Rogers, J. M., Daston, G. P., Oteiza, P., and Uriu-Adams, J. Y. (2003). The plausibility of micronutrient deficiencies being a significant contributing factor to the occurrence of pregnancy complications. *J Nutr.* **133**: 1597S-1605S.
- Kim, J., Paik, H. Y., Joung, H., Woodhouse, L. R., Li, S., and King, J. C. (2007). Effect of dietary phytate on zinc homeostasis in young and elderly Korean women. *J Am Coll Nutr.* **26**: 1-9.
- King, J. C. (1986). Assessment of techniques for determining human zinc requirements. *J Am Diet Assoc.* **86**: 1523-1528.
- King, J. C. (1990). Assessment of zinc status. [Review] [28 refs]. *Journal of Nutrition.* **120**: Suppl-9.

- Larsson, M., Rossander-Hulthén, L., Sandström, B., and Sandberg, A. S. (1996). Improved zinc and iron absorption from breakfast meals containing malted oats with reduced phytate content. *In: The British journal of nutrition*, pp. 677-688.
- Liuzzi, J. P., and Cousins, R. J. (2004). Mammalian zinc transporters. [Review] [108 refs]. *Annual Review of Nutrition*. **24**: 151-172.
- Lowe, N. M., Fekete, K., and Decsi, T. (2009). Methods of assessment of zinc status in humans: a systematic review. [Review] [74 refs]. *American Journal of Clinical Nutrition*. **89**: 2040S-2051S.
- Mackenzie, G. G., Zago, M. P., Keen, C. L., and Oteiza, P. I. (2002). Low intracellular zinc impairs the translocation of activated NF-kappa B to the nuclei in human neuroblastoma IMR-32 cells. *J Biol Chem*. **277**: 34610-34617.
- Maret, W., and Sandstead, H. H. (2008). Possible roles of zinc nutriture in the fetal origins of disease. *Experimental Gerontology*. **43**: 378-381.
- Mariani, E., Neri, S., Cattini, L., Mocchegiani, E., Malavolta, M., Dedoussis, G. V., Kanoni, S., Rink, L., Jajte, J., and Facchini, A. (2008). Effect of zinc supplementation on plasma IL-6 and MCP-1 production and NK cell function in healthy elderly: interactive influence of +647 MT1a and -174 IL-6 polymorphic alleles. *Experimental Gerontology*. **43**: 462-471.
- Matthys, C., Bucchini, L., Busstra, M. C., Cavelaars, A. E., Eleftheriou, P., Garcia-Alvarez, A., Fairweather-Tait, S., Gurinovic, M., van, O. B., and Contor, L. (2010). EURRECA: development of tools to improve the alignment of micronutrient recommendations. *European Journal of Clinical Nutrition*. **64**: Suppl-31.
- Matthys C, v. T. V. P., de Groot L, Hooper L, Cavelaars AE, Collings R, Donutske-Rutten R, Harvey LJ, Casgrain A, Rollin F, Contor L. (2011). EURRECA's Approach for Estimating Micronutrient Requirements. *Int J Vitam Nutr Res*. **81**: 256-263.

Ministry of Health Labour and Welfare Japan (2005). Dietary Reference Intakes for Japanese.

In: The Report from the Scientific Committee of "Dietary Reference Intakes for Japanese- Recommended Dietary Allowance", p. 91.

Mocchegiani, E., Giacconi, R., Costarelli, L., Muti, E., Cipriano, C., Tesei, S., Pierpaoli, S., Giuli, C., Papa, R., Marcellini, F., Gasparini, N., Pierandrei, R., Piacenza, F., Mariani, E., Monti, D., Dedoussis, G., Kanoni, S., Herbein, G., Fulop, T., Rink, L., Jajte, J., and Malavolta, M. (2008). Zinc deficiency and IL-6 -174G/C polymorphism in old people from different European countries: effect of zinc supplementation. ZINCAGE study. *Experimental Gerontology*. **43**: 433-444.

Mocchegiani, E., Malavolta, M., Marcellini, F., and Pawelec, G. (2006). Zinc, oxidative stress, genetic background and immunosenescence: Implications for healthy ageing. *Immunity and Ageing*. **3**.

Nordic Council of Ministers (2004). Nordic Nutrition Recommendations: Integrating Nutrition and Physical Activity. pp. 397-402. Nordic Council of Ministers, Copenhagen.

Novakovic R, C. A., Bekkering E G, Roman B, Ngo J, Gurinovic M et al. (Submitted 2011). Micronutrients intake and status in Central and Eastern Europe as compared to other European countries, results from the EURRECA network.

Novakovic R, C. A., Geelen A, Nikolic M, Iglesia Altaba I, Vinas Roman B et al (Submitted 2012). Socioeconomic determinants of micronutrient intake and status in Europe: a systematic review

Petterson, D. S., Sandström, B., and Cederblad, A. (1994). Absorption of zinc from lupin (*Lupinus angustifolius*)-based foods. **In:** The British journal of nutrition, pp. 865-871.

- Reszka, E., Wasowicz, W., and Gromadzinska, J. (2007). Antioxidant defense markers modulated by glutathione S-transferase genetic polymorphism: Results of lung cancer case-control study. *Genes and Nutrition*. **2** (3): 287-294.
- Reszka, E., Wasowicz, W., Gromadzinska, J., Winnicka, R., and Szymczak, W. (2005). Evaluation of selenium, zinc and copper levels related to GST genetic polymorphism in lung cancer patients. *Trace Elements and Electrolytes*. **22** (1): 23-32.
- Richards, M., Hardy, R., and Wadsworth, M. E. (2002). Long-term effects of breast-feeding in a national birth cohort: educational attainment and midlife cognitive function. *Public Health Nutrition*. **5**: 631-635.
- Roman-Vinas, B., Ortiz-Andrellucchi, A., Mendez, M., Sanchez-Villegas, A., Quintana, L. P., Aznar, L. A. M., Hermoso, M., and Serra-Majem, L. (2010). Is the food frequency questionnaire suitable to assess micronutrient intake adequacy for infants, children and adolescents? *Maternal and Child Nutrition*. **6**: 112-121.
- Sandstead, H. H., and Smith, J. C., Jr. (1996). Deliberations and evaluations of approaches, endpoints and paradigms for determining zinc dietary recommendations. *J Nutr*. **126**: 2410S-2418S.
- Sandström, B., Arvidsson, B., Cederblad, A., and Björn-Rasmussen, E. (1980). Zinc absorption from composite meals. I. The significance of wheat extraction rate, zinc, calcium, and protein content in meals based on bread. **In:** The American journal of clinical nutrition, pp. 739-745.
- Sandstrom, B., Bugel, S., McGaw, B. A., Price, J., and Reid, M. D. (2000). A high oat-bran intake does not impair zinc absorption in humans when added to a low-fiber animal protein-based diet. *Journal of Nutrition*. **130**: 594-599.

- Sandstrom, B., and Sandberg, A. S. (1992). Inhibitory effects of isolated inositol phosphates on zinc absorption in humans. *Journal of Trace Elements and Electrolytes in Health and Disease*. **6**: 99-103.
- Scientific Committee for Food (1993). Nutrient and energy intakes for the European Community. Chapter 28: Selenium. **In**: Reports of the Scientific Committee for Food, Thirty First Series. Nutrition. , pp. 202-207. European Commission, Luxembourg.
- Seiler, W. O., Itin, P., and Stahelin, H. B. (2002). Zinc deficiency, a problem of old age frequently not recognized. *Ernahrungs-Umschau*. **49**: 260-+.
- Serra-Majem L, N. J., Roman-Vinas B. (2009). Micronutrient Intake Assessment in Europe: Best evidence and practice. . *British Journal of Nutrition*. **Vol 101**.
- Shankar, A. H., and Prasad, A. S. (1998). Zinc and immune function: The biological basis of altered resistance to infection. *American Journal of Clinical Nutrition*. **68**: 447S-463S.
- Silvia Bel-Serrat, A.-L. S., Marisol Warthon-Medina, Iris Iglesia-Altaba, Maria Hermoso, Luis A. Moreno, Nicola M. Lowe (In Progress). Factorial Approach to assessing zinc requirements and the affect of bioavailability in adults and elderly people: A systematic review and meta-analysis from the Eurreca Network.
- Singh, R. B., Niaz, M. A., Rastogi, S. S., Bajaj, S., Zhang, G. L., and Zhu, S. M. (1998). Current zinc intake and risk of diabetes and coronary artery disease and factors associated with insulin resistance in rural and urban populations of north India. *Journal of the American College of Nutrition*. **17**: 564-570.
- Solomons, N., and Slavin, J. L. (2001). What impact does stage of physiological development and/or physiological state have on the bioavailability of dietary supplements? Summary of workshop discussion. *J Nutr*. **131**: 1392S-1395S.

Souverein, O. W., Dullemeyer, C., van, T. V. P., and van der Voet, H. (2012).

Transformations of summary statistics as input in meta-analysis for linear dose-response models on a logarithmic scale: a methodology developed within EURRECA.

BMC Med Res Methodol. **12**: 57.

Swanson, C. A., Turnlund, J. R., and King, J. C. (1983). Effect of dietary zinc sources and

pregnancy on zinc utilization in adult women fed controlled diets. *Journal of Nutrition.* **113**: 2557-2567.

Tamura, T., Johanning, G. L., Goldenberg, R. L., Johnston, K. E., and DuBard, M. B. (1996).

Effect of angiotensin-converting enzyme gene polymorphism on pregnancy outcome, enzyme activity, and zinc concentration. *Obstetrics & Gynecology.* **88**: 497-502.

Taylor, C. M., Bacon, J. R., Aggett, P. J., and Bremner, I. (1991). Intestinal absorption and

losses of copper measured using ⁶⁵Cu in zinc-deprived men. *Eur J Clin Nutr.* **45**: 187-194.

Turnlund, J. R., King, J. C., and Keyes, W. R. (1984). A stable isotope study of zinc

absorption in young men: Effects of phytate and alpha-cellulose. *American Journal of Clinical Nutrition.* **40**: 1071-1077.

Van Biervliet, S., Van Biervliet, J. P., Vande Velde, S., and Robberecht, E. (2007). Serum

zinc concentrations in cystic fibrosis patients aged above 4 years: a cross-sectional evaluation. *Biological Trace Element Research.* **119**: 19-26.

Vinas, B. R., Barba, L. R., Ngo, J., Gurinovic, M., Novakovic, R., Cavelaars, A., de Groot, L.

C., van't Veer, P., Matthys, C., and Majem, L. S. (2011). Projected prevalence of inadequate nutrient intakes in Europe. *Ann Nutr Metab.* **59**: 84-95.

WHO Conclusions of the Joint WHO/UNICEF/IAEA/IZiNCG

Interagency Meeting on Zinc Status Indicators

<http://www.who.int/nutrition/publications/micronutrients/FNBvol28N3supsep07.pdf>.

Zheng, J., Mason, J., Rosenberg, I., and Wood, R. (1993). Measurement of zinc bioavailability from beef and a ready-to-eat high- fiber breakfast cereal in humans: application of a whole-gut lavage technique. *The American journal of clinical nutrition*. **58**: 902-907.

Figure 1. Flow diagram showing the results of the systematic review of studies investigating zinc bioavailability.




Figure 2 . Forest plot of high and low phytate meals and zinc absorption as % of the diet.





Figure 3. Serum/plasma zinc concentration ($\mu\text{mol/L}$) as a function of dietary zinc intake (mg/day), estimated by random-effects meta-analyses of RCTs. In figure 3a the data are presented on natural log scale, where $Y=0.11 \cdot x + 2.28$.

a



b.





Figure 4 Serum/plasma zinc concentration ($\mu\text{mol/L}$) as a function of dietary zinc intake (mg/day), estimated by random-effects meta-analyses of RCTs of pregnant (solid line) and lactating (dashed line) women. In figure 4a the data are presented on natural log scale.

4a



4b.




Table 1. - Selected recommended intake levels for zinc (mg)

DATA SOURC E	Gend er	Special condition s	POPULATION GROUP						
			Infant s	Childr en	Adolesce nts	Adult s	Elderl y	Lactati on	Pregnan cy
WHO / FAO	<i>Male</i>	High bioavailabil ity	0-6m	1-3y	10-18y	19- 65y	>65y	-	-
			1.1	2.4	5.1	4.2			
			7- 12m	4-6y		4.2			
			0.8	7-9y					
				3.3					
	<i>Femal e</i>	Low bioavailabil ity	0-6m	1-3y	10-18y	19- 50y	>65y	0-3m	< 3m.
			6.6	8.3	14.4	9.8	19	11	
			7- 12m	4-6y		9.8			
			8.4	7-9y					
				11.2					
Nordic (Norwa y)	<i>Male</i>	-	<6m	2-5y	14-17y	18- 30y	≥75y	-	-
			NUL	6	12	9			
			L	6-9y		9			
			6- 11m	7		31- 60y			
			5	11		9			
	<i>Femal e</i>	-	<6m	2-5y	14-17y	18- 30y	≥75y	-	-
			NUL	6	9	7			
			L	6-9y		7			
			6- 11m	7		31- 60y			
			5	8		7			
Australi a / NZ	<i>Male</i>	-	0-6m	1-3y	14-18y	19- 30y	>70y	-	-
			2	3	13	14			
			7- 12m	4-8y		14			
			3	4		31- 50y			
				9-13y		14			
	<i>Femal e</i>	-	0-6m	1-3y	14-18y	19- 30y	>70y	14-18y	14-18y
			2	3	7	8	11	10	

			7- 12m 3	4-8y 4 9-13y 6		8 31- 50y 8		19-50y 12	19-30y 11
DACH (German y)	<i>Male</i>	-	0-3m 1 4- 11m 2	1-3y 3 4-6y 5 7-9y 7 10-12y 7	13-14y 9.5 15-18y 10		≥65y 10	-	-
	<i>Femal</i>	-	0-3m 1 4- 11m 2	1-3y 3 4-6y 5 7-9y 7 10-12y 7	13-14y 7 15-18y 7	19- 64y 7	≥65y 7	10	>4m 10
	<i>e</i>								
EC	<i>Male</i>	-	6- 11m 4	1-3y 4 4-6y 6 7-10y 7	11-14y 9 15-17y 9	≥18y 9.5	≥18y 9.5	-	-
	<i>Femal</i>	-	6- 11m 4	1-3y 4 4-6y 6 7-10y 7	11-14y 9 15-17y 7	≥18y 7	≥18y 7	12	7
	<i>e</i>								
IOM (US/ Canada)	<i>Male</i>	-	0-6m 2 7- 12m 3	1-3y 3 4-8y 5 9-13y 8	14-18y 11	19- 70y 11	>70y 11	-	-
	<i>Femal</i>	-	0-6m 2 7- 12m 3	1-3y 3 4-8y 5 9-13y 8	14-18y 9	19- 70y 11	>70y 11	14-18y	19-30y
	<i>e</i>								

United Kingdom	Male	-	2	3		70y	8	12	11
			7-	4-8y		8		19-30y	31-50y
			12m	5				12	11
			3	9-13y					
				8					
	Female	-	0-3m	1-3y	11-14y	9	19-	>50y	-
			4	5			50y	9.5	
			4-6m	4-6y			9.5		
			4	6.5					
				7-10y					
	Male	-	0-3m	1-3y	14-18y	9	19-	>50y	NULL
			4	5			50y	7	NULL
			4-6m	4-6y			7		
			4	6.5					
				7-10y					

Data presented in the Table was obtained from online EURRECA web resource

NutriRecQuest: <http://www.serbianfood.info/eurreca/> (Cavelaars et al., 2010) from original source documents (Australian Government Department of Health and Aging New Zealand Ministry of Health and National Health and Medical Research Council, 2005; Department of Health, 1991; Institute of Medicine, 2000; Ministry of Health Labour and Welfare Japan, 2005; Nordic Council of Ministers, 2004; Scientific Committee for Food, 1993)

[†] RDA, Recommended Dietary Allowance (USA, Japan); equivalent to RNI, Reference Nutrient Intake (UK); PRI, Population Reference Intake (EFSA); RNI, Recommended Nutrient Intake (WHO/FAO); RDI, Recommended Dietary Intake (AU/NZ); RI, Recommended Intake (Nordic)

*upper level of safe intake for European Community from EC2006, other EC data from 1993

[#]Safe upper limit of selenium intake for UK from Expert Group on Vitamins and Minerals 2003, Safe Upper Levels for Vitamins and Minerals Food Standards Agency (full document available at <http://www.food.gov.uk/multimedia/pdfs/vitmin2003.pdf>)

'DACH' refers to reference intakes for Germany, Austria and Switzerland

'Nordic countries' refers to Denmark, Finland, Norway and Sweden.

'EC' refers to European Community.

Table 2. Biomarkers identified in systematic review (Lowe et al., 2009).

Potentially useful	<ul style="list-style-type: none"> • Plasma/serum zinc concentration • Hair zinc concentration • Urinary zinc excretion
Not useful	<ul style="list-style-type: none"> • Erythrocyte zinc concentration • Mononuclear cell zinc concentration • Polymorphonuclear cell zinc concentration • Platelet zinc concentration • Plasma alkaline phosphatase activity
Inconclusive due to lack of data	<ul style="list-style-type: none"> • Aminolevulinic acid dehydratase • Erythrocyte metallothionein • Monocyte metallothionein cDNA • Salivary zinc • Salivary-sediment zinc • Mixed saliva zinc • Plasma extracellular superoxide dismutase • Lymphocyte zinc • Lymphocyte ecto-5'-nucleotidase • Nail zinc • Plasma angiotensin-converting enzyme • Neutrophil zinc • T lymphocyte metallothionein -2A mRNA • Plasma 5'-nucleotidase • Endogenous zinc excretion • Plasma zinc flux • Exchangeable zinc pool • Carbonic anhydrase • Fecal Zinc • Neutrophil α-D-mannosidase • Neutrophil alkaline phosphatase • Erythrocyte membrane zinc • Erythrocyte membrane alkaline phosphatase • Erythrocyte membrane neutral phosphatase

Table 3. Priority health outcomes associated with inadequate zinc intake for each population group.

Infants	Children & Adolescent	Pregnant & lactating women	Adults/Elderly
Growth	Growth	<u>Foetus</u>	Immune function
Immune response to vaccination	Immune function	Foetal growth	Cognitive function
Neurodevelopment	Cognitive functions and Psychomotor development	Foetal malformation	Dermatitis
	Dermatitis	<u>Mother</u>	Anorexia
		Preeclampsia	Hypogeusia
		Preterm delivery	Ischemic heart disease
			Depression
			Diabetes Mellitus
			Carcinogenesis

Table 4: Results of the Meta-analysis of intake-status in all populations groups.

Population group	Overall Beta	95% CI's	I ²	p
Adults and Elderly	0.09	0.07,0.12	79.1%	P<0.0001
Pregnant women	0.04	0.02, 0.07	55%	p=0.002
Lactating women	0.02	-0.01,0.05	0%	p=0.28
Children and adolescents	0.12	0.04, 0.20	97.6%	p<0.005
Infants	0.09	0.06,0.12	95%	P<0.00001

Table 5. Results of the systematic search for studies of the effect of gene polymorphisms on zinc metabolism.

Author	Year	Reference	Gene with significant interaction with Zn
(Tamura et al.)	1996	Obstet Gynecol 88:497-502	ACE insertion/deletion DD, DI and II
(Mariani et al.)	2008	Experimental Gerontology 43:462-471 Eur J Clin Investigation 29:637-642	MT1a +674 A/C transition and Interleukin 6 IL-6 +174C/G transition
(Gonzalez et al.)	1999	Biogerontology 6:407-413	Epsilon 4 apoE
(Giacconi et al.)	2005	Nutrition and Cancer 60(5):585-591	MT2A rs1610216 AA and AG, 246bp, 131 and 115bp
(Dar et al.)	2008	Trace Elements and Electrolytes 22(1):23-32	TP53 mutation at exon 5-8
(Reszka et al.)	2005	Cancer Epidemiology 32:182-187	GSTP1, GSTT1 and GSTM1
(Jin et al.)	2011	Genes Nutr 2:287-294	CYP1A1 mspl Aa or aa and GSTM1null
(Reszka et al.)	2007	Experimental Gerontology 43:433-444	GSTM1 and GSTT1
(Mocchegiani et al.)	2008	Southeast Asian J Trop Med Public Health 28(4):877-880	IL-6 +174G/C
(Kajanachumpol et al.)	1997	Trace Elements and Electrolytes 14(2):69-71	HbH, β -thal/HbE, β -thal major
(Ajayi)	1997	Trace Elements and Electrolytes 23(4):266-269	HbAA, HbSS, HbAS and HbAC
(Akar et al.)	2006		SLC30A4 ZNT4 915T-C at exon 5