

Central Lancashire Online Knowledge (CLoK)

Title	A chemical analysis of the salt content of sandwiches purchased from independent outlets
Type	Article
URL	https://clock.uclan.ac.uk/13037/
DOI	https://doi.org/10.1017/S0029665115004036
Date	2015
Citation	Watts, Emilea Sarah, Davies, I.G., Sinclair, Jonathan Kenneth and Dillon, Stephanie (2015) A chemical analysis of the salt content of sandwiches purchased from independent outlets. <i>Proceedings of the Nutrition Society</i> , 74 (OCE5). ISSN 0029-6651
Creators	Watts, Emilea Sarah, Davies, I.G., Sinclair, Jonathan Kenneth and Dillon, Stephanie

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1017/S0029665115004036>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>



Summer Meeting, 6–9 July 2015, The future of animal products in the human diet: health and environmental concerns

A chemical analysis of the salt content of sandwiches purchased from independent outlets

E.S. Watts¹, I.G. Davies², J.K. Sinclair¹ and S.A. Dillon¹

¹Division of Sport Exercise and Nutritional Sciences, School of Sport, Tourism and The Outdoors, University of Central Lancashire, Preston, PR1 2HE, UK and ²Faculty of Education, Health and Community, IM Marsh Campus, Liverpool John Moores University, Liverpool, L17 6BD, UK

In recent years there has been an increase in eating away from home and sandwiches are popular choices which are often viewed as the healthy option^{1–2}. However, foods eaten away from home are characterised as having a high salt content and are generally nutritionally inadequate^{3–6}. Diet is widely recognized as a modifiable determinant of health and considering the prevailing problem of non-communicable disease⁷ the nutritional quality of food served from independent sandwich outlets is of considerable public health concern. Therefore the present pilot study investigated the salt content of sandwiches purchased from six independent outlets in the UK. A convenience sample of three popular categories of sandwiches (cheese, tuna, and chicken-tikka) were analysed and the results compared against national guidelines. Samples (n = 36) were analysed for sodium by atomic absorption spectrometry and values were converted to salt using a standard conversion factor (2.54)⁸. The calculated salt content of sandwiches (per portion as consumed) from independent outlets was high (Cheese: 3.00 ± 0.97 g, Chicken: 2.88 ± 0.74 g, Tuna: 2.78 ± 0.46 g). In comparison to current guidelines for intake of salt per meal for an adult (1.8 g) these levels equate to over-consumption by an average of 160 %. Regular consumption of sandwiches purchased from independent outlets, particularly in combination with other high salt foods, may contribute to the development of hypertension. Given the popularity of commercially prepared sandwiches, addressing the salt content would have a major public health benefit.

1. Tangari A, Burton S, Howlett E, Cho Y, Thyroff A. (2010) Weighing in on fast food consumption: the effects of meal and calorie disclosures on consumer fast food evaluations. *J Consumer Affairs* 44(3), 431–462.
2. Hwang J, Cranage D. (2010) Customer health perceptions of selected fast-food restaurants according to their nutritional knowledge and health consciousness. *J Foodservice Business Res* 13(2), 68–84.
3. Stender S, Dyerberg J, Astrup A. (2007) Fast food: unfriendly and unhealthy. *Int J Obesity* 31(6), 887–890.
4. Jaworowska A, Blackham T, Stevenson L, Davies IG. (2012) Determination of salt content in hot takeaway meals in the United Kingdom. *Appetite* 59(2), 517–522.
5. Nielsen S, Popkin B. (2003) Patterns and trends in food portion sizes, 1977–1998. *JAMA* 289(4), 450–453.
6. Prentice A, Jebb S. Fast foods, energy density and obesity: a possible mechanistic link. *Obesity Rev* 2003; 4(4), 187–194.
7. World Health Organization. (2010) *Global status report on noncommunicable diseases*. WHO.
8. Food Standard Agency. 2017 UK salt reduction targets. <http://www.food.gov.uk/scotland/scotnut/salt/saltreduction> (accessed 4th May 2014).