

## ASTRONOMY

# Shape of a slowly rotating star measured by asteroseismology

Laurent Gizon,<sup>1,2,3,4\*</sup> Takashi Sekii,<sup>3</sup> Masao Takata,<sup>5</sup> Donald W. Kurtz,<sup>6</sup> Hiromoto Shibahashi,<sup>5</sup> Michael Bazot,<sup>4</sup> Othman Benomar,<sup>4,5</sup> Aaron C. Birch,<sup>1</sup> Katepalli R. Sreenivasan<sup>4,7</sup>

Stars are not perfectly spherically symmetric. They are deformed by rotation and magnetic fields. Until now, the study of stellar shapes has only been possible with optical interferometry for a few of the fastest-rotating nearby stars. We report an asteroseismic measurement, with much better precision than interferometry, of the asphericity of an A-type star with a rotation period of 100 days. Using the fact that different modes of oscillation probe different stellar latitudes, we infer a tiny but significant flattening of the star's shape of  $\Delta R/R = (1.8 \pm 0.6) \times 10^{-6}$ . For a stellar radius  $R$  that is 2.24 times the solar radius, the difference in radius between the equator and the poles is  $\Delta R = 3 \pm 1$  km. Because the observed  $\Delta R/R$  is only one-third of the expected rotational oblateness, we conjecture the presence of a weak magnetic field on a star that does not have an extended convective envelope. This calls to question the origin of the magnetic field.

## INTRODUCTION

According to Clairaut's theorem, slowly rotating stars are oblate spheroids (1, 2). Other factors may affect the shapes of stars, such as magnetic fields, thermal asphericities, large-scale flows, or strong stellar winds. Global poloidal magnetic fields tend to make stars oblate, whereas toroidal magnetic fields tend to make them prolate (3, 4). The tidal interaction of a star with a close stellar companion or a giant planet is yet another cause of stellar deformation (5). Thus, measuring the asphericity of stars can place constraints on a wide range of phenomena beyond the standard model of stellar structure and evolution. Direct imaging of the deformed shapes of nearby stars requires a resolution better than a milli-arc second. The elongated projected shape of the rapidly rotating A star Altair has been observed with infrared interferometry to have a flattening  $\Delta R/R = 0.14 \pm 0.03$  (6, 7). Vega, another rapidly rotating A star, has an apparent deformation that is too small to be measured because it is seen almost pole-on (8). Here, we present with unprecedented precision the first measurement of stellar asphericity by means of asteroseismology (9), for the star KIC 11145123, which has an equatorial velocity two orders of magnitude smaller than either Altair or Vega. This work is motivated by helioseismology's ability to probe the Sun's asphericities and their temporal variations with the 11-year solar magnetic cycle (10, 11).

## RESULTS

The star KIC 11145123 belongs to the class of hybrid pulsators (12). It oscillates both in a high-frequency band (15 to 25  $\text{day}^{-1}$ ) and in a low-frequency band (below 5  $\text{day}^{-1}$ ). The observed modes of oscillation are acoustic (p), gravity (g), and mixed (p and g) modes. Modes with dominant p-mode character are seen in the high-frequency band.

<sup>1</sup>Max-Planck-Institut für Sonnensystemforschung, 37077 Göttingen, Germany. <sup>2</sup>Institut für Astrophysik, Georg-August-Universität Göttingen, 37077 Göttingen, Germany. <sup>3</sup>National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan. <sup>4</sup>Center for Space Science, NYUAD Institute, New York University Abu Dhabi, P.O. Box 129188, Abu Dhabi, UAE. <sup>5</sup>Department of Astronomy, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan. <sup>6</sup>Jeremiah Horrocks Institute, University of Central Lancashire, Preston PR1 2HE, U.K. <sup>7</sup>New York University, New York, NY 10012, USA.

\*Corresponding author. Email: gizon@mps.mpg.de

These modes oscillate throughout most of the star, with larger oscillation amplitudes near the surface. They are labeled with the radial order  $n$ , which counts the number of nodes in the radial direction with a positive sign for nodes in the p-mode cavity and a negative sign for nodes in the g-mode cavity. Most known hybrid pulsators, including KIC 11145123, belong to the  $\gamma$  Doradus- $\delta$  Scuti class (13). Oscillations in these stars are likely to be excited by the opacity (p and mixed modes) and the convective-blocking (g modes) mechanisms.

Oscillations of KIC 11145123 were observed in intensity over a period of  $T = 3.94$  years by Kepler (14). Because the oscillations are purely harmonic, random errors in the inferred mode frequencies scale as  $T^{-3/2}$  times the noise-to-signal ratio of the periodic oscillations (15), and thus, the mode frequencies can be determined with astounding precision. In the p-mode frequency band, Kurtz *et al.* (12) report frequency errors between  $5 \times 10^{-7} \text{ day}^{-1}$  and  $10^{-4} \text{ day}^{-1}$ . The stellar model that best explains the observed mode frequencies implies that KIC 11145123 is a terminal-age main sequence A star. It has a small convective core ( $r < 0.04 R$ ), in which the fraction of hydrogen content is less than 5%. Outside this convective core, energy is transported by radiation up to the surface layers of the star. In the top few thousand kilometers, there are very thin convective layers associated with the ionization of helium and hydrogen. See Table 1 for a summary of the basic stellar parameters.


In spherically symmetric stars, the eigenfunctions of stellar oscillations are proportional to spherical harmonics  $Y_l^m(\theta, \varphi)$  of degree  $l$  and azimuthal order  $m = -l, -l+1, \dots, l$ , where  $\theta$  is the colatitude and  $\varphi$  is the longitude. Internal rotation and departures from spherical symmetry lift the  $(2l+1)$ -fold degeneracy in  $m$  of the mode frequencies,  $\nu_{nlm}$ . The antisymmetric component of the frequency splittings in a multiplet,  $\delta\nu_{nlm} = (\nu_{nlm} - \nu_{nl,-m})/2$ , is a weighted average over the stellar volume of the stellar angular velocity (16). KIC 11145123 is one of a very few stars in which these rotational splittings have unambiguously been detected in both the p-mode and g-mode bands. The observed frequency splittings imply an internal rotation period of more than 105 days and a surface rotation period of less than 99 days, showing that the star rotates a little more quickly at the surface than in the core (12). Internal angular momentum transfer or external accretion mechanisms have spun up the atmosphere, a result of theoretical interest (17).

2016 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).

Downloaded from <http://advances.sciencemag.org/> on December 22, 2016

**Table 1. Parameters of the star KIC 11145123 and the best-fit seismic model.**

	Photometry (12, 18)	Seismology (12)
Spectral type	A	Late main sequence
Kepler visual magnitude	13	
Effective temperature, $T_{\text{eff}}$	$8050 \pm 200$ K	7032 K
Surface gravity, $\log g$ (cgs)	$4.0 \pm 0.2$	3.904
Surface rotation rate		$\Omega/2\pi = 0.01 \text{ day}^{-1}$
Mass		$M = 1.46 M_{\odot}$
Radius		$R = 2.24 R_{\odot}$
Initial abundances		$(X, Z) = (0.65, 0.010)$
Radius of convective core		$0.04 R$
Hydrogen core abundance		$X_c = 0.033$



**Fig. 1. Latitudinal dependence of mode kinetic energy density.** Dipole ( $l = 1$ , left panel) and quadrupole ( $l = 2$ , right panel) modes of oscillation. The arrow points along the stellar rotation axis. Scalar eigenfunctions of stellar oscillation are proportional to  $P_l^m(\cos\theta) e^{im\phi}$ , where  $P_l^m$  are associated Legendre functions. Polar plots of the kinetic energy density  $E_m(\theta) = c_{lm} |P_l^m(\cos\theta)|^2 \sin\theta$ , where  $\theta$  is the colatitude, for the modes with azimuthal orders  $m = 0$  (black),  $m = 1$  (red), and  $m = 2$  (blue) are shown. The constants of normalization,  $c_{lm}$  are such that  $\int_0^\pi d\theta E_m(\theta) = 1$  for each  $(l, m)$ . For dipole modes, we see that  $E_{10}$  is maximum at latitude  $\lambda = \pi - \theta = \pm 63^\circ$  and  $E_{11}$  is maximum at the equator. For quadrupole modes,  $E_{20}$  peaks at  $\lambda = 0^\circ$  and  $\pm 59^\circ$ ,  $E_{21}$  is maximum at  $\lambda = \pm 39^\circ$ , and  $E_{22}$  is maximum at the equator. For reference, the dashed curves show a highly distorted (oblate) stellar shape of the form  $r(\theta) = 1 - 0.15 P_2(\cos\theta)$ , where  $P_2(x) = (3x^2 - 1)/2$  is the second-order Legendre polynomial.

Stellar asphericity is measured through the symmetric component of the splittings (9)

$$s_{nlm} = (v_{nlm} + v_{nl,-m})/2 - v_{nl0}, \text{ for } 1 \leq m \leq l \quad (1)$$

This differential measurement exploits the different sensitivities in latitude of modes with different values of  $|m|$  at fixed  $l$  and  $n$ . As shown in Fig. 1, modes with larger values of  $|m|$  are confined to lower latitudes. For p modes, the mean frequencies  $\bar{v}_{nlm} = (v_{nlm} + v_{nl,-m})/2$  are

not sensitive to rotation at first order and inform us about the inverse acoustic stellar radius at specific latitudes, with increased sensitivity at lower latitudes for larger  $|m|$ . For a spherical star,  $\bar{v}_{nlm} = v_{nl0}$  and  $s_{nlm} = 0$  for all  $m$ . Latitudinal variations in stellar shape or wave speed will cause a nonzero  $s_{nlm}$ . The  $s_{nlm}$  values are positive for prolate spheroids and negative for oblate spheroids. Latitudinal variations in the wave speed may result from variations in a magnetic field or chemical composition.

In the p-mode frequency band of KIC 11145123, five multiplets have been identified (12) and assigned values of  $(n, l, m)$  by comparison with the best-fit seismic model. These are two dipole ( $l = 1$ ) and three quadrupole ( $l = 2$ ) multiplets, for which all  $2l + 1$  azimuthal modes are identified. The measured values of  $s_{nlm}$  are tabulated in Table 2. Among these, the quadrupole multiplet near  $23.5 \text{ day}^{-1}$  does not provide frequencies with sufficient precision to affect the results of this study. The  $s_{nlm}$  values are plotted in Fig. 2 for the four multiplets of interest. The average of all values,  $\langle s_{nlm} \rangle = (-1.4 \pm 0.5) \times 10^{-5} \text{ day}^{-1}$ , is negative (3 standard deviations away from zero); therefore, the star is oblate.

As mentioned in the Introduction, several physical mechanisms can make a star aspherical to stellar oscillations. One mechanism that must be present is rotational oblateness, which is relatively easy to compute when rotation is slow. The centrifugal force distorts the equilibrium structure of a rotating star. The corresponding perturbation to the mode frequencies scales as the ratio of centrifugal to gravitational forces

$$\epsilon = \Omega^2 R^3 / (GM) \quad (2)$$

where  $\Omega$  is the star's surface angular velocity;  $R$  and  $M$  are the radius and mass of the star, respectively; and  $G$  is the universal constant of gravity. Using  $\Omega/2\pi = 0.01 \text{ day}^{-1}$  for KIC 11145123, we have  $\epsilon = 1.34 \times 10^{-6} (R/R_{\odot})^3 (M/M_{\odot})^{-1}$ , where  $R_{\odot}$  and  $M_{\odot}$  are reference solar values (19). The mass and radius of the star are not known to the same level of confidence as the rotation. Our best-fit seismic model (12) has a metallicity of  $Z = 0.01$ , a mass of  $1.46 M_{\odot}$ , and a radius of  $2.24 R_{\odot}$ . For this stellar model, the ratio of the centrifugal to gravitational forces becomes

$$\epsilon = 1.0 \times 10^{-5} \quad (3)$$

This is a very small number, but it is not small compared to the relative errors of the most precise frequencies in the p-mode range from Table 2. Note that  $\epsilon$  is roughly half the solar value ( $\epsilon_{\odot} = 1.8 \times 10^{-5}$ ).

For slow rotators, rotational oblateness is described by a quadrupole distortion of the stellar structure. To leading order, the contribution of rotational oblateness to  $s_{nlm}$  can be written as (16, 20)

$$s_{nlm}^{\text{rot}} = -\epsilon m^3 v_{nl} \Delta_{nl} (2l - 1)^{-1} (2l + 3)^{-1} \quad (4)$$

where the dependence on  $m$  and  $l$  is due to the latitudinal sensitivity of the modes of oscillation (Fig. 1). The amplitude of the effect is proportional to the degenerate mode frequency  $v_{nl}$  of the nonrotating reference model and to the numbers  $\Delta_{nl}$  which are mode-weighted

**Table 2. Mode frequencies and frequency shifts.** Values of  $s_{nlm}$ , as defined by Eq. 1, are given for the dipole and quadrupole multiplets in the p-mode range of KIC 11145123. The frequency shifts expected from rotational oblateness,  $s_{nlm}^{\text{rot}}$ , are computed using Eq. 4. Mode identification is according to the best-fit stellar model, with  $R = 2.24 R_{\odot}$  and  $M = 1.46 M_{\odot}$  (12). Mode amplitudes are measured to a precision of 0.01 mmag.

$n$	$l$	$m$	$\nu_{nlm} \pm \sigma_{nlm}$ (day <sup>-1</sup> )	Amplitude (mmag)	$s_{nlm}$ (10 <sup>-5</sup> day <sup>-1</sup> )	$\Delta_{nl}$	$s_{nlm}^{\text{rot}}$ (10 <sup>-5</sup> day <sup>-1</sup> )	Index
-1	2	-2	16.7258824 ± 0.0000017	2.33				
		-1	16.7339455 ± 0.0000186	0.21				
		0	16.7420104 ± 0.0000075	0.53				
		1	16.7500755 ± 0.0000110	0.36	0.1 ± 1.3	0.40	-1.0	1
		2	16.7580083 ± 0.0000504	0.08	-6.4 ± 2.6	0.40	-3.9	2
2	1	-1	18.3558305 ± 0.0000029	1.36				
		0	18.3660001 ± 0.0000135	0.29				
		1	18.3761210 ± 0.0000034	1.13	-2.3 ± 1.4	0.68	-7.7	3
0	2	-2	18.9869603 ± 0.0000078	0.50				
		-1	18.9967001 ± 0.0000060	0.65				
		0	19.0064482 ± 0.0000056	0.69				
		1	19.0161736 ± 0.0000101	0.39	-1.1 ± 0.8	0.39	-1.1	4
		2	19.0259102 ± 0.0000086	0.45	-1.4 ± 0.8	0.39	-4.3	5
3	1	-1	21.9933315 ± 0.0000064	0.61				
		0	22.0018915 ± 0.0000286	0.14				
		1	22.0104220 ± 0.0000064	0.61	-1.5 ± 2.9	0.68	-9.2	6
2	2	-2	23.5455350 ± 0.0000303	0.13				
		-1	23.5553428 ± 0.0000582	0.07				
		0	23.5651835 ± 0.0000480	0.08				
		1	23.5749885 ± 0.0000607	0.06	-1.7 ± 6.4	0.20	-0.6	7
		2	23.5847898 ± 0.0000163	0.24	-2.2 ± 5.1	0.20	-2.2	8

radial averages of the stellar distortion (see Table 2 and Materials and Methods). The numerical values of  $s_{nlm}^{\text{rot}}$  are listed in Table 2 and are overplotted in Fig. 2 for the available modes. We find that the theoretical  $s_{nlm}^{\text{rot}}$  values are of the same sign and same order of magnitude as the measured  $s_{nlm}$ . As illustrated in Fig. 3, a good representation of the measurements is

$$s_{nlm} = (0.35 \pm 0.12) s_{nlm}^{\text{rot}} \quad (5)$$

Hence, the star is more round than rotational oblateness would imply. Equation 5 also implies that the modes of oscillation see a quadrupole distortion of the shape of the star. The amplitude of the distortion is smaller than would be expected from rotation alone; therefore, an additional physical ingredient is needed.


## DISCUSSION

The flattening of the stellar surface due to rotation alone would be  $(\Delta R/R)^{\text{rot}} = \epsilon/2 = 5 \times 10^{-6}$ , where  $\Delta R$  is the difference between the equatorial and polar radii. Here, the effective flattening of the stellar surface implied by the seismic measurements (Eq. 5) is only


$$\Delta R/R = (1.8 \pm 0.6) \times 10^{-6} \quad (6)$$

To our knowledge, KIC 411145123 is the most spherical natural object ever measured, more spherical than the quiet Sun (21).

Using  $R = 2.24 R_{\odot} = 1.56 \times 10^6$  km, we have  $\Delta R = 2.7 \pm 0.9$  km. This is an astonishing illustration of the precision of the asteroseismic diagnostic and a direct consequence of the very long lifetime of the oscillations under study. However, there is a limitation in accuracy mainly due to the uncertainty in the radius of the stellar model. We



**Fig. 2. Symmetric component of observed frequency splittings  $s_{nlm}$ .** Observations are plotted as red circles with error bars. Each value is associated with the mode index given in the last column of Table 2. The data points are labeled by the values of  $(l, m)$ . The theoretical values for rotational oblateness alone,  $s_{nlm}^{rot}$ , are plotted as open squares. Note that the last two values of  $s_{nlm}$  from Table 2 are not plotted here because they are associated with errors that are too large to provide additional constraints.



**Fig. 3. Ratios of observed  $s_{nlm}$  to theoretical prediction for rotational oblateness  $s_{nlm}^{rot}$ .** The horizontal solid line and the gray area indicate the average and the  $1-\sigma$  bounds,  $\langle s_{nlm}/s_{nlm}^{rot} \rangle = 0.35 \pm 0.12$ . Each value is associated with the mode index given in the last column of Table 2. The distributions of the data points and their errors are consistent with a single value for the ratio of  $s_{nlm}/s_{nlm}^{rot}$ .

may incorporate the uncertainty in the stellar radius in the error for  $\Delta R$ ; the conservative assumption of a systematic error of one solar radius implies a combined error of 1.5 km on  $\Delta R$ . We emphasize that the uncertainty in the stellar radius is a systematic error that does not change the  $3-\sigma$  significance level of the result; it only changes the absolute value of  $\Delta R$ .

Guided by the well-established results of helioseismology (11, 22), we suggest that a weak surface magnetic field (much weaker than the

Sun’s surface magnetic field at solar maximum; see Materials and Methods) is a possible explanation for the reduced oblateness of KIC 11145123: Waves propagate faster in magnetized regions, so surface magnetic fields at low latitudes will make a star appear less oblate to acoustic waves. We note that observations of photometric variability have led to the speculation that a large fraction of A stars without star-spots (23). However, the origin of magnetic fields in stars without deep convective envelopes is a matter of debate (24). Dynamo action may take place in the core of the star or in the very thin convective layers near the surface, or the magnetic field may have a fossil origin.

Other than a magnetic field, there are few alternative explanations for the reduced oblateness. At this level of precision, the physics of stellar oscillations may need to be studied in more detail. In particular, it is not quite excluded that nonlinear (amplitude) effects could play a role; this should be investigated further. On the other hand, nonadiabatic effects are spherically symmetric and will not affect  $s_{nlm}$  to leading order.

Nearly all slowly rotating A stars have overabundances of certain metal elements (25). The fact that KIC 11145123 is not a chemically peculiar star (Am or Ap) is surprising, hence the speculation by Kurtz *et al.* (12) about possible blue straggler mass transfer. An enhancement or deficiency of metals in the atmosphere would only affect seismic asphericity if the abundances were nonuniformly distributed in latitude. This could happen in magnetic Ap stars, but a Subaru high-resolution spectrum does not show Ap abundances and shows a metal deficiency of 0.7 dex. Although we cannot rule out a latitudinal gradient in chemical composition, this explanation is more involved than a weak magnetic field.

Because stars more massive than the Sun are more likely to harbor giant planets (26), one may also ask whether KIC 11145123 could be deformed by tidal interaction. In the linear regime, only the equilibrium tidal deformation should be considered. However, it is smaller than the rotational deformation by a factor proportional to the ratio of the mass of the planet to the mass of the star (20, 27); thus, it is negligible for Jupiter mass planets. Furthermore, a planetary companion (or a stellar companion) in the equatorial plane of the star would make the star look more oblate to the acoustic modes, not less oblate as required by the observations.

This work is a first step in the study of stellar shapes through asteroseismology. The method demonstrated here will be applied to other stars, including more rapidly rotating stars and stars with stronger magnetic fields, where deformations will be greater. Because of the unprecedented high precision and long time span of the Kepler observations, an important field of theoretical astrophysics is now also observational.

**MATERIALS AND METHODS**

**Mode frequency measurements**

The frequencies of the modes of oscillation were measured using Kepler light curves for quarters Q0 to Q16, spanning a total of 51 months of data. The mode frequencies were determined by nonlinear least squares in the time domain assuming Gaussian uncorrelated errors; see the study of Kurtz *et al.* (12) for a full description of the data reduction. We tested a new frequency solution on Kepler quarters Q0 to Q17 and found that there were no significant changes compared to the published Q0 to Q16 analysis. For our work to be easily tested and reproduced by others, we used published data (12).

Downloaded from <http://advances.sciencemag.org/> on December 22, 2016



The frequency errors were determined using an estimate of the variance around each mode frequency. Because many nearby frequency peaks may contribute to this variance, the frequency errors are conservative. Had all significant peaks been removed from the amplitude spectrum, the error estimates would have been smaller.

The Kepler data were averaged over consecutive 29.4-min time intervals, that is, a significant fraction of the shortest p-mode periods (~1 hour). This effect reduces the observed amplitudes of the modes but does not affect the mode frequencies. The only effect is a reduced signal-to-noise ratio (compared to shorter integration times); this ratio was taken into account in the estimation of the errors on mode frequencies.

### Effect of centrifugal distortion on mode frequencies

The effect of the centrifugal force on mode frequencies can be evaluated using the second-order perturbation theory, either in the spherical geometry of the reference model (28) or in the distorted geometry of the oblate spheroid (20, 29). It consists of several terms, which account for geometrical distortion, change in wave speed, and first-order perturbation to the mode eigenfunctions. In the p-mode frequency range, for which  $[\Omega/(2\pi\nu_{nl})]^2 \ll \epsilon$ , the effect of rotation on mode frequencies is well approximated by (16)

$$\nu_{nlm} \approx \nu_{nl} + m\Omega/(2\pi) + \epsilon\nu_{nl}\Delta_{lm}Q_{2lm} \quad (7)$$

where  $\nu_{nl}$  is the mode frequency in the nonrotating reference stellar model

$$Q_{2lm} = \int_0^\pi d\theta P_2(\cos\theta)E_{lm}(\theta) = [l(l+1) - 3m^2](2l-1)^{-1}(2l+3)^{-1} \quad (8)$$

is the latitudinal average of quadrupole distortion weighted by latitudinal mode energy density  $E_{lm}$  (see Fig. 1), and the dimensionless number

$$\Delta_{nl} = 4/3 \int_0^R dr (r/R)^3 \xi_{nl}^2(r) \rho r^2 \quad (9)$$

is the radial average of the centrifugal distortion weighted by mode energy density (fig. S1), which depends on the (normalized) radial mode displacement  $\xi_{nl}$ . For modes with pure p-mode character,  $\Delta_{nl} \approx 0.7$ . For the modes under consideration here,  $\Delta_{nl}$  ranges from 0.2 to 0.7 (Table 2), where the smaller values are for the mixed modes (fig. S2). The above expression (Eq. 9) assumes a rigid body rotation and neglects the perturbation to the gravitational potential related to the star's quadrupole moment; both approximations are at the level of a few percent (30) and are thus acceptable for the purpose of estimating the contribution of rotational oblateness to  $s_{nlm}$ . Combining the definition of  $s_{nlm}$  (Eq. 1) and Eq. 7, we obtain Eq. 4.

### Alternative stellar model

We note that the effective temperature of the best-fit seismic model is not consistent with the photometric value (Table 1). This prompted Kurtz *et al.* (12) to consider an alternative stellar model with  $M = 2.05 M_\odot$  and  $R = 2.82 R_\odot$ , whose effective temperature is within error bars. However, this model is a worse fit to the p-mode frequencies, making mode identification more difficult. In particular, in Table 2, only the  $l = 2$  mode at  $16.7 \text{ day}^{-1}$  and the  $l = 1$  mode at  $18.4 \text{ day}^{-1}$

can be identified. For the alternative model, we have  $\epsilon = 1.5 \times 10^{-5}$ , which is 50% larger than for the best-fit seismic model. Should this alternative stellar model be preferred, the estimates of  $s_{nlm}^{\text{rot}}$  should be scaled appropriately so that  $s_{nlm} = (0.23 \pm 0.08) s_{nlm}^{\text{rot}}$  and  $\Delta R = 3.4 \pm 1.1 \text{ km}$ . We note that this stellar model and the best-fit seismic model referred to in this study were obtained from a stellar evolutionary code that does not include rotation or magnetic fields.

### Helioseismology and upper limit on magnetic field

In helioseismology, azimuthal mode frequencies in a multiplet are traditionally expanded as  $a$  coefficients on a basis of Clebsch-Gordan polynomials (31). The odd  $a$  coefficients,  $a_{2k+1}$ , are measures of differential rotation, whereas the even  $a$  coefficients,  $a_{2k}$ , are measures of asphericity ( $k = 0, 1, 2, \dots$ ). The  $s_{nlm}$  are related to the even  $a$  coefficients. In particular, for dipole modes,  $s_{n11} = 3a_2$  informs us about the  $P_2$  component of distortion. The effect of solar rotational oblateness is too small ( $a_2 \sim -10 \text{ nHz}$  for  $l \leq 2$  modes) to be measured on individual multiplets (11). However, solar asphericity measurements are possible by averaging over sets of intermediate degree modes ( $l < 150$ ). When the Sun's magnetic activity is very low,  $a_2$  coefficients are negative but of smaller magnitude than implied by rotational oblateness (11). During maxima of solar activity, the solar  $a_2$  coefficients become positive ( $a_2 \sim 100 \text{ nHz}$  for  $l < 5$  modes) (22), and the Sun appears prolate to the acoustic modes: They sense magnetic activity at mid to low latitudes ( $< 40^\circ$ ). At the solar surface, the quadrupole components of the solar magnetic field vary by less than 10 G with the sunspot cycle (32). Baldner *et al.* (22) used measurements of solar  $a_2$  coefficients to infer toroidal and poloidal magnetic field components below the solar surface, at the level of a few hundred gauss.

In light of the solar observations, a possible explanation for KIC 11145123's reduced oblateness is a magnetic perturbation (9). Let us consider the dipole mode at  $18.3 \text{ day}^{-1}$ , for which  $\Delta a_2 = a_2 - a_2^{\text{rot}} = 1.8 \times 10^{-5} \text{ day}^{-1} = 0.2 \text{ nHz}$ . By comparison with the solar observations and given that  $\Delta a_2$  is expected to scale like the square of the magnetic field, we infer that a much smaller level of magnetism than the Sun's would be needed to explain the observations. However, it is difficult to be more specific because  $\Delta a_2$  sensitively depends on the geometrical configuration of the magnetic field (22).

### SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/2/11/e1601777/DC1>

fig. S1. Mode kinetic energy density.

fig. S2. Radial dependence of centrifugal distortion.

### REFERENCES AND NOTES

1. A. C. Clairaut, *Théorie de la figure de la Terre, tirée des principes de l'hydrostatique* (Durand, 1743).
2. S. Chandrasekhar, *Ellipsoidal Figures of Equilibrium* (Yale Univ. Press, 1969).
3. S. Chandrasekhar, E. Fermi, Problems of gravitational stability in the presence of a magnetic field. *Astrophys. J.* **118**, 116 (1953).
4. D. G. Wentzel, On the shape of magnetic stars. *Astrophys. J.* **133**, 170 (1961).
5. G. I. Ogilvie, D. N. C. Lin, Tidal dissipation in rotating solar-type stars. *Astrophys. J.* **661**, 1180 (2007).
6. G. T. van Belle, D. R. Ciardi, R. R. Thompson, R. L. Akeson, E. A. Lada, Altair's oblateness and rotation velocity from long-baseline interferometry. *Astrophys. J.* **559**, 1155 (2001).
7. J. D. Monnier, M. Zhao, E. Pedretti, N. Thureau, M. Ireland, P. Muirhead, J.-P. Berger, R. Millan-Gabet, G. Van Belle, T. ten Brummelaar, H. McAlister, S. Ridgway, N. Turner,

- L. Sturmann, J. Sturmann, D. Berger, Imaging the surface of Altair. *Science* **317**, 342–345 (2007).
8. D. M. Peterson, C. A. Hummel, T. A. Pauls, J. T. Armstrong, J. A. Benson, G. C. Gilbreath, R. B. Hindsley, D. J. Hutter, K. J. Johnston, D. Mozurkewich, H. R. Schmitt, Vega is a rapidly rotating star. *Nature* **440**, 896–899 (2006).
  9. L. Gizon, Prospects for detecting stellar activity through asteroseismology. *Astron. Nachr.* **323**, 251–253 (2002).
  10. K. G. Libbrecht, M. F. Woodard, Solar-cycle effects on solar oscillation frequencies. *Nature* **345**, 779–782 (1990).
  11. H. M. Antia, S. M. Chitre, M. J. Thompson, The Sun's acoustic asphericity and magnetic fields in the solar convection zone. *Astron. Astrophys.* **360**, 335–344 (2000).
  12. D. W. Kurtz, H. Saio, M. Takata, H. Shibahashi, S. J. Murphy, T. Sekii, Asteroseismic measurement of surface-to-core rotation in a main-sequence A star, KIC 11145123. *Mon. Not. R. Astron. Soc.* **444**, 102–116 (2014).
  13. A. Grigahcène, V. Antoci, L. Balona, G. Catanzaro, J. Daszyńska-Daszkiewicz, J. A. Guzik, G. Handler, G. Houdek, D. W. Kurtz, M. Marconi, M. J. P. F. G. Monteiro, A. Moya, V. Ripepi, J.-C. Suárez, K. Uytterhoeven, W. J. Borucki, T. M. Brown, J. Christensen-Dalsgaard, R. L. Gilliland, J. M. Jenkins, H. Kjeldsen, D. Koch, S. Bernabei, P. Bradley, M. Breger, M. Di Criscienzo, M.-A. Dupret, R. A. García, A. García Hernández, J. Jackiewicz, A. Kaiser, H. Lehmann, S. Martín-Ruiz, P. Mathias, J. Molenda-Zakowicz, J. M. Nemeč, J. Nuspl, M. Paparo, M. Roth, R. Szabó, M. D. Suran, R. Ventura, Hybrid  $\gamma$  Doradus- $\delta$  Scuti pulsators: New insights into the physics of the oscillations from *Kepler* observations. *Astrophys. J.* **713**, L192 (2010).
  14. W. J. Borucki, D. Koch, G. Basri, N. Batalha, T. Brown, D. Caldwell, J. Caldwell, J. Christensen-Dalsgaard, W. D. Cochran, E. DeVore, E. W. Dunham, A. K. Dupree, T. N. Gautier III, J. C. Geary, R. Gilliland, A. Gould, S. B. Howell, J. M. Jenkins, Y. Kondo, D. W. Latham, G. W. Marcy, S. Meibom, H. Kjeldsen, J. J. Lissauer, D. G. Monet, D. Morrison, D. Sasselov, J. Tarter, A. Boss, D. Brownlee, T. Owen, D. Buzasi, D. Charbonneau, L. Doyle, J. Fortney, E. B. Ford, M. J. Holman, S. Seager, J. H. Steffen, W. F. Welsh, J. Rowe, H. Anderson, L. Buchhave, D. Ciardi, L. Walkowicz, W. Sherry, E. Horch, H. Isaacson, M. E. Everett, D. Fischer, G. Torres, J. A. Johnson, M. Endl, P. MacQueen, S. T. Bryson, J. Dotson, M. Haas, J. Kolodziejczak, J. Van Cleve, H. Chandrasekaran, J. D. Twicken, E. V. Quintana, B. D. Clarke, C. Allen, J. Li, H. Wu, P. Tenenbaum, E. Verner, F. Bruhweiler, J. Barnes, A. Prsa, *Kepler* planet-detection mission: Introduction and first results. *Science* **327**, 977–980 (2010).
  15. T. Stahn, L. Gizon, Fourier analysis of gapped time series: Improved estimates of solar and stellar oscillation parameters. *Sol. Phys.* **251**, 31–52 (2008).
  16. C. Aerts, J. Christensen-Dalsgaard, D. W. Kurtz, *Asteroseismology* (Springer, 2010).
  17. T. M. Rogers, On the differential rotation of massive main-sequence stars. *Astrophys. J.* **815**, L30 (2015).
  18. D. Huber, V. S. Aguirre, J. M. Matthews, M. H. Pinsonneault, E. Gaidos, R. A. García, S. Hekker, S. Mathur, B. Mosser, G. Torres, F. A. Bastien, S. Basu, T. R. Bedding, W. J. Chaplin, B.-O. Demory, S. W. Fleming, Z. Guo, A. W. Mann, J. F. Rowe, A. M. Serenelli, M. A. Smith, D. Stello, Revised stellar properties of *Kepler* targets for the quarter 1–16 transit detection run. *Astrophys. J. Suppl. S.* **211**, 2 (2014).
  19. E. E. Mamajek, A. Prsa, G. Torres, P. Harmanec, M. Asplund, P. D. Bennett, N. Capitaine, J. Christensen-Dalsgaard, E. Depagne, W. M. Folkner, M. Haberreiter, S. Hekker, J. L. Hilton, V. Kostov, D. W. Kurtz, J. Laskar, B. D. Mason, E. F. Milone, M. M. Montgomery, M. T. Richards, J. Schou, S. G. Stewart, IAU 2015 resolution B3 on recommended nominal conversion constants for selected solar and planetary properties. arXiv:1510.07674 (2015).
  20. H. Saio, Rotational and tidal perturbations of nonradial oscillations in a polytropic star. *Astrophys. J.* **244**, 299–315 (1981).
  21. J. R. Kuhn, R. Bush, M. Emilio, I. F. Scholl, The precise solar shape and its variability. *Science* **337**, 1638–1640 (2012).
  22. C. S. Baldner, H. M. Antia, S. Basu, T. P. Larson, Solar magnetic field signatures in helioseismic splitting coefficients. *Astrophys. J.* **705**, 1704 (2009).
  23. L. A. Balona, Rotational light variations in *Kepler* observations of A-type stars. *Mon. Not. R. Astron. Soc.* **415**, 1691–1702 (2011).
  24. J. Braithwaite, H. C. Spruit, Magnetic fields in non-convective regions of stars. arXiv:1510.03198 (2015).
  25. H. A. Abt, N. I. Morrell, The relation between rotational velocities and spectral peculiarities among A-type stars. *Astrophys. J. Suppl. S.* **99**, 135 (1995).
  26. J. A. Johnson, C. Clanton, A. W. Howard, B. P. Bowler, G. W. Henry, G. W. Marcy, J. R. Crepp, M. Endl, W. D. Cochran, P. J. MacQueen, J. T. Wright, H. Isaacson, Retired A stars and their companions: Eighteen new Jovian planets. *Astrophys. J. Suppl. S.* **197**, 26 (2011).
  27. S. Chandrasekhar, The equilibrium of distorted polytropes. (III) The double star problem. *Mon. Not. R. Astron. Soc.* **93**, 462–471 (1933).
  28. W. A. Dziembowski, P. R. Goode, Effects of differential rotation on stellar oscillations—A second-order theory. *Astrophys. J.* **394**, 670–687 (1992).
  29. D. O. Gough, M. J. Thompson, The effect of rotation and a buried magnetic field on stellar oscillations. *Mon. Not. R. Astron. Soc.* **242**, 25–55 (1990).
  30. L. Paterno, S. Sofia, M. P. di Mauro, The rotation of the Sun's core. *Astron. Astrophys.* **314**, 940–946 (1996).
  31. J. Schou, J. Christensen-Dalsgaard, M. J. Thompson, On comparing helioseismic two-dimensional inversion methods. *Astrophys. J.* **433**, 389–416 (1994).
  32. M. L. DeRosa, A. S. Brun, J. T. Hoeksema, Solar magnetic field reversals and the role of dynamo families. *Astrophys. J.* **757**, 96 (2012).

**Acknowledgments:** We acknowledge useful discussions with H. Saio and T. Hoeksema.

**Funding:** L.G. acknowledges support through a visiting professorship at the National Astronomical Observatory of Japan (Tokyo, Japan). L.G., M.B., O.B., and K.R.S. acknowledge research funding from the New York University Abu Dhabi Institute under grant no. G1502. M.T. acknowledges financial support from the Japan Society for the Promotion of Science KAKENHI under grant no. 26400219. **Author contributions:** L.G., T.S., and M.T. designed the research. D.W.K. performed multiple tests to validate mode frequencies and associated errors. L.G. drafted the paper, and all authors contributed to the final manuscript. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. The mode eigenfunctions may be requested from the authors.

Submitted 1 August 2016  
Accepted 12 October 2016  
Published 16 November 2016  
10.1126/sciadv.1601777

**Citation:** L. Gizon, T. Sekii, M. Takata, D. W. Kurtz, H. Shibahashi, M. Bazot, O. Benomar, A. C. Birch, K. R. Sreenivasan, Shape of a slowly rotating star measured by asteroseismology. *Sci. Adv.* **2**, e1601777 (2016).

This article is published under a Creative Commons license. The specific license under which this article is published is noted on the first page.

For articles published under [CC BY](#) licenses, you may freely distribute, adapt, or reuse the article, including for commercial purposes, provided you give proper attribution.

For articles published under [CC BY-NC](#) licenses, you may distribute, adapt, or reuse the article for non-commercial purposes. Commercial use requires prior permission from the American Association for the Advancement of Science (AAAS). You may request permission by clicking [here](#).

**The following resources related to this article are available online at <http://advances.sciencemag.org>. (This information is current as of December 22, 2016):**

**Updated information and services**, including high-resolution figures, can be found in the online version of this article at:

<http://advances.sciencemag.org/content/2/11/e1601777.full>

**Supporting Online Material** can be found at:

<http://advances.sciencemag.org/content/suppl/2016/11/14/2.11.e1601777.DC1>

This article **cites 27 articles**, 7 of which you can access for free at:

<http://advances.sciencemag.org/content/2/11/e1601777#BIBL>

*Science Advances* (ISSN 2375-2548) publishes new articles weekly. The journal is published by the American Association for the Advancement of Science (AAAS), 1200 New York Avenue NW, Washington, DC 20005. Copyright is held by the Authors unless stated otherwise. AAAS is the exclusive licensee. The title *Science Advances* is a registered trademark of AAAS