

Central Lancashire Online Knowledge (CLoK)

Title	Effects of prophylactic knee bracing on patellar tendon loading parameters during functional sports tasks in recreational athletes
Type	Article
URL	https://clock.uclan.ac.uk/21080/
DOI	##doi##
Date	2017
Citation	Sinclair, Jonathan Kenneth orcid iconORCID: 0000-0002-2231-3732, Richards, James orcid iconORCID: 0000-0002-4004-3115 and Taylor, Paul John orcid iconORCID: 0000-0002-9999-8397 (2017) Effects of prophylactic knee bracing on patellar tendon loading parameters during functional sports tasks in recreational athletes. Sport Sciences for Health, 14 . pp. 151-160. ISSN 1824-7490
Creators	Sinclair, Jonathan Kenneth, Richards, James and Taylor, Paul John

It is advisable to refer to the publisher's version if you intend to cite from the work. ##doi##

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

1 **Effects of prophylactic knee bracing on patellar tendon loading parameters during**
2 **functional sports tasks in recreational athletes.**

3 **Keywords:** Biomechanics, knee brace, patellar tendon, tendinopathy

4 **Word count:** 3200

5 **Conflict statement:** No conflict of interest to declare.

6
7 **Abstract**

8 **PURPOSE:** This study investigated the effects of prophylactic knee bracing on patellar
9 tendon loading parameters.

10 **METHODS:** Twenty recreational athletes (10 male & 10 female), from a different athletic
11 disciplines performed run, cut and single leg hop movements under two conditions
12 (prophylactic knee brace/ no-brace). Lower extremity kinetics and kinematics were examined
13 using a piezoelectric force plate, and three-dimensional motion capture system. Patellar
14 tendon loading was explored using a mathematical modelling approach, which accounted for
15 co-contraction of the knee flexors. Tendon loading parameters were examined using 2
16 (*brace*)*3 (*movement*)*2 (*sex*) mixed ANOVA's.

17 **RESULTS:** Tendon instantaneous load rate was significantly reduced in female athletes, in
18 the run (brace = 289.14BW/s no-brace = 370.06BW/s) and cut (brace = 353.17BW/s/ no-
19 brace = 422.01BW/s) conditions whilst wearing the brace.

20 **CONCLUSIONS:** Female athletes may be able to attenuate their risk from patellar
21 tendinopathy during athletic movements, through utilization of knee bracing, although further
22 prospective research into the prophylactic effects of knee bracing is required before this can
23 be clinically substantiated.

24
25 **Introduction**

26 Chronic patellar tendinopathy is an extremely common musculoskeletal condition in both
27 recreational and elite athletes, and has previously been reported to account for as many as
28 25% of all soft tissue injuries (1). Patellar tendinopathy is characterized by pain localized at
29 the lower pole of the patella, and pain symptoms that are augmented by activities which place
30 high demands on the knee extensors, notably in physical disciplines which repeatedly store
31 and release elastic energy in the tendon itself (2). Patellar tendinopathy is more common in
32 skeletally mature individuals, and there remains disagreement as to whether this condition is
33 most common in male or female athletes (3). Chronic patellar tendinopathy is established
34 after 1-3 months, as degenerative alterations occur in the tendon itself (4). Degenerative
35 alterations at the tendon are mediated primarily by the absence of inflammatory cells within
36 the tendon itself, which reduces healing of the tendon and ultimately leads to decreased
37 tensile strength and disorganization of the collagen fibers (5). Patellar tendinopathy can be
38 debilitating; Cook et al., (6) showed that 1/3 of athletes with patellar tendinopathy are unable
39 to return to physical activity within 6 months, and it has also been evidenced that 53% of
40 athletes who present with this condition were forced to permanently cease physical activities.

41

42 Knee braces are utilized extensively in both recreationally active and competitive athletes, in
43 order to attenuate their risk from knee pathology (7). Knee braces are external devices which
44 are designed to improve the alignment of the knee joint (8). Prophylactic knee braces aim to
45 protect athletes from sustaining injury, whilst being minimally restrictive, allowing athletes to
46 utilize full knee range of motion during their physical activities (9). Recently, the effects of
47 prophylactic knee braces on the biomechanics of the knee joint during dynamic sports tasks
48 have received significant attention in clinical literature. Sinclair et al., (7), examined the
49 effects of knee bracing on knee joint kinetics and kinematics in netball specific movements.
50 They showed that the brace did not alter knee kinetics but did reduce range of motion in the

51 transverse plane. Ewing et al., (10), examined muscle kinetics with and without the presence
52 of a prophylactic knee brace during double limb drop landings. Hamstring and vasti muscles
53 produced significantly greater flexion and extension torques, and greater peak muscle forces
54 in the brace condition. Lee et al., (11), analyzed the effects of a prophylactic bilateral hinge
55 brace, fitted with torque transducers during four functional sports tasks; drop vertical jump,
56 pivot, stop vertical jump and cut. Their results showed that the knee brace hinges absorbed up
57 to 18% of the force and 2.7% of the torque at the knee, during the different athletic motions.
58 Which they concluded, was minimal evidence that the brace was able to reduce the
59 mechanical load at the knee. Although knee braces have been studied in terms of both their
60 therapeutic and prophylactic effects, there is currently no literature which has considered
61 their role in the prevention of patellar tendinopathy.

62

63 Therefore, the aim of the current investigation was to investigate the effects of a prophylactic
64 knee brace on patellar tendon loading parameters linked to the aetiology of patellar
65 tendinopathy, in male and female recreational athletes. Research of this nature may provide
66 important clinical information, regarding the potential role of prophylactic knee bracing for
67 the prevention of patellar tendinopathy.

68

69 **Methods**

70 *Participants*

71 Twenty participants (10 male; age = 26.70 ± 4.24 , mass = 73.90 ± 5.3 , stature = $176.50 \pm$
72 4.25 & BMI = 23.73 ± 1.80 & and 10 female age = 27.60 ± 4.72 , mass = 60.40 ± 7.86 , stature
73 = 166.50 ± 5.06 & BMI = 21.86 ± 2.21), volunteered to take part in the current investigation.
74 Participants were all recreational level athletes who came from squash, netball, basketball and
75 association football athletic backgrounds, with a minimum of 2 years of experience in their

76 chosen discipline. In addition, all were free from lower extremity pathology at the time of
77 data collection, and had not previously experienced an injury to the patellar tendon. Written
78 informed consent was provide,d in accordance with the declaration of Helsinki and the rights
79 of all participants were protected. The procedure was approved by the Universities Science,
80 Technology, Engineering, Medicine and Health ethics committee, with the reference STEMH
81 295.

82

83 *Knee Brace*

84 A single knee brace was utilized in this investigation, (Trizone, DJO USA), which was worn
85 on the dominant limb in all participants. The brace examined in the current investigation
86 represents a compression sleeve reinforced with silicone designed to support the knee joint
87 and improve proprioception.

88

89 *Procedure*

90 Participants were required to complete five repetitions of three sports specific movements';
91 jog, cut and single leg hop, with and without presence of the brace. The order that
92 participants performed in the movement/ brace conditions was counterbalanced. To quantify
93 lower extremity segments, the calibrated anatomical systems technique was utilized (12).
94 Retroreflective markers (19 mm), were positioned unilaterally allowing the; foot, shank and
95 thigh to be defined. The foot was defined via the 1st and 5th metatarsal heads, medial and
96 lateral malleoli and tracked using the calcaneus, 1st metatarsal and 5th metatarsal heads. The
97 shank was defined via the medial and lateral malleoli and medial and lateral femoral
98 epicondyles and tracked using a cluster positioned onto the shank. The thigh was defined via
99 the medial and lateral femoral epicondyles and the hip joint centre and tracked using a cluster
100 positioned onto the thigh. To define the pelvis additional markers were positioned onto the

101 anterior (ASIS) and posterior (PSIS) superior iliac spines and this segment was tracked using
102 the same markers. The hip joint centre was determined using a regression equation, which
103 uses the positions of the ASIS markers (13). The centers of the ankle and knee joints were
104 delineated as the mid-point between the malleoli and femoral epicondyle markers (14, 15).
105 Each tracking cluster comprised four retroreflective markers, mounted onto a rigid piece of
106 lightweight carbon-fibre. Static calibration trials were obtained allowing for the anatomical
107 markers to be referenced in relation to the tracking markers/ clusters. The Z (transverse) axis
108 was oriented vertically from the distal segment end to the proximal segment end. The Y
109 (coronal) axis was oriented in the segment from posterior to anterior. Finally, the X (sagittal)
110 axis orientation was determined using the right hand rule and was oriented from medial to
111 lateral.

112

113 Data were collected during run, cut and jump movements using the protocol below:

114

115 *Run*

116 Participants ran at $4.0 \text{ m}\cdot\text{s}^{-1} \pm 5\%$, and struck the force platform with their right (dominant)
117 limb. The average velocity of running was monitored using infra-red timing gates
118 (SmartSpeed Ltd UK). The stance phase of running, was defined as the duration over $> 20 \text{ N}$
119 of vertical force was applied to the force platform (16).

120

121 *Cut*

122 Participants completed 45° sideways cut movements, using an approach velocity of $4.0 \text{ m}\cdot\text{s}^{-1}$
123 $\pm 5\%$ striking the force platform with their right (dominant) limb. In accordance with McLean
124 et al., (17), cut angles were measured from the centre of the force plate and the corresponding
125 line of movement was delineated using masking tape, so that it was clearly evident to

126 participants. The stance phase of the cut-movement was similarly defined as the duration over
127 > 20 N of vertical force was applied to the force platform (16).

128

129 *Hop*

130 Participants began standing by on their dominant limb; they were then requested to hop
131 forward maximally, landing on the force platform with same leg without losing balance. The
132 arms were held across the chest to remove arm-swing contribution. The hop movement was
133 defined as the duration from foot contact (defined as > 20 N of vertical force applied to the
134 force platform), to maximum knee flexion. The hop distance was recorded and maintained
135 throughout data collection.

136

137 *Processing*

138 Dynamic trials were processed using Qualisys Track Manager, and then exported as C3D
139 files. Ground reaction force and marker data were filtered at 50 Hz and 15 Hz respectively
140 using a low-pass Butterworth 4th order filter, and processed using Visual 3-D (C-Motion,
141 Germantown, MD, USA). Internal moments were computed using Newton-Euler inverse-
142 dynamics, allowing net knee joint moments to be calculated. Angular kinematics of the knee
143 joint were calculated using an XYZ (sagittal, coronal and transverse) sequence of rotations,
144 allowing sagittal angles at footstrike and peak flexion angles to be extracted.

145

146 A commonly utilized mathematical model for the quantification of patellar tendon loading is
147 that developed by Janssen et al., (18). Whereby the Patellar tendon load is determined by
148 dividing the knee extensor moment by the estimated patellar tendon moment arm. This
149 algorithm has been successfully utilized previously, to resolve differences in patellar tendon

150 kinetics during different movements (18), different footwear conditions (19), and also
151 between sexes (20).

152

153 However, a limitation of the aforementioned model is that the knee extensor moment does
154 not account for co-contraction of the knee flexor musculature. In order to account for this, we
155 also calculated hamstring and gastrocnemius force in accordance with the procedures
156 described by DeVita and Hortobagyi (21). To summarize, the hamstring force was calculated
157 using the hip extensor moment, hamstrings and gluteus maximus cross-sectional areas (22),
158 and by fitting a 2nd order polynomial curve to the data of Nemeth & Ohlsen, (23) who
159 provided muscle moment arms at the hip as a function of hip flexion angle. The
160 gastrocnemius force, was calculated firstly by quantifying the ankle plantarflexor force,
161 which was resolved by dividing the plantarflexion moment by the Achilles tendon moment
162 arm. The Achilles tendon moment arm was calculated by fitting a 2nd order polynomial curve
163 to the ankle plantarflexion angle in accordance with Self and Paine (24). The quantity of
164 plantarflexion force accredited to the gastrocnemius muscles, was calculated via the cross-
165 sectional area of this muscle relative to the triceps surae (22).

166

167 The hamstring and gastrocnemius forces were multiplied by their estimated muscle moment
168 arms to the knee joint in relation to the knee flexion angle (25), and then added together to
169 estimate the knee flexor moment. The derived knee flexor moment was added to the net knee
170 extensor moment quantified using inverse dynamics, and then divided by the moment arm of
171 the patellar tendon, generating the patellar tendon force. The tendon moment arm was
172 quantified as a function of the sagittal plane knee angle, by fitting a 2nd order polynomial
173 curve to the data provided by Herzog & Read, (26), showing patellar tendon moment arms at
174 different knee flexion angles.

175

176 All patellar tendon load parameters were normalized by dividing the net values by
177 bodyweight (BW). Patellar tendon instantaneous load rate (BW/s), was quantified as the peak
178 increase in patellar tendon force between adjacent data points. In addition, we also calculated
179 the total patellar tendon force impulse (BW·s) during each movement using a trapezoidal
180 function.

181

182 *Statistical analyses*

183 Descriptive statistics of means, standard deviations and 95% confidence intervals (95% CI)
184 were obtained for each outcome measure. Shapiro-Wilk tests were used to screen the data for
185 normality. Differences in patellar tendon loading parameters between conditions, were
186 examined using 2 (*brace*) * 3 (*movement*) * 2 (*sex*) mixed ANOVA's. Statistical significance
187 was accepted at the P<0.05 level. Effect sizes for all significant findings were calculated
188 using partial Eta² (η^2). Post-hoc pairwise comparisons were conducted on all significant
189 main effects. Significant interactions were further evaluated by performing simple main
190 effect examinations on each level of the interaction, in the event of a significant simple main
191 effect pairwise comparisons were performed. All statistical actions were conducted using
192 SPSS v22.0 (SPSS Inc, Chicago, USA).

193

194 **Results**

195 Tables 1-4 and figure 1 present patellar tendon loading parameters as a function of *brace*,
196 *movement* and *sex*.

197

198 @@@ **FIGURE 1 NEAR HERE** @@@

199 @@@ **FIGURE 2 NEAR HERE** @@@

200 @@@ TABLE 1 NEAR HERE @@@

201 @@@ TABLE 2 NEAR HERE @@@

202 @@@ TABLE 3 NEAR HERE @@@

203 @@@ TABLE 4 NEAR HERE @@@

204

205 *Peak patellar tendon force*

206 A significant main effect ($P < .05$, $\eta^2 = .20$) was found for *movement*. Post-hoc pairwise
207 comparisons showed that peak patellar tendon force was significantly larger in the cut
208 movement compared to the hop ($P = .046$) and run ($P = .008$) conditions.

209

210 In addition a significant main effect ($P < .05$, $\eta^2 = .31$) was observed for *brace*. Post-hoc
211 pairwise comparisons showed that peak patellar tendon force was significantly larger in the
212 no-brace ($P = .013$) condition compared to wearing the brace.

213

214 *Patellar tendon instantaneous load rate*

215 A significant main effect ($P < .05$, $\eta^2 = .29$) was found for *movement*. Post-hoc pairwise
216 comparisons showed that patellar tendon instantaneous load rate was significantly larger in
217 the cut ($P = .032$) and hop ($P = .003$) conditions compared to the run movement. In addition a
218 significant main effect ($P < .05$, $\eta^2 = .45$) was observed for *brace*, with patellar tendon
219 instantaneous load rate being significantly in the no-brace condition compared to wearing the
220 brace.

221

222 Finally a significant ($P < .05$, $\eta^2 = .19$) *brace * movement * sex* interaction was shown.
223 Follow up analyses using simple main effects showed for males that there was a significant
224 main effect ($P < .05$, $\eta^2 = .21$) for *movement*, with the hop ($P = .01$) and cut ($P = .04$)

225 movements being associated with a greater instantaneous load rate than the run movement.
226 For females there was a significant main effect ($P < .05$, $\eta^2 = .86$) for *movement*, with the hop
227 ($P = .00001$) and cut ($P = .002$) movements being associated with a greater instantaneous load
228 rate than the run movement. In addition there was also a main effect ($P < .05$, $\eta^2 = .57$) for
229 *brace* with instantaneous load rate being significantly ($P = .018$) larger in the no-brace
230 condition. Finally a significant ($P < .05$, $\eta^2 = .42$) *brace * movement* interaction was found for
231 females. Follow up analyses showed that there were main effects for the run ($P < .05$, $\eta^2 =$
232 $.89$) and cut ($P < .05$, $\eta^2 = .72$) movements, with instantaneous load rate being significantly
233 greater in the no-brace condition for both movements (cut – $P = .004$ & run – $P = .00001$). No
234 differences were shown for the hop condition.

235

236 *Patellar tendon impulse*

237 A significant main effect ($P < .05$, $\eta^2 = .20$) was found for *movement*. Post-hoc pairwise
238 comparisons showed that peak tendon impulse was significantly larger in the cut ($P = .0002$)
239 and hop ($P = .048$) movements compared to the run condition.

240

241 In addition a significant main effect ($P < .05$, $\eta^2 = .19$) was observed for *brace*, with patellar
242 tendon impulse was significantly larger in the no-brace ($P = .042$) condition compared to
243 wearing the brace.

244

245 Finally, a significant ($P < .05$, $\eta^2 = .19$) *brace * movement * sex* interaction was shown.
246 Follow up analyses using simple main effects showed for males that there was a significant
247 main effect ($P < .05$, $\eta^2 = .35$) for *movement*, with the hop ($P = .001$) and cut ($P = .023$)
248 movements being associated with a greater impulse than the run movement. For females there
249 was a significant main effect ($P < .05$, $\eta^2 = .22$) for *movement*, with the cut ($P = .01$) being

250 associated with a greater impulse than the run movement. Finally a significant ($P < .05$, $\eta^2 =$
251 $.56$) *brace * movement* interaction was found for females. Follow up analyses showed that
252 there was a main effect for the run ($P < .05$, $\eta^2 = .89$) movement, with impulse being
253 significantly ($P = .0004$) greater in the no-brace condition.

254

255 *Sagittal knee kinematics*

256 For the knee flexion angle at footstrike, a significant main effect ($P < .05$, $\eta^2 = .36$) was
257 observed for *brace*, with knee flexion being reduced in the brace condition. For the peak
258 flexion angle, a significant main effect ($P < .05$, $\eta^2 = .28$) was observed for *brace*, with peak
259 flexion being reduced in the brace condition. In addition, a significant main effect ($P < .05$,
260 $\eta^2 = .60$) was observed for *movement*. Post-hoc pairwise comparisons indicated that peak
261 flexion was significantly greater in the cut ($P = .000008$) and hop ($P = .0000009$) movement in
262 comparison to the run and also in the hop compared to the cut ($P = .02$). Finally, a significant
263 *brace * sex* ($P < .05$, $\eta^2 = .22$) interaction was found. Follow up analyses showed that in
264 female athletes only peak knee flexion was significantly reduced in the brace condition for
265 the run ($P < .05$, $\eta^2 = .37$) and hop ($P < .05$, $\eta^2 = .66$) movements.

266

267 **Discussion**

268 The aim of the current investigation was to investigate the effects of a prophylactic knee
269 brace on patellar tendon loading parameters linked to the aetiology of patellar tendinopathy,
270 in male and female recreational athletes. To the authors' knowledge, this represents the first
271 investigation to examine the effects of prophylactic knee bracing in relation to the aetiology
272 patellar tendinopathy.

273

274 A key finding from the current study is that indices of patellar tendon instantaneous load rate
275 and impulse were found to be significantly reduced in female athletes during the run and cut
276 movements when wearing the knee brace. This observation is interesting in that female
277 athletes exhibited significant reductions in patellar tendon loading parameters as a function of
278 the prophylactic brace, yet in male athletes there were no statistical alterations. The
279 mechanisms responsible for this observation are unknown at this stage. However, previous
280 analyses have shown that female's exhibit diminished knee joint proprioception in relation to
281 males (27-30). Prophylactic knee sleeves, such as that used in the current investigation are
282 proposed to promote stimulation of type δ sensory fibres within skin mechanoreceptors (31),
283 and clinical research into their efficacy has shown that they are associated with improvements
284 in knee joint proprioception (32-34). It can be speculated upon that there may be more scope
285 for proprioceptive benefits in females, and that the positive effect of the knee brace in female
286 athletes was mediated by a proprioceptive effect, which may have been responsible for the
287 alterations in peak knee flexion that were evident only in female participants. Reductions in
288 knee flexion are associated with lengthening of the moment arm of the patellar tendon itself,
289 which leads to a reduction in tendon loading. Nonetheless, further mechanistic investigations
290 into the specific effects of prophylactic knee sleeves on joint position sense at the knee are
291 required before this notion can be recognized.

292

293 As stated previously, the aetiology of patellar tendinopathy in athletic populations, relates to
294 the storage and release of energy by the tendon during sports movements (2). Therefore given
295 the increased rate at which the tendon was loaded in the no-brace condition, this observation
296 may have clinical significance. It can be conjectured that female athletes may be able to
297 attenuate their risk from patellar tendinopathy during specific athletic movements through

298 utilization of prophylactic knee bracing. However, further prospective research into the
299 prophylactic effects of knee bracing is required before this can be clinically substantiated.

300

301 A further important observation from this investigation, is that for both male and female
302 athletes, patellar tendon loading was significantly greater in the cut and hop movements in
303 relation to the run condition. It is proposed that this observation relates to the ballistic nature
304 of cut and single leg hop movements, in relation to the run condition, placing greater
305 demands on the knee extensors. It has been shown through epidemiological analyses, that the
306 aetiology of patellar tendinopathy is related to the magnitude of the loads experienced by the
307 tendon itself (2). Importantly, cutting is one of the key abilities of sports games (35) and
308 cutting actions are functionally specific to a range of different individual and team events
309 including but not limited to; association football (36), American football (37), netball (4),
310 tennis (38), squash (16) and basketball (39). In addition, single leg hop landings are similarly
311 common in multidirectional sports including but not limited to; association football (40),
312 American football (41), gymnastics (42), netball (7) and basketball (39). The findings from
313 the current investigation indicate that cut and hop motions may place athletes at increased
314 risk from patellar tendon pathology, therefore conservative prophylactic measures such as
315 knee bracing may be important apparatuses in athletic disciplines and their associated training
316 regimens whereby these movements are common. Future prospective research is clearly
317 required to investigate the longitudinal prophylactic effects of different conservative
318 modalities, in sports which place high mechanical demands on the patellar tendon.

319

320 A potential drawback to the current investigation is that patellar tendon loading parameters
321 were quantified via a musculoskeletal driven model. Although this approach represents an
322 advancement in relation to previous mechanisms, further progression is needed to improve

323 the efficacy of musculoskeletal modeling of patellar tendon kinetics. Although muscle driven
324 simulations of musculoskeletal loading require a range of mechanical assumptions, they have
325 developed significantly in recent years. Thus, musculoskeletal simulations have the potential
326 to become useful tools for clinical analyses in the field of biomechanics.

327

328 In conclusion, whilst previous analyses have investigated the therapeutic and prophylactic
329 effects of knee bracing, the current knowledge with regards to the effects of prophylactic
330 knee bracing on the patellar tendon in functional athletic movements is limited. The current
331 investigation therefore addresses this, by examining the effects of wearing a prophylactic
332 knee brace on patellar tendon loading parameters during run, cut and jump movements in
333 male and female athletes. The current study showed firstly that patellar tendon loading
334 parameters were significantly reduced in female athletes in the run and cut conditions whilst
335 wearing the brace. In addition, for both males and females the cut and hop movements were
336 associated with significantly greater tendon loading in relation to the run motion. Given the
337 association between patellar tendon loading and the aetiology of patellar tendinopathy, this
338 observation may be clinically important. It can be conjectured that female athletes may be
339 able to attenuate their risk from tendinopathy during specific athletic movements through
340 utilization of knee bracing, although further prospective research into the prophylactic effects
341 of knee bracing is required before this can be clinically substantiated.

342

343 **References**

- 344 1. Lian, Ø.B., Engebretsen, L., and Bahr, R. Prevalence of jumper's knee among elite
345 athletes from different sports a cross-sectional study. American Journal of Sports
346 Medicine. 2005; 33: 561-567.

- 347 2. Rudavsky, A., and Cook, J. Physiotherapy management of patellar tendinopathy.
348 Journal of Physiotherapy. 2014; 60: 122-129.
- 349 3. Witvrouw, E., Bellemans, J., Lysens, R., Danneels, L., and Cambier, D. Intrinsic risk
350 factors for the development of patellar tendinitis in an athletic population: a two-year
351 prospective study. American Journal of Sports Medicine. 2001; 29: 190–195.
- 352 4. Maffulli, N., Wong, J., and Almekinders, L.C. Types and epidemiology of
353 tendinopathy. Clinical Journal of Sports Medicine. 2003; 22: 675–692.
- 354 5. Cook, J.L., Khan, K.M., Harcourt, P.R., Grant, M., Young, D.A., and Bonar, S.F. A
355 cross sectional study of 100 athletes with jumper’s knee managed conservatively and
356 surgically. The Victorian Institute of Sport Tendon Study Group. British Journal of
357 Sports Medicine. 1997; 31: 332-336.
- 358 6. Cook, J.L., Khan, K.M, and Purdam, C.R. Conservative treatment of patellar
359 tendinopathy. Physical Therapy in Sport. 2001; 35: 291–294.
- 360 7. Sinclair, J., Vincent, H., Richards, J. Effects of prophylactic knee bracing on knee
361 joint kinetics and kinematics during netball specific movements. Physical Therapy in
362 Sport. 2017; 23: 93–98.
- 363 8. Paluska, S.A., and McKeag, D.B. Knee braces: current evidence and clinical
364 recommendations for their use. American Family Physician. 2000; 61: 411-418.
- 365 9. Warden, S.J., Hinman, R.S., Watson, M.A., Avin, K.G., Bialocerkowski, A.E., and
366 Crossley, K.M. Patellar taping and bracing for the treatment of chronic knee pain: A
367 systematic review and meta-analysis. Arthritis Care & Research. 2008; 59: 73-83.
- 368 10. Ewing, K.A., Fernandez, J.W., Begg, R.K., Galea, M.P., and Lee, P.V. Prophylactic
369 knee bracing alters lower-limb muscle forces during a double-leg drop landing.
370 Journal of Biomechanics. 2016; 49; 3347-3354.

- 371 11. Lee, H., Ha, D, Kan, Y.S., and Park, H.S. Biomechanical Analysis of the Effects of
372 Bilateral Hinged Knee Bracing. *Frontiers in Bioengineering and Biotechnology*. 2016;
373 4: 50-55.
- 374 12. Cappozzo, A., Catani, F., Leardini, A., Benedetti, M.G., and Della C.U. Position and
375 orientation in space of bones during movement: Anatomical frame definition and
376 determination. *Clinical Biomechanics*. 1995; 10: 171-178.
- 377 13. Sinclair, J., Taylor, P.J., Currigan, G., and Hobbs, S.J. The test-retest reliability of
378 three different hip joint centre location techniques. *Movement & Sport Sciences*.
379 2014; 83: 31-39.
- 380 14. Sinclair, J., Hebron, J., and Taylor, P.J. The Test-retest Reliability of Knee Joint
381 Center Location Techniques. *Journal of Applied Biomechanics*. 2015; 31: 117-121.
- 382 15. Graydon, R., Fewtrell, D., Atkins, S., and Sinclair, J. The test-retest reliability of
383 different ankle joint center location techniques. *Foot & Ankle Online Journal*. 2015;
384 8: 1-11.
- 385 16. Sinclair, J., Hobbs, S.J., Protheroe, L., Edmundson, C.J., and Greenhalgh, A.
386 Determination of gait events using an externally mounted shank accelerometer.
387 *Journal of Applied Biomechanics*. 2013; 29: 118-122.
- 388 17. McLean, S.G., Huang, X., Su, A., and Van Den Bogert, A.J. Sagittal plane
389 biomechanics cannot injure the ACL during sidestep cutting. *Clinical Biomechanics*.
390 2004; 19: 828-838.
- 391 18. Janssen, I., Steele, J.R., Munro, B.J., and Brown, N.A. Predicting the patellar tendon
392 force generated when landing from a jump. *Medicine & Science in Sport & Exercise*.
393 2013; 45: 927-934.

- 394 19. Sinclair, J, and Taylor, PJ. Effects of court specific and minimalist footwear on
395 patellar tendon loading during a maximal change of direction task. *Baltic Journal of*
396 *Physical Activity in Health* (In Press).
- 397 20. Sinclair, J., and Taylor, P.J. Sex variation in patellar tendon kinetics during running.
398 *Human Movement*. 2015; 16: 60-63.
- 399 21. DeVita, P., and Hortobagyi, T. Functional knee brace alters predicted knee muscle
400 and joint forces in people with ACL reconstruction during walking. *Journal of*
401 *Applied Biomechanics*. 2001; 17: 297–311.
- 402 22. Ward, S.R., Eng, C.M., Smallwood, L.H., and Lieber, R.L. Are current measurements
403 of lower extremity muscle architecture accurate? *Clinical Orthopaedics and Related*
404 *Research*. 2009; 467: 1074–1082.
- 405 23. Nemeth, G., and Ohlsen, H. In vivo moment arm lengths for hip extensor muscles at
406 different angles of hip flexion. *Journal of Biomechanics*. 1985; 18: 129–140.
- 407 24. Self, B.P., and Paine, D. Ankle biomechanics during four landing techniques.
408 *Medicine & Science in Sports & Exercise*. 2001; 33: 1338-1344.
- 409 25. Spoor, C.W., van Leeuwen, J.L. Knee muscle moment arms from MRI and from
410 tendon travel. *Journal of Biomechanics*. 1992; 25: 201–206.
- 411 26. Herzog, W., and Read, L.J. Lines of action and moment arms of the major force-
412 carrying structures crossing the human knee joint. *Journal of Anatomy*. 1993; 182:
413 213-230.
- 414 27. Nagai, T., Sell, T.C., Abt, J.P., and Lephart, S.M. Reliability, precision, and gender
415 differences in knee internal/external rotation proprioception measurements. *Physical*
416 *Therapy in Sport*. 2012; 13: 233-237.

- 417 28. Muaidi, Q.I. Does gender make a difference in knee rotation proprioception and range
418 of motion in healthy subjects?. *Journal of Back and Musculoskeletal Rehabilitation*.
419 2017; (In press).
- 420 29. Rozzi, S.L., Lephart, S.M., Gear, W.S., and Fu, F.H. Knee joint laxity and
421 neuromuscular characteristics of male and female soccer and basketball players. *The*
422 *American Journal of Sports Medicine*. 1999; 27: 312-319.
- 423 30. Karkousha, R.N. Sex differences of knee joint repositioning accuracy in healthy
424 adolescents. *Bulletin of Faculty of Physical Therapy*. 2016; 21: 56-60.
- 425 31. Callaghan, M.J., Selfe, J., McHenry, A., and Oldham, J. A. Effects of patellar taping
426 on knee joint proprioception in patients with patellofemoral pain syndrome. *Manual*
427 *Therapy*. 2008; 13: 192-199.
- 428 32. Baltaci, G., Aktas, G., Camci, E., Oksuz, S., Yildiz, S., and Kalaycioglu, T. The effect
429 of prophylactic knee bracing on performance: balance, proprioception, coordination,
430 and muscular power. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2011; 19:
431 1722-1728.
- 432 33. Herrington, L., Simmonds, C., and Hatcher, J. The effect of a neoprene sleeve on knee
433 joint position sense. *Research in Sports Medicine*. 2005; 13: 37-46.
- 434 34. McNair, P.J., Stanley, S.N., and Strauss, G.R. Knee bracing: effects on
435 proprioception. *Archives of Physical Medicine and Rehabilitation*. 1996; 77: 287-289.
- 436 35. Sheppard, J.M., and Young, W.B. Agility literature review: classifications, training
437 and testing. *Journal of Sports Sciences*. 2006; 24: 919-932.
- 438 36. Strutzenberger, G., Cao, H. M., Koussev, J., Potthast, W., and Irwin, G. Effect of turf
439 on the cutting movement of female football players. *Journal of Sport and Health*
440 *Science*. 2014; 3: 314-319.

- 441 37. Carson, D.W., Myer, G.D., Hewett, T.E., Heidt, R. S., and Ford, K.R. Increased
442 plantar force and impulse in American football players with high arch compared to
443 normal arch. *The Foot* 2012; 22: 310-314.
- 444 38. Stacoff, A., Steger, J., Stuessi, E., and Reinschmidt, C. Lateral stability in sideward
445 cutting movements. *Medicine & Science in Sport & Exercise*. 1996; 28: 350-358.
- 446 39. Cowley, H.R., Ford, K.R., Myer, G.D., Kernozek, T.W., and Hewett, T.E. Differences
447 in neuromuscular strategies between landing and cutting tasks in female basketball
448 and soccer athletes. *Journal of Athletic Training*. 2006; 41: 67-73.
- 449 40. Lyle, M.A., Valero-Cuevas, F.J., Gregor, R.J., and Powers, C.M. Control of dynamic
450 foot-ground interactions in male and female soccer athletes: females exhibit reduced
451 dexterity and higher limb stiffness during landing. *Journal of Biomechanics*. 2014; 47:
452 512-517.
- 453 41. Masters, C., Johnstone, J., and Hughes, G. The Effect of Arm Position on Lower
454 Extremity Kinematics during a Single Limb Drop Landing: A Preliminary Study.
455 *Journal of Functional Morphology and Kinesiology*. 2016; 1: 282-288.
- 456 42. Chaudhari, A.M., Hearn, B.K., and Andriacchi, T.P. Sport-dependent variations in
457 arm position during single-limb landing influence knee loading. *The American
458 journal of Sports Medicine*. 2005; 33: 824-830.

459 Table 1: Patellar tendon load parameters (means, standard deviations and 95% confidence intervals) as a function of *brace* and *movement*
 460 conditions in male athletes.

	Male																	
	Run						Cut						Hop					
	Brace			No-Brace			Brace			No-Brace			Brace			No-Brace		
	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI
Peak patellar tendon load (BW)	7.03	1.25	6.24 - 7.83	7.48	1.48	6.54 - 8.42	8.08	2.03	6.80 - 9.37	8.30	1.46	7.37 - 9.22	7.76	1.67	6.69 - 8.82	8.07	1.22	7.30 - 8.85
Patellar tendon instantaneous load rate (BW/s)	335.41	115.57	261.98 - 408.84	358.54	114.05	286.07 - 431.01	445.64	162.25	342.55 - 548.73	457.89	153.72	360.22 - 555.56	442.39	184.86	324.94 - 559.85	518.55	270.58	346.63 - 690.49
Patellar tendon impulse (BW·s)	0.61	0.13	0.52 - 0.69	0.82	0.25	0.66 - 0.97	1.01	0.31	0.81 - 1.21	0.98	0.30	0.79 - 1.17	1.01	0.50	0.69 - 1.32	0.96	0.38	0.72 - 1.20

461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473

474 Table 2: Patellar tendon load parameters (means, standard deviations and 95% confidence intervals) as a function of brace and movement
 475 conditions in female athletes.

	Female																	
	Run						Cut						Hop					
	Brace			No-Brace			Brace			No-Brace			Brace			No-Brace		
	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI
Peak patellar tendon load (BW)	7.69	0.76	7.05 - 8.32	9.42	1.03	8.56 - 10.29	8.79	1.14	7.84 - 9.73	9.26	1.93	7.64 - 10.87	7.88	0.76	7.24 - 8.52	8.70	2.38	6.72 - 10.69
Patellar tendon instantaneous load rate (BW/s)	289.14	65.59	234.31 - 343.98	370.06	93.67	291.75 - 488.40	353.17	116.46	255.81 - 450.54	422.01	142.91	302.54 - 541.49	484.43	63.87	431.0 - 537.83	487.58	115.96	390.64 - 584.53
Patellar tendon impulse (BW·s)	0.79	0.10	0.70 - 0.87	1.00	0.07	0.94 - 1.05	0.95	0.12	0.89 - 1.05	1.05	0.19	0.90 - 1.25	0.84	0.09	0.76 - 0.91	0.99	0.42	0.64 - 1.34

476

477

478 Table 3: Knee flexion parameters (means, standard deviations and 95% confidence intervals) as a function of brace and movement conditions in
 479 male athletes.

	Male																	
	Run						Cut						Hop					
	Brace			No-Brace			Brace			No-Brace			Brace			No-Brace		
	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI
Angle at footstrike (°)	10.92	4.34	8.16-16.68	13.30	5.98	9.50-17.10	10.26	4.48	7.42-13.11	12.67	5.76	9.01-16.32	12.94	6.29	8.95-16.94	13.70	3.16	11.70-15.71
Peak flexion (°)	36.55	2.64	34.87-38.23	39.05	4.06	36.47-41.63	44.45	4.18	41.79-47.10	43.92	3.82	41.50-46.35	45.26	6.60	41.07-49.46	45.00	5.79	41.32-48.68

480 Table 4: Knee flexion parameters (means, standard deviations and 95% confidence intervals) as a function of brace and movement conditions in
 481 female athletes.

	Female																	
	Run						Cut						Hop					
	Brace			No-Brace			Brace			No-Brace			Brace			No-Brace		
	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI
Angle at footstrike (°)	11.46	2.66	9.24-13.69	16.44	4.94	12.31-20.57	13.16	3.98	9.83-16.49	17.87	4.53	14.09-21.65	12.49	3.14	9.86-15.12	17.99	6.27	12.74-23.23
Peak flexion (°)	36.64	1.92	35.04-38.25	41.12	3.84	37.91-44.33	44.35	2.12	42.85-46.12	45.71	3.12	43.10-48.32	49.74	8.48	42.65-56.83	53.39	11.50	43.78-63.00

482

483

484

485 **List of figures**

486 Figure 1: Patellar tendon forces as a function of brace and movement conditions – black = no-brace & grey = brace (a. = male run, b. = female

487 run, c. = male cut, d. = female cut, e. = male hop and f. = female hop).