

Central Lancashire Online Knowledge (CLoK)

Title	On convoluters on Lp-spaces
Type	Article
URL	https://clock.uclan.ac.uk/21777/
DOI	https://doi.org/10.4064/sm170601-24-11
Date	2018
Citation	Daws, Matthew and Spronk, Nico (2018) On convoluters on Lp-spaces. <i>Studia Mathematica</i> , 245. pp. 15-31. ISSN 0039-3223
Creators	Daws, Matthew and Spronk, Nico

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.4064/sm170601-24-11>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

On convoluters on L^p -spaces

Matthew Daws

Leeds, United Kingdom

E-mail: matt.daws@cantab.net

Nico Spronk

Department of Pure Mathematics

University of Waterloo

Waterloo, Ontario, N2L3G1, Canada

E-mail: nspronk@uwaterloo.ca

November 29, 2017

Abstract

We prove two theorems about convolution operators on $L^p(G)$ for a locally compact group G . First, if G has the approximation property, then the algebra of convoluters is the algebra of pseudo-measures. Second, the bicommutant of the algebra of pseudo-measures is the algebra of convoluters.

1 Introduction

Let G be a locally compact group and $L^p(G)$ denote the L^p -space with respect to left Haar measure, for $p \in [1, \infty]$. The algebra of bounded operators $\mathcal{B}(L^p(G))$ for $p \in (1, \infty)$ admits a natural predual (which we specify in Section 2, below) and hence admits a specified weak* topology which is finer than the weak operator topology. We consider the two subalgebras, the *pseudo-measures* and *convoluters*, which are given by

$$PM_p(G) = \overline{\text{lin}}^{w*} \lambda_p(G) \quad (\text{weak* closed linear span})$$

$$CV_p(G) = \rho_p(G)' \quad (\text{commutant})$$

2010 *Mathematics Subject Classification*: Primary 43A15; Secondary 22D12, 47L10.
Key words and phrases: convoluter, pseudo-measure, approximation property.

where the right and left regular representations $\lambda_p, \rho_p : G \rightarrow \mathcal{B}(L^p(G))$ are given by

$$\lambda_p(s)f(t) = f(s^{-1}t), \quad \rho_p(s)f(t) = \Delta(s)^{1/p}f(ts)$$

for all s and a.e. t . Here Δ is the Haar modular function. Since $\lambda_p(G) \subseteq \rho_p(G)'$ and commutant algebras are weak operator closed, hence weak* closed, we have that $PM_p(G) \subseteq CV_p(G)$. Notice, moreover, that the invertible isometry U in $\mathcal{B}(L^p(G))$, given by $Uf(t) = \Delta^{1/p}(t^{-1})f(t^{-1})$, is self-inverse and intertwines λ_p and ρ_p , which shows we may interchange the roles of left and right.

The *approximation property* was defined by Haagerup and Kraus [9]. A detailed definition is provided in Section 3, below.

Theorem 1.1. *If G has the approximation property, then*

$$CV_p(G) = PM_p(G).$$

A complete description of connected groups with the approximation property is provided by Haagerup, Knudby and de Laat [10]: any simple Lie quotient must have real rank 1. This builds on an enormous body of work, including [18, 11, 12]. The approximation property passes to closed subgroups, extension groups, and is passed up from lattices [9]. In particular, countably generated free groups enjoy this property, even the stronger property of *weak amenability*, defined by de Cannière and Haggerup [7]. The approximation property does not pass to general quotients: the finitely generated lattice $SL_3(\mathbb{Z})$ in $SL_3(\mathbb{R})$ is a quotient of some free group \mathbb{F}_n . In fact, we have the following implications of properties of G :

$$\text{amenable} \Rightarrow \text{weakly amenable} \Rightarrow \text{approximation property}.$$

The first implication is given by Leptin [19], i.e. a bounded approximate identity in the Fourier algebra $A_2(G)$ satisfies the definition of weak amenability of [7]. For amenable G , Theorem 1.1, is proved by Herz [15]; see Remark 2.9.

A certain p -approximation property, which is implied by the approximation property was defined by An, Lee and Ruan [1], and studied vigorously by Vergara [23]. Using this, Vergara strengthens our Theorem 1.1, but relies heavily on our methods. See Remark 3.3, below.

On a related note, we also give an elementary proof of the following. Recall that the intertwiner U is given above

Theorem 1.2. *We have commutation results:*

$$PM_p(G)' = U CV_p(G)U \text{ and } (U CV_p(G)U)' = CV_p(G).$$

In particular, we have bicommutant $PM_p(G)'' = CV_p(G)$.

For $p = 2$ the comutation result of convoluteds was proved by Dixmier [8] using left Hilbert algebras. Recently, Pham [21] gave an elementary proof in the $p = 2$ case. This proof is similar to the one offered here, but we wish to note that ours was conducted independently and posted on [arXiv](#) in 2016.

The present article is an updated version of our work [6], the main body of which was first posted to the [arXiv](#) in 2013. Both theorems appear to be folklore — see Cowling [3], and Herz [15] — but we have been unable to track down complete proofs. Theorem 1.1 was used in recent work of Öztop and the second named author [20] to obtain, for a pair of groups G and H with approximation property, a tensor product description of $A_p(G \times H)$. Given recent developments related to this work and interest in it, in particular [21, 23], we have elected to update our results and submit them for publication.

2 On preduals of pseudo-measures and convoluteds

The entire goal of the present section is to present background for the proofs of our two main theorems. We also redevelop some methods of Cowling [3], who gave an innovative and insightful description of a predual of $CV_p(G)$. The present methods give a new perspective and help to illuminate the role of the Herz-Schur multipliers which we discuss in 2.2, below.

We let p' be the conjugate index to p , so $\frac{1}{p} + \frac{1}{p'} = 1$. We recall that $\mathcal{B}(L^p(G))$ is the dual of the space $N^p(G) = L^{p'}(G) \hat{\otimes} L^p(G)$ by way of dual pairing $\langle T, \xi \otimes f \rangle = \langle \xi, Tf \rangle = \int_G \xi(t)[Tf](t) dt$. Elements $\omega = \sum_{n=1}^{\infty} \xi_n \otimes f_n$ of $N^p(G)$ may be viewed as functions $\omega(s, t) = \sum_{n=1}^{\infty} \xi_n(s)f_n(t)$ up to marginal almost everywhere (m.a.e.) equivalence; i.e. excepting marginally null sets, $(N_1 \times G) \cup (G \times N_2)$, where N_1 and N_2 are Haar null sets.

We let $P : N^p(G) \rightarrow A_p(G) \subseteq C_0(G)$ be given for s in G by

$$P \left(\sum_{n=1}^{\infty} \xi_n \otimes f_n \right) (s) = \sum_{n=1}^{\infty} \langle \xi_n, \lambda_p(s) f_n \rangle = \sum_{n=1}^{\infty} \xi_n * \check{f}_n$$

where $\check{f}(t) = f(t^{-1})$ a.e., i.e. $P\omega(s) = \int_G \omega(t, s^{-1}t) dt$. Hence $A_p(G)$ is normed as the coimage of P , i.e. $A_p(G) \cong N^p(G)/\ker P$ isometrically. It

is straightforward to see that $PM_p(G)$ and $CV_p(G)$ admit the following pre-annihilators in $N^p(G)$:

$$\begin{aligned} {}^\perp PM_p(G) &= \ker P \\ {}^\perp CV_p(G) &= \overline{\text{lin}}\{\rho_{p'}(t^{-1})\xi \otimes f - \xi \otimes \rho_p(t)f : \\ &\quad t \in G, \xi \in L^{p'}(G), f \in L^p(G)\} \end{aligned}$$

where we note that adjoint of these right translation operators is given by $\rho_p(t)^* = \rho_{p'}(t^{-1})$.

It is standard that

$$PM_p(G) \cong A_p(G)^* \text{ and } CV_p(G) \cong [N^p(G)/{}^\perp CV_p(G)]^*$$

which gives us distinguished preduals of these algebras of operators.

2.1 $N^p(G)$ as an algebra

Herz [13] showed that $A_p(G)$ is always a subalgebra of $C_0(G)$, and further showed ([14]) that it has Gelfand spectrum given by evaluations at points of G . We note that $A_p(G)$ is known as the *Figà-Talamanca–Herz algebra*. Herz’s technique for showing that $A_p(G)$ is an algebra lifts naturally to $N^p(G)$. We note that the amplification $T \mapsto T \otimes I : \mathcal{B}(L^p(G)) \rightarrow \mathcal{B}(L^p(G; L^p(G))) \cong \mathcal{B}(L^p(G \times G))$ is an isometry. We consider the *fundamental isometry* W_p in $L^p(G \times G)$ given for a.e. (s, t) by

$$W_p f(s, t) = f(s, st).$$

We note that W_p is invertible with $W_p^* = W_{p'}^{-1}$. We define a *co-product* on $\mathcal{B}(L^p(G))$, $\Gamma : \mathcal{B}(L^p(G)) \rightarrow \mathcal{B}(L^p(G \times G))$ by

$$\Gamma(T) = W_p^{-1}(T \otimes I)W_p.$$

This spatially implemented map is evidently weak*-weak* continuous, and hence admits a preadjoint. We note that $N^p(G) \otimes N^p(G)$ comprises a dense subspace of $N^p(G \times G)$ given on elementary tensors of elementary tensors (“really elementary tensors”) by $(\xi \otimes f) \otimes (\eta \otimes g) \mapsto (\xi \otimes \eta) \otimes (f \otimes g)$. Let us compute Γ_* on these really elementary tensors. We have

$$\begin{aligned} \langle T, \Gamma_*((\xi \otimes f) \otimes (\eta \otimes g)) \rangle &= \langle W_p^{-1}(T \otimes I)W_p, (\xi \otimes f) \otimes (\eta \otimes g) \rangle \\ &= \langle W_{p'}(\xi \otimes \eta), (T \otimes I)W_p(f \otimes g) \rangle \\ &= \int_G \int_G \xi(s)\eta(st)(T \otimes I)[(s, t) \mapsto f(s)g(st)] ds dt \\ &= \int_G \langle \xi \eta_t, T(fg_t) \rangle dt \end{aligned}$$

where $\eta_t(s) = \eta(st)$, for example. Hence we see that

$$(2.1) \quad \Gamma_*((\xi \otimes f) \otimes (\eta \otimes g)) = \int_G (\xi \eta_t) \otimes (f g_t) dt.$$

The equation

$$\int_G \int_G (\xi \eta_t \vartheta_s) \otimes (f g_t h_s) dt ds = \int_G \int_G (\xi \eta_t \vartheta_{ts}) \otimes (f g_t h_{ts}) ds dt$$

shows that Γ_* is an associative product. We note that though $\Gamma(\lambda_p(r)) = \lambda_p(r) \otimes \lambda_p(r)$, which implies that $\Gamma|_{PM_p(G)}$ is cocommutative, we have for a multiplication operator M_φ , where $\varphi \in L^\infty(G)$, that $\Gamma(M_\varphi) = M_\varphi \otimes I$, which shows that Γ is not generally cocommutative. Hence Γ_* is not commutative.

Let $\mathcal{K}(G)$ denote the family of all compact neighbourhoods of the identity in G . For K in $\mathcal{K}(G)$ we let $L^p(K) = 1_K L^p(G)$, which is a 1-complemented subspace. Likewise we define $L^{p'}(K)$ and

$$N^p(K) = L^{p'}(K) \hat{\otimes} L^p(K)$$

which is a subspace of $N^p(G)$.

Proposition 2.1. *The space ${}^\perp PM_p(G) = \ker P$ is a Γ_* -ideal in $N^p(G)$, while each of ${}^\perp CV_p(G)$ and $N^p(K)$, for K in $\mathcal{K}(G)$, are right ideals.*

Proof. That $P : N^p(G) \rightarrow C_0(G)$ is Γ_* -pointwise multiplicative is shown in [13], hence ${}^\perp PM_p(G) = \ker P$ is an ideal. Let us consider the case of ${}^\perp CV_p(G)$, manually. On really elementary tensors we have

$$\begin{aligned} & \Gamma_*((\rho_{p'}(s^{-1})\xi \otimes f - \xi \otimes \rho_p(s)f) \otimes (\eta \otimes g)) \\ &= \int_G [(\rho_{p'}(s^{-1})\xi)\eta_t \otimes f g_t - \xi \eta_t \otimes (\rho_p(s)f)g_t] dt \\ &= \int_G \rho_{p'}(s^{-1})(\xi \eta_{st}) \otimes f g_t dt - \int_G \xi \eta_t \otimes \rho_p(s)(f g_{s^{-1}t}) dt \\ &= \int_G [\rho_{p'}(s^{-1})(\xi \eta_{st}) \otimes f g_t - \xi \eta_{st} \otimes \rho_p(s)(f g_t)] dt \end{aligned}$$

which is an integral of elements in ${}^\perp CV_p(G)$. That each $N^p(K)$ is a right ideal follows readily from (2.1). Indeed if $\xi \otimes f \in N^p(K)$ then

$$\begin{aligned} (1_K \otimes 1_K)\Gamma_*((\xi \otimes f) \otimes (\eta \otimes g)) &= \int_G 1_K \xi \eta_t \otimes 1_K f g_t dt \\ &= \Gamma_*((\xi \otimes f) \otimes (\eta \otimes g)) \end{aligned}$$

so $\Gamma_*((\xi \otimes f) \otimes (\eta \otimes g)) \in N^p(K)$. □

It will be convenient below, to consider the following space of elements. We let $L^1_{\text{loc}}(G)$ denote the space of locally a.e. equivalence classes of measurable functions on G which are integrable on any compact set. An application of Hölder's inequality shows that $L^q(G) \subseteq L^1_{\text{loc}}(G)$ for any q in $[1, \infty]$. If $h \in L^1_{\text{loc}}(G)$ and $g \in C_c(G)$ then $h * g$ is well-defined as an element of $L^1_{\text{loc}}(G)$. We let

$$CV_p^1(G) = \{h \in L^1_{\text{loc}}(G) : \sup\{\|h * g\|_p : g \in C_c(G), \|g\|_p \leq 1\} < \infty\}.$$

Each element h of $CV_p^1(G)$ defines an operator $\lambda_p(h)$ in $\mathcal{B}(L^p(G))$. By testing on elements $\rho_{p'}(s^{-1})g \otimes f - g \otimes \rho_p(s)f$ where $f, g \in C_c(G)$, and applying a standard density argument, we see that $\lambda_p(h) \in CV_p(G)$.

The next result is really a repackaging of the main result of Cowling [3], and is used extensively through the rest of this note. The second part is a localization theorem.

Theorem 2.2. (i) *Each T in $CV_p(G)$ may be approximated in the strong operator topology by a bounded net of elements $\lambda_p(h_i)$ where each $h_i \in CV_p^p(G) = CV_p^1 \cap L^p(G)$.*

(ii) *For each K in $\mathcal{K}(G)$, we have that*

$${}^\perp PM_p(G) \cap N^p(K) = {}^\perp CV_p(G) \cap N^p(K).$$

Proof. (i) Let $T \in CV_p(G)$. Let $(f_i) \subset C_c(G)$ be a contractive approximate identity for $L^1(G)$. Then for g in $C_c(G)$ we have

$$T(f_i * g) = T(\rho_p(\Delta^{-1/p}\check{g})f_i) = \rho_p(\Delta^{-1/p}\check{g})T(f_i) = T(f_i) * g$$

where $\check{g}(t) = g(t^{-1})$ a.e. Notice then, that

$$\begin{aligned} \|T(f_i) * g\|_p &= \|T(f_i * g)\|_p \leq \|T\| \|f_i * g\|_p \leq \|T\| \|g\|_p \\ \text{and } \|T(f_i) * g - Tg\|_p &= \|T(f_i * g - g)\|_p \xrightarrow{i} 0. \end{aligned}$$

It follows that $h_i = T(f_i)$ gives the desired net of elements in $CV_p^p(G)$.

(ii) Since ${}^\perp PM_p(G) \supseteq {}^\perp CV_p(G)$, it suffices to show that ${}^\perp PM_p(G) \cap N^p(K) \subseteq {}^\perp CV_p(G)$. From (i), above, it suffices to show that for any h in $CV_p^1(G)$, that $\lambda_p(h)$ annihilates ${}^\perp PM_p(G) \cap N^p(K)$.

Now let $\xi \otimes f \in N^p(K)$, and let (f_n) be a sequence in $C_c(G)$ whose supports are in the neighbourhood KK^{-1} of K and converges to f in $L^p(G)$.

Then we have that

$$\begin{aligned}
\langle \xi, \lambda_p(h)f \rangle &= \lim_n \langle \xi, \lambda_p(h)f_n \rangle = \lim_n \int_K \xi(t)h * f_n(t) dt \\
&= \lim_n \int_K \xi(t) \int_{K^2K^{-1}} h(s)f_n(s^{-1}t) ds dt \\
&= \lim_n \int_{K^2K^{-1}} h(s) \int_K \xi(t)f_n(s^{-1}t) dt ds \\
&= \lim_n \int_{K^2K^{-1}} h(s)P(\xi \otimes f_n)(s) ds \\
&= \int_{K^2K^{-1}} h(s)P(\xi \otimes f)(s) ds
\end{aligned}$$

where each interchange of integrals is justified by Fubini's theorem, as $(\xi \otimes h)1_{K \times K^2K^{-1}}$ is integrable and each f_n is bounded, and the limits are justified by L^p -convergence of f_n , then uniform convergence of $P(\xi \otimes f_n)$. Hence, if $\omega = \sum_{n=1}^{\infty} \xi_n \otimes f_n \in {}^\perp PM_p(G) \cap N^p(K)$, then uniform convergence of $\sum_{n=1}^{\infty} P(\xi_n \otimes f_n) = P\omega = 0$ provides that

$$\langle \lambda_p(h), \omega \rangle = \sum_{n=1}^{\infty} \langle \xi_n, \lambda_p(h)f_n \rangle = \sum_{n=1}^{\infty} \int_{K^2K^{-1}} h(s)P(\xi_n \otimes f_n)(s) ds = 0$$

which is the desired annihilation condition. \square

We note that proving Theorem 1.1 amounts, in effect, to showing that $\lambda_p(CV_p^1(G)) \subseteq PM_p(G)$, even that $\lambda_p(CV_p^p(G)) \subseteq PM_p(G)$. Attacking this directly seems very delicate; see Cowling's approach [2] for certain simple connected Lie groups with real rank 1. We will proceed through a different sequence of observations.

Corollary 2.3. (i) *The space $\bigcup_{K \in \mathcal{K}(G)} {}^\perp PM_p(G) \cap N^p(K)$ is dense in ${}^\perp CV_p(G)$.*

(ii) *Every Γ_* -commutator is an element of ${}^\perp CV_p(G)$. Hence ${}^\perp CV_p(G)$ is a two-sided Γ_* -ideal.*

Proof. (i) We observe that the family of elements

$$\bigcup_{K \in \mathcal{K}(G)} \{ \rho_{p'}(s^{-1})\xi \otimes f - \xi \otimes \rho_p(s)f : s \in K, \xi \in L^{p'}(K), f \in L^p(K) \}$$

is contained in $\bigcup_{K \in \mathcal{K}(G)} {}^\perp CV_p(G) \cap N^p(KK^{-1} \cup K^2)$, and its linear span is dense in ${}^\perp CV_p(G)$.

(ii) Any commutator $\Gamma_*(\omega \otimes \mu - \mu \otimes \omega)$ can be approximated by elements of the form $\Gamma_*([1_K \otimes 1_K]\omega \otimes [1_K \otimes 1_K]\mu - [1_K \otimes 1_K]\mu \otimes [1_K \otimes 1_K]\omega)$ where $K \in \mathcal{K}(G)$. Such elements are in ${}^\perp PM_p(G) \cap N^p(K)$, hence in ${}^\perp CV_p(G)$.

Now, if $\omega \in {}^\perp CV_p(G)$ and $\mu \in N^p(G)$ we have

$$\Gamma_*(\mu \otimes \omega) = \Gamma_*(\omega \otimes \mu) + \Gamma_*(\mu \otimes \omega - \omega \otimes \mu) \in {}^\perp CV_p(G)$$

which establishes the second fact. \square

Hence Γ_* induces a commutative product on

$$\overline{A}_p(G) = N^p(G)/{}^\perp CV_p(G)$$

which is a predual of $CV_p(G)$. We let $\overline{P} : N^p(G) \rightarrow \overline{A}_p(G)$ denote the quotient map. We note that there is a further natural quotient map $\overline{P}(\omega) \mapsto P\omega : \overline{A}_p(G) \rightarrow A_p(G)$. It is shown by Cowling [3] that this is the Gelfand transform. In particular, semisimplicity of $\overline{A}_p(G)$ is equivalent to having ${}^\perp CV_p(G) = {}^\perp PM_p(G)$, i.e. $CV_p(G) = PM_p(G)$.

Now we let for K in $\mathcal{K}(G)$

$$A_p(K) = P(N^p(K)).$$

We remark that if K is an open subgroup of G , then our definition of $A_p(K)$ coincides with the usual definition. For a in $A_p(K)$

$$\|a\|_{A_p(K)} = \inf\{\|\omega\|_{N^p(G)} : \omega \in N^p(K), a = P(\omega)\}$$

is a norm on $A_p(K)$.

The set $\bigcup_{K \in \mathcal{K}(G)} A_p(K)$ is the algebra $A_{p,c}(G)$ of compactly supported elements of $A_p(G)$. Indeed, if u in $A_{p,c}(G)$ is supported on compact S , let $L \in \mathcal{K}(G)$ and $u_L = \frac{1}{m(L)} 1_{SL} * \check{1}_L$ is 1 on S , hence $u = uu_L \in A_p(K)$ whenever $SL \cup L \subseteq K$.

The following will play a critical role in the proof of Lemma 3.2, and hence in the proof of Theorem 1.1.

Corollary 2.4. *Let $N_c^p(G) = \bigcup_{K \in \mathcal{K}} N^p(K)$. Then $\overline{P}(N_c^p(G))$ is a dense ideal in $\overline{A}_p(G)$ which is algebraically isomorphic to $A_{p,c}(G)$. Furthermore, for ω in $N_c^p(G)$ we have*

$$\|\overline{P}(\omega)\|_{\overline{A}_p(G)} = \inf\{\|P(\omega)\|_{A_p(K)} : \omega \in N^p(K)\}$$

and the pairing with $CV_p(G)$ depends only on $P(\omega)$; i.e. if $T \in CV_p(G)$ we may write

$$(2.2) \quad \langle T, \omega \rangle = \langle T, P\omega \rangle.$$

Proof. It is evident that $N_c^p(G)$ is a dense right Γ_* -ideal in $N^p(G)$, so $\overline{P(N_c^p(G))}$ is a dense ideal in $\overline{A_p(G)}$, by part (ii) of the prior corollary. We use part (i) of the prior corollary to see that

$$\begin{aligned} \|\overline{P}(\omega)\|_{\overline{A_p(G)}} &= \text{dist}(\omega, {}^\perp CV_p(G)) \\ &= \inf_{K \in \mathcal{K}(G)} \text{dist}(\omega, {}^\perp PM_p(G) \cap N^p(K)) \\ &= \inf\{\|P(\omega)\|_{A_p(K)} : K \in \mathcal{K}(G), \omega \in N^p(K)\} \end{aligned}$$

where we note that the norm on $A_p(K)$ is given as the coimage: $A_p(K) \cong N^p(K)/\ker P|_{N^p(K)}$, where $\ker P|_{N^p(K)} = {}^\perp PM_p(G) \cap N^p(K)$. Hence if, further, $P\omega = 0$ then for T in $CV_p(G)$ we have

$$|\langle T, \omega \rangle| \leq \|T\| \|\overline{P}(\omega)\|_{\overline{A_p(G)}} = 0$$

which establishes (2.2). \square

We record the following localization principle. In terminology of Herz [15], (b), below, is the condition of being “formally of compact support”.

Theorem 2.5. *The following are equivalent:*

- (a) $CV_p(G) = PM_p(G)$;
- (b) for each u in $A_{p,c}(G)$ we have

$$\|u\|_{A_p(G)} = \inf\{\|u\|_{A_p(K)} : K \in \mathcal{K}(G), u \in A_p(K)\}; \text{ and}$$

- (c) there is $C > 0$ such that for each u in $A_{p,c}(G)$ we have

$$\inf\{\|u\|_{A_p(K)} : K \in \mathcal{K}(G), u \in A_p(K)\} \leq C\|u\|_{A_p(G)}.$$

Proof. That (a) implies (b) is immediate from the last corollary, whilst that (b) implies (c) is obvious for $C \geq 1$. If (c) holds, then on the dense subspace $A_{p,c}(G)$, which we may identify with $\overline{P(N_c^p(G))}$, we have that $\|\cdot\|_{A_p(G)}$ and $\|\cdot\|_{\overline{A_p(G)}}$ are equivalent norms, so the map $\omega + {}^\perp CV_p(G) \mapsto P(\omega)$ is injective, which implies that ${}^\perp CV_p(G) = {}^\perp PM_p(G)$. The latter fact gives (a). \square

2.2 The action of Herz-Schur multipliers

Let $[\text{SQ}_p]$ denote the class of spaces which are isometrically isomorphic to a subspace of a quotient of an L^p -space. Kwapien [17] noted that for E in $[\text{SQ}_p]$ that the ampliation $T \mapsto T \otimes I_E : \mathcal{B}(L^p(G)) \rightarrow \mathcal{B}(L^p(G; E))$ is an

isometry. In fact, he showed that this property characterizes elements of the class $[\text{SQ}_p]$. See Dales et al [4] for an exposition on this. Hence, by duality, the map $\gamma_E : L^{p'}(G; E^*) \hat{\otimes} L^p(G; E) \rightarrow N^p(G)$ given on elementary tensors m.a.e. by

$$(2.3) \quad \gamma_E(\Xi \otimes F)(s, t) = \langle \Xi(s), F(t) \rangle_{E^*, E}$$

is a quotient map, in particular a contraction. Also see Herz [14]. Let us define the p -Herz-Schur multiplier space by

$$\begin{aligned} \mathcal{M}_p(G) = \{ \varphi : G \rightarrow \mathbb{C} \mid \varphi(st^{-1}) = \langle \beta(s), \alpha(t) \rangle \text{ where each} \\ \alpha : G \rightarrow E, \beta : G \rightarrow E^* \text{ is continuous} \\ \text{and bounded for some } E \text{ in } [\text{SQ}_p] \} \end{aligned}$$

These were essentially described by Herz [15, 16], and the specific version we are using here was given by the first named author [5]. As shown in [5], $\mathcal{M}_p(G)$ is the algebra of “ p -completely bounded multipliers” of $A_p(G)$, and a Banach algebra with respect to the norm

$$\|\varphi\|_{\mathcal{M}_p} = \inf \{ \|\beta\|_{\infty} \|\alpha\|_{\infty} : \varphi(st^{-1}) = \langle \beta(s), \alpha(t) \rangle \text{ as above} \}.$$

Proposition 2.6. (i) $N^p(G)$ is a Banach $\mathcal{M}_p(G)$ -module via the action given m.a.e. by

$$\varphi \cdot \omega(s, t) = \varphi(st^{-1})\omega(s, t).$$

(ii) Each of ${}^{\perp}PM_p(G)$ and $N^p(K)$ ($K \in \mathcal{K}(G)$) are $\mathcal{M}_p(G)$ -submodules, hence so too is ${}^{\perp}CV_p(G)$.

Proof. (i) This is an immediate application of the map γ_E in (2.3), above. If $\varphi \in \mathcal{M}_p(G)$ with $\varphi(st^{-1}) = \langle \beta(s), \alpha(t) \rangle$, and $\xi \otimes f$ is an elementary tensor in $N^p(G)$, then let $\xi\beta(s) = \xi(s)\beta(s)$ in $L^{p'}(G; E^*)$ and $f\alpha(t) = f(t)\alpha(t)$ in $L^p(G; E)$. Notice that $\|\xi\beta\|_{p'; E^*} \leq \|\xi\|_{p'} \|\beta\|_{\infty}$ and $\|f\alpha\|_{p; E} \leq \|f\|_p \|\alpha\|_{\infty}$. Then

$$\varphi \cdot (\xi \otimes f) = \gamma_E(\xi\beta \otimes f\alpha) \in N^p(G)$$

and hence

$$\|\varphi \cdot (\xi \otimes f)\|_{N^p} \leq \|\xi\beta\|_{p'; E^*} \|f\alpha\|_{p; E} \leq \|\beta\|_{\infty} \|\alpha\|_{\infty} \|\xi\|_{p'} \|f\|_p$$

from which it follows that $\|\varphi \cdot (\xi \otimes f)\|_{N^p} \leq \|\varphi\|_{\mathcal{M}_p} \|\xi \otimes f\|_{N^p}$.

(ii) If $\varphi \in \mathcal{M}_p(G)$ and $\omega \in \ker P = {}^{\perp}PM_p(G)$ then

$$P(\varphi \cdot \omega)(s) = \int_G \varphi(t(s^{-1}t)^{-1})\omega(t, s^{-1}t) dt = \varphi(s)P\omega(s) = 0.$$

It is facile that each $N^p(K)$ is an $\mathcal{M}_p(G)$ -submodule. Thus each ${}^\perp PM_p(G) \cap N^p(K)$ is an $\mathcal{M}_p(G)$ -submodule, and the result for ${}^\perp CV_p(G)$ follows from the density result, Corollary 2.3 (i), above. \square

It is immediate that the action in (ii), above, induces actions of $\mathcal{M}_p(G)$ on each of $A_p(G)$ and $A_p(K)$ (each by pointwise multiplication) and on $\overline{A_p(G)}$, making each a continuous module. The spaces $A_p(K)$ are shown to be Banach $A_p(G)$ -modules in [20], by a method quite similar to that above. We note that the natural map $\overline{P}(\omega) \mapsto P\omega : \overline{A_p(G)} \rightarrow A_p(G)$ is an $\mathcal{M}_p(G)$ -module map.

We shall have need to consider the adjoint action on the dual spaces. That $CV_p(G)$ is a $\mathcal{M}_p(G)$ -module seems also to be shown by Herz [16].

Corollary 2.7. *The algebras $CV_p(G)$ and $PM_p(G)$ are dual Banach $\mathcal{M}_p(G)$ -modules:*

$$\langle \varphi \cdot T, \omega \rangle = \langle T, \varphi \cdot \omega \rangle.$$

If $\varphi \in \mathcal{M}_p(G)$ and $h \in CV_p^1(G)$, then $\varphi h \in CV_p^1(G)$ and

$$\varphi \cdot \lambda_p(h) = \lambda_p(\varphi h).$$

Proof. The first statement being an evident adjoint module operation, we are left only to inspect the second statement. However, if $\xi \otimes f$ is an elementary tensor in $N_c^p(G)$, i.e. in $N^p(K)$ for some K , then arguments just as in the proof of Theorem 2.2 (ii) provide that

$$\begin{aligned} \langle \varphi \cdot \lambda_p(h), \xi \otimes f \rangle &= \langle \lambda_p(h), \varphi \cdot (\xi \otimes f) \rangle \\ &= \int_{KK^{-1}} h(s) P(\varphi \cdot (\xi \otimes f))(s) ds \\ &= \int_{KK^{-1}} \varphi(s) h(s) \int_K \xi(t) f(s^{-1}t) dt ds \\ &= \langle \xi, (\varphi h) * f \rangle \end{aligned}$$

where the last interchange of integrals is justified as

$$(s, t) \mapsto |\varphi(s)h(s)\xi(t)f(s^{-1}t)|$$

is integrable thanks to Hölder's inequality and Tonelli's theorem. If $f \in C_c(G)$ then

$$\begin{aligned} \|(\varphi h) * f\|_p &= \sup\{|\langle \varphi \cdot \lambda_p(h), \xi \otimes f \rangle| : \xi \in L^p(K), K \in \mathcal{K}(G), \|\xi\|_p \leq 1\} \\ &\leq \|\varphi \cdot \lambda_p(h)\| \|f\|_p. \end{aligned}$$

Then taking supremum over such f with $\|f\|_p \leq 1$, reveals that $\varphi h \in CV_p^1(G)$, and further gives the desired formula. \square

We note that $A_p(G) \subseteq \mathcal{M}_p(G)$, is an ideal within the latter space, and the inclusion is a contractive map. Indeed if $a = \sum_{n=1}^{\infty} \langle \xi_n, \lambda_p(\cdot) f_n \rangle$ in $A_p(G)$, with $\sum_{n=1}^{\infty} \|\xi_n\|_{p'} \|f_n\|_p < \infty$ and each $\|\xi_n\|_{p'} \|f_n\|_p > 0$, let $\xi'_n = \|\xi_n\|_{p'}^{-1/p} \|f_n\|_p^{1/p'} \xi_n$ and $f'_n = \|\xi_n\|_{p'}^{1/p} \|f_n\|_p^{-1/p'} f_n$. Then $\beta(s) = (\lambda_{p'}(s^{-1}) \xi'_n)_{n=1}^{\infty}$ in $\ell^{p'}(\mathbb{N}, L^{p'}(G))$ and $\alpha(t) = (\lambda_p(t^{-1}) f'_n)_{n=1}^{\infty}$ in $\ell^p(\mathbb{N}, L^p(G))$ show that $a \in \mathcal{M}_p(G)$, with $\|a\|_{\mathcal{M}_p} \leq \|a\|_{A_p(G)}$.

Hence we get the following crucial result which will help us in the proof of Theorem 1.1.

Corollary 2.8. *Let $T \in CV_p(G)$ and $a \in A_p(G)$. Then $a \cdot T \in PM_p(G)$.*

Proof. By a norm estimate, we may assume that $a \in A_{p,c}(G)$. Furthermore, Theorem 2.2 (i) provides that T is a weak* limit of $\lambda_p(h_i)$ for some h_i in $CV_p^1(G)$. Then the prior corollary provides that each $a \cdot \lambda_p(h_i) = \lambda_p(ah_i)$, where $ah_i \in L^1_{\text{loc}}(G)$ with compact support, hence $ah_i \in L^1(G)$, so $\lambda_p(ah_i) \in PM_p(G)$. Thus $a \cdot T = \lim_i a \cdot \lambda_p(h_i) = \lim_i \lambda_p(ah_i) \in PM_p(G)$. \square

Remark 2.9. We say that G is *p-weakly amenable* whenever $A_p(G)$ – if we like $A_{p,c}(G)$ – admits a net (φ_i) of elements which is bounded in $\mathcal{M}_p(G)$ and tends to 1 uniformly on compact sets. For $p = 2$, this is the property of weak amenability of de Cannière and Haagerup [7]. Remarks in the proof of Theorem 1.1, in Section 3 below, show that the assumption of 2-weak amenability is sufficient to obtain general p -weak amenability.

Suppose that G is p -weakly amenable. Using the density of $N_c^p(G)$ in $N^p(G)$, it is easy to conclude that $\varphi_i \cdot \omega$ tends in norm to ω , for any ω in $N^p(G)$. Hence, by the last corollary, any T in $CV_p(G)$ is approximated weak* by the (bounded) net of elements $\varphi_i \cdot T$ from $PM_p(G)$. Hence we have proved Theorem 1.1 for such groups. For amenable groups, this is the proof of Herz [15]. For weakly amenable groups, this proof is hinted at by Cowling [3].

3 Proof of the approximation result, Theorem 1.1

We recall that $\mathcal{M}_2(G)$ is a dual space. We identify $L^1(G)$ with its evident image in $\mathcal{M}_2(G)^*$, acting by integration, and let $Q(G) = \overline{L^1(G)}^{\mathcal{M}_2(G)^*}$, the closure of $L^1(G)$ in $\mathcal{M}_2(G)^*$. De Cannière and Haagerup [7] show that $\mathcal{M}_2(G) \cong Q(G)^*$, and thus we define a weak*-topology on $\mathcal{M}_2(G)$.

As in Haagerup and Kraus [9], we say that G admits the *approximation property* if $1 \in \overline{A_p(G)}^{w*}$, i.e. there is a net $(\varphi_i) \subset A_2(G)$ – we may suppose, in fact that $(\varphi_i) \subset A_{2,c}(G)$ – for which $\langle 1, q \rangle = \lim_i \langle \varphi_i, q \rangle$ for q in $Q(G)$.

The following claim is implicit in [9] (see notes prior to Proposition 1.3, there), but the proof offered seems incomplete since it is not known if translations on $\mathcal{M}_2(G)$ are continuous in the translating variable.

Lemma 3.1. *If $f \in L^1(G)$ and $\varphi \in \mathcal{M}_2(G)$, then $f * \varphi \in \mathcal{M}_2(G)$.*

Proof. Since we have a contractive embedding $\mathcal{M}_2(G) \hookrightarrow L^\infty(G)$, we have that for g in $L^1(G)$ that $\|g\|_Q \leq \|g\|_1$. Further $\mathcal{M}_2(G)$ is closed under left translations, and translation operators are isometries, so $Q(G)$ is a homogeneous space for left translation, and thus admits left convolution by elements of $L^1(G)$. Also, $\mathcal{M}_2(G)$ consists of weakly almost periodic functions (see the observation of Xu [24]) and hence of uniformly continuous functions, so $f * \varphi$ makes sense as a uniformly continuous function. Now we have

$$\begin{aligned} & \sup \left\{ \left| \int_G f * \varphi(s)g(s) ds \right| : g \in L^1(G), \|g\|_Q \leq 1 \right\} \\ &= \sup \left\{ \left| \int_G \varphi(s)\tilde{f} * g(s) ds \right| : g \in L^1(G), \|g\|_Q \leq 1 \right\} \\ &\leq \|f\|_1 \sup \left\{ \left| \int_G \varphi(s)g(s) ds \right| : g \in L^1(G), \|g\|_Q \leq 1 \right\} = \|f\|_1 \|\varphi\|_{\mathcal{M}_2} \end{aligned}$$

where $\tilde{f}(t) = \Delta(t^{-1})f(t^{-1})$. Hence $f * \varphi \in \mathcal{M}_2(G)$. □

The fact that Hilbert spaces, which are exactly the $[\text{SQ}_2]$ -spaces, are also $[\text{SQ}_p]$ spaces (see [17]) gives a contractive embedding $\mathcal{M}_2(G) \hookrightarrow \mathcal{M}_p(G)$.

The following construction is modelled after Proposition 1.3 of [9].

Lemma 3.2. *Fix a in $A_{p,c}(G)$ with $a \geq 0$ and $\int_G a = 1$. For T in $CV_p(G)$ and ω in $N^p(G)$ let*

$$q_{T,\omega} \in \mathcal{M}_2(G)^* \text{ be given by } q_{T,\omega}(\varphi) = \langle T, (a * \varphi) \cdot \omega \rangle.$$

Then $q_{T,\omega} \in Q(G)$.

Proof. It is evident that $\|q_{T,\omega}\|_{\mathcal{M}_2^*} \leq \|T\| \|\omega\|_{N^p(G)}$. Hence by norm approximation, it suffices to show that if $\omega \in N_c^p(G)$, say $\omega \in N^p(K)$ for some K in $\mathcal{K}(G)$, then $q_{T,\omega}(\varphi) = \int_G \varphi g$ for some g in $L^1(G)$. We may further assume K is so large that $a \in A_p(K)$. With these assumptions, let

$S = \text{supp}(a)^{-1}\text{supp}(P\omega)$, and we have

$$\begin{aligned} (a * \varphi)(s)P\omega(s) &= \int_G a(t)\varphi(t^{-1}s)P\omega(s) dt \\ &= \int_G a(t)1_S(t^{-1}s)\varphi(t^{-1}s)P\omega(s) dt \\ &= \int_G 1_S(t^{-1})\varphi(t^{-1})a_t(s)P\omega(s) dt. \end{aligned}$$

Let $L = KK^{-1}S^{-1} \cup KK^{-1}$ in $\mathcal{K}(G)$, which contains the support of each $a_t P\omega$. We note that $t \mapsto a_t : S^{-1} \rightarrow A_p(L)$ is continuous. Indeed, on elementary tensors $\xi \otimes f$ in $N^p(K)$, we have that $(\xi * \check{f})_t = \xi * (\lambda_p(t)f)^\vee$ and left translation is continuous on elements of $L^p(G)$. Hence we realize

$$(a * \varphi)P\omega = \int_G 1_S(t^{-1})\varphi(t^{-1})[a_t P\omega] dt$$

as a Bochner integral in $A_p(L)$, i.e. respecting the norm $\|\cdot\|_{A_p(L)}$. Then let

$$g(t) = 1_S(t)\Delta(t^{-1})\langle T, a_{t^{-1}} P\omega \rangle$$

which defines an element of $L^1(G)$. We may thus use the dual pairing (2.2) to compute

$$\begin{aligned} \int_G \varphi g &= \int_G 1_S(t^{-1})\varphi(t^{-1})\langle T, a_t P\omega \rangle dt \\ &= \left\langle T, \int_G 1_S(t^{-1})\varphi(t^{-1})[a_t P\omega] \right\rangle = \langle T, (a * \varphi) \cdot \omega \rangle. \end{aligned}$$

This establishes the desired claim. \square

Proof of Theorem 1.1. Let a be as in Lemma 3.2 and $(\varphi_i) \subset A_{2,c}(G)$ be a net converging weak* to 1. Note that $A_{2,c}(G) \subseteq \mathcal{M}_2(G) \subseteq \mathcal{M}_p(G)$, and hence $A_{2,c}(G)$ is a space of compactly supported elements of $\mathcal{M}_p(G)$. The argument just before Corollary 2.4 tells us that any compactly supported element of $\mathcal{M}_p(G)$ is in $A_{p,c}(G)$. Since $A_p(G)$ is a homogeneous space for left translations, we find that each $a * \varphi_i \in A_p(G)$.

Given T in $CV_p(G)$, let $q_{T,\omega}$ be as in Lemma 3.2. Then we see that

$$\begin{aligned} \langle (a * \varphi_i) \cdot T, \omega \rangle &= \langle T, (a * \varphi_i) \cdot \omega \rangle = q_{T,\omega}(\varphi_i) \\ &\xrightarrow{i} q_{T,\omega}(1) = \langle T, (a * 1) \cdot \omega \rangle = \langle T, \omega \rangle \end{aligned}$$

since $a * 1 = [\int_G a] 1 = 1$, by assumption on a . But Corollary 2.8 provides that each $(a * \varphi_i) \cdot T \in PM_p(T)$, and hence so too the weak* limit of this net, T , is also in $PM_p(G)$. \square

Hints of the Theorem 1.1 are given by Cowling [3]. Given how heavily we have exploited his results, i.e. Theorem 2.2, we suspect he knew how to prove this.

Remark 3.3. The space $\mathcal{M}_p(G)$ generally has a predual $Q_p(G)$, as was verified by T. Miao in an unpublished manuscript. Using this, An, Lee and Ruan [1] defined the *p-approximation property* (*p-AP*), which is having 1 in the weak* closure of $A_p(G)$ in $\mathcal{M}_p(G)$. As verified by Vergara [23], *p-AP* is equivalent to *p'-AP*, and if $2 \leq p \leq q < \infty$, then *p-AP* implies *q-AP*. Hence *p-AP* for $p \neq 2$ is ostensibly more general than approximation property (*2-AP*). However, it is shown in [23] that certain canonical higher rank simple Lie groups fail *p-AP* for any p – the same ones known to Lafforgue and de la Salle [18] and Haagerup and de Laat [11] – giving convincing evidence that, at least for connected groups and lattices within, that *p-AP* is equivalent to *2-AP*. No examples are known of groups admitting *p-AP* for some p , but not *2-AP*.

Nonetheless, it is simple to modify the Lemmas of this section to accommodate the assumption of *p-AP*. We refer the reader [23] for details.

4 Proof of the commutation results, Theorem 1.2

Theorem 2.2, above, provides a very useful approximation of a convoluter T in $CV_p(G)$ by a bounded net of operators $(\lambda_p(h_i))$ where each $h_i \in CV_p^p(G)$. If we were conducting approximations for $p = 2$, then we need only consider self adjoint $T = T^*$ and we need not worry about determining the structure of $\lambda_p(h_i)^*$. For $p \neq 2$ we have no such luck. We overcome this difficulty below. Furthermore, in the spirit of Theorem 2.2, we give a careful pointwise approximation of a convoluter, on a dense subspace, by convolutions with elements of $L^1(G)$.

We note that $f \mapsto \tilde{f}$, where $\tilde{f}(t) = \Delta(t^{-1})f(t^{-1})$ for (locally) a.e. s , defines an isometric involution on $L^1(G)$, and clearly defines a linear map on $L_{\text{loc}}^1(G)$ which satisfies $(h * f)^\sim = \tilde{f} * \tilde{h}$ for any convolvable pair of elements. Hence if (f_i) is a contractive approximate identity on $L^1(G)$, then so too is (\tilde{f}_i) . Furthermore, if $f \in L^1(G)$, then $\lambda_p(f)^* = \lambda_{p'}(\tilde{f})$, as is standard to check. We also observe that since $\rho_p(s)^* = \rho_{p'}(s^{-1})$ for all s in G , we have $\{T^* : T \in CV_p(G)\} = CV_{p'}(G)$.

Theorem 4.1. *Let $T \in CV_p(G)$.*

(i) Then there is a net of elements (k_i) in $CV_p^p(G)$ for which $(\lambda_p(k_i))$ is a bounded net in $CV_p(G)$ converging strong operator to T , and also (\tilde{k}_i) is a net in $CV_{p'}^{p'}(G)$ for which $(\lambda_{p'}(\tilde{k}_i))$ is bounded in $CV_{p'}(G)$ and converges to T^* .

(ii) Given g, f in $C_c(G)$ and $\varepsilon > 0$, there is k in $C_c(G)$ for which

$$\|Tf - \lambda_p(k)f\|_p < \varepsilon \text{ and } \|T^*g - \lambda_{p'}(\tilde{k})g\|_{p'} < \varepsilon.$$

Here k is dependant upon the choices of g, f and ε .

Proof. (i) As in Theorem 2.2 we let $(f_i) \subset C_c(G)$ be a contractive approximate identity for $L^1(G)$ and set $h_i = Tf_i$. We note that

$$\lambda_p(h_i) = \lambda_p(Tf_i) = T\lambda_p(f_i).$$

Let $g, f \in C_c(G)$. Since $\tilde{h}_i \in L_{\text{loc}}^1(G)$, $\tilde{h}_i * g \in L_{\text{loc}}^1(G)$ and, letting $S = \text{supp}f \cup \text{supp}g$, we have

$$\begin{aligned} \int_G \tilde{h}_i * g(s)f(s) ds &= \int_S \int_{SS^{-1}} \Delta(t^{-1})h_i(t^{-1})g(t^{-1}s)f(s) dt ds \\ &= \int_{SS^{-1}} \int_S g(s)h_i(t)f(t^{-1}s) ds dt \\ &= \langle g, \lambda_p(h_i)f \rangle = \langle g, T\lambda_p(f_i)f \rangle = \langle \lambda_{p'}(\tilde{f}_i)T^*g, f \rangle. \end{aligned}$$

Hence by density it follows that $\tilde{h}_i * g = \lambda_{p'}(\tilde{f}_i)T^*g$, and hence $\tilde{h}_i \in CV_{p'}^1(G)$ with

$$\lambda_p(\tilde{h}_i) = \lambda_{p'}(\tilde{f}_i)T^*.$$

We do not know of a means to see that $\tilde{h}_i \in CV_{p'}^{p'}(G)$. We use another convolution to overcome this difficulty.

Let $k_i = f_i * h_i$. It is clear that $k_i \in CV_p^p(G)$ with $\lambda_p(k_i) = \lambda_p(f_i)\lambda_p(h_i)$, and hence $\|\lambda_{p'}(\tilde{k}_i)\| \leq \|f_i\|_1 \|\lambda_p(h_i)\|$. Likewise $\tilde{k}_i = \tilde{h}_i * \tilde{f}_i \in CV_{p'}^1(G)$ with $\lambda_{p'}(\tilde{k}_i) = \lambda_{p'}(\tilde{h}_i)\lambda_p(\tilde{f}_i)$, with $\|\lambda_{p'}(\tilde{k}_i)\| \leq \|\lambda_{p'}(\tilde{h}_i)\| \|f_i\|_1$. Moreover, as $T^* \in CV_{p'}(G)$ we have that $T^*(\tilde{f}_i) \in CV_{p'}^{p'}(G)$ and

$$\lambda_{p'}(\tilde{k}_i) = \lambda_{p'}(\tilde{h}_i)\lambda_p(\tilde{f}_i) = \lambda_{p'}(\tilde{f}_i)T^*\lambda_p(\tilde{f}_i) = \lambda_{p'}(\tilde{f}_i * T^*\tilde{f}_i)$$

so $\tilde{k}_i = \tilde{f}_i * T^*(\tilde{f}_i) \in CV_{p'}^{p'}(G)$ too. We have that

$$\begin{aligned} \|\lambda_p(k_i)f - Tf\|_p &= \|f_i * [Tf_i * f] - Tf\|_p \\ &\leq \|f_i * [Tf_i * f] - f_i * Tf\|_p + \|f_i * Tf - Tf\|_p \\ &\leq \|f_i\|_1 \|Tf_i * f - Tf\|_p + \|f_i * Tf - Tf\|_p \end{aligned}$$

which tends to zero. By uniform boundedness of the operators $\lambda_p(k_i)$ and density of $C_c(G)$ in $L^p(G)$, we see that $\lambda_p(k_i)$ tends to T in the strong operator topology. Similarly, the net $(\lambda_{p'}(\tilde{k}_i))$ is bounded and tends to T^* in the strong operator topology.

(ii) Let $\delta > 0$. Let i be such that

$$\|Tf - k_i * f\|_p < \delta \text{ and } \|T^*g - \tilde{k}_i * g\|_{p'} < \delta.$$

Inner regularity of the Haar measure provides K in $\mathcal{K}(G)$ for which

$$\|k_i - 1_K k_i\|_p < \delta \text{ and } \|\tilde{k}_i - 1_{K^{-1}} \tilde{k}_i\|_{p'} < \delta.$$

Let $k = 1_K k_i$ which is in $L^1 \cap L^p(G)$, and note that $\tilde{k} = 1_{K^{-1}} \tilde{k}_i$. Since $l * f = \rho_p(\Delta^{-1/p} \check{f})l$ for l in $L^p(G)$, we have that

$$\begin{aligned} \|Tf - k * f\|_p &\leq \|Tf - k_i * f\|_p + \|k_i * f - k * f\|_p \\ &< \delta + \|\rho_p(\Delta^{-1/p} \check{f})\| \delta. \end{aligned}$$

Likewise we have that

$$\|T^*g - \tilde{k} * g\|_{p'} < \delta + \|\rho_{p'}(\Delta^{-1/p'} \check{g})\| \delta.$$

Hence an obvious choice of δ yields the desired inequalities. \square

We note that the sorts of global estimates found in (i), above, were required to obtain the pointwise estimates of (ii).

Remark 4.2. As noted in the introduction, there is a self-inverting isometry U on $L^p(G)$, which intertwines λ_p and ρ_p : $U\lambda_p(s) = \rho_p(s)U$ for s in G . Hence U^* intertwines $\lambda_{p'}$ and $\rho_{p'}$. We let $CV'_p(G) = \lambda_p(G)'$ so

$$CV'_p(G) = (U\rho_p(G)U)' = U\rho_p(G)'U = UCV_p(G)U.$$

If $S \in CV'_p(G)$ and $g, f \in C_c(G)$, then $U^*g, Uf \in C_c(G)$. Hence, given $\varepsilon > 0$ Theorem 4.1 (ii) provides k' in $L^1(G)$ for which

$$\|USUUf - \lambda_p(k')Uf\|_p < \varepsilon \text{ and } \|(USU)^*U^*g - \lambda_{p'}(\tilde{k}')U^*g\|_{p'} < \varepsilon$$

hence multiplying the arguments of the expressions by the respective isometries U and U^* we obtain

$$\|Sf - \rho_p(k')f\|_p < \varepsilon \text{ and } \|S^*g - \rho_{p'}(\tilde{k}')g\|_{p'} < \varepsilon.$$

The key reason we have Theorem 4.1, is that for k, k' in $L^1(G)$ we have $\lambda_p(k)\rho_p(k') = \rho_p(k')\lambda_p(k)$. We do not know how to verify directly that $\lambda_p(h)\rho_p(h') = \rho_p(h')\lambda_p(h)$ for h, h' in $CV'_p(G)$.

Proof of Theorem 1.2. Let $PM'_p(G) = U PM_p(G) U = \overline{\text{lin}}^{w*} \rho_p(G)$. We have that

$$PM_p(G)' = \lambda_p(G)' = CV'_p(G) \supseteq PM'_p(G) \text{ and } PM'_p(G)' = CV_p(G).$$

Thus it follows that

$$PM_p(G)'' = CV'_p(G)' \subseteq PM'_p(G)' = CV_p(G).$$

Hence we will be done once we show that $CV_p(G) \subseteq CV'_p(G)'$, i.e. if $T \in CV_p(G)$ and $S \in CV'_p(G)$, we wish to see that $TS = ST$.

To this end, fix g, f in $C_c(G)$. As in Theorem 4.1 (ii) and Remark 4.2, given $\varepsilon > 0$, find k, k' in $L^1(G)$ for which

$$\begin{aligned} \|Tf - \lambda_p(k)f\|_p &< \varepsilon \text{ and } \|T^*g - \lambda_{p'}(\tilde{k})g\|_{p'} < \varepsilon \\ \|Sf - \rho_p(k')f\|_p &< \varepsilon \text{ and } \|S^*g - \rho_{p'}(\tilde{k}')g\|_{p'} < \varepsilon. \end{aligned}$$

Then

$$\begin{aligned} &|\langle g, TSf \rangle - \langle g, \lambda_p(k)\rho_p(k')f \rangle| \\ &\leq |\langle T^*g, Sf - \rho_p(k')f \rangle| + |\langle T^*g - \lambda_{p'}(\tilde{k})g, \rho_p(k')f \rangle| \\ &\leq \|T\| \|g\|_{p'} \varepsilon + \varepsilon \|\rho_p(k')f\|_p \leq (\|T\| \|g\|_{p'} + \|S\| \|f\|_p + \varepsilon) \varepsilon \end{aligned}$$

and, by a similar calculation, we see that

$$|\langle g, STf \rangle - \langle g, \rho_p(k')\lambda_p(k)f \rangle| \leq (\|S\| \|g\|_{p'} + \|T\| \|g\|_p + \varepsilon) \varepsilon.$$

Since $\lambda_p(k)\rho_p(k') = \rho_p(k')\lambda_p(k)$ for any k, k' in $L^1(G)$, we can take ε to 0 and see that

$$\langle g, TSf \rangle = \langle g, STf \rangle.$$

Hence by density we see that $\langle \xi, TSf \rangle = \langle \xi, STf \rangle$ for any ξ in $L^{p'}(G)$ and f in $L^p(G)$, so $TS = ST$, as desired. \square

We obtain an “elementary” proof of Dixmier’s result [8], i.e. without use of left Hilbert algebras. The common algebra in the result below is usually denoted $VN(G)$ and called the *group von Neuman algebra*.

Corollary 4.3. *We have that*

$$CV_2(G) = CV'_2(G)' = PM_2(G) = PM'_2(G)'.$$

Proof. Here $\lambda_2(G)$ is a self-adjoint subset of $\mathcal{B}(L^2(G))$, so $VN(G) = PM_2(G)$ is self-adjoint. The bicommutant theorem of von Neumann, [22, II.3.9] for example, tells us that $PM_2(G)''$ is the σ -strong* closure of $PM_2(G)$. As a convex subspace of $\mathcal{B}(L^2(G))$ is weak*-closed if and only if it is σ -strong* closed (see [22, II.2.6]), we see that $PM_2(G)'' = PM_2(G)$. The rest follows from the theorem above. \square

Acknowledgements

The second named researcher was partially supported by an NSERC Discovery Grant.

References

- [1] G. An, J.-J. Lee and Z.-J. Ruan, *On p -approximation properties for p -operator spaces*, J. Funct. Anal. 259 (2010), no. 4, 933–974.
- [2] M. Cowling, *La syntèse des convoluteurs de L^p de certains groupes pas moyennables*, Boll. Un. Mat. Ital. A (5) 14 (1977), no. 3, 551–555.
- [3] M. Cowling, *The predual of the space of convolutors on a locally compact group*, Bull. Austral. Math. Soc. 57 (1998), no. 3, 409–414.
- [4] H. G. Dales, N. J. Laustsen, T. Oikberg and V. G. Troitsky, *Multinorms and Banach lattices*, Dissertationes Math. (Rozprawy Mat.), 524 (2017), 115 pages.
- [5] M. Daws, *p -operator spaces and Figà-Talamanca–Herz algebras*, J. Operator Theory 63 (2010), no. 1, 47–83.
- [6] M. Daws and N. Spronk, *The approximation property implies that convolvers are pseudo-measures*, unpublished, see [arXiv:1308.1073v3](https://arxiv.org/abs/1308.1073v3) [math.FA].
- [7] J. De Cannière and U. Haagerup, *Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups*, Amer. J. Math. 107 (1985), no. 2, 455–500.
- [8] J. Dixmier, *von Neumann algebras*, North-Holland Mathematical Library, 27. North-Holland Publishing Co., Amsterdam-New York, 1981.
- [9] U. Haagerup and J. Kraus, *Approximation properties for group C^* -algebras and group von Neumann algebras*, Trans. Amer. Math. Soc. 344 (1994), no. 2, 667–699.
- [10] U. Haagerup, S. Knudby and T. De Laat, *A complete characterization of connected Lie groups with the approximation property*, Ann. Sci. Éc. Norm. Supér. (4) 49 (2016), no. 4, 927–946.

- [11] U. Haagerup and T. de Laat, *Simple Lie groups without the Approximation Property*, Duke Math. J. 162 (2013), no. 5, 925–964.
- [12] U. Haagerup and T. de Laat, *Simple Lie groups without the Approximation Property II*, Trans. Amer. Math. Soc. 368 (2016), no. 6, 3777–3809.
- [13] C. S. Herz, *Le rapport entre l’algèbre A_p d’un groupe et d’un sous-groupe*, C. R. Acad. Sci. Paris Sér. A-B 271 (1970), A244–A246.
- [14] C. S. Herz, *The theory of p -spaces with an application to convolution operators*, Trans. Amer. Math. Soc. 154 (1971), 69–82.
- [15] C. S. Herz, *Harmonic synthesis for subgroups*, Ann. Inst. Fourier (Grenoble) 23 (1973), no. 3, 91–123.
- [16] C. S. Herz, *Une généralisation de la notion de transformée de Fourier-Stieltjes*, Ann. Inst. Fourier (Grenoble) 24 (1974), no. 3, 145–157.
- [17] S. Kwapien, *On operators factorizable through L_p space*, Actes du Colloque d’Analyse Fonctionnelle de Bordeaux (Univ. de Bordeaux, 1971), pp. 215–225, Bull. Soc. Math. France, Mém. No. 3–32, Soc. Math. France, Paris, 1972.
- [18] V. Lafforgue and M. de la Salle, *Noncommutative L^p -spaces without the completely bounded approximation property*, Duke Math. J. 160 (2011), no. 1, 71–116.
- [19] H. Leptin, *Sur l’algèbre de Fourier d’un groupe localement compact*, C. R. Acad. Sci. Paris sér. A-B 266 (1968), A1180–A1182.
- [20] S. Öztop and N. Spronk, *p -operator space structure on Feichtinger–Figà-Talamanca–Herz Segal algebras*, J. Operator Theory 74 (2015), no. 1, 45–74.
- [21] H. L. Pham, *A short self-contained proof of the commutation theorem*, Studia Math. 237 (2017), no. 2, 195–200.
- [22] M. Takesaki, *Theory of operator algebras. I*, reprint of the first (1979) edition, Encyclopaedia of Mathematical Sciences, 124, Springer-Verlag, Berlin, 2002.

- [23] I. Vergara, *The p -approximation property for simple Lie groups with finite center*, preprint, see [arXiv1609.07632](https://arxiv.org/abs/1609.07632).
- [24] G. Xu, *Amenability and uniformly continuous functionals on the algebras $A_p(G)$ for discrete groups*, Proc. Amer. Math. Soc. 123 (1995), no 11, 3425–3429.