

Central Lancashire Online Knowledge (CLoK)

Title	Effects of fluoxetine on functional outcomes after acute stroke (FOCUS): a pragmatic, double-blind, randomised, controlled trial
Type	Article
URL	https://clock.uclan.ac.uk/24808/
DOI	https://doi.org/10.1016/S0140-6736(18)32823-X
Date	2019
Citation	Hackett, Maree, Dennis, Martin, Forbes, John, Graham, Catriona, Hankey, Graeme, House, Allan, Lewis, Stephanie, Lundström, Erik, Sandercock, Peter et al (2019) Effects of fluoxetine on functional outcomes after acute stroke (FOCUS): a pragmatic, double-blind, randomised, controlled trial. <i>The Lancet</i> , 393 (10168), pp. 265-274. ISSN 0140-6736
Creators	Hackett, Maree, Dennis, Martin, Forbes, John, Graham, Catriona, Hankey, Graeme, House, Allan, Lewis, Stephanie, Lundström, Erik, Sandercock, Peter, Mead, Gillian and Watkins, Caroline Leigh

It is advisable to refer to the publisher's version if you intend to cite from the work.
[https://doi.org/10.1016/S0140-6736\(18\)32823-X](https://doi.org/10.1016/S0140-6736(18)32823-X)

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Effects of fluoxetine on functional outcomes after acute stroke (FOCUS): a pragmatic, double-blind, randomised, controlled trial



FOCUS Trial Collaboration*

Summary

Background Results of small trials indicate that fluoxetine might improve functional outcomes after stroke. The FOCUS trial aimed to provide a precise estimate of these effects.

Methods FOCUS was a pragmatic, multicentre, parallel group, double-blind, randomised, placebo-controlled trial done at 103 hospitals in the UK. Patients were eligible if they were aged 18 years or older, had a clinical stroke diagnosis, were enrolled and randomly assigned between 2 days and 15 days after onset, and had focal neurological deficits. Patients were randomly allocated fluoxetine 20 mg or matching placebo orally once daily for 6 months via a web-based system by use of a minimisation algorithm. The primary outcome was functional status, measured with the modified Rankin Scale (mRS), at 6 months. Patients, carers, health-care staff, and the trial team were masked to treatment allocation. Functional status was assessed at 6 months and 12 months after randomisation. Patients were analysed according to their treatment allocation. This trial is registered with the ISRCTN registry, number ISRCTN83290762.

Findings Between Sept 10, 2012, and March 31, 2017, 3127 patients were recruited. 1564 patients were allocated fluoxetine and 1563 allocated placebo. mRS data at 6 months were available for 1553 (99·3%) patients in each treatment group. The distribution across mRS categories at 6 months was similar in the fluoxetine and placebo groups (common odds ratio adjusted for minimisation variables 0·951 [95% CI 0·839–1·079]; $p=0\cdot439$). Patients allocated fluoxetine were less likely than those allocated placebo to develop new depression by 6 months (210 [13·43%] patients vs 269 [17·21%]; difference 3·78% [95% CI 1·26–6·30]; $p=0\cdot0033$), but they had more bone fractures (45 [2·88%] vs 23 [1·47%]; difference 1·41% [95% CI 0·38–2·43]; $p=0\cdot0070$). There were no significant differences in any other event at 6 or 12 months.

Interpretation Fluoxetine 20 mg given daily for 6 months after acute stroke does not seem to improve functional outcomes. Although the treatment reduced the occurrence of depression, it increased the frequency of bone fractures. These results do not support the routine use of fluoxetine either for the prevention of post-stroke depression or to promote recovery of function.

Funding UK Stroke Association and NIHR Health Technology Assessment Programme.

Copyright © 2018 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

Introduction

Each year, stroke affects around 9 million people worldwide for the first time and results in long-term disability for around 6·5 million people.¹ Fluoxetine, a selective serotonin reuptake inhibitor (SSRI), is used to treat depression and emotional lability after stroke. Many clinical and preclinical studies have suggested that SSRIs might improve outcomes after stroke through a range of mechanisms, which include enhancing neuroplasticity and promoting neurogenesis. In 2011, the results of the FLAME (FLuoxetine for motor recovery After acute ischaemic stroke) trial indicated that fluoxetine enhanced motor recovery.² In this double-blind, placebo-controlled, multicentre trial, 118 patients with ischaemic stroke and unilateral motor weakness, and a median National Institutes of Health Stroke Scale (NIHSS) score of 13, were randomly allocated between 5 and 10 days after stroke onset to receive fluoxetine 20 mg daily or placebo for

3 months. At day 90, the improvement from baseline in the Fugl-Meyer motor score was significantly greater in the fluoxetine group than in the placebo group. Additionally, the proportion of patients who were independent in daily living (with a modified Rankin Scale [mRS] score of 0–2) was significantly higher in the fluoxetine group than in the placebo group (26% vs 9%, $p=0\cdot015$). More participants were free from depression at 3 months in the fluoxetine group than in the placebo group (93% vs 71%; $p=0\cdot002$). A subsequent Cochrane systematic review³ of SSRIs for stroke recovery identified 52 randomised controlled trials of SSRIs versus controls (in 4060 patients), but no others tested the effect of fluoxetine on functional outcomes measured with the mRS. The findings of the Cochrane review suggested that SSRIs might reduce post-stroke disability, although this estimate was based on a meta-analysis done across various measures of function and greater effects were seen if studies with increased risk of



Published Online

December 5, 2018

[http://dx.doi.org/10.1016/S0140-6736\(18\)32823-X](http://dx.doi.org/10.1016/S0140-6736(18)32823-X)

See Online/Comment

[http://dx.doi.org/10.1016/S0140-6736\(18\)32983-0](http://dx.doi.org/10.1016/S0140-6736(18)32983-0)

*Members of the writing group are listed at the end of the Article; all members of the FOCUS Trial Collaboration are listed in the appendix

Correspondence to:

Prof Martin Dennis, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh EH16 4SB, UK
martin.dennis@ed.ac.uk

See Online for appendix

Research in context

Evidence before this study

We searched the literature in July, 2018, using the same search strategy as that of a 2012 Cochrane review. In addition to the FLAME (Fluoxetine for motor recovery After acute ischaemic stroke) trial we identified three other small, randomised, placebo-controlled trials of fluoxetine, which enrolled patients who did not have depression at recruitment and which reported the modified Rankin Scale (mRS) during follow-up. These three trials recruited a total of 154 patients and reported improvements in the mRS in those allocated fluoxetine, but two trials (n=122) did not publish their mRS data in a format that would facilitate a meta-analysis. The FLAME trial indicated that fluoxetine, when given to patients with a recent ischaemic stroke, a motor deficit, and a median National Institutes of Health Stroke Scale (NIHSS) of 13, improved recovery in motor function as measured by the Fugl-Meyer motor score at about 3 months. In a published post-hoc analysis, the proportion of patients who were independent in daily living (mRS 0–2) was significantly higher in the fluoxetine group than in the placebo

group (26% vs 9%, $p=0.015$). However, an ordinal analysis of the mRS data did not show a significant difference between groups (common odds ratio 1.501 [95% CI 0.757–2.974]; $p=0.2446$).

Added value of this study

The results of the Fluoxetine Or Control Under Supervision (FOCUS) trial suggest that fluoxetine 20 mg given orally daily for 6 months after acute stroke does not improve functional outcomes. Although the treatment might lead to a reduction in the occurrence of depression, it also seems to increase the frequency of bone fractures. These results do not support the routine use of fluoxetine either for prevention of post-stroke depression or to promote recovery of function.

Implications of all the available evidence

Ongoing trials might be able to confirm the external generalisability of these findings to different populations, and a planned individual patient data meta-analysis could clarify whether any subgroups might benefit from fluoxetine and provide more precise estimates of any harms.

bias were retained and patients with depression were included. Although promising, data from the FLAME trial and the Cochrane review were not sufficiently compelling to alter stroke treatment guidelines or to alleviate concerns that any possible benefits might be offset by serious adverse reactions.⁴

The primary aim of the Fluoxetine Or Control Under Supervision (FOCUS) trial was to ascertain whether patients with a clinical stroke diagnosis would have improved functional outcomes with a 6-month course of fluoxetine compared with placebo. Important secondary aims were to identify any other benefits or harms and to assess whether any benefits persisted from the end of the treatment period to 12 months after stroke.

Methods

Study design and patients

FOCUS was a pragmatic, multicentre, parallel group, double-blind, randomised, placebo-controlled trial done at 103 hospitals in the UK. The protocol and statistical analysis plan were published before completion of follow-up.^{5,6}

Patients were eligible if they were aged 18 years or older; had a clinical diagnosis of acute stroke with brain imaging compatible with intracerebral haemorrhage or ischaemic stroke (including a normal brain scan); were randomly assigned between 2 days and 15 days after stroke onset; and had a persisting focal neurological deficit at the time of randomisation that was severe enough to warrant 6 months of treatment from the patient's or carer's perspective.

Patients were excluded if they had subarachnoid haemorrhage except where secondary to a primary intracerebral haemorrhage; they were unlikely to be

available for follow-up for the following 12 months; they were unable to speak English and had no close family member available to help with follow-up; they had another life-threatening illness (eg, advanced cancer) that would make 12-month survival unlikely; they had a history of epileptic seizures; they had a history of allergy to fluoxetine; they had contraindications to fluoxetine, including hepatic impairment (alanine aminotransferase more than three times the upper normal limit) or renal impairment (creatinine $>180 \mu\text{mol/L}$); they were pregnant or breastfeeding, or women of childbearing age not taking contraception; they had a previous drug overdose or attempted suicide; they were already enrolled into a controlled trial of an investigational medicinal product; they had current or recent (within the last month) depression treated with an SSRI; or they were taking or had, in the past 5 weeks, taken medications that have a potentially serious interaction with fluoxetine.

Patients (or their carers or relatives if patients had mental incapacity) provided written informed consent.

We monitored the quality and integrity of the accumulating clinical data according to a protocol agreed with the study sponsors (the Academic and Clinical Central Office for Research and Development [ACCORD] representing the University of Edinburgh and NHS Lothian), which involved central statistical monitoring, supplemented by onsite monitoring and detailed source data verification in the coordinating centre and triggered visits when patterns in the data at a centre seemed anomalous. All FOCUS monitoring procedures were compliant with requirements of the study sponsors, the ethics committee and regulatory agencies, and they met all appropriate regulatory and good clinical practice requirements. All baseline data, inpatient data, and

6-month and 12-month outcome data were subject to verification checks built into the randomisation and data management system.

During recruitment, interim analyses of baseline and follow-up data were supplied, in strict confidence at least once every year, to the chairman of the data monitoring committee. In light of these analyses, the data monitoring committee advised the chairman of the trial steering committee whether, in their view, the randomised comparisons provided “proof beyond reasonable doubt” that for all, or some, patients the treatment was clearly indicated or contraindicated, and evidence that might reasonably be expected to materially influence future patient management.

The protocol was approved by the Scotland A Multicentre Research Ethics Committee (Dec 21, 2011). The study was jointly sponsored by the University of Edinburgh and NHS Lothian. The full protocol is available in the appendix.

Randomisation and masking

Patients were randomly assigned in a 1:1 ratio to receive fluoxetine or placebo, by use of a centralised randomisation system. The clinician entered the patient's baseline data into a secure web-based randomisation system hosted by the University of Edinburgh. After the data were checked for completeness and consistency, the system generated a unique study identification number and a treatment pack number, which corresponded to either fluoxetine or placebo. A minimisation algorithm⁷ was used to achieve optimum balance between treatment groups for the following factors: delay since stroke onset (2–8 days vs 9–15 days), computer-generated prediction of 6-month outcome (probability of mRS⁸ 0 to 2 was ≤ 0.15 vs > 0.15 based on the six simple variable [SSV] model⁹), and presence of a motor deficit or aphasia (according to the NIHSS).¹⁰ The SSV model includes the patient's age; whether the patient is independent in activities of daily living before the stroke; whether they are living alone before the stroke; whether they are able to lift both arms off the bed; whether they are able to walk unassisted; whether they are able to talk, and whether they are not confused.⁹ The randomising clinicians in each centre had received training and certification in the application of the NIHSS. The system also incorporated an element of randomisation over and above the minimisation algorithm, so that it allocated patients to the treatment group that minimised the difference between groups with a probability of 0.8 rather than 1.0.⁷

Patients, their families, and the health-care team including the pharmacist, staff in the coordinating centre, and anyone involved in outcome assessments were all masked to treatment allocation by use of a placebo capsule that was visually identical to the fluoxetine capsules even when broken open. An emergency unblinding system was available but was designed so that

those in the coordinating centre and those doing follow-up remained masked to treatment allocation.

Procedures

Fluoxetine 20 mg or placebo were administered to patients orally once daily for 6 months. The study medication (active and placebo) was manufactured by Unichem (Goa, India), imported by Niche Generics Ltd (Hitchin, UK), purchased from Discovery Pharmaceuticals Ltd (Castle Donington, UK), and quality assured, packaged, labelled, and distributed by Sharp Clinical Services. Patients were supplied with 186 capsules and were prescribed the study medication (20 mg capsules of fluoxetine or placebo capsule) to be taken daily. If a patient was unable to swallow capsules and had an enteral feeding tube in place, the capsules were broken open and the contents put down the tube according to accepted methods.¹¹ We measured adherence to the study medication by recording the date of the first and last dose taken, the number of missed doses while in hospital, capsule counts when unused capsules were returned, and estimated adherence at the 6-month follow-up. We recorded the reasons for stopping the study medication early. Our primary measure of adherence was the best estimate of the interval between the first and last dose based on all the information available. Therefore, for a particular patient a capsule count might lead us to modify the estimate of the timing of the last dose.

Outcomes

The primary outcome was functional status, measured with the mRS, at the 6-month follow-up. We used the simplified mRS questionnaire (smRSq) delivered by post.^{8,12,13} Among those without a complete postal questionnaire, a telephone interview was done for any further clarification, for completion of missing items, or for the whole questionnaire. Those doing telephone assessments (chief investigators or other staff at the coordinating centre) were trained in their use.

Secondary outcomes were survival at 6 and 12 months, functional status at 12 months (mRS), and health status with the Stroke Impact Scale (SIS; for each of nine domains on which the patient scores 0–100).^{14–18} Arm, hand, leg, and foot strength; hand function; mobility; communication and understanding; memory and thinking; mood and emotions; daily activities; and participation in work, leisure, and social activities were assessed by a Likert scale. Overall rating of recovery was assessed on a visual analogue scale. Mood was assessed with the Mental Health Inventory (MHI-5).^{19,20} Fatigue was measured on the Vitality subscale of SF36.^{21,22} Health-related quality of life was measured with the EuroQoL-5 Dimensions-5 Levels (EQ5D-5L) to generate utilities.²³ The following adverse events and safety outcomes were systematically recorded: recurrent stroke including ischaemic and haemorrhagic strokes, acute coronary syndromes, epileptic seizures, hyponatraemia (< 125 mmol/L), upper

gastrointestinal bleeding, other major bleeding (lower gastrointestinal, extracranial, subdural, extradural, and subarachnoid), poorly controlled diabetes including hyperglycaemia (>22 mmol/L) and symptomatic hypoglycaemia, falls resulting in injury, bone fractures, new depression (including a diagnosis made by their treating clinician and initiation of a new antidepressant prescription), and self-harm.

The recruiting hospitals monitored adherence, identified adverse events in hospital, and completed the follow-up form at hospital discharge or death in hospital. National coordinating centre staff followed up patients at 6 months and 12 months to measure the primary and secondary outcomes. Data on adverse events and medications were also collected from patients' general practitioners at 6 months and 12 months.

Our protocol stipulated that if patients developed depression that a clinician wished to treat with an antidepressant during the treatment period, then the clinician should continue the study medication and avoid

the use of an SSRI if possible, and instead use either mirtazapine, trazadone, or a tricyclic antidepressant. We monitored the use of all antidepressants during follow-up.

Statistical analysis

We aimed to recruit at least 3000 patients. We estimated that this sample size would allow us to identify a treatment effect size of fluoxetine in the FOCUS trial that we thought would be important to patients and health and social care services. This effect size would also justify a 6-month course of treatment. FOCUS had 90% power to identify an increase in the proportion of patients with good outcomes (ie, mRS of 0–2) from 39.6% to 44.7% (ie, an absolute difference of 5.1 percentage points), based on an ordinal analysis expressed as a common odds ratio (OR) of 1.23.

The unmasked trial statistician (C Graham) prepared analyses of the accumulating data, which the data monitoring committee reviewed in strict confidence at least once a year. No other members of the trial team, trial steering committee, or patients had access to these analyses. Before recruitment was completed, and without input from the unmasked trial statistician or reference to the unblinded data, the trial steering committee prepared a detailed statistical analysis plan that was then published.⁶ For all primary analyses, including our primary analyses of adverse events and safety outcomes, we retained patients in the treatment group to which they were randomly allocated irrespective of the treatment they had actually received. We did a secondary safety analysis according to the treatment patients actually received rather than what they were randomly allocated (comparing those who received some fluoxetine in the first 6 months and those who received no fluoxetine).

Inevitably, some patients withdrew from the trial and were lost to follow-up. Some did not return follow-up questionnaires or left items blank. We excluded patients who had no follow-up data from the analyses, and did sensitivity analyses to assess the effect of these exclusions on the results.

For our primary outcome we did an ordinal analysis expressing the result as a common OR and 95% CI, where a common OR in favour of placebo is less than 1.0, adjusted with logistic regression for the variables in the minimisation algorithm. We did Cox proportional hazards modelling to analyse the effect of treatment on survival up to 12 months, also adjusting for variables included in our minimisation algorithm. We compared the frequency of outcome events by calculating the differences in proportions between treatment groups with their 95% CIs and p values. We present the median scores on the SIS, MHI-5, and the Vitality subscale of the SF36, and EQ5D-5L with the IQRs and p value derived by non-parametric methods (Mann-Whitney test). For all these scales, higher values represent better outcomes.




Figure 1: Trial profile
mRS=modified Rankin Scale. *1544 inpatients with discharge form; 20 recruited as outpatients. †1536 inpatients with discharge form; 27 recruited as outpatients.

	Fluoxetine (n=1564)	Placebo (n=1563)
Sex		
Women	589 (38%)	616 (39%)
Men	975 (62%)	947 (61%)
Age		
Age ≤70 years	666 (43%)	664 (42%)
Age >70 years	898 (57%)	899 (58%)
Mean age, years	71.2 (12.4)	71.5 (12.1)
Ethnicity		
Asian	30 (2%)	31 (2%)
Black	35 (2%)	29 (2%)
Chinese	0 (0%)	1 (0%)
White	1495 (96%)	1493 (96%)
Other	4 (0%)	9 (1%)
Marital status		
Married	879 (56%)	846 (54%)
Partner	93 (6%)	91 (6%)
Divorced or separated	109 (7%)	100 (6%)
Widowed	337 (22%)	354 (23%)
Single	124 (8%)	150 (10%)
Other	22 (1%)	22 (1%)
Living arrangement		
Living with someone else	1057 (68%)	1034 (66%)
Lives alone	485 (31%)	516 (33%)
Living in an institution	10 (1%)	4 (0%)
Other	12 (1%)	9 (1%)
Employment status		
Full-time employment	287 (18%)	258 (17%)
Part-time employment	76 (5%)	70 (4%)
Retired	1122 (72%)	1134 (73%)
Unemployed or disabled	53 (3%)	60 (4%)
Other	26 (2%)	41 (3%)
Independent before stroke	1431 (92%)	1435 (92%)
Previous medical history		
Coronary heart disease	281 (18%)	300 (19%)
Ischaemic stroke or TIA	274 (18%)	294 (19%)
Diabetes	337 (22%)	303 (19%)
Hyponatraemia	19 (1%)	26 (2%)
Intracranial bleed	27 (2%)	23 (1%)
Upper gastrointestinal bleed	25 (2%)	26 (2%)
Bone fractures	241 (15%)	256 (16%)
Depression	130 (8%)	123 (8%)
Stroke diagnosis		
Non-stroke (final diagnosis)	2 (0%)	2 (0%)
Ischaemic stroke	1410 (90%)	1406 (90%)
Intracerebral haemorrhage	154 (10%)	157 (10%)
OCSF classification of ischaemic strokes ²⁴		
Total anterior circulation infarct	318 (20%)	317 (20%)
Partial anterior circulation infarct	561 (36%)	553 (35%)
Lacunar infarct	307 (20%)	283 (18%)
Posterior circulation infarct	191 (12%)	230 (15%)
Uncertain	33 (2%)	23 (2%)

(Table 1 continues in next column)

	Fluoxetine (n=1564)	Placebo (n=1563)
(Continued from previous column)		
Cause of stroke, modified TOAST classification ²⁵		
Large artery disease	278 (18%)	234 (15%)
Small vessel disease	252 (16%)	218 (14%)
Embolism from heart	377 (24%)	411 (26%)
Another cause	38 (2%)	35 (2%)
Unknown or uncertain	465 (30%)	508 (33%)
Predictive variables		
Able to walk at time of randomisation	435 (28%)	412 (26%)
Able to lift both arms off bed	924 (59%)	935 (60%)
Able to talk and not confused	1166 (75%)	1164 (74%)
Predicted 6-month outcome based on SSV		
Probability of being alive and independent	0.28 (0.07–0.63)	0.26 (0.07–0.63)
0.00 to ≤0.15	592 (38%)	591 (38%)
>0.15 to 1.00	972 (62%)	972 (62%)
Neurological deficits		
NIHSS	6 (3–11)	6 (3–11)
Presence of a motor deficit	1361 (87%)	1361 (87%)
Presence of aphasia	457 (29%)	449 (29%)
Depression at baseline		
Current diagnosis of depression (patient or proxy reported)	26 (2%)	18 (1%)
Taking a non-SSRI antidepressant	65 (4%)	77 (5%)
Current mood, PHQ-2 ²⁶		
2 yes responses	81 (5%)	60 (4%)
1 yes response	136 (9%)	130 (8%)
0 yes responses	1347 (86%)	1373 (88%)
Delay (days) since stroke onset at randomisation		
Mean delay	6.9 (3.6)	7.0 (3.6)
2–8 days	1070 (68%)	1072 (69%)
9–15 days	494 (32%)	491 (31%)
Details of enrolment		
Enrolled as a hospital inpatient (not outpatient clinic)	1544 (99%)	1536 (98%)
Patient consented	1136 (73%)	1118 (72%)
Proxy consented	428 (27%)	445 (28%)
Data are n (%), mean (SD), or median (IQR). TIA=transient ischaemic attack. OCSF=Oxfordshire Community Stroke Project. SSV=six simple variable. NIHSS=National Institutes of Health Stroke Scale. SSRI=selective serotonin reuptake inhibitor. PHQ-2=Patient Health Questionnaire 2.		

Table 1: Patient characteristics at randomisation by allocated treatment

Prespecified subgroup analyses were the effect of treatment allocation on the primary outcome subdivided by key baseline variables described in our published statistical analysis plan,⁶ including the probability of being alive and independent (0.00 to ≤0.15 vs >0.15 to 1.00); delay from stroke onset to randomisation (2–8 days vs 9–15 days), motor deficit (present or absent) or aphasia (present or absent), pathological type of stroke (ischaemic vs haemorrhagic), and age (≤70 years




Figure 2: Primary outcome of disability on the modified Rankin Scale at 6 months by treatment group
 Ordinal analysis of the modified Rankin Scale (mRS) adjusted with logistic regression for the variables included in our minimisation algorithm. 1553 patients had mRS data available in each group; 11 patients in the fluoxetine group and ten in the placebo group had missing mRS data. Common odds ratio 0.951 (95% CI 0.839–1.079), $p=0.439$; adjusted for baseline variables.

vs >70 years); ability to consent for themselves (yes or no); whether or not mood was assessable at baseline, and whether the patient was or was not depressed at baseline. Subgroup analyses were done by observing the change in log-likelihood when the interaction between the treatment and the subgroup was added into a logistic regression model. Statistical analyses were done with SAS, version 9.2.

The study is registered with the ISRCTN registry, number ISRCTN83290762.

Role of the funding source

None of the funding organisations had any role in study design, data collection, data analysis, data interpretation, or writing of this report, or the decision to publish. The corresponding author had full access to all data in the study and had final responsibility for the decision to submit for publication.

Results

Between Sept 10, 2012, and March 31, 2017, 3152 patients consented and 3127 were enrolled. 25 patients were not enrolled; 15 were identified as ineligible between obtaining consent and randomisation and in nine cases the patients, their carer or family member, or their treating clinician changed their mind about participation in the trial (figure 1). Of the 3127 patients enrolled, 1564 were allocated

fluoxetine and 1563 allocated placebo. 11 patients did not meet our eligibility criteria after randomisation: two in each group had a final diagnosis other than stroke, and seven others (three in the fluoxetine group and four in the placebo group) were identified as having exclusion criteria (eg, a history of epilepsy, self-harm, or some other contraindication to fluoxetine). Ineligible patients were retained in our intention-to-treat analyses. Baseline characteristics of the two treatment groups were well balanced (table 1) and were similar to those of unselected patients with stroke admitted to UK hospitals (appendix). 1393 (49%) of 2847 6-month follow-up assessments were obtained by postal questionnaire (693 in the fluoxetine group and 700 in the placebo group); the remainder required a telephone reminder or were completed by telephone interview (appendix). The emergency unblinding procedure was done in only three patients, all allocated fluoxetine (one at the request of a coroner, after the patient died, one for a suspected unexpected serious adverse reaction, and one because the responsible clinician felt that knowledge of the treatment would substantially alter their management of the patient).

The primary measure of adherence was the estimated duration of study medication (interval in days from first to last dose of study medication) based on all available data, including a capsule count, which was available in 398 (25%) of 1564 patients allocated fluoxetine, and 410 (26%) of 1563 allocated placebo. Patients returned a median of 32 capsules (IQR 10–135) in the fluoxetine group and 33 (11–139) in the placebo group. Our primary measure of adherence was available in 1417 (91%) patients in each group. The median duration of treatment was 185 days (IQR 149–186) in the fluoxetine group, and 183 days (136–186) in the placebo group. The median delay between randomisation and first dose was 0 days (IQR 0–1) in both treatment groups. 1519 (97%) patients in the fluoxetine group and 1494 (96%) in the placebo group received their first dose by day 2 after randomisation. The number and proportion of patients meeting our eligibility criteria and with different levels of adherence to the study medication are shown in the appendix. 143 (9%) patients in the fluoxetine group stopped the trial medication because of perceived adverse effects within the first 90 days compared with 122 (8%) in the placebo group. Around two-thirds of patients took the study medication for at least 150 days.

The primary outcome, an ordinal comparison of the distribution of patients across the mRS categories at 6 months, adjusted for variables included in the minimisation algorithm, was similar in the two groups (common OR 0.951 [95% CI 0.839–1.079]; $p=0.439$; figure 2). The unadjusted analysis provided similar results (common OR 0.961 [95% CI 0.848–1.089]; $p=0.531$). The ordinal analysis was done with the assumption of proportional odds, in the model of mRS by treatment. This assumption was found to hold in the score test for proportional odds assumption ($p=0.9947$).

Comparison of the mRS dichotomised into 0–2 vs 3–6 similarly showed no significant difference between the groups (adjusted OR 0.955 [95% CI 0.812–1.123], $p=0.576$; unadjusted OR 0.957 [0.827–1.107], $p=0.352$).

The results of our prespecified subgroup analyses are shown in the appendix. No significant interactions were observed between the prespecified subgroups and the effect of treatment on the primary outcome.

The appendix shows the effect of fluoxetine on our primary outcome in subgroups defined by the eligibility criteria and increasing degrees of adherence to the study medication; we did a series of prespecified per-protocol analyses, which sequentially excluded subgroups of patients who either did not meet our eligibility criteria or had incomplete adherence to the study medication.⁶ We did not observe greater benefit in patients with greater adherence.

Secondary outcomes at 6 months are shown in table 2 and adverse events at 6 months shown in table 3. Patients allocated fluoxetine were less likely than those allocated placebo to be diagnosed with new depression at 6 months (210 [13.43%] patients vs 269 [17.21%]; difference in proportions 3.78% [95% CI 1.26–6.30]; $p=0.0033$) and had better mood measured on MHI-5 at the 6-month follow-up (median 76 [IQR 60–88] vs 72 [56–88]; $p=0.0100$). Those allocated fluoxetine had an increased risk of bone fractures compared with those allocated placebo (45 [2.88%] patients vs 23 [1.47%]; difference in proportions 1.41% [95% CI 0.38–2.43]; $p=0.0070$). There were no significant differences in any other secondary outcomes at 6 months, including any of the nine domains of the SIS, the Vitality subscale of SF36, and EQ5D-5L (table 2) or other recorded adverse reactions (table 3). The appendix shows the progress through the trial for patients who received any fluoxetine and those who received no fluoxetine by 6 months. Adverse events and safety outcomes in patients who received any fluoxetine within the first 6 months and those who received no fluoxetine, irrespective of the group to which they were allocated, are shown in the appendix.

The appendix shows secondary outcomes at 12 months. The difference in MHI-5 at 6 months was not sustained at 12 months, and the difference between the two treatment groups in the cumulative number of patients diagnosed with new depression over 12 months was no longer significant. More patients had been started on antidepressants in the placebo group than in the fluoxetine group, but some were started on antidepressants for indications other than depression. There were no significant differences between treatment groups in any other secondary outcomes at 12 months, including survival (hazard ratio 0.929 [95% CI 0.756–1.141]; $p=0.4819$; appendix).

We assessed the effect of treatment among the subgroup with motor deficit at baseline ($n=2702$) but found no evidence of an effect on the mRS (common OR 0.919 [95% CI 0.803–1.051]; $p=0.2172$) or on motor score based

	Fluoxetine	Placebo	p value*
SIS			
Strength	56.25 (31.25–81.25)	62.50 (37.50–81.25)	0.7008
Hand ability	45.00 (0.00–90.00)	50.00 (0.00–90.00)	0.4824
Mobility	63.89 (36.11–86.11)	63.89 (33.33–88.89)	0.5486
Motor†	54.86 (27.31–83.33)	56.78 (28.75–82.64)	0.5125
Daily activities	62.50 (37.50–90.00)	65.00 (35.00–90.00)	0.6235
Physical function‡	56.77 (30.38–84.31)	58.82 (30.56–84.10)	0.5154
Memory	82.14 (57.14–96.43)	82.14 (57.14–96.43)	0.3070
Communication	89.29 (67.86–100)	92.86 (71.43–100.0)	0.1919
Emotion	75.00 (58.33–88.89)	75.00 (58.33–88.89)	0.4687
Participation	62.50 (37.50–87.50)	65.63 (40.63–87.50)	0.2595
Recovery (VAS)	60.00 (40.00–80.00)	60.00 (40.00–80.00)	0.9820
Vitality	56.25 (37.50–75.00)	56.25 (43.75–75.00)	0.6726
MHI-5	76.00 (60.00–88.00)	72.00 (56.00–88.00)	0.0100
EQ5D-5L	0.56 (0.21–0.74)	0.56 (0.19–0.75)	0.5866

Data were only available for those who survived and who completed sufficient questions to derive a score. The number of patients with missing scores was similar in the two treatment groups. The number of survivors with missing data across both treatment groups varied from 16 for EQ5D-5L and Mobility to 71 for Emotion. Data are median (IQR). EQ5D-5L=EuroQoL-5 Dimensions-5 Levels (where 1 indicates perfect health, and <0=worse than death). VAS=visual analogue scale. MHI-5=Mental Health Inventory 5 (where higher scores are better). SIS=Stroke Impact Scale (where higher scores are better). *Mann-Whitney test. †Mean of the Strength, Hand ability, and Mobility domains. ‡Mean of the Strength, Hand ability, Mobility, and Daily activities domains.

Table 2: Secondary outcomes at 6 months by allocated treatment

	Fluoxetine (n=1564)	Placebo (n=1563)	Difference (95% CI)	p value
Any stroke	56 (3.58%)	64 (4.09%)	-0.51% (-1.90 to 0.80)	0.4543
All thrombotic events	78 (4.99%)	92 (5.89%)	-0.90% (-2.49 to 0.69)	0.2677
Ischaemic stroke	43 (2.75%)	45 (2.88%)	-0.13% (-1.30 to 1.00)	0.8264
Other thrombotic events	20 (1.28%)	27 (1.73%)	-0.45% (-1.30 to 0.40)	0.3025
Acute coronary events	15 (0.96%)	23 (1.47%)	-0.51% (-1.28 to 0.26)	0.1910
All bleeding events	41 (2.62%)	38 (2.43%)	0.19% (-0.91 to 1.29)	0.7346
Haemorrhagic stroke	7 (0.45%)	9 (0.58%)	-0.13% (-0.60 to 0.37)	0.6153
Upper gastrointestinal bleed	21 (1.34%)	16 (1.02%)	0.32% (-0.44 to 1.08)	0.4094
Other major bleeds	13 (0.83%)	14 (0.90%)	-0.06% (-0.71 to 0.58)	0.8454
Epileptic seizures	58 (3.71%)	40 (2.56%)	1.15% (-0.07 to 2.37)	0.0651
Fall with injury	120 (7.67%)	94 (6.01%)	1.66% (-0.11 to 3.43)	0.0663
Fractured bone	45 (2.88%)	23 (1.47%)	1.41% (0.38 to 2.43)	0.0070
Hyponatraemia <125 mmol/L	22 (1.41%)	14 (0.90%)	0.51% (-0.24 to 1.26)	0.1805
Hyperglycaemia	23 (1.47%)	16 (1.02%)	0.45% (-0.33 to 1.22)	0.2602
Symptomatic hypoglycaemia	23 (1.47%)	13 (0.83%)	0.64% (-0.11 to 1.39)	0.0940
New depression	210 (13.43%)	269 (17.21%)	-3.78% (-6.30 to -1.26)	0.0033
New antidepressant	280 (17.90%)	357 (22.84%)	-4.94% (-7.76 to -2.12)	0.0006
Attempted or actual suicide	3 (0.19%)	2 (0.13%)	0.06% (-0.02 to 0.34)	0.6550

Data are n (%), unless otherwise stated.

Table 3: Adverse events at 6 months by treatment group

on the mean of SIS Strength, Hand, and Mobility domains (fluoxetine median 48.43 [IQR 24.98–78.84] vs placebo 52.66 [25.28–77.22]; $p=0.4714$). Additionally, in patients with aphasia at baseline and an SIS communication domain score available at 6 months ($n=894$) we found no difference in median SIS communication domain

scores (fluoxetine 64·29 [IQR 32·14–89·29] vs placebo 64·29 [35·71–89·29]; $p=0\cdot4971$).

Discussion

The results of the FOCUS trial show that fluoxetine 20 mg given daily for 6 months after an acute stroke does not significantly improve patients' functional outcome or survival at 6 and 12 months. However, fluoxetine decreased the occurrence of depression and increased bone fractures at 6 months.

The strengths of the study, supporting the internal validity of the results, are that bias was minimised by central randomisation without any prospect of foreknowledge; patients, carers, and outcome assessment were masked (with only three episodes of unmasking); there were few losses to follow-up (<1%), and pre-specified intention-to-treat analyses were done. The small difference in the numbers of patients stopping the trial medication for perceived adverse effects suggests that unmasking because of adverse effects was unlikely to have had a significant effect on our results. In any case, expectation bias would normally be expected to bias the result in favour of active treatment. Random error was also minimised by the large sample size and high rates of follow-up, which provided greater statistical power than in previous similar trials. The external validity of the results, at least for the UK stroke population, is supported by the large number of participating hospitals throughout the UK. Compared with unselected patients with stroke admitted to UK hospitals (appendix), there were few differences in the baseline characteristics of patients enrolled in the FOCUS trial.^{27,28} Patients enrolled in FOCUS had slightly more severe strokes than unselected patients (NIHSS 6 vs 4), which probably reflected inclusion criteria that required patients to have a neurological deficit persisting at the time of enrolment. Also, 60% of enrolled patients were men compared with a UK average of 50%—an unexplained but common observation in stroke trials.²⁹ Enrolled patients were slightly younger than the UK average (71 years vs 77 years), which might partly explain the male preponderance, with older women being under-represented. Many studies included in the previously published systematic review of randomised controlled trials of fluoxetine were from China, whereas non-white patients comprised less than 5% of those recruited in FOCUS. The ongoing AFFINITY trial is recruiting in Vietnam and will include a larger proportion of Asian patients.⁵

The validity of our results is also supported by the observed reduction in the occurrence of new post-stroke depression at 6 months with fluoxetine, which is consistent with its known antidepressant effects and the results of the FLAME trial. A previous systematic review of five randomised controlled trials (two of fluoxetine, two of sertraline, and one of escitalopram), including FLAME, in patients with stroke and no depression tested

whether SSRIs prevented the development of post-stroke depression.³⁰ In a pooled analysis, 23 (9·3%) of 248 patients treated with an SSRI developed post-stroke depression compared with 59 (24·4%) of 242 treated with a placebo (OR 0·37 [95% CI 0·22–0·61]; $p=0\cdot001$). The rate of depression in the placebo groups of these trials was much higher than that in FOCUS, which might have reflected the characteristics of the patients (as they tended to have had more severe strokes than those enrolled in FOCUS) or the different methods of diagnosing depression. Although this observation is consistent with our findings in terms of the direction (but not the magnitude) of treatment effect, it does not take into account the possible excess risk of adverse effects (such as bone fractures), which might offset any benefits of reducing the frequency of post-stroke depression.

The observed 1·4% absolute excess risk of bone fractures at 6 months with fluoxetine in FOCUS is also consistent with previous reports from large case-control and cohort studies.³¹ The magnitude of the increased risk in previous observational studies tended to be greater than in FOCUS, but this difference might be attributable to the inherent confounding by treatment indication in observational studies. The rates of serious adverse reactions to fluoxetine referred to in the summary of product characteristics, which we included as secondary outcomes in this trial (eg, epileptic seizures, falls, hyponatraemia, uncontrolled diabetes, and upper gastrointestinal bleeding), were higher in the fluoxetine group than in the placebo group, but the absolute differences were small and not significant. Despite concerns about the effects of fluoxetine on platelet function and interactions with antiplatelet and anticoagulant medications, we observed no effect on bleeding or thrombotic adverse events.

The main limitation of FOCUS was the moderate adherence to the trial medication, which might have led us to under-estimate any treatment effect. However, adherence measured in FOCUS was superior to that reported in routine clinical practice, and did not differ substantially between the treatment groups.³² Differences in adherence between the fluoxetine and placebo groups were more likely if reduced adherence resulted from possible adverse reactions or perceived change (or no change) in patients' conditions. We repeated the analysis of our primary outcome after sequentially excluding patients with different reasons for, and different degrees of, adherence. Such per-protocol analyses can increase the risk of bias, usually in favour of the active treatment. However, our analyses (shown in the appendix) did not show any increased benefit from fluoxetine in patients with greater adherence.

Our use of the smRSq as the primary outcome measure could be perceived as a limitation. However, the smRSq is a valid, reliable, and patient-centred measure of functional outcome, thus ensuring our results are relevant to

patients and their families.^{8,12,13} Additionally, local, face-to-face assessments of outcomes might be more prone to unmasking than those done through postal and telephone follow-up because of patients reporting adverse effects of trial medication. We used patient-reported outcomes, the Patient Health Questionnaire 2 (PHQ-2) at baseline and the smRSq, MHI-5, and SIS motor score at follow-up by postal and telephone questionnaires. Other limitations of FOCUS include the absence of a standardised psychiatric assessment at baseline or follow-up and absence of a structured neurological examination during follow-up, which were impractical to include in this large, pragmatic, multicentre trial.

We cannot definitively exclude an effect of fluoxetine on a directly measured neurological deficit—such as the Fugl-Meyer motor score, which was measured in the FLAME trial. However, we have shown that a resulting improvement in functional status measured with the mRS or SIS is unlikely.

Other trials of similar design to FOCUS, but with smaller recruitment targets, are ongoing.^{5,6} These studies should allow us to confirm the effects of fluoxetine on post-stroke depression and bone fractures, and provide more precise estimates of the benefits and harms of early fluoxetine, to guide its use in patients with stroke and perhaps other older people with comorbidities. These ongoing trials will also establish the external validity of the FOCUS trial in stroke populations with different ethnic groups and health-care backgrounds—for example, with different intensities of physical rehabilitation.

In summary, the results of the FOCUS trial show that fluoxetine 20 mg given daily for 6 months after an acute stroke did not influence patients' functional outcomes but did decrease the occurrence of depression and increase the occurrence of bone fractures. These results do not support the routine use of fluoxetine either for the prevention of post-stroke depression or to promote recovery of function. Ongoing trials and a planned individual patient data meta-analysis are needed to confirm or refute a more modest benefit, either overall or in particular subgroups, and to provide more precise estimates of any harms.

Contributors

MD was Co-Chief Investigator, participated in the steering committee, was involved in the design of the trial, and collected, verified, and analysed data and drafted this report. JF participated in the steering committee, was involved in the design of the trial, and analysed health economic data. CG participated in the steering committee, was involved in the design of the trial, wrote the first draft of the statistical analysis plan, and verified and analysed data. MH was involved in the trial design and helped conduct relevant systematic reviews. GJH was involved in the trial design and helped conduct relevant systematic reviews. AH was involved in the trial design and advised on the management of depression within the trial. SL was involved in the trial design and advised on the statistical analysis plan. EL was involved in the design of the trial. PS chaired the steering committee of the initial phase. GM was Co-Chief Investigator, participated in the steering committee, was involved in the design of the trial and data collection, and coordinated the systematic review of the randomised controlled trials. All members of the writing committee listed here have

commented on the analyses and drafts of this report and have seen and approved the final version of the report.

Writing group of the FOCUS Trial Collaboration

Martin Dennis (Chair), John Forbes, Catriona Graham, Maree Hackett, Graeme J Hankey, Allan House, Stephanie Lewis, Erik Lundström, Peter Sandercock, Gillian Mead.

Declaration of interests

We declare no competing interests.

Data sharing

The study protocol and statistical analysis plan have been published.^{5,6} A fully anonymised trial dataset with individual participant data and a data dictionary will be made available to other researchers after the publication of the full trial report in the *Health Technology Assessment* journal in 2019. Requests should first be directed to Martin Dennis (Co-Chief Investigator). Written proposals will be assessed by the FOCUS trial team and a decision made about the appropriateness of the use of data. A data sharing agreement will be put in place before any data will be shared.

Acknowledgments

The start-up phase of the FOCUS trial was funded by the UK Stroke Association (TSA 2011101) and the main phase funded by the NIHR Health Technology Assessment Programme (project number 13/04/30). The views and opinions expressed herein are those of the authors and do not necessarily reflect those of the NIHR Health Technology Assessment Programme. Recruitment and follow-up was supported by the NIHR-funded UK Stroke Research Network and the Scottish Stroke Research Network, which was supported by NHS Research Scotland (NRS).

References

- 1 GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; **388**: 1545–602.
- 2 Chollet F, Tardy J, Albuher JF, et al. Fluoxetine for motor recovery after acute ischaemic stroke (FLAME): a randomised placebo-controlled trial. *Lancet Neurol* 2011; **10**: 123–30.
- 3 Mead GE, Hsieh CF, Lee R, et al. Selective serotonin reuptake inhibitors (SSRIs) for stroke recovery. *Cochrane Database Syst Rev* 2012; **11**: CD009286.
- 4 Schambra H, O'Dell MW. Should this patient with ischemic stroke receive fluoxetine? *PM R* 2015; **7**: 1294–99.
- 5 Mead G, Hackett ML, Lundstrom E, Murray V, Hankey GJ, Dennis M. The FOCUS, AFFINITY and EFFECTS trials studying the effect(s) of fluoxetine in patients with a recent stroke: a study protocol for three multicentre randomised controlled trials. *Trials* 2015; **16**: 369.
- 6 Graham C, Lewis S, Forbes J, et al. The FOCUS, AFFINITY and EFFECTS trials studying the effect(s) of fluoxetine in patients with a recent stroke: statistical and health economic analysis plan for the trials and for the individual patient data meta-analysis. *Trials* 2017; **18**: 627.
- 7 Altman DG, Bland JM. Treatment allocation by minimisation. *BMJ* 2005; **330**: 843.
- 8 Bruno A, Shah N, Lin C, et al. Improving modified Rankin scale assessment with a simplified questionnaire. *Stroke* 2010; **41**: 1048–50.
- 9 Counsell C, Dennis M, McDowall M, Warlow C. Predicting outcome after acute stroke: development and validation of new models. *Stroke* 2002; **33**: 1041–47.
- 10 The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995; **333**: 1581–87.
- 11 White R, Bradnam V. Handbook of drug administration via enteral feeding tubes, 2nd edn. London: Pharmaceutical Press, 2010.
- 12 Bruno A, Akinwuntan AE, Lin C, et al. Simplified modified Rankin scale questionnaire: reproducibility over the telephone and validation with quality of life. *Stroke* 2011; **42**: 2276–79.
- 13 Dennis M, Mead G, Doubal F, Graham C. Determining the modified Rankin score after stroke by postal and telephone questionnaires. *Stroke* 2012; **43**: 851–53.
- 14 Duncan PW, Wallace D, Lai SM, Johnson D, Embretson S, Laster LJ. The stroke impact scale version 2.0: evaluation of reliability, validity and sensitivity to change. *Stroke* 1999; **30**: 2131–40.

For more on the NIHR Health Technology Assessment Programme see <http://www.hta.ac.uk/project>

- 15 Duncan PW, Bode R, Min Lai S, Perera S, Glycine Antagonist in Neuroprotection Americans Investigators. Rasch analysis of a new stroke specific outcome scale: the Stroke Impact Scale. *Arch Phys Med Rehabil* 2003; **84**: 950–63.
- 16 Duncan P, Reker D, Horner R, et al. Performance of a mail-administered version of a stroke specific outcome measure, the Stroke Impact Scale. *Clin Rehabil* 2002; **16**: 493–505.
- 17 Duncan P, Reker D, Kwon S, et al. Measuring stroke impact with the stroke impact scale: telephone versus mail administration in veterans with stroke. *Med Care* 2005; **43**: 507–15.
- 18 Kwon S, Duncan P, Studenski S, Perera S, Lai SM, Reker D. Measuring stroke impact with SIS: construct validity of SIS telephone administration. *Qual Life Res* 2006; **15**: 367–76.
- 19 Hoeymans N, Garssen AA, Westert GP, Verhaak PF. Measuring mental health of the Dutch population: a comparison of the GHQ-12 and the MHI-5. *Health Qual Life Outcomes* 2004; **2**: 23.
- 20 McCabe CJ, Thomas KJ, Brazier JE, Coleman P. Measuring the mental health status of a population: a comparison of the GHQ-12 and the SF-36 (MHI-5). *Br J Psychiatry* 1996; **169**: 516–21.
- 21 Mead GE, Lynch J, Greig CA, Young A, Lewis SJ, Sharpe M. Evaluation of fatigue scales in stroke. *Stroke* 2007; **38**: 2090–95.
- 22 Mead GE, Graham C, Dorman P, et al, on behalf of UK Collaborators of IST. Fatigue after stroke: baseline predictors and influence on survival. Analysis of data from UK Patients Recruited in the International Stroke Trial. *PLoS One* 2011 **6**: e16988.
- 23 Herdman M, Gudex C, Lloyd A, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res* 2011; **20**: 1727–36.
- 24 Bamford J, Sandercock P, Dennis MS, Burn J, Warlow C. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet* 1991; **337**: 1521–26.
- 25 Adams HP Jr, Bendixen BH, Kappelle LJ, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of ORG 10172 in acute stroke treatment. *Stroke* 1993; **24**: 35–41.
- 26 Whooley MA, Avins AL, Miranda J, Browner WS. Case-finding instruments for depression. Two questions are as good as many. *J Gen Intern Med* 1997; **12**: 439–45.
- 27 Bray B D, Cloud GC, James MA, et al. Weekly variation in health-care quality by day and time of admission: a nationwide, registry-based, prospective cohort study of acute stroke care. *Lancet* 2016; **388**: 170–77.
- 28 NHS National Services Scotland. Scottish Stroke Improvement Programme. 2017 report. <https://www.strokeaudit.scot.nhs.uk/Downloads/docs/2017-07-11-SCCA-Report.pdf> (accessed July 19, 2018).
- 29 Tsivgoulis G, Katsanos AH, Caso V. Under-representation of women in stroke randomized controlled trials: inadvertent selection bias leading to suboptimal conclusions. *Ther Adv Neurol Disord* 2017; **10**: 241–44.
- 30 Salter KL, Foley NC, Zhu L, Jutai JW, Teasell RW. Prevention of poststroke depression: does prophylactic pharmacotherapy work? *J Stroke Cerebrovasc Dis* 2013; **22**: 1243–51.
- 31 Wadhwa R, Kumar M, Talegaonkar S, Vohora D. Serotonin reuptake inhibitors and bone health: a review of clinical studies and plausible mechanisms. *Osteoporosis Sarcopenia* 2017; **3**: 75–81.
- 32 Sansone RA, Sansone LA. Antidepressant adherence: are patients taking their medications? *Innov Clin Neurosci* 2012; **9**: 41–46.