

Central Lancashire Online Knowledge (CLoK)

Title	Interventions for treating urinary incontinence after stroke in adults
Type	Article
URL	https://clock.uclan.ac.uk/28346/
DOI	##doi##
Date	2019
Citation	Thomas, Lois Helene orcid iconORCID: 0000-0001-5218-6546, Coupe, Jacqueline orcid iconORCID: 0000-0002-2025-5620, Cross, Lucy orcid iconORCID: 0000-0002-7272-9445, Tan, Aidan L and Watkins, Caroline Leigh orcid iconORCID: 0000-0002-9403-3772 (2019) Interventions for treating urinary incontinence after stroke in adults. Stroke, 50 (8). e226-e227. ISSN 0039-2499
Creators	Thomas, Lois Helene, Coupe, Jacqueline, Cross, Lucy, Tan, Aidan L and Watkins, Caroline Leigh

It is advisable to refer to the publisher's version if you intend to cite from the work. ##doi##

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Interventions for treating urinary incontinence after stroke in adults

Key words: stroke, urinary incontinence, rehabilitation

Corresponding author:

Professor Lois Thomas PhD, BA(Hons), RN

Professor of Health Services Research

Brook Building room 416

Faculty of Health and Wellbeing

University of Central Lancashire

PRESTON PR1 2HE

+44 (0)1772 893643

lhthomas@uclan.ac.uk

Other authors:

Jacqueline Coupe MPH, BSc (Hons)

Research Associate, Faculty of Health and Wellbeing, University of Central Lancashire, UK
(until November 2018)

Lucy D Cross BA(Hons)

Research Assistant, Faculty of Health and Wellbeing, University of Central Lancashire, UK

Aidan L Tan MPH, MBBS, GDip(Occ Med)

Department of Preventive Medicine, National University Hospital, Singapore

Caroline L Watkins PhD, BA(Hons), RN

Professor of Stroke and Older People' Care, Faculty of Health and Wellbeing, University of Central Lancashire, UK

Social media handles: @UCLanHealth; @UCLan; @UCLanResearch;
@ukstrokenursing

Background

Urinary incontinence can affect 40% to 60% of people admitted to hospital after a stroke; 25% and 15% still have problems at discharge and one year respectively.

Objectives

To assess the effects of interventions for treating urinary incontinence after stroke in adults at least one-month post-stroke.

Search methods

We searched the Cochrane Incontinence and Cochrane Stroke Specialised Registers (searched 30 October 2017 and 1 November 2017 respectively) and hand searched journals and conference proceedings.

Selection criteria

We included randomised or quasi-randomised controlled trials.

Data collection and analysis

Two review authors independently undertook data extraction, risk of bias assessment and implemented GRADE.

Main results

We included 20 trials (reporting 21 comparisons) with 1338 participants. Interventions in the included trials included behavioural interventions, specialised professional input interventions, complementary therapy interventions, pharmacotherapy interventions, and physical therapy interventions. Findings for our main comparison, **intervention versus no intervention/usual care**, are presented below.

Findings

Behavioural interventions: Low-quality evidence suggests behavioural interventions may reduce the mean number of incontinent episodes in 24 hours (mean difference (MD) -1.00 , 95% confidence interval (CI) -2.74 to 0.74 ; 1 trial; 18 participants; $P = 0.26$). Further, low-quality evidence from two trials suggests that behavioural interventions may make little or no difference to quality of life (SMD -0.99 , 95% CI -2.83 to 0.86 ; 55 participants).

Specialised professional input interventions: One trial of moderate-quality suggested structured assessment and management by continence nurse practitioners probably made little or no difference to the number of people continent three months after treatment (risk

ratio (RR) 1.28, 95% CI 0.81 to 2.02; 121 participants; equivalent to an increase from 354 to 453 per 1000, 95% CI 287 to 715).

Complementary therapy: Five trials assessed complementary therapy using traditional acupuncture, electroacupuncture and ginger-salt-partitioned moxibustion plus routine acupuncture. Low-quality evidence from five trials suggested complementary therapy may increase the number of participants continent after treatment; participants in the treatment group were three times more likely to be continent (RR 2.82, 95% CI 1.57 to 5.07; 524 participants; equivalent to an increase from 193 to 544 per 1000, 95% CI 303 to 978).

Physical therapy: Two trials reporting three comparisons suggest that physical therapy using transcutaneous electrical nerve stimulation (TENS) may reduce the mean number of incontinent episodes in 24 hours (MD -4.76, 95% CI -8.10 to -1.41; 142 participants; low-quality evidence). One trial of TENS reporting two comparisons found that the intervention probably improves overall functional ability (MD 8.97, 95% CI 1.27 to 16.68; 81 participants; moderate-quality evidence).

Conclusions

There is insufficient evidence to guide continence care of adults in the rehabilitative phase of stroke. As few trials tested the same intervention, conclusions are drawn from few, usually small, trials. Many trials were limited by poor reporting, making it impossible to judge the extent to which they were prone to bias.

More appropriately powered, multi-centre trials of promising interventions (e.g. TENS) are required to provide robust evidence.

Disclosures:

None.

Funding sources:

- Faculty of Health and Wellbeing, University of Central Lancashire, UK.
- National Institute of Health Research (NIHR), UK.

This project was supported by the National Institute for Health Research, via the Cochrane Incentive Scheme and Infrastructure funding to Cochrane Incontinence.

Reference and Copyright requirements:

a) This paper is based on a Cochrane Review published in The Cochrane Library 2019, Issue 2 (see www.thecochranelibrary.com for information). Cochrane Reviews are regularly updated as new evidence emerges and in response to feedback, and The Cochrane Library should be consulted for the most recent version of the review.

b) Thomas LH, Coupe J, Cross LD, Tan AL, Watkins CL. Interventions for treating urinary incontinence after stroke in adults. Cochrane Database of Systematic Reviews 2019, Issue 2. Art. No.: CD004462. DOI: 10.1002/14651858.CD004462.pub4.

c) <http://dx.doi.org/10.1002/14651858.CD004462>

