

Central Lancashire Online Knowledge (CLOK)

Title	Non-clinical interventions to reduce unnecessary caesarean sections: WHO recommendations
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/31213/
DOI	https://doi.org/10.2471/blt.19.236729
Date	2019
Citation	Opiyo, Newton, Kingdon, Carol, Oladapo, Olufemi T, Souza, João Paulo, Vogel, Joshua P, Bonet, Mercedes, Bucagu, Maurice, Portela, Anayda, McConville, Frances et al (2019) Non-clinical interventions to reduce unnecessary caesarean sections: WHO recommendations. Bulletin of the World Health Organization, 98 (1). pp. 66-68. ISSN 0042-9686
Creators	Opiyo, Newton, Kingdon, Carol, Oladapo, Olufemi T, Souza, João Paulo, Vogel, Joshua P, Bonet, Mercedes, Bucagu, Maurice, Portela, Anayda, McConville, Frances, Downe, Soo, Gülmezoglu, Ahmet Metin and Betrán, Ana Pilar

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.2471/blt.19.236729>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Non-clinical interventions to reduce unnecessary caesarean sections: WHO recommendations

Newton Opiyo,^a Carol Kingdon,^b Olufemi T Oladapo,^a João Paulo Souza,^a Joshua P Vogel,^a Mercedes Bonet,^a Maurice Bucagu,^c Anayda Portela,^c Frances McConville,^c Soo Downe,^b Ahmet Metin Gülmezoglu^a & Ana Pilar Betrán^a

Caesarean sections can be a lifesaving procedure for mother and baby, but rates beyond 10% of live births are not associated with reductions in maternal and newborn mortality.¹ Caesarean section rates at national level vary between around 2% in Chad, Burkina Faso, Ethiopia or Madagascar and above 50% in Brazil, Dominican Republic or Egypt.² The trend towards overuse of caesarean sections is a major concern globally, given the risks to the mother and her child associated with unnecessary caesarean birth. These risks include avoidable maternal complications such as infections, haemorrhage, complications related to use of anaesthesia or blood transfusion, and infant morbidity, for example, respiratory problems, asthma and obesity in children. Caesarean sections can also lead to added complications for the mother in subsequent pregnancies, including uterine rupture, placental implantation problems and need for hysterectomy.³ High rates of caesarean sections are also associated with substantial health-care costs, which can pose a considerable burden on health systems.

Multiple factors are driving increases in caesarean section rates. Clinical reasons for growing rates include increases in the incidence of maternal obesity, multiple pregnancies and a higher maternal age at birth. These factors alone are unlikely to explain the extent of the rise in caesarean section rates or the substantial variations among health-care providers, hospitals and regions. Studies have shown associations between caesarean section rates and non-clinical factors such as differences in health provider practices, fear of malpractice litigation and organizational, economic, social and cultural factors.⁴ A growing

proportion of caesarean sections globally are not medically indicated and could have been avoided.

To address the rising rates worldwide and prevent the harm to women and newborns resulting from overuse of this procedure, in 2018 the World Health Organization (WHO) published new recommendations on non-clinical interventions to reduce unnecessary caesarean sections.⁵ In this guideline, non-clinical interventions are defined as those interventions that are applied outside of the routine clinical interactions between a provider and pregnant woman. The interventions may target women (for instance, birth preparation classes), health-care providers (clinical practice guidelines) or health organizations (different payment systems for caesarean sections).

The recommendations (Box 1), are grouped according to the target of interventions and address major determinants of caesarean section rates. The recommendations are intended to inform the development of national and subnational policies and protocols to reduce unnecessary caesarean births in high-, middle- and low-income countries. The new recommendations should be integrated and implemented with other related WHO guidelines, such as WHO recommendations on antenatal and intrapartum care for a positive childbirth experience.

Implementation strategies

Countries should use a combination of interventions and strategies tailored to local determinants of inappropriate caesarean section use, for example the beliefs about the impact, professional norms and incentives, and economic

and organizational factors influencing caesarean sections.

During implementation, monitoring and assessing caesarean section rates and maternal and perinatal outcomes is essential. This analysis should be done in a standardized and action-oriented manner, considering the specific characteristics of the populations served. For this purpose, WHO recommends the Robson Classification system as a global standard.⁶

Several factors may hinder uptake and scale-up of these new recommendations. These factors may be related to the behaviours or preferences of women and their families, the behaviour of health-care providers, the organization of care, health service delivery or financial arrangements. The barriers were identified from multiple sources: (i) qualitative reviews undertaken for this guideline;^{7–9} (ii) case studies and systematic reviews exploring factors affecting the implementation of interventions to reduce caesarean section rates;^{10,11} and (iii) Cochrane overviews of reviews of health-system implementation, care delivery arrangements and financial strategies.^{12–14} The barriers and proposed implementation strategies to overcome them are outlined below.

Patient factors

These factors involve the lack of understanding of the value of recommended practices among women seeking maternity care, families or communities. To address this gap, a suggested strategy is to undertake community-level sensitization to disseminate information about the risks of unnecessary caesarean sections and the benefits of adhering to recommended practices. Such strategy should be delivered in a clear and con-

^a UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), Department of Reproductive Health and Research, World Health Organization, Avenue Appia 20, 1211 Geneva 27, Switzerland.

^b School of Community Health and Midwifery, University of Central Lancashire, Preston, England.

^c Department of Maternal, Newborn, Child and Adolescent Health and Ageing, World Health Organization, Geneva, Switzerland.

Correspondence to Newton Opiyo (email: opiyon@who.int).

(Submitted: 16 May 2019 – Revised version received: 22 August 2019 – Accepted: 23 August 2019 – Published online: 29 November 2019)

sistent format using locally appropriate information, education and communication materials and activities.

Health-care providers factors

Several factors can hinder implementation of the guideline's recommendations. First, resistance of health-care providers to changing their practices coupled with a lack of understanding of the value of newly-recommended interventions. Suggested strategies to address this barrier include: involving local opinion leaders or champions to promote the implementation of the guideline's recommendations; providing information or education that helps the targeted health-care providers to fit the recommended behaviour into their current practice; and involving training institutions and professional bodies in implementing the guideline so that pre-service and in-service training curricula can be updated with the recommendations. Successful implementation strategies should be documented and shared as examples to other implementers. Second, the working arrangements and numbers of senior clinicians (obstetrician-gynaecologists) may hinder the implementation of mandatory second opinion for caesarean section indication. This challenge can be addressed with a strategy that involves developing local case-specific protocols to ensure timely and appropriate senior clinician review for caesarean section indication and by organizing teams with defined roles and a shared goal towards reducing unnecessary caesarean sections. Third, patients may make demands for caesarean sections that hinder adherence to recommended evidence-based practices, which could be addressed by providing tailored patient education materials and training health-care professionals to provide patient education. Fourth, financial incentives may hinder adherence (for example higher pay for caesarean sections compared with vaginal births). One strategy to address this could be to remove or modify financial incentive by involving key stakeholder groups, including policy-makers and medical insurance agencies. Fifth, dysfunctional teamwork among health-care providers, such as lack of communication between maternity and operating room staff, and on occasion, difficult relationships between obstetricians, midwives and family doctors can hinder implementation of guideline recom-

Box 1. Summary list of WHO recommendations on non-clinical interventions to reduce unnecessary caesarean sections

Interventions targeted at women

Recommendation 1. Health education for women is an essential component of antenatal care. The following educational interventions and support programmes are recommended to reduce caesarean births only with targeted monitoring and evaluation. (*Context-specific recommendation, low-certainty evidence*)

- Childbirth training workshops: Content includes sessions about childbirth fear and pain, pharmacological pain-relief techniques and their effects, non-pharmacological pain-relief methods, advantages and disadvantages of caesarean sections and vaginal delivery, indications and contraindications of caesarean sections, among others.
- Nurse-led applied relaxation training programme: Content includes group discussion of anxiety and stress-related issues in pregnancy and purpose of applied relaxation, deep breathing techniques, among other relaxation techniques.
- Psychosocial couple-based prevention programme: Content includes emotional self-management, conflict management, problem solving, communication and mutual support strategies that foster positive joint parenting of an infant. Couple in this recommendation includes couples, people in a primary relationship or other close people.
- Psychoeducation for women with fear of pain: Content includes information about fear and anxiety, fear of childbirth, normalization of individual reactions, stages of labour, hospital routines, birth process, and pain relief led by a therapist and midwife, among other topics.

When considering the educational interventions and support programmes, no specific format (pamphlet, videos, role play education) is recommended as more effective. (*Low- to moderate-certainty evidence*)

Interventions targeted at health-care professionals

Recommendation 2.1. Implementation of evidence-based clinical practice guidelines combined with structured, mandatory second opinion for caesarean section indication is recommended to reduce caesarean births in settings with adequate resources and senior clinicians able to provide second opinion for caesarean section indication. (*Context-specific recommendation, high-certainty evidence*)

Recommendation 2.2. Implementation of evidence-based clinical practice guidelines, caesarean section audits and timely feedback to health-care professionals are recommended to reduce caesarean births. (*Recommended, high-certainty evidence*)

Interventions targeted at health organizations or facilities

Recommendation 3.1. For the sole purpose of reducing caesarean section rates, collaborative midwifery-obstetrician model of care (that is, a model of staffing based on care provided primarily by midwives, with 24-hour back-up from an obstetrician who provides in-house labour and delivery coverage without other competing clinical duties) is recommended only in the context of rigorous research. This model of care primarily addresses intrapartum caesarean sections. (*Context-specific recommendation, low-certainty evidence*)

Recommendation 3.2. For the sole purpose of reducing unnecessary caesarean sections, financial strategies (such as insurance reforms equalizing physician fees for vaginal births and caesarean sections) for health-care professionals or health-care organizations are recommended only in the context of rigorous research. (*Context-specific recommendation, very low-certainty evidence*)

WHO: World Health Organization.

mendations. A suggested strategy would be to organize teams with defined roles and a shared goal towards reducing unnecessary caesarean sections.

Health-care organization factors

Four factors belong to this category. First, lack of staff with the necessary expertise and skills to implement, supervise and support recommended practices. To address this shortage, a possible strategy would be to redistribute health-care resources, where and when possible, and to consider task shifting of roles where appropriate. Strategic long-term planning and budgeting to

provide the necessary resources should also be contemplated. Second, lack of physical space, for example a venue for birth preparation classes, counselling sessions and training workshops. A proposed strategy to address this barrier is to adapt implementation strategies to work within the constraints of the existing systems and engage in strategic long-term planning and budgeting to provide the necessary resources. Third, lack of essential supplies including locally adapted information, education and communication materials to support training. Devising strategies to improve supply chain management

according to local requirements, such as developing protocols for obtaining and maintaining the stock of supplies, is one of the suggested solutions. Fifth, lack of health information management systems designed to document and monitor recommended practices, which could be addressed by providing appropriate incentives to record the needed information. In this case as well, strategic long-term planning and budgeting to provide the necessary resources for health information management systems should be undertaken.

Future research

The evidence underpinning the new recommendations was drawn from single studies with largely small samples. Therefore, the reported effect estimates and applicability of interventions should be confirmed in other settings, preferably in multicentre randomized trials. Future studies should be preceded by formative research to define locally relevant determinants of caesarean births that can be targeted by tailored interventions. The formative research would thus provide an important opportunity to modify and adapt the interventions to address key determinants of inappropriate caesarean use and achieve best

fit with national policies and health system capacities, for example resources available.

Combinations rather than single-standalone interventions to reduce unnecessary caesarean sections are preferred, given the multiple determinants and stakeholders involved in decision-making about the mode of birth. Understanding of views, values and preferences, and active engagement of all stakeholders through participatory approaches is crucial throughout the process.

During pregnancy, women and the health-care providers need to interact, occasions which offer multiple opportunities to support informed decision-making around choice of the mode of birth. More research is needed to understand determinants of birth choices, so that the content and format of educational interventions can be tailored to relevant determinants of caesarean births. Prioritized research gaps are outlined in the full guideline.⁴

Conclusions

Reducing unnecessary caesarean sections requires interventions that address both the clinical and non-clinical drivers of overuse. Clinical interventions that

could help to reduce caesarean section rates have been addressed in previous WHO guidelines. The new guideline, targeting non-clinical drivers, should be implemented alongside existing WHO-recommended clinical interventions as part of an integrated approach to optimize the use of caesarean sections. Multifaceted strategies selected based on local determinants of caesarean section practices and health system capacities are recommended to address barriers and help improve successful implementation of the recommendations. ■

Acknowledgements

We thank the people and organizations involved in the development of this guideline.

Funding: The United States Agency for International Development and the United Nations Development Fund / United Nations Population Fund / United Nations Children's Fund / WHO / the World Bank Special Programme of Research and Development and Research Training in Human Reproduction funded the development of this guideline.

Competing interests: None declared.

References

- Betrán AP, Torloni MR, Zhang J, Ye J, Mikolajczyk R, Deneux-Tharaux C, et al. What is the optimal rate of caesarean section at population level? A systematic review of ecologic studies. *Reprod Health*. 2015 06 21;12(1):57. doi: <http://dx.doi.org/10.1186/s12978-015-0043-6> PMID: 26093498
- Betrán AP, Ye J, Moller AB, Zhang J, Gülmezoglu AM, Torloni MR. The Increasing Trend in Caesarean Section Rates: Global, Regional and National Estimates: 1990-2014. *PLoS One*. 2016;11(2):e0148343.
- Keag OE, Norman JE, Stock SJ. Long-term risks and benefits associated with cesarean delivery for mother, baby, and subsequent pregnancies: Systematic review and meta-analysis. *PLoS Med*. 2018 01 23;15(1):e1002494. doi: <http://dx.doi.org/10.1371/journal.pmed.1002494> PMID: 29360829
- Lin HC, Xirasagar S. Institutional factors in cesarean delivery rates: policy and research implications. *Obstet Gynecol*. 2004 Jan;103(1):128-36. doi: <http://dx.doi.org/10.1097/01.AOG.0000102935.91389.53> PMID: 14704256
- WHO recommendations on non-clinical interventions to reduce unnecessary caesarean births. Geneva: World Health Organization; 2018. Available from: <https://www.who.int/reproductivehealth/publications/non-clinical-interventions-to-reduce-cs/en/> [cited 2019 Nov 16].
- WHO statement on caesarean section rates. Geneva: World Health Organization; 2015. Available from: https://apps.who.int/iris/bitstream/handle/10665/161442/WHO_RHR_15.02_eng.pdf?sequence=1 [cited 2019 Dec 12]
- Kingdon C, Downe S, Betrán AP. Women's and communities' views of educational interventions targeted at them to reduce unnecessary caesarean section: a qualitative evidence synthesis. *Reprod Health*. 2018;15:130. doi: <http://dx.doi.org/10.1186/s12978-018-0570-z> PMID: 30041661
- Kingdon C, Downe S, Betrán AP. Interventions targeted at health professionals to reduce unnecessary caesarean sections: a qualitative evidence synthesis. *BMJ Open*. 2018 12 16;8(12):e025073. doi: <http://dx.doi.org/10.1136/bmjopen-2018-025073> PMID: 30559163
- Kingdon C, Downe S, Betrán AP. Non-clinical interventions to reduce unnecessary caesarean section targeted at organisations, facilities and systems: Systematic review of qualitative studies. *PLoS One*. 2018 09 4;13(9):e0203274. doi: <http://dx.doi.org/10.1371/journal.pone.0203274> PMID: 30180198
- Bermúdez-Tamayo C, Fernández Ruiz E, Pastor Moreno G, Maroto-Navarro G, García-Mochon L, Pérez-Ramos FJ, et al. Barriers and enablers in the implementation of a program to reduce cesarean deliveries. *Reprod Health*. 2017 08 29;14(1):106. doi: <http://dx.doi.org/10.2471/BLT.06.039289> PMID: 18038061
- Chaillet N, Dubé E, Dugas M, Francoeur D, Dubé J, Gagnon S, et al. Identifying barriers and facilitators towards implementing guidelines to reduce caesarean section rates in Quebec. *Bull World Health Organ*. 2007 Oct;85(10):791-7. doi: <http://dx.doi.org/10.2471/BLT.06.039289> PMID: 18038061
- Pantoja T, Opiyo N, Lewin S, Paulsen E, Ciapponi A, Wiyongse CS, et al. Implementation strategies for health systems in low-income countries: an overview of systematic reviews. *Cochrane Database Syst Rev*. 2017 09 12;9(9):CD011086. PMID: 28901005
- Ciapponi A, Lewin S, Herrera CA, Opiyo N, Pantoja T, Paulsen E, et al. Delivery arrangements for health systems in low-income countries: an overview of systematic reviews. *Cochrane Database Syst Rev*. 2017 09 13;9(9):CD011083. PMID: 28901005
- Wiyongse CS, Paulsen E, Lewin S, Ciapponi A, Herrera CA, Opiyo N, et al. Financial arrangements for health systems in low-income countries: an overview of systematic reviews. *Cochrane Database Syst Rev*. 2017 09 11;9(9):CD011084. PMID: 28891235