



## Article

# Delivering motivational interviewing early post stroke: standardisation of the intervention

Patel, Kulsum, Auton, Malcolm Frederick, Watkins, Caroline Leigh, Sutton, Chris J, Benedetto, Valerio, Hackett, Maree, Holland, Emma-Joy and Lightbody, Catherine Elizabeth

Available at <http://clock.uclan.ac.uk/36233/>

*Patel, Kulsum, Auton, Malcolm Frederick ORCID: 0000-0002-8173-8159, Watkins, Caroline Leigh ORCID: 0000-0002-9403-3772, Sutton, Chris J ORCID: 0000-0002-6406-1318, Benedetto, Valerio ORCID: 0000-0002-4683-0777, Hackett, Maree ORCID: 0000-0003-1211-9087, Holland, Emma-Joy ORCID: 0000-0003-3029-7573 and Lightbody, Catherine Elizabeth ORCID: 0000-0001-5016-3471 (2022) Delivering motivational interviewing early post stroke: standardisation of the intervention. Disability and Rehabilitation, 44 (14). pp. 3453-3458. ISSN 0963-8288*

It is advisable to refer to the publisher's version if you intend to cite from the work.  
<http://dx.doi.org/10.1080/09638288.2020.1864035>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

## **Implications for Rehabilitation**

- Motivational Interviewing (MI) can help reduce depression in stroke survivors when delivered early after stroke.
- The effectiveness of our MI intervention depends on the delivery of high quality MI; in particular, interactions with low levels of MI-inconsistency, and high global MI ratings, ideally delivered over more than one session, each lasting at least 30 minutes.
- Provided high quality MI is being delivered, the intervention can still have a beneficial effect on participant outcome, even with flexibility and variation in therapist characteristics, and duration and number of sessions, which may be inevitable in a clinical context.