

Central Lancashire Online Knowledge (CLoK)

Title	The Radio Luminosity-Risetime Function of Core-Collapse Supernovae
Туре	Article
URL	https://clok.uclan.ac.uk/id/eprint/36815/
DOI	https://doi.org/10.3847/1538-4357/abccd9
Date	2021
Citation	Bietenholz, Michael F., Bartel, N., Argo, Megan, Dua, R., Ryder, S. and Soderberg, A. (2021) The Radio Luminosity-Risetime Function of Core- Collapse Supernovae. Astrophysical Journal, 908 (1). ISSN 0004-637X
Creators	Bietenholz, Michael F., Bartel, N., Argo, Megan, Dua, R., Ryder, S. and Soderberg, A.

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.3847/1538-4357/abccd9

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <u>http://clok.uclan.ac.uk/policies/</u>

The Radio Luminosity-Risetime Function of Core-Collapse Supernovae

M. F. BIETENHOLZ,^{1,2} N. BARTEL,² M. ARGO,³ R. DUA,⁴ S. RYDER,^{5,6} AND A. SODERBERG⁷

¹Hartebeesthoek Radio Astronomy Observatory, PO Box 443, Krugersdorp, 1740, South Africa

²Department of Physics and Astronomy, York University, Toronto, M3J 1P3, Ontario, Canada

³ Jeremiah Horrocks Institute, University of Central Lancashire, Preston PR1 2HE, UK

⁴Birla Institute of Technology and Science, Pilani Campus, Pilani, India

⁵Dept. of Physics & Astronomy, Macquarie University, NSW 2109, Australia

⁶Astronomy, Astrophysics and Astrophotonics Research Centre, Macquarie University, Sydney, NSW 2109, Australia

⁷ formerly at Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

(Accepted for publication in the Atrophysical Journal, January 18, 2021)

ABSTRACT

We assemble a large set of 2–10 GHz radio flux density measurements and upper limits of 294 different supernovae (SNe), from the literature and our own and archival data. Only 31% of SNe were detected. We characterize the SN radio lightcurves near the peak using a two-parameter model, with $t_{\rm pk}$ being the time to rise to a peak and $L_{\rm pk}$ the spectral luminosity at that peak. Over all SNe in our sample at D < 100 Mpc, we find that $t_{\rm pk} = 10^{1.7\pm0.9}$ d, and that $L_{\rm pk} = 10^{25.5\pm1.6}$ erg s⁻¹ Hz⁻¹, and therefore that generally, 50% of SNe will have $L_{\rm pk} < 10^{25.5}$ erg s⁻¹ Hz⁻¹. These $L_{\rm pk}$ values are ~30 times lower than those for only detected SNe. Types I b/c and II (excluding IIn's) have similar mean values of $L_{\rm pk}$ but the former have a wider range, whereas Type IIn SNe have ~ 10 times higher values with $L_{\rm pk}=10^{26.5\pm1.1}$ erg s⁻¹ Hz⁻¹. As for $t_{\rm pk}$, Type I b/c have $t_{\rm pk}$ of only $10^{1.1\pm0.5}$ d while Type II have $t_{\rm pk}=10^{1.6\pm1.0}$ and Type IIn the longest timescales with $t_{\rm pk}=10^{3.1\pm0.7}$ d. We also estimate the distribution of progenitor mass-loss rates, \dot{M} , and find the mean and standard deviation of $\log_{10}(\dot{M}/[M_{\odot} \text{ yr}^{-1}])$ are -5.4 ± 1.2 (assuming $v_{\rm wind}=1000 \text{ km s}^{-1}$) for Type I b/c SNe, and -6.9 ± 1.4 (assuming $v_{\rm wind}=10 \text{ km s}^{-1}$) for Type II SNe excluding Type IIn.

Keywords: Core-collapse supernovae, radio transient sources

1. INTRODUCTION

Core collapse supernova (SNe) can produce bright radio emission. The chief source of this emission is the interaction of the rapidly expanding ejecta with the circumstellar medium (CSM), which usually consists of the stellar wind of the SN progenitor, but may also have a significant contribution from mass-stripping in binary systems. Shocks are formed in this interaction, which serve to accelerate particles to relativistic velocities and amplify the magnetic field, resulting in synchrotron radio emission.

The radio emission provides us with a probe of the CSM, as well as for the outer, highest-velocity portion of the SN ejecta, for which few other observational probes are available. SNe are much less luminous in the radio

than in the optical, with typical radio luminosities $< 10^{-4}$ of those in the optical. Compared to the thousands of SNe detected in the optical, only ~100 SNe have been detected in the radio. Furthermore, only core-collapse SNe have been detected to date, and as yet no Type Ia SN (for recent limits on the radio emission of Type Ia SNe, see Lundqvist et al. 2020). In this paper, therefore, we consider only core-collapse SNe, that is SN of Types Ib, Ic and II, and whenever we use the term "SN" we are referring only to ones of the core-collapse variety.

The radio emission from SNe is synchrotron emission. It generally displays a high brightness temperature, and a non-thermal spectrum. Their radio lightcurves follow a general pattern with a rise to a maximum, which can occur days to years after the SN explosion. The peak is followed by a decay, often of an approximately power-law form, with $S_{\nu} \propto t^{\beta}$, where S_{ν} is the flux density at frequency ν , t the time since the explosion, and β is



Figure 1. An example radio lightcurve of a core-collapse SN. We show the lightcurve of the Type IIb SN 1993J (data from Bartel et al. 2002; see also Martí-Vidal et al. 2011), which is one of the most intensively observed radio SNe. We plot the flux density, S, at 8.4 GHz against the time, t, since the explosion, with both axes being logarithmic. The errorbars are mostly smaller than the plotted points. The typical features are a rapid rise to a peak, which occurs at $t_{\rm pk} \simeq 130$ d with $S_{\rm pk} = 123$ mJy, followed by a decline. In the case of SN 1993J the decline is of an approximately power-law form $(S \propto t^{-\beta})$, until $t \simeq 2500$ d, at which time there is a distinct steepening of the slope of the logarithmic decline. For our purposes here, we concentrate only on the region of the lightcurve near the peak and do not try to fit features such as the change in slope in the decay.

usually in the range of -1 to -3. As an illustration, we show the 8.4 GHz lightcurve of SN 1993J in Figure 1 (data from Bartel et al. 2002, and our own unpublished measurements). SN 1993J shows the typical rise and then power-law decay, although in this case, there is a distinct change in the slope of the decay after about 7 yr.

The radio lightcurves of SNe vary over a large range. Although the brightest SNe reach peak spectral luminosities, $L_{\rm pk}{}^1 = 10^{29}$ erg s⁻¹ Hz⁻¹ (e.g., SN 1998bw, SN 2009bb), a considerable fraction of even nearby SNe are never detected in the radio, and must have $L_{\rm pk}$ values at least 3 orders of magnitude lower, or $< 10^{26}$ erg s⁻¹ Hz⁻¹. Indeed, the $L_{\rm pk}$ of SN 1987A was another 2 orders of magnitude lower at $\lesssim 10^{24} \text{ erg s}^{-1} \text{ Hz}^{-1}$. Similarly, the risetimes $t_{\rm pk}$, have a very wide range. Some SNe, such as SN 1987A, have a very short $t_{\rm pk} \simeq 1$ d, while others, such as SN 1986J, can take several years to reach their peak.

A considerable number of radio flux density measurements of individual SNe have been published over the years. Much effort has also gone into parameterizing and modeling the radio lightcurves for the subset of SNe for which densely-sampled lightcurves are available (see e.g., Weiler et al. 2002). However, there has been rather less examination of the population as a whole. In this paper, we will explore, in a largely empirical way, the radio luminosity function of supernovae, and attempt first to answer the questions: how bright do we expect a core-collapse supernova to get in the radio, and how long do we expect it to take to reach this peak?

Our approach is as follows: we will adopt a simple parameterization of a supernova radio lightcurve, with only two parameters: $t_{\rm pk}$, the time between the explosion and L_{pk} , the peak spectral luminosity at that time. The challenge is to find the values of t_{pk} and L_{pk} . For a lightcurve with many flux density measurements as depicted in Figure 1 this can be done straightforwardly and relatively unambiguously. However, if there is only a single flux density measurement available, then the determination of t_{pk} and L_{pk} is ambiguous but as we will quantify later, the range of possible values of t_{pk} and L_{pk} is still well constrained. In the case of only upper limits on the flux density, the values of t_{pk} and L_{pk} are also ambiguous, but nonetheless still constrained, although generally less so than in the case of a single measurement. In this paper we use all our measurements to derive statistically meaningful results.

Many SNe in fact show behavior more complex than assumed in our simple model, with modulated lightcurves and anomalous rises at late times (for example, SN 1993J, already shown in Figure 1; but also SN 1979C, Bartel & Bietenholz 2008; SN 1986J, Bietenholz et al. 2002; SN 1987A, Zanardo et al. 2010; Cendes et al. 2018; SN 2001em Bietenholz & Bartel 2005; and SN 2001ig Ryder et al. 2004). However, most SNe do show an initial rise to a peak brightness and then a subsequent decay, so our model should suffice for giving us some insight into the population as a whole. For those SNe, such as SN 1987A, which showed a late-time rise in the radio emission, we use only the measurements for the first rise and subsequent decay.

We divide SNe into different Types such as Types I b/c or II and determine the difference in the luminosityrisetime function for different SN Types. We use the following three main classifications: Type I b/c, Type IIn, and then the remainder of the Type II's. In what fol-

¹ More formally, $L_{\rm pk}$ should be denoted $L_{\nu,{\rm pk}}$ since the peak spectral luminosity will depend on the observing frequency, ν . We omit the ν subscript on $L_{\nu,{\rm pk}}$ and $L_{\nu,{\rm pk,obs}}$ for clarity. We expect in any case that the dependence on ν will not be large, since we restrict ourselves to frequencies, $4 < \nu < 10$ GHz, with the exception of SN 1987A. Indeed, Weiler et al. (2002) found that the dependence of the $L_{\rm pk}$ on ν was not large for a variety of SNe of Type Ib/c and II.

lows, when we mention Type II, we always mean Type II excluding the Type IIn. In addition, we examine separately the subset of Type I b/c SNe which has broad optical lines, which we call "BL", and the Type IIb subset of Type II's.

Type IIn SNe are those with narrow optical lines. They constitute $\sim 12\%$ of all Type II SNe (Smith et al. 2011). Examples are SN 1986J and SN 1998S. These SNe are thought to be due to interaction with a dense CSM, which produces the narrow lines, and often strong radio emission. The radio evolution of Type IIn SNe is quite different from that of normal Type II SNe, which is why we treat then as a separate group. Occasionally Type Ib SNe are also observed to have narrow lines, and classed as Type Ibn. Our sample, however, contained only a single Type Ibn SN, SN 2015G, which was not detected, therefore we do not discuss the Ibn subtype separately.

We also discuss the subset of Type I b/c SNe which have broad optical lines, indicating high ejection velocities, (BL) as a group. This subtype has been of special interest because it is associated with gamma-ray bursts (Woosley & Bloom 2006; Cano et al. 2017).

Finally, we also discuss the Type IIb subset of Type II SNe, of which SN 1993J is the most famous example. These SNe initially have H in their spectra and are therefore classified as Type II, but transition subsequently to having He-dominated spectra more characteristic of Type Ib. They constitute $\sim 14\%$ of all Type II's (Smith et al. 2011).

An important caveat must be mentioned here. The classification of SNe into Types is based on features in the optical spectrum. Since such features can vary as the SN evolves, there is the possibility that a SN may appear as different Types at different stages in its evolution. Indeed, we just mentioned the Type IIb SNe whose spectra changes from Type II to one resembling Type Ib.

The classification of Type IIn SNe is also occasionally time-variable. The interaction with the dense CSM giving rise to the narrow lines and the "n" characteristics can start only some time after the explosion, so some SNe might first appear to be normal Type I or II, and then develop the "n" characteristics. SN 2014C is a prominent example of this behavior, which started as a Type Ib but developed IIn characteristics after about 1 yr (Milisavljevic et al. 2015). SN 2001em is the other example of this behaviour in our sample. Since the dominant part of the radio lightcurve for both SN 2001em and SN 2014C occurs at later times, when the optical spectrum was of Type IIn, we classify both SNe as Type IIn. Given the possible time-variability of the spectral characteristics, and therefore the non-uniqueness of the SN Type classification, our division by the SN Types is not completely unique. However, since only a small fraction of SNe show such time-variable spectral characteristics, our statistical results should not be greatly affected by their occurrence.

The remainder of this paper is organized as follows. First in Section 2, we briefly describe the observations and data reduction for the new data in this paper. Then, in Section 3, we describe our collection of radio measurements of 294 SNe. In Section 4 we describe the model of a SN radio lightcurve we fit to our measurements, which is characterized by only two parameters, $t_{\rm pk}$ and $L_{\rm pk}$. For many SNe, the measurements are not sufficient to uniquely determine the values of $t_{\rm pk}$ and $L_{\rm pk}$, for example, if there is only a single measurement, or only upper limits. In Section 5, we combine these constraints over all our SNe, and determine the likelihood of different values of t_{pk} and L_{pk} given our measurements. We then parameterize the distribution of L_{pk} and t_{pk} , finding that lognormal form is the most likely, and proceed to determine the particular lognormal distributions for $t_{\rm pk}$ and $L_{\rm pk}$ which are most compatible with our measurements. We also examine various SN subtypes, such as Type I b/c and Type II, to ask whether the distribution of $t_{\rm pk}$ and $L_{\rm pk}$ differs for different SN Types. In Section 6, we use our distrutions of t_{pk} and L_{pk} to estimate the distribution of mass-loss rates. In Section 7, we discuss the implications of our results, and finally in Section 8 we summarize them and give our conclusions.

2. OBSERVATIONS AND DATA REDUCTION

We discuss our complete data-set which includes both published and previously unpublished values in the next section. Here we give a brief summary of the observations and data reduction of the 296 previously unpublished SN observations.

We re-reduced a number of archival observations of SNe from the Karl G. Jansky Very Large Array (VLA). This was done in a standard manner, using the Astronomical Image Processing System (AIPS; Associated Universities 1999) for observations from the VLA before about 2011, and Common Astronomy Software Application (CASA; International Consortium Of Scientists 2011) The flux density calibration was done using observations of 3C 48, 3C 138 or 3C 286. Phase selfcalibration was done on the supernova observations in cases where the signal-to-noise ratio was adequate, but no amplitude self-calibration was done. In most of the archival data sets, the supernova was not detected, so no self-calibration was done. The flux densities were determined by fitting to the images elliptical Gaussians, fixed to the dimensions of the restoring beam, along with a zero-level to account for any extended emission from the host galaxies. The total uncertainties (in Table 1) include a 5% uncertainty on the flux-density calibration, and in some cases a contribution from the uncertainty in separating the SN from the background emission, added in quadrature to the image background rms.

All observations with the Australia Telescope Compact Array (ATCA) used the 2 GHz bandwidth CABB system (Wilson et al. 2011) and were processed and measured using the MIRIAD package (Sault et al. 1995), as described in Bufano et al. (2014). The primary flux density calibrator was PKS B1934-638, and no selfcalibration was applied.

Observations with the Multi-Element Radio-Linked Interferometer Network (MERLIN) used the e-Merlin pipeline (Argo 2014) using 512 MHz bandwidth. The primary flux density calibrator was 3C 286, and no selfcalibration was done.

3. THE DATA-SET

We avail ourselves of as many of the published results as possible, taking care to include any published upper limits in the cases of non-detection. To keep our data-set as uniform as possible, we restricted ourselves to measurements between 4 and 10 GHz since the most commonly used observing frequencies are 4.8 and 8.4 GHz, making an exception for SN 1987A, where only a very few measurements are available in the first years at those frequencies and we therefore use the more complete 2.3 GHz lightcurve. We add to the previously published values a number of previously unpublished measurements, which are listed in Table 1.

Our previously unpublished values include new measurements from the ATCA and MERLIN, as well as a number of results from re-reduced data from the Karl G. Jansky Very Large Array (VLA) available in the National Radio Astronomy Observatory (NRAO)² data archive. There are a considerable number of such observations which were never published. Many SNe, even relatively nearby ones, are never detected in the radio. Such non-detections are much less likely to be published, therefore the sample of published values is likely to be biased towards detections and thus higher radio luminosities. We have therefore re-reduced a significant number of unpublished archival measurements, the majority of which are indeed non-detections.

Finally, we include values from the website of the late Kurt W. Weiler. Dr. Weiler obtained many radio observations of SNe during his illustrious career — see, for example, Weiler et al. (2002). Some of these were made available for a time on his website at the U. S. Naval Observatory, but were never formally published. We had retrieved some of those values from the website, which we now include also in our data set and in Table 1.

While the largest fraction of our assembled observations are from the VLA and ATCA, we also have measurements from a number of other telescopes including MERLIN, the Westerbork Synthesis Radio Telescope, the European VLBI Network, the Urumqi radio telescope and the Parkes-Tidbinbilla Interferometer.

In total we have 1475 measurements the flux density, or upper limits on it, for 294 SNe. For well observed SNe, such as SN 1993J (Figure 1) or SN 1986J, we have very well sampled lightcurves, with many measurements (n = 29 and 39 respectively), allowing t_{pk} and L_{pk} to be accurately determined. For the majority of SNe, however, only one or two measurements are available, which thus provide only weak constraints on t_{pk} or L_{pk} . In fact, in many cases, the observations yielded only upper limits on the SN's flux density.

Of our 294 SNe, only 31% (n = 90) are detected. For the remaining 69% (n = 204) we have only upper limits on the flux density. The average number of measurements or limits per SN, detected or not, is 5.0. However, this number is skewed by the 9% (n = 26) of wellobserved SNe which have more than 12 measurements each. In fact, 35% (n = 104) of our SNe have only a single measurement or limit. At least three observations are required to uniquely determine the peak of the lightcurve (one near, one before and one after the peak). Only 27% (n = 79) of our SNe have three or more measurements or limits, although in many of those cases, they all occur after the peak, so that the peak is not determined.

Given this relatively modest number of measurements, compared to what is available in the optical, and the fact that our sample is of necessity heterogeneous and incomplete, we cannot provide a definitive radio luminosity function for supernovae. Nonetheless, we have a larger data set than has ever previously been assembled, and sufficiently large that some reasonably robust inferences can be drawn. It is crucial for this purpose to consider the non-detections as well as the published detections.

Table 2 gives some details of the SNe in our database. In order to determine luminosities, we need the dis-

 $^{^2}$ The NRAO, is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

NOTE—Table 1 is published in its entirety in the machine-readable format. Only a portion is shown here for guidance regarding its form and content.

```
Title: The Radio Luminosity-Risetime Function of Core-Collapse Supernovae
Authors: Bietenholz M.F., Bartel N., Argo M., Dua R., Ryder S., Soderberg A.
Table: Supernova flux densities or limits from radio observations
_____
Byte-by-byte Description of file: datafile1.txt
  Bytes Format Units Label Explanations
  1- 9 A9
               ____
                      ID
                             SN identifier
               ____
     11 A1
                      Limit [L] Limit flag on Flux (1)
  13- 16 I4
               yr
                      Obs.Y UT Year of observation midpoint
  18- 19 I2
                      Obs.M UT Month of observation midpoint
               month
  21- 25 F5.2
               d
                      Obs.D UT Day of observation midpoint
  27- 33 A7
               ____
                      Tel
                             Telescope identifier (2)
  36- 40 F5.2
               GHz
                      Freq
                            Observed frequency
  42- 49 F8.4
                      Flux
                             Measured flux density at Freq (3)
               mJy
  51- 56 F6.4
                     e_Flux
                             Uncertainty in Flux
               mJy
  58- 63 A6
               ___
                      Com
                             Additional comment (4)
                      _____
                             _____
                                                                _____
Note (1): "L" indicates a limit, blank indicates a measured value.
Note (2):
   VLA = Very Large Array, USA; if known, the VLA configuration
         is appended, e.g. VLA-A;
 MERLIN= the Multi-Element Radio-Linked Interferometer Network, UK;
   ATCA= Australia Telescope Compact Array, Australia.
Note (3): A negative value indicates a limit, with the magnitude of
         the value being the 3-sigma upper limit
Note (4): "Weiler" indicates that this value was retrieved from the
         website of the late Kurt Weiler.
SN19800
        L 1988 2 1.32 VLA-AB
                                  4.86 -0.360 0.120
SN1982F
         L 1984 8 31.00 VLA-D
                                  4.86 -1.160
                                               0.390
         L 1984 12 23.00 VLA-A
                                       -0.180
SN1982F
                                  4.86
                                               0.060
SN1985F
         L 1985 3 18.00 VLA
                                  4.86
                                       -0.189
                                               0.064
        L 1985 7 31.00 VLA
                                  4.86
                                       -0.330
SN1985F
                                               0.110
SN1985G
         L 1985 5 7.3 VLA
                                  4.86
                                       -0.212
                                               0 071
SN1985G
         L 1985 9 1.00 VLA
                                  4.86
                                        -0.675
                                               0.225
SN1985G
         L 1986 12 15.00 VLA
                                  4.86
                                       -0.623
                                               0.208
SN1993N
        L 1994 2 18.32 VLA
                                       -0.110 0.037
                                  8.44
SN1993N
        L 1997 1 23.00 VLA
                                  8.46
                                       -0.186 0.062 Weiler
. . . .
                                        2.19
SN2010as
           2010 4 16.7 ATCA
                                  9.00
                                               0.11
           2010 4 25.5 ATCA
                                  9.00
SN2010as
                                        3.10
                                               0.9
. . . .
```

tances, D, for our SNe. In most cases, we calculated D from the recession velocity for the parent galaxy from the NASA/IPAC Extragalactic Database (NED)³, using the value corrected for our motion with respect to the cosmic microwave background, and infall to the Virgo cluster, to the Great Attractor and to the Shapley supercluster (Mould et al. 2000).

We use the latest values from the Planck collaboration, which are $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_{\text{matter}} = 0.315$ and $\Omega_{\Lambda} = 0.685$ (Planck Collaboration et al. 2020). Since our most distant (SN 2010ay) is at $D \simeq 300 \text{ Mpc}$, and most SNe (89%) are at D < 100 Mpc, the precise values adopted for the cosmological parameters do not significantly affect our results. For SNe closer than 30 Mpc, we use the mean of the redshift-independent distances from NED when available in preference to those calculated from the recession velocity.

SN Name	$_{\mathrm{Type}}a$	Galaxy	$_{ m Distance}b$	Explosion	Number of	Detected	$_{\rm References} e$
			$(D; \mathrm{Mpc})$	$_{ m date}c$	$_{\rm measurements}d$		
SN 1979C	IIL	NGC 4321	16.2	$1979 \ 04 \ 06$	67	Υ	1, 2
SN 1980K	IIb-L	NGC 6946	5.5	$1980 \ 10 \ 25$	69	Υ	3, 4
SN 1980O	II	NGC 1255	17.9	$1980\ 12\ 30$	2		5, 6
SN 1981A	II	NGC 1532	17.9	$1981 \ 02 \ 28$	1		5
SN 1981K	II	NGC 4258	7.3	$1981 \ 07 \ 31$	30	Υ	1
SN 1982F	IIP	NGC 4490	6.2	$1982 \ 02 \ 24$	2		6
SN 1982aa	?	NGC 6052	80.5	$1979 \ 08 \ 16$	11	Υ	7
SN 1983I	Ic	NGC 4051	13.7	$1983\ 04\ 25$	2		8
SN 1983K	II	NGC 4699	19.7	$1983\ 06\ 22$	3		9
SN 1983N	Ib	NGC 5236	4.9	$1983\ 06\ 29$	15	Υ	10
SN 1984E	IIL	NGC 3169	22.4	$1984 \ 03 \ 29$	4		9
SN 1984L	Ib	NGC 991	8.8	$1984 \ 08 \ 10$	3	Υ	11
SN 1985F	Ib/c	NGC 4618	7.2	$1984 \ 03 \ 30$	2		6
SN 1985G	IIP	NGC 4451	20.9	$1985\ 03\ 17$	3		6
SN 1985H	II	NGC 3359	16.0	$1985 \ 04 \ 12$	2		6
SN 1985L	IIL	NGC 5033	16.5	$1985\ 06\ 13$	7	Υ	12
SN 1986E	IIL	NGC 4302	16.8	$1986 \ 03 \ 28$	7	Υ	13
SN 1986J	IIn	NGC 891	10.0	$1983\ 03\ 14$	39	Υ	14, 15
SN 1987A	IIf	LMC	0.051	$1997 \ 02 \ 23$	8	Υ	16
SN 1987F	IIn:	NGC 4615	79.6	$1987 \ 03 \ 22$	4		6
SN 1987K	IIb	NGC 4651	16.5	$1987 \ 07 \ 31$	2		6
SN 1988I	IIn	Leda 86944	178	$1988 \ 03 \ 07$	1		9
SN 1988Z	IIn	MCG+03-28-22	111	$1988\ 12\ 01$	26	Υ	6, 17
SN 1989C	IIP	UGC 5249	32.1	$1989\ 02\ 01$	1		9
SN 1989L	II	NGC 7339	22.0	$1989\ 05\ 04$	3		6
SN 1989R	IIn	UGC 2912	80.1	$1989 \ 09 \ 15$	1		9
SN 1990B	Ic	NGC 4568	17.4	$1990\ 01\ 18$	8	Υ	18
SN 1990K	II	NGC 150	23.4	$1990\ 05\ 14$	2		6
SN 1991G	IIP	NGC 4088	13.9	$1991 \ 01 \ 23$	2		6
SN 1991N	Ic	NGC 3310	18.1	$1991 \ 04 \ 02$	2		8, 19
SN 1991ae	IIn	MCG+11-19-18	138	$1991 \ 05 \ 15$	2		6, 9
SN 1991av	IIn	Anon J215601+0059	288	$1991 \ 09 \ 15$	3		9
SN 1992H	II	NGC 5377	35.1	$1992\ 02\ 11$	2		6
SN 1992ad	II	NGC 4411B	22.4	$1992 \ 06 \ 30$	5	Υ	6, 20
SN 1992bd	II	NGC 1097	16.9	$1992 \ 10 \ 12$	5		6
SN 1993G	IIL	NGC 3690	53.1	$1993\ 02\ 24$	1		6
SN 1993J	IIb	M81	3.7	$1993 \ 03 \ 28$	29	Υ	21
SN 1993N	IIn	UGC 5695	50.2	$1993 \ 04 \ 15$	2		6, 22
SN 1993X	II	NGC 2276	40.5	$1993 \ 08 \ 22$	1		6
SN 1994I	Ic	M51	7.9	$1994 \ 03 \ 31$	39		6, 19, 23
SN 1994P	II	UGC 6983	19.6	$1994 \ 01 \ 20$	3		6
SN 1994W	IIn-P	NGC 4041	25.4	$1994\ 07\ 30$	3		6, 22
SN 1994Y	IIn	NGC 5371	46.4	$1994 \ 07 \ 09$	1		6
SN 1994ai	Ic	NGC 908	15.6	$1994 \ 12 \ 20$	2		6, 19
SN 1994ak	IIn	NGC 2782	43.1	$1994 \ 12 \ 24$	1		6
SN 1995N	IIn	MCG-02-38-17	31.4	$1994\ 07\ 04$	18	Υ	24
SN 1995X	П	UGC 12160	25.5	1995 08 03	4		22

 Table 2.
 Supernovae Observed in Radio

Table 2 continued

(D; Mpc) date ^C measurements ^d SN 1995ad II NGC 2139 27.0 1995 09 22 1 22 SN 1996L IIn ESO 266-G10 157 1996 03 12 1 22 SN 1996N Ib NGC 1398 19.8 1996 03 09 3 Y 6, 19,	25
SN 1995ad II NGC 2139 27.0 1995 09 22 1 22 SN 1996L IIn ESO 266-G10 157 1996 03 12 1 22 SN 1996N Ib NGC 1398 19.8 1996 03 09 3 Y 6, 19,	25
SN 1996L IIn ESO 266-G10 157 1996 03 12 1 22 SN 1996N Ib NGC 1398 19.8 1996 03 09 3 Y 6, 19,	25
SN 1996N Ib NGC 1398 19.8 1996 03 09 3 Y 6, 19,	25
SN 1996W II NGC 4027 12.2 1996 04 10 3 6, 22	
SN 1996ae IIn NGC 5775 19.9 1996 01 27 4 6, 22	
SN 1996an II NGC 1084 19.1 1996 05 30 2 22	
SN 1996aq Ic NGC 5584 21.8 1996 08 17 4 6, 19	26
SN 1996bu IIn NGC 3631 10.3 1996 11 14 2 6	
SN 1996bw II NGC 664 79.0 1996 11 30 1 22	
SN 1996cb IIb NGC 3510 13.9 1996 12 12 3 Y 6, 27	
SN 1996cr IIn: Circinus 3.8 1995 03 01 11 Y 28	
SN 1997W II NGC 664 79.0 1997 02 01 2 6, 22	
SN 1997X Ib/c NGC 4691 21.3 1997 01 25 3 Y 6, 19	
SN 1997ab IIn Anon J095100+2004 53.9 1996 04 11 2 22	
SN 1997db II UGC 11861 18.9 1997 08 02 3 6, 22	
SN 1997dn II NGC 3451 27.1 1997 10 29 1 6	
SN 1997dq IcBL NGC 3810 15.7 1997 10 13 3 22, 10	
SN 1997ef IbBL UGC 4107 55.9 1997 11 20 2 6, 19	
SN 1997eg IIn NGC 5012 47.6 1997 12 04 3 Y 29	
SN 1997ei Ic NGC 3963 48.8 1997 11 20 1 22	
SN 1998S IIn NGC 3877 14.9 1998 02 28 8 Y 6, 22	30
SN 1998bm II IC 2458 24.7 1998 04 21 2 6	
SN 1998bw IcBL ESO 184-82 41.4 1998 04 25 31 Y 31	
SN 1998dl IIP NGC 1084 19.1 1998 08 02 2 22	
SN 1998dn II NGC 337A 13.7 1998 08 19 2 22	
SN 1999B II UGC 7189 31.2 1999 01 14 1 6	
SN 1999D II NGC 3690 52.6 1999 01 16 2 6, 22	
SN 1999E IIn Anon J131716-1833 119 1998 09 10 1 22	
SN 1999cn Ic MCG+02-38-43 111 1999 06 14 1 22	
SN 1999dn Ib NGC 7714 29.1 1999 08 15 1 6, 19	
SN 1999eb IIn NGC 664 79.0 1999 10 02 1 6	
SN 1999eh Ib NGC 2770 28.6 1999 07 26 2 8, 19	
SN 1999el IIn NGC 6951 23.1 1999 10 20 2 6	
SN 1999em IIP NGC 1637 11.5 1999 10 24 5 Y 6, 22	32
SN 1999ev IIP NGC 4724 13.9 1999 11 07 1 6	
SN 1999ex Ic IC 5179 53.3 1999 11 01 1 33	
SN 1999gi IIP NGC 3184 12.4 1999 12 06 3 6	
SN 1999go II NGC 1376 60.4 1999 12 18 1 6	
SN 1999gg IIP NGC 4523 16.7 1999 12 23 1 Y 6	
SN 2000C Ic NGC 2415 59.4 2000 01 01 1 19.3	
SN 2000F Ic IC 302 86.1 2000 01 29 1 19	
SN 2000P IIn NGC 4965 30.2 2000 03 08 2 22	
SN 2000S Ic MCG-01-27-20 138 1999 10 09 1 19	
SN 2000cr Ic NGC 5395 61.3 2000 06 21 1 33	
SN 2000ds Ib/c NGC 2768 20.5 2000 05 28 3 8.19	
SN 2000ew Ic NGC 3810 15.7 2000 11.21 1 6	
SN 2000fm Ib NGC 2526 72.3 2000 11 09 1 33	
SN 2000ft ? NGC 7469 73.5 2000 07.19 7 Y 34	
SN 2001B Ib IC 391 27 4 2000 12 31 3 V 19 3	
SN 2001M Ic NGC 3240 57 3 2001 01 17 1 33	
SN 2001ai Ic NGC 5278 191 2001 01 17 1 33	
SN 2001bb Ic IC 4319 82.0 2001 04 29 2 10 35	
SN 2001ch Ic MCG-01-54-16 46.8 2001 03 24 1 19	

Table 2 (continued)

SN Name	Type	Galaxy	Distance	Explosion	Number of	Detected	References
	- J F -	C			1		
			(D; Mpc)	$date^{C}$	measurements u		
SN 2001ci	Ic	NGC 3079	16.4	2001 04 21	3	Y	6, 19, 33
SN 2001ef	Ic	IC 381	40.2	$2001 \ 09 \ 04$	2		19, 33
SN 2001ej	Ib	UGC 3829	62.8	2001 09 09	2		19, 33
SN 2001em	$_{\rm IIn}f$	UGC 11794	89.7	2001 09 12	8	Υ	35, 36, 37
SN 2001gd	IIb	NGC 5033	17.5	2001 09 03	11	Υ	38
SN 2001ig	IIb	NGC 7424	9.3	2001 12 03	23	Υ	39
SN 2001is	Ib	NGC 1961	61.2	2001 12 19	1		19
SN 2002ap	IcBL	NGC 628	8.9	2001 02 28	9	Υ	19, 40
SN 2002bl	IcPecBL	UGC 5499	77.5	2002 02 23	2		19, 33
SN 2002cj	Ic	ESO 582-05	113	2002 04 16	1	Υ	33
SN 2002cp	Ib/c	NGC 3074	82.9	2002 04 20	2		19, 33
SN 2002dg	Ib	Anon J145716+0554	225	2002 05 29	2		19, 33
SN 2002dn	Ic	IC 5145	112	2002 06 08	1		19, 33
SN 2002gy	Ib/c:	UGC 2701	107	2002 10 13	1		33
SN 2002hf	Ic	MCG-05-03-20	82.9	2002 10 26	2		19, 33
SN 2002hh	II	NGC 6946	5.6	2002 10 31	8	Υ	22, 41
SN 2002hn	Ic	NGC 2532	82.1	2002 10 26	1		33
SN 2002ho	Ic	NGC 4210	47.2	2002 11 01	2		19, 33
SN 2002hy	IbPec	NGC 3464	60.7	2002 10 28	2		19, 33
SN 2002hz	Ib	UGC 12044	82.9	2002 11 07	2		19, 33
SN 2002ji	Ic	NGC 3655	30.3	2002 10 19	3		19, 33, 42
SN 2002jj	Ic	IC 340	60.6	2002 10 13	2		19, 33
SN 2002jp	Ic	NGC 3313	59.6	2001 11 15	2		19, 33
SN 2002jz	Ic	UGC 2984	22.9	2001 12 14	1		33
SN 2003H	IbPec	NGC 2207	22.3	2003 01 08	3		6, 42
SN 2003L	Ic	NGC 3506	104	2001 01 01	40	Υ	43
SN 2003bg	IcPecBL	MCG -05-10-15	19.3	2003 02 22	41	Υ	44
SN 2003bu	Ic	NGC 5953	105	2003 03 03	2		8
SN 2003dr	Ib/c	NGC 5714	41.7	2004 04 10	3		8, 19, 42
SN 2003 dv	IIn	UGC 9638	33.9	$2004 \ 04 \ 16$	1		6
SN 2003ed	II	NGC 5303A	25.2	2003 04 30	4	Υ	6, 45
SN 2003el	Ic	NGC 5000	93.4	2003 05 11	1		19
SN 2003gd	IIP	NGC 628	8.6	2003 03 17	5		6
SN 2003gk	Ib	NGC 7460	48.5	$2003 \ 06 \ 15$	1		8
SN 2003ie	IIP	NGC 4051	13.7	2003 09 19	4		6, 22
SN 2003jd	IcPecBL	MCG -01-59-2	84.6	2003 10 10	4		8, 19
SN 2003jg	Ib/c	NGC 2997	9.0	$2003 \ 10 \ 01$	3		8, 42
SN 2003lo	IIn	NGC 1376	60.4	$2003 \ 12 \ 31$	1		6
SN 2004A	IIP	NGC 6207	17.0	$2004 \ 01 \ 06$	6		6
SN 2004C	Ic	NGC 3683	32.6	2003 12 23	3	Υ	6, 8
SN 2004am	IIP	NGC 3034	3.8	$2003 \ 11 \ 07$	3		22, 46
SN 2004ao	Ib	UGC 10862	26.8	$2004 \ 02 \ 21$	1		8
SN 2004bm	Ic	NGC 3437	24.4	$2004 \ 04 \ 17$	2		6, 42
SN 2004bu	IcBL	UGC 10089	92.1	$2004 \ 05 \ 14$	1		8
SN 2004cc	Ic	NGC 4568	17.4	$2004 \ 05 \ 23$	8	Υ	47
SN 2004dj	IIP	NGC 2403	3.4	2004 07 13	40	Υ	48
SN 2004 dk	Ib	NGC 6118	20.8	$2004 \ 07 \ 30$	10	Υ	47
SN 2004 et	IIP	NGC 6946	5.6	$2004 \ 09 \ 22$	19	Υ	22, 49
SN 2004gq	Ib	NGC 1832	24.3	$2004 \ 12 \ 08$	21	Υ	47
SN 2004gt	Ib/c	NGC 4038	21.1	$2004 \ 11 \ 27$	2		8, 42, 60
SN 2005E	Ib/c	NGC 1032	38.8	$2005 \ 01 \ 04$	1		8
SN 2005U	IIb	NGC 3690	53.1	2005 01 28	2		6
SN 2005V	Ib/c	NGC 2146	19.6	$2005 \ 01 \ 01$	5		6, 8, 42

 Table 2 continued

Table 2 (continued)

SN Name	Typea	Galaxy	Distance	Explosion	Number of	Detected	$\overline{\mathrm{References}}e$
			$(D; \mathrm{Mpc})$	$_{ m date}c$	$_{\rm measurements}d$		
SN 2005aj	Ic	UGC 2411	41.1	$2005 \ 02 \ 09$	2		8, 42
SN 2005at	Ic	NGC 6744	7.2	$2005 \ 03 \ 05$	2		50
SN 2005ay	IIP	NGC 3938	12.7	$2005 \ 03 \ 21$	4		6
SN 2005cs	IIP	M51	7.9	$2005 \ 06 \ 27$	5		51
SN 2005ct	Ic	NGC 207	58.6	$2005 \ 05 \ 29$	1		8
SN 2005cz	Ib	NGC 4589	35.7	$2005 \ 06 \ 17$	1		8
SN 2005da	IcBL	UGC 11301	74.4	$2005 \ 06 \ 25$	3		8
SN 2005dl	II	NGC 2276	20.4	$2005 \ 08 \ 25$	2		6
SN 2005ek	Ic	UGC 2526	73.0	$2005 \ 09 \ 22$	1		52
SN 2005gl	IIn	NGC 266	68.8	$2005 \ 10 \ 26$	1		22
SN 2005ip	IIn	NGC 2906	36.5	$2005 \ 10 \ 27$	3	Υ	53
SN 2005kd	IIn	2MFGC 3318	69.4	2005 11 10	4	Υ	6, 22, 54, 55
SN 2005kl	Ic	NGC 4369	29.7	$2005 \ 11 \ 01$	1		6
SN 2006aj	IcBL	2XMM J032139.6+165202	153	$2006 \ 02 \ 18$	17	Υ	56
SN 2006be	II	IC 4582	40.6	2006 03 13	1		57
SN 2006bp	IIP	NGC 3953	16.6	2006 04 09	4		22, 58
SN 2006gy	IIn	NGC 1260	85.0	2005 08 20	8		6, 59, 60
SN 2006jd	IIn	UGC 4179	83.7	2006 10 07	11	Υ	61
SN 2006my	IIP	NGC 4651	16.5	2006 08 01	2		22
SN 2006ov	IIP	NGC 4303	14.6	2006 10 26	2		6, 22
SN 2007C	Ib	NGC 4981	22.7	2006 12 28	2	Y	6
SN 2007Y	IbPec	NGC 1187	16.8	2007 02 14	7		42, 62
SN 2007ak	IIn	UGC 3293	69.6	2007 03 10	1		22
SN 2007bg	IcBL	Anon J114926+5149	155	2007 04 16	18	Y	63
SN 2007gr	Ib/c	NGC 1058	5.2	2007 08 13	9	Y	64
SN 2007iq	IcBL	UGC 3416	62.5	2007 08 01	2		8, 42
SN 2007ke	Ib	NGC 1129	76.7	2007 09 02	1		8
SN 2007ki	Ib/c	NGC 7803	79.3	2007 09 14	1		8
SN 2007pk	IInPec	NGC 579	73.4	2007 11 08	1		65
SN 2007rt	IIn	UGC 6109	107	2007 09 05	1		66
SN 2007ru	IcBL	UGC 12381	70.3	2007 11 25	2		8
SN 2007rz	Ic	NGC 1590	57.1	2007 11 19	2		8, 42
SN 2007uv	Ib	NGC 2770	28.6	2007 12 27	16	Y	67
SN 2008B	IIn	NGC 5829	94.5	2008 01 02	1		68
SN 2008D	Ib	NGC 2770	28.6	2008 01 09	21	Y	69
SN 2008X	IIP	NGC 4141	35.4	2008 01 14	2		6, 70
SN 2008ai	IIn	MCG+06-30-34	122	2008 02 12	1		71
SN 2008ax	IIb	NGC 4490	6.2	2008 03 03	24	Y	22. 72
SN 2008be	IIn	NGC 5671	142	2008 03 12		-	73
SN 2008bk	IIP	NGC 7793	3.9	2008 03 07	- 1		74
SN 2008bm	IIn	Leda 45053	155	2008 03 29	- 1		75
SN 2008bo	IIh	NGC 6643	19.1	2008 03 27	7	v	22 76
SN 2008du	Ic	NGC 7422	72.4	2008 06 30	1	1	42
SN 2008dy	Ic	NGC 1343	10.5	2008 05 26	1		42 8 42
SN 2008ew	Ic	IC1236	99.5	2008 08 06	- 1		8
SN 2008gm	IIn	NGC 7530	53.0	2008 10 02	1		77
SN 2008bb	Ic	IC 112	85.0	2008 11 04	1		8
SN 2008hp	Ic	NCC 2545	54.2	2008 11 04	1		8
SN 2008ii	п	NGC 6643	10.1	2000 11 12	1		78
SN 2000ij	њ	UCC 2006	19.1	2000 12 19	1		2
SN 2008in	IID	NGC 4303	40.2	2000 12 10 2008 12 22	1		6 70
SN 2008in	IIn	NGC 4846	14.0	2000 12 22	2		0, 19 80
SN 2008iz	?	M82	3.8	2008 02 20	25	Y	81

Table 2 (continued)

SN Name Type ^{<i>a</i>} Galaxy		Distance	Explosion	Number of	Detected	References	
			(D; Mpc)	$_{ m date}c$	$_{\rm measurements}d$		
SN 2008jb	II	ESO 302-14	9.3	$2008\ 11\ 11$	1	Υ	6
SN 2009E	IIP	NGC 4141	35.4	$2008 \ 01 \ 01$	1		6
SN 2009H	II	NGC 1084	19.1	$2009 \ 01 \ 02$	2		82
SN 2009N	IIP	NGC 4487	17.2	$2009 \ 01 \ 24$	2		82
SN 2009au	IIn	ESO 443-21	36.5	$2009 \ 03 \ 07$	1		83
SN 2009bb	IcBL	NGC 3278	43.5	$2009 \ 03 \ 19$	17	Υ	84
SN 2009dd	II	NGC 4088	13.9	$2009\ 04\ 12$	3		6, 85
SN 2009eo	IIn	Leda 53491	212	$2009\ 04\ 13$	1		86
SN 2009fs	IIn	UGC 11205	256	$2009 \ 06 \ 01$	1		87
SN 2009gj	IIb	NGC 134	16.8	$2009 \ 05 \ 31$	3	Υ	88
SN 2009hd	II	NGC 3627	9.6	$2009 \ 06 \ 19$	1		6
SN 2009ip	IIn	NGC 7259	28.1	$2009 \ 09 \ 13$	5	Υ	89
SN 2009kn	IIn	MCG-03-21-06	74.6	$2009\ 10\ 11$	1		90
SN 2009mk	IIb	ESO 293-34	20.3	$2009\ 12\ 15$	4		91
SN 2010O	Ib	NGC 3690	53.1	$2010 \ 01 \ 24$	2		92
SN 2010P	?	NGC 3690	53.1	$2010 \ 01 \ 10$	7	Υ	92
SN 2010ah	IcBL	Anon J114403+5541	230	$2010 \ 02 \ 21$	4		93
SN 2010al	IInPec	UGC 4286	80.6	$2010 \ 03 \ 07$	1		94
SN 2010as	IIb	NGC 6000	27.4	$2010 \ 03 \ 16$	10	Υ	60, 95
SN 2010ay	IcBL	Anon J123527+2704	314	$2010\ 02\ 22$	3		96
SN 2010bh	IcBL	Anon J071031-5615	276	$2010 \ 03 \ 16$	7	Υ	97
SN 2010br	Ib/c	NGC 4051	13.7	$2010 \ 04 \ 10$	1		98
PTF10vgv	IcBL	2MASX J22160156 + 4052065	63.8	$2010 \ 09 \ 13$	1		99
SN 2010id	II	NGC 7483	74.1	$2010 \ 09 \ 15$	1		100
SN 2010jl	IIn	UGC 5189A	53.3	$2010 \ 10 \ 01$	11	Υ	101
SN 2010jp	IIn	Anon J061630-2124	44.8	$2010\ 11\ 13$	2		102
SN 2010kp	II	Anon J040341+7045	22.3	$2010\ 11\ 30$	2		6, 103
PTF10abyy	II	galaxy unknown	133	$2010\ 12\ 06$	1		104
SN 2011cb	IIb	IC 5249	36.0	$2011 \ 04 \ 18$	4	Υ	60, 105
SN 2011dh	IIb	M51	7.9	$2011 \ 05 \ 31$	16	Υ	106
PTF11iqb	IIn	NGC 151	55.1	$2011 \ 07 \ 20$	1		107
PTF11qcj	IcBL	Leda 2295826	135	2011 10 08	20	Υ	108
SN 2011ei	II	NGC 6925	28.7	$2011 \ 07 \ 25$	11	Υ	109
SN 2011hp	Ic	NGC 4219	22.1	$2011 \ 11 \ 04$	1		110
SN 2011hs	IIb	IC 5267	21.3	$2011 \ 11 \ 06$	10	Υ	111
SN 2011ja	IIP	NGC 4945	4.2	2011 12 12	2	Y	112
SN 2012A	IIP	NGC 3239	9.7	2012 01 07	2		6
SN 2012ap	IcBL	NGC 1729	53.6	$2012 \ 02 \ 05$	3	Y	113
SN 2012au	Ib	NGC 4790	22.9	2012 03 03	3	Y	114
SN 2012aw	IIP	NGC 3351	10.0	2012 03 15	9	Y	115
PTF 12gzk	Ic	SDSS J221241.53+003042.7	63.4	2012 07 24	3	Y	116
SN 2013df	IIb	NGC 4414	18.1	$2013 \ 06 \ 04$	5	Y	117
SN 2013ej	IIP	NGC 628	8.6	2013 07 28	2	Y	6
SN 2013fs	IIP	NGC 7610	53.4	2013 10 06	2		118
SN 2013ge	Ib/c	NGC 3287	15.4	2013 11 07	3		119
iPTF13bvn	lb f	NGC 5806	24.7	2013 06 16	2		120
SN 2014C	IIn^{J}	NGC 7331	13.4	$2013\ 12\ 30$	14	Υ	121
SN 2014ad	IcBL	Mrk 1309	28.9	$2014 \ 03 \ 09$	6		122
SN 2014bc	IIP	NGC 4258	14.1	$2014 \ 04 \ 08$	1		6, 123
SN 2014bi	IIP	NGC 4096	11.5	$2014 \ 04 \ 22$	2		6, 123
SN 2014eh	Ic	NGC 6907	51.8	$2014 \ 10 \ 29$	1		124
AT 2014 ge	Ib	NGC 4080	15.5	$2014 \ 09 \ 26$	5	Υ	125
SN 2015G	Ibn	NGC 6951	23.1	$2015 \ 02 \ 27$	3		126

Table 2 (continued)

SN Name	$_{\mathrm{Type}}a$	Galaxy	Distance	Explosion	Number of	Detected	${}^{\rm References}e$
			$(D; \mathrm{Mpc})$	$_{ m date}c$	$_{\rm measurements}d$		
SN 2015J	IIn	Anon J073505-6907	24.1	$2015 \ 04 \ 26$	5	Y	60, 127
iPTF15eqv	IIb/Ib	NGC 3430	26.5	$2015 \ 08 \ 18$	4		128
ASASSN-15oz	IIL	HIPASS J1919-33	34.6	$2015 \ 08 \ 27$	2	Υ	129
PSN J22460504-1059484	Ib	NGC 7371	41.4	$2015 \ 07 \ 10$	1	Υ	130
PSN J14102342-4318437	Ib	NGC 5483	18.5	$2015\ 12\ 03$	1	Υ	131
SN 2016aqf	II	NGC 2101	16.1	$2016\ 02\ 24$	2	Υ	132
SN 2016bas	IIb	ESO 163-11	42.4	$2016 \ 03 \ 02$	8	Υ	60, 133
SN 2016bau	Ib	NGC 3631	10.3	$2016 \ 03 \ 12$	2	Υ	6, 134
SN 2016coi	IcBL	UGC 11868	18.1	$2016 \ 05 \ 24$	7	Υ	135
SN 2016cvk	IIn-pec	ESO 344-21	50.4	$2016 \ 06 \ 13$	1		136
SN 2016gfy	II	NGC 2276	20.4	$2016 \ 09 \ 10$	1		6
Spirits 16tn	?	NGC 3556	10.0	$2016 \ 05 \ 05$	2		137
SN 2017ahn	II	NGC 3318	39.8	$2017 \ 02 \ 08$	1		138
SN 2017eaw	IIP	NGC 6946	5.6	$2017 \ 05 \ 12$	4	Υ	139
SN 2017gax	Ib/c	NGC 1672	11.8	$2017 \ 08 \ 12$	1		140
SN 2018ec	Ic	NGC 3256	40.3	$2017 \ 12 \ 27$	1		60
SN 2018ie	IcBL	NGC 3456	70.6	$2018 \ 01 \ 05$	1		141
SN 2018if	IcBL	SDSS J091423.85+493533.4	141	$2018 \ 01 \ 19$	1		141
SN 2018bvw	IcBL	SDSS J115244.11+254027.1	258	$2018 \ 04 \ 25$	4	Υ	142
SN 2018cow	Icpec	CGCG 137-068	72.7	$2018 \ 06 \ 16$	7	Υ	143
SN 2018gep	IcBL	SDSS J164348.22+410243.3	144	$2018 \ 09 \ 09$	3	Υ	144
SN 2018lab	II	IC 2163	21.0	$2018\ 12\ 29$	1		145
SN 2019eez	II	NGC 2207	22.3	$2019 \ 04 \ 26$	1		146
SN 2019ehk	Ib	NGC 4321	16.2	$2019 \ 04 \ 28$	5		147
SN 2019ejj	II	ESO 430-20	11.5	$2019 \ 04 \ 29$	1		146
SN 2019esa	IIn	ESO 035-18	25.9	$2019 \ 05 \ 05$	1		146
SN 2019fcn	II	ESO 430-20	11.5	$2019 \ 05 \ 03$	1		146
SN 2019mhm	IIP	NGC 6753	50.6	$2019\ 10\ 09$	1		148
SN 2019qar	Ib/c-pec	NGC 7083	48.5	$2019 \ 09 \ 10$	1		149
SN 2020ad	II	IC 4351	28.8	$2019\ 12\ 03$	1		150
SN 2020oi	Ic	NGC 4321	16.2	$2020 \ 01 \ 07$	9	Υ	151
SN 2020bvc	IcBL	UGC 09379	122	$2020 \ 02 \ 04$	2	Υ	152
SN 2020fqv	Ib/c	NGC 4568	21.0	$2020 \ 03 \ 31$	1		153
SN 2020fsb	II	ESO 515-04	35.2	$2020\ 04\ 02$	1		153
SN 2020llx	II	NGC 7140	46.7	2020 05 29	1		154

 Table 2 continued

Table 2 (continued)

$_{(D; \text{ Mpc})} \text{date}^c \text{measurements}^d$	SI	N Name	Type ^a	Galaxy	Distance	Explosion	Number of	Detected	References
					$(D; \mathrm{Mpc})$	$_{ m date}c$	$_{\rm measurements}d$		

^a The Type of the SN. "BL" stands for "broad-lined", "Pec" for "peculiar", a ":" means the Type is somewhat uncertain, and "?" means the SN Type is unknown, because no optical spectrum was available. We do not include the unknown-Type SNe in either our I b/c or II groups.

^b The (luminosity) distance to the SN, derived from the NED database (see text for details).

^c The explosion date, t_0 , is taken from the literature. If the maximum-light time is known, but there is no other estimate of the explosion date, we take t_0 to be two weeks prior to maximum light. If maximum light time is also not known we use the discovery date for t0, in most of these cases, the radio observations occur only several months later and the exact value of t0 will have relatively little effect.

^d The number of measurements refers to those used in this work. For each SN we picked one of 4-8 GHz (C-band) or 8-12 GHz (X-band), whichever had more or better measurements, with the exception of SN 1987A where we picked 2.3 GHz.

^e References: 1 Weiler et al. (1986); 2 Weiler et al. (1991); Montes et al. (2000); 3 Weiler et al. (1992); 4 Montes et al. (1998); 5 Weiler et al. (1989); 6 re-reduced archival data; 7 Yin (1994); 8 Soderberg (2007); 9 van Dyk et al. (1996d); 10 Sramek et al. (1984); 11 Panagia et al. (1986); 12 van Dyk et al. (1998); 13 Montes et al. (1997); 14 Weiler et al. (1990); 15 Bietenholz et al. (2002); Bietenholz & Bartel (2017a); 16 Turtle et al. (1987); 17 van Dyk et al. (1993b); Williams et al. (2002); 18 van Dyk et al. (1993a); 19 Soderberg et al. (2006b); 20 van Dyk et al. (1996b); 21 Bartel et al. (2002); 22 measurements retrieved from the website of the late Kurt W. Weiler; 23 Weiler et al. (2011); 24 Chandra et al. (2009); 25 van Dyk et al. (1996c); 26 Stockdale et al. (2009f); 27 van Dyk et al. (1996a); 28 Bauer et al. (2008); 29 Lacey et al. (1998); 30 van Dyk et al. (1999); 31 Kulkarni et al. (1998); Wieringa et al. (1999); 32 Lacey et al. (1999); 33 Berger et al. (2003); 34 Alberdi et al. (2006); Pérez-Torres et al. (2009); 35 Schinzel et al. (2009), interpolated between the measured 22 GHz and 5 GHz values; 36 Stockdale et al. (2004); 37 Bietenholz & Bartel (2005, 2007b); 38 Stockdale et al. (2007); Chandra et al. (2002); 39 Ryder et al. (2004); 40 Berger et al. (2002); 41 Beswick et al. (2005); 42 Bietenholz et al. (2014); 43 Soderberg et al. (2005); 44 Soderberg et al. (2006a); 45 Stockdale et al. (2003); 46 Beswick et al. (2004); 47 Wellons et al. (2012); 48 Nayana et al. (2018); 49 Martí-Vidal et al. (2007); 50 Kankare et al. (2014); 51 Stockdale et al. (2005); 52 Drout et al. (2013); 53 Smith et al. (2017), and Charles Kilpatrick, private communication; 54 Chandra & Soderberg (2007a); 55 Dwarkadas et al. (2016); 56 (Soderberg et al. 2006c); 57 Argo (2007); 58 Kelley et al. (2006); 59 Argo et al. (2007); Bietenholz & Bartel (2007a, 2008a,b); 60 this paper; 61 Chandra et al. (2012); 62 Stritzinger et al. (2009); 63 Salas et al. (2013); 64 Soderberg et al. (2010); 65 Chandra & Soderberg (2007b); 66 Chandra & Soderberg (2008c); 67 van der Horst et al. (2011); Roy et al. (2013); 68 Chandra & Soderberg (2008b); 69 Soderberg et al. (2008); Bietenholz et al. (2009); 70 Chandra & Soderberg (2008e); 71 Chandra & Soderberg (2008d); 72 Argo et al. (2008); Stockdale et al. (2008c); Roming et al. (2009); 73 Soderberg & Chandra (2008); 74 Stockdale et al. (2008e); 75 Chandra & Soderberg (2008a); 76 Stockdale et al. (2008b,a); 77 Soderberg (2008); 78 Stockdale et al. (2009c); 79 Stockdale et al. (2008d, 2009a); 80 Chandra & Soderberg (2009a); 81 Marchili et al. (2010); Brunthaler et al. (2010); Kimani et al. (2016); 82 Stockdale et al. (2009c,d); 83 Chandra & Soderberg (2009d); 84 Bietenholz et al. (2010b); 85 Stockdale et al. (2009b); 86 Chandra & Soderberg (2009e); 87 Chandra & Soderberg (2009b); 88 Stockdale et al. (2009e); 89 Margutti et al. (2014); 90 Chandra & Soderberg (2009c); 91 Ryder et al. (2010b); 92 Romero-Cañizales et al. (2014); 93 Corsi et al. (2011); 94 Chandra et al. (2010); 95 Ryder et al. (2010a); 96 Sanders et al. (2012); 97 Margutti et al. (2013); 98 van der Horst et al. (2010); 99 Corsi et al. (2012); 100 Kasliwal et al. (2010b); 101 Chandra et al. (2015); 102 Smith et al. (2012); 103 Kasliwal et al. (2010a); 104 Kasliwal et al. (2010c); 105 Ryder et al. (2011a); 106 Krauss et al. (2012); torial, (2013); 102 United and (2014); 103 International Control, 104 International Control, 105 Inter (2014); 116 Horesh et al. (2013c); 117 Kamble et al. (2016a); Perez-Torres et al. (2015b); 118 Yaron et al. (2017); 119 Drout et al. (2016); 120 Kamble & Soderberg (2013); Horesh et al. (2013a); 121 Margutti et al. (2017); Bietenholz et al. (2018); 122 Marongiu et al. (2019); 123 Bietenholz (2017); 124 Kamble et al. (2014); 125 Chandra et al. (2019); 126 Shivvers et al. (2017); 127 Ryder et al. (2015); 128 Milisavljevic et al. (2017); 129 Bostroem et al. (2019); 130 Kamble et al. (2015); 131 Hancock & Horesh (2016); 132 Ryder et al. (2016c); 133 Ryder et al. (2016a); 134 Kamble et al. (2016b); 135 Argo et al. (2016); Terreran et al. (2019); 136 Ryder et al. (2016b); 137 Jencson et al. (2018); 138 Ryder et al. (2017); 139 Argo et al. (2017a,b); 140 Bannister et al. (2017); 141 Corsi et al. (2018); 142 Ho et al. (2020b); 143 Dobie et al. (2018a,b,c); Margutti et al. (2019) 144 Ho et al. (2019); 145 Ryder et al. (2019b); 146 Ryder et al. (2019a); 147 Jacobson-Galán et al. (2020); 148 Kundu & Ryder (2019); 149 Ryder et al. (2019c); 150 Kundu et al. (2020a); 151 Horesh et al. (2020); 152 Ho et al. (2020a); 153 Ryder et al. (2020); 154 Kundu et al. (2020b)

^f SN 2001em and SN 2014C were initially classified as Type Ic and Ib, respectively, but both developed the spectral characteristics of a Type IIn later in their evolution. Since the bright radio emission occurred at later times corresponding to the IIn spectra, we classify both as IIn

3.1. Observed Radio Lightcurves

We plot the observed values in the form of radio lightcurves (i.e., spectral luminosity curves), including any upper limits, for all our SNe with known Types in Figure 2. We then also separate the SNe by Type, and restrict our sample to those SNe at D < 100 Mpc (except as noted below), and plot values for Type I b/c SNe in Figure 3, those for Type II SNe (excluding IIn's) in Figure 4, and those for Type IIn SNe in Figure 5.

The subtype IIb seem to have brighter radio emission than the remainder of the Type II's, and we plot the Type IIb's separately from the other Type II's in Figure 6.

Finally we plot the values for the "broad-lined" (BL) TYpe Ic SNe separately from the remainder of the Type

I b/c in Figure 7. Since SNe-BL are rare, and we have only 6 detected examples at D < 100 Mpc, we plot all 27 BL SNe in our sample, regardless of D.

A number of things are apparent from these figures. First, it can be seen that the lightcurves vary over a large range. $L_{\rm pk}$ can vary over more than 5 orders of magnitude, from ~ 10^{29} erg s⁻¹ Hz⁻¹ for SN 1998bw (Kulkarni et al. 1998), which is associated with GRB 980425, and SN 2009bb (Bietenholz et al. 2010b), to < 10^{24} erg s⁻¹ Hz⁻¹ for SN 1987A (Turtle et al. 1987). Similarly, some SNe, such as SN 1987A peak at $t_{\rm pk} \leq 2$ d, while others such as SN 1986J have $t_{\rm pk} > 1000$ d (Bietenholz et al. 2002), almost 3 orders of magnitude larger.



Figure 2. A plot of the measurements and lightcurves for all the 289 SNe in our sample with known SN Types. We plot the spectral luminosity against the time since the explosion, with both axes being logarithmic. The 129 Type I b/c SNe are plotted in red. The 53 SNe of Types IIn are plotted in green, and 107 remaining Type II's in blue. The lines are lightcurves in the case of multiple detections, while the round points are single detections, and the pale triangles show upper limits. The lines do not show any fit, they just connect the available measurements. All measurements were between 4 and 10 GHz with the exception of SN 1987A. For SN 1987A, which is the lowest-luminosity curve in the plot, there were only very few early measurements available above 2.3-GHz, and we therefore use the more complete 2.3 GHz lightcurve.

It can also be seen that the lightcurves exhibit a wide variety of forms. While generally they do show an initial rise and a subsequent decay of approximately power-law form, various "bumps" and changes in the slope of the power-law decay are seen.

Figure 2 shows that Type I b/c (red) reach the highest peak luminosities, followed by the Type IIn (green), while those of Type II SNe (blue) are lower. Type I b/c's are more likely to peak earlier, while the Type II's are likely to peak later and the Type IIn even later. This pattern has been noted earlier, for example in Weiler et al. (2002), but with only a relatively small sample of SNe. While we only have a single example detected at a low value of $L_{\rm pk} < 10^{24}$ erg s⁻¹ Hz⁻¹ (SN 1987A, at D only ~50 kpc), which was of Type II, the distribution of upper limits for Type I b/c SNe is not obviously different than that for Type II's, implying that low values of $L_{\rm pk} < 10^{25}$ erg s⁻¹ Hz⁻¹ likely occur for both Type I b/c and II SNe. Type IIb SNe tend to have higher values of $L_{\rm pk}$ than the remainder of the Type II's, and are therefore more likely to be detected. The Type Ic-BL SNe also tend to have high values of $L_{\rm pk}$ but note that some Ic-BL SNe, such as SN 2002ap and SN 2014ad have fairly low values of $L_{\nu} \lesssim 10^{25.5}$ erg s⁻¹ Hz⁻¹.

4. LIGHTCURVE MODELING



Figure 3. As Figure 2, but showing only the 110 SNe of Type I b/c and showing only those at D < 100 Mpc. Different colors are used for the different SNe. The lightcurves for several SNe are labeled in the corresponding colors.

As mentioned, in our model for the lightcurves, the spectral luminosity, L_{ν} , of the SN rises to a peak, and then decays in a power-law fashion with $L_{\nu} \propto t^{\beta}$, where we take $\beta = -1.5$. Our model has only two free parameters, $t_{\rm pk}$, the time from the explosion to reach the peak, and $L_{\rm pk}$, the peak spectral luminosity.

The rise in the lightcurve is caused by an optical depth, τ , which decreases as a function of time. This optical depth could be due to either external free-free absorption or internal synchrotron self-absorption, or a combination of the two. The peak in the lightcurve occurs approximately when $\tau = 1$. We take $\tau \propto t^{-\delta}$ and $\delta = 1$, which is a value which fits most SNe moderately well, although we explore different rise parameterizations in Section 7.4 below.

We fixed the slope of the power-law decay at $\beta = -1.5$ for all SNe. Different well-observed SNe do in fact show different values of β : For example, SN 1993J has a flatter decay particularly during the first ~1000 d (Figure 1 and Bartel et al. 2002), while SN 1986J shows a steeper decay (Bietenholz & Bartel 2017b). However, for our purposes, an average value of $\beta = -1.5$ gives a reasonable fit near the peak of the lightcurve.

Our model lightcurve, normalized so that it reaches $L_{\rm pk}$ at $t_{\rm pk}$, therefore has the form

$$L(t) = L_{\rm pk} \cdot 4.482 \cdot e^{-1.5(t_{\rm pk}/t)} \cdot (t/t_{\rm pk})^{-1.5}.$$

As can be seen in Figure 2, the lightcurves of individual SNe are often more complex than our simple model. However, our model gives an adequate fit to the peak in the lightcurve, and thus serves our purpose here of providing an approximate, but sufficient, parameterization of SN lightcurves in general.

While more complex models are certainly warranted for studying individual SNe, and would likely yield more accurate values for $t_{\rm pk}$ and $L_{\rm pk}$, our purpose here is to examine the distribution of $t_{\rm pk}$ and $L_{\rm pk}$ over all SNe, so the approximate values obtained from our simple model



Figure 4. As Figure 2, but showing only the 106 SNe of Type II (excluding Type IIn) at D < 100 Mpc. Different colors are used for the different SNe. The lightcurves for several SNe, including SN 1987A, are labeled in the corresponding colors. The lightcurve for SN 1987A is at 2.3 GHz, unlike all the others which are at 4 to 10 GHz.

are adequate. In particular, the fitted distributions of $t_{\rm pk}$ and $L_{\rm pk}$ depend only very weakly on the choice of parameterization for the rise and fall of the lightcurve, so even in cases where the shape of the actual lightcurve differs from the model, our fitted values for $t_{\rm pk}$ and $L_{\rm pk}$ should be adequate to our purpose.

In cases where we have many measurements, clearly those near to $t_{\rm pk}$ provide the best constraints on $t_{\rm pk}$ and $L_{\rm pk}$. Values that are either much earlier or much later than $t_{\rm pk}$ and well below $L_{\rm pk}$ provide little additional constraint on $t_{\rm pk}$ and $L_{\rm pk}$, and could drive the fitted values to deviate from the peak in the actual lightcurve in cases where our model is not a good match for the actual lightcurve shape. To minimize this effect, for any given SN, we downweight any measurements that are at < 10% of the observed peak by treating them as upper limits. Note that we downweight measurements in this way only in cases where we have better measurements available for the same SN, that is those with $> 10 \times$ higher flux density. The effect of this is two-fold: firstly any "bumps" in the lightcurve that happen well below the peak have little effect on our fitted values of $t_{\rm pk}$ and $L_{\rm pk}$, and secondly, it serves to smooth the likelihood function in the $t_{\rm pk}$ - $L_{\rm pk}$ plane slightly, which reduces the effect of our relatively coarse sampling in this plane.

An example of this can be seen in the case of SN 1993J, where the slope of the decay changes. Figure 1 shows the full set of 8.4-GHz measurements for SN 1993J, while the left panel of Figure 9 below shows the values that we used to fit $t_{\rm pk}$ and $L_{\rm pk}$ in this case, with the flux-densities < 10% of the peak treated as upper limits.

4.2. Estimates of t_{pk} and L_{pk}

For many of our SNe, particularly if only upper limits were obtained, the measurements do not determine a unique set of values of $t_{\rm pk}$ and $L_{\rm pk}$. Instead, some



Figure 5. As Figure 2, but showing only the 41 SNe of Type IIn at D < 100 Mpc. Different colors are used for the different SNe. The lightcurves for a few SNe are labeled in the corresponding colors. Since we treat values below 10% of the peak as upper limits (see Section 3), the trailing part of the SN 1986J lightcurve is indicated here as upper limits, though in fact the spectral luminosities are well measured (Bietenholz & Bartel 2017b).



Figure 6. As Figure 2, but showing only the 19 SNe of Type IIb at D < 100 Mpc. Different colors are used for the different SNe. The lightcurves for several SNe are labeled in the corresponding colors.

ranges of values are allowed and others excluded. In order to establish the *distribution* of $t_{\rm pk}$ and $L_{\rm pk}$ over our sample, we proceed in a Bayesian fashion as follows. We define a 2-dimensional array of possible values of $t_{\rm pk}$ and $L_{\rm pk}$. We choose logarithmically spaced values of $t_{\rm pk}$ and $L_{\rm pk}$ in view of the large range these quantities can take on. Then, for each SN, we calculate the likelihood of



Figure 7. As Figure 2, but showing only the 27 broad-lined (BL) SNe (including those at D > 100 Mpc). Different colors are used for the different SNe. We highlight the limits for SN 2014ad (Marongiu et al. 2019) with larger, lime-green triangles, since for that SN the measurements constrain the radio emission to low levels over a wide range of times.

obtaining the flux-density measurements for that SN as a function of $t_{\rm pk}$ and $L_{\rm pk}$ (assuming the distance given in Table 2). If the likelihood is high for some particular pair of values $t_{\rm pk}$ and $L_{\rm pk}$, then a lightcurve characterized by those values of $t_{\rm pk}$ and $L_{\rm pk}$ represents a good fit to the measurements of the spectral luminosity.

Some values of $t_{\rm pk}$ and $L_{\rm pk}$ are un-physical: the frequency at which the spectrum turns over due to synchrotron self-absorption (SSA) depends only on the luminosity and size of the source. A lower limit on the size of the source can therefore be estimated from the observing frequency and value of $L_{\rm pk}$ (see Chevalier & Fransson 2006). Assuming a spherical source, this size can be expressed as a radius, which we call the SSA-radius, $r_{\rm SSA}$. In the case that absorbing mechanisms other than SSA are active, for instance free-free absorption (FFA) in the CSM, the turnover frequency could be higher, so the source could be larger, but not smaller than calculated assuming only SSA, so r_{SSA} , is a lower limit on the physical radius. The speed, $v_{\rm SSA} = r_{\rm SSA}/t_{\rm pk}$ is therefore a lower limit on the source's expansion speed. Projection effects do allow apparent velocities somewhat larger than c in the case of relativistic SNe, as were observed in SN 2003dh / GRB 030329 (Pihlström et al. 2007), but highly superluminal values are not expected. To exclude physically unlikely cases where highly relativistic expansion would be required, we therefore assign a likelihood of 0 to all points in the t_{pk} , L_{pk} plane for which $v_{\rm SSA} > 2c$. Although we use a non-relativistic calculation for $v_{\rm SSA}$, which will not provide accurate values when $v_{\rm SSA} \gtrsim c$, our cut at $v_{\rm SSA} > 2c$ should nonetheless serve to exclude the majority of the physically unlikely combinations of $t_{\rm pk}$ and $L_{\rm pk}$. (Indeed, there are no welldetermined values of $t_{\rm pk}$ and $L_{\rm pk}$ in this part of the plane.)

We show three examples of these likelihood arrays in Figure 8, and three examples of the possible lightcurves in Fig 9.

The first example is for a well-sampled case like SN 1993J (e.g., Bartel et al. 2002), Figure 8 left. The many luminosity measurements allow for only one specific fit of our model, which narrowly constrains the possible pairs of values of $t_{\rm pk}$ and $L_{\rm pk}$ and only one specific pair, corresponding to a single pixel in the $t_{\rm pk}$, $L_{\rm pk}$ plane, has a significantly non-zero likelihood. Only a single lightcurve fits the measurements in Figure 9 left.

The second example is for a supernova with only a single detection like PSN J22460504-1059484 (Kamble et al. 2015), shown in Figure 8 center. In this case many lightcurves are possible, all of them going through the sole luminosity measurement but some having the measured luminosity on the rising part and some on the falling part of the model lightcurve. In this case the allowed pairs of values of $t_{\rm pk}$ and $L_{\rm pk}$ are constrained to a thin curve. A family of related lightcurves, all passing through the single measurement, fit in Figure 9 center.

The third example is for a case where only one single upper limit of a luminosity measurement is available, like for SN 2017gax (Bannister et al. 2017), shown in Figure 8 right. Here the range of lightcurves with high likelihood is the largest, with many points in the t_{pk} - L_{pk} plane having almost the same high likelihood, but still a portion of the plane is excluded. A range of lightcurves, constrained only by having to go below the observed limit, fit in Figure 9 right.

5. THE RADIO LUMINOSITY-RISETIME FUNCTION, OR THE DISTRIBUTION OF $t_{\rm pk}$ AND $L_{\rm pk}$

5.1. The Distribution of the Observed Values of $t_{\rm pk}$ and $L_{\rm pk}$

We want to determine the distribution of $t_{\rm pk}$ and $L_{\rm pk}$, which is the radio luminosity-risetime function for corecollapse SNe. To guide our investigation, we start first with the subset of SNe that have well-determined values of $t_{\rm pk}$ and $L_{\rm pk}$, which is the subset of examples similar to SN 1993J in Figures 8 and 9. We adopt simple observational values of $t_{\rm pk}$ and $L_{\rm pk}$ here, where $L_{\rm pk,obs}$ is the L_{ν} corresponding to the highest measured flux density, provided that the highest value was not either the first or the last measurement, and $t_{\rm pk,obs}$ is the time since the explosion of that measurement. Note that these observational values of $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$ will generally not be identical to the values of $t_{\rm pk}$ and $L_{\rm pk}$ that have the highest likelihood from the previous section, since the latter are influenced by all the measured values, not just the single highest measurement. However, the maximum likelihood values of $t_{\rm pk}$ and $L_{\rm pk}$ should be similar to $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$. We will return below to the fitted values of $t_{\rm pk}$ and $L_{\rm pk}$, which are required for the majority of SNe for which $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$ are not determined. First however, we plot a scattergram of the observed values of $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$ in Figure 10. As already noted in Figure 2, SNe of Type I b/c (shown in red) tend to have higher values of $L_{\rm pk}$ and lower values of $t_{\rm pk}$ than do Type II.

In Figure 11 we plot the histograms showing distributions of $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$. For both, the values are scattered relatively uniformly in logarithmic space, suggesting that parameterizing the distributions of $t_{\rm pk}$ and $L_{\rm pk}$ in logarithmic space. Only for 57 SNe, (19% of our total of 294), can the values of $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$ be determined.

It is important to note that the histograms in Figure 11 represent only the population of well-observed, detected SNe, and are not representative of the overall population at D < 100 Mpc, of which 69% was never detected and 80% do not have well-defined values of $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$.

The most obvious bias is in the distribution of $L_{\rm pk,obs}$: If one were to take into account the 69% of SNe for which only upper limits on L_{ν} were ever obtained, many of them would be at $L_{\rm pk} < 10^{25}$ erg s⁻¹ Hz⁻¹, and the distribution of $L_{\rm pk,obs}$ must therefore be biased towards higher values than the distribution of $L_{\rm pk}$ over all SNe. Indeed, only for SN 1987A could a value of $\log_{10}(L_{\rm pk}) <$ 25 have been observed.

As far as $t_{\rm pk}$ is concerned, very few SNe are observed at all at times < 1 week, therefore many SNe could lie in the range $t_{\rm pk} < 10$ d, and the distribution in Figure 11 may be significantly biased here also.

5.2. The Distribution Function for L_{pk} and t_{pk} From All SNe

We now turn to incorporating the 80% of our sample for which $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$ were not defined, which includes the 69% of SNe for which only upper limits on L_{ν} could be determined. Although the observations for these SNe do not determine $t_{\rm pk}$ or $L_{\rm pk}$ uniquely, they do provide some constraints on their possible values. We incorporate them by examining the likelihood of various values of $t_{\rm pk}$ and $L_{\rm pk}$ given the observations.

In Section 4.2, we calculated the likelihoods for each SN for different pairs of values of t_{pk} and L_{pk} , with



Figure 8. Plots of the likelihood of pairs of t_{pk} and L_{pk} values for three example SNe. The likelihoods are normalized to a maximum value of unity. The horizontal axis is t_{pk} , while the vertical one is L_{pk} , and the likelihood is shown in color. The values of t_{pk} and L_{pk} that imply an apparent expansion speed > 2c are excluded, which results in the region above the dashed line, at the top left of the plots, always having zero likelihood. Left: SN 1993J, for which many measurements tightly constrain the possible values of t_{pk} and L_{pk} to a region smaller than our resolution in the t_{pk} , L_{pk} plane, and thus to a single pixel in the image, which is indicated by the red arrow. Middle: PSN J22460504-1059484, for which there was only a single measurement, but the SN was detected, thus constraining the possible locations in the t_{pk} - L_{pk} plane to the thin curved line, occupying only a small part of the plane. The pixellation of the curved region is an artefact of our relatively low resolution in the t_{pk} , L_{pk} plane, but should not significantly affect our results. Right: SN 2017gax, for which a single measurement yielded only an upper limit to the flux density. Many parts of the t_{pk} - L_{pk} plane are therefore almost equally likely.



Figure 9. Plots of the possible lightcurves for the three example SNe. The darkness of the line indicates the relative likelihood of the measurements for that particular lightcurve. The errorbars indicate the p = 68% (1 σ) confidence limits in the case of both measured values (blue) and limits (orange). The lightcurves (in mJy) are calculated using the distances given in Table 2. Left: SN 1993J. The measurements (including the limits) have a high likelihood only for a single lightcurve defined by a particular set of $t_{\rm pk}$, $L_{\rm pk}$ values. In this case, one can see that modeled lightcurves do not match the measurements precisely, with the measurements suggesting a slightly slower rise, as well as a flatter decay, especially at t > 1000 d, than our simple two-parameter model. However, the model reasonably reproduces the peak of the lightcurve. Note also that the measurements plotted as lower limits here were in fact detections (see Figure 1). As we explain in Section 3, we treat all values below 10% of the brightest observed value as upper limits so as to not unduly influence the fits near the peak. Middle: PSN J22460504-1059484. The measurements have a high likelihood for a range of related lightcurves, in some cases (with small values of $t_{\rm pk}$) placing the single measurement during the rise, and in others (with larger values of $t_{\rm pk}$) placing it during the decay. Right: SN 2017gax. The measurements have a high likelihood for a wide range of lightcurves, but nonetheless some lightcurves, e.g., those having $S_{\rm peak} \gtrsim 0.1$ mJy and $t_{\rm pk} \sim 5$ d, are excluded by the measurements.



Figure 10. The observed values of the \log_{10} of the peak spectral luminosity, $L_{pk,obs}$, against the time in days at which it occurs, $t_{\rm pk,obs}$, for all 54 SNe for which these values were determined. These values are not derived from any lightcurve fit to the measurements, but for each SN are merely the largest value of L_{ν} that was observed and the time it was observed. We do not plot SNe for which the highest observed value was the earliest (or only) one, since in those cases the peak cannot be determined. The interacting SNe, of Type IIn are shown as green triangles, while the remainder of the Type I b/c SNe are shown as red circles and the remainder of the Type II SNe are shown as blue squares. The isolated square at the lower left corner is SN 1987A. Many SNe for which only upper limits on the flux density could be determined would likely fall in the range below $L_{\rm pk}$ $< 10^{25} \text{ erg s}^{-1} \text{ Hz}^{-1}.$

examples being shown in Figure 8. If we normalize these likelihood functions, they become the probability, $p_i(t_{\rm pk}, L_{\rm pk})$, of SN number *i*, having some particular pair of $t_{\rm pk}$ and $L_{\rm pk}$ values (in Bayesian terms, this is equivalent to incorporating a flat prior for $t_{\rm pk}$ and $L_{\rm pk}$ to form the posterior probability). If we then sum these arrays over all of our SNe and divide by our total number of SNe (294), we arrive at the probability for particular pairs of values of $t_{\rm pk}$, $L_{\rm pk}$ over all of our SNe, $p_{\rm tot}(t_{\rm pk}, L_{\rm pk})$. We show $p_{\rm tot}(t_{\rm pk}, L_{\rm pk})$ in Figure 12.

The probability of different values of $t_{\rm pk}$ and $L_{\rm pk}$ is hard to interpret from Figure 8. On the one hand, there are a small number of SNe that have well-determined $t_{\rm pk}$ and $L_{\rm pk}$ (those in Figure 10 that produce a small number of high-probability pixels in Figure 8). As mentioned, these constitute an almost certainly biased subset of only 19% of our sample. On the other hand, there are many SNe for which the sparse measured values or limits mean that large areas of the $t_{\rm pk}$ - $L_{\rm pk}$ plane have low, but significantly non-zero probability. Pairs of $t_{\rm pk}$ - $L_{\rm pk}$ values which are physically unlikely, such as $\log_{10}(t_{\rm pk}) = 4$, $\log_{10}(L_{\rm pk}) = 30$, have non-zero probabil-



Figure 11. Histograms of the observed values of the \log_{10} of the time in days at which the observed peak occurred, $t_{\rm pk,obs}$, (top), and the spectral luminosity of that peak, $L_{\rm pk,obs}$ (bottom) for all 57 SNe for which these values were determined. See Figure 10 for the description of $t_{\rm pk,obs}$ and $L_{\rm pk,obs}$. These histograms represent only the population of detected, wellobserved, SNe, and are not representative of the overall population, of which 69% is not detected. In particular, the distribution of $L_{\rm pk,obs}$ is strongly biased by exclusion of SNe for which only limits on L_{ν} were obtained. The mean of $t_{\rm pk,obs}$ was 1.88, and the standard devotion was 0.67, while the corresponding values for $L_{\rm pk,obs}$ were 27.09 and 1.09.

ity because for many SNe they are not excluded by the measurements.

To proceed we want to impose some reasonable constraints on the distributions of t_{pk} and L_{pk} , for example considering extreme values unlikely even if they are allowed by our measurements. So, instead of attempting to estimate the probability distributions of $t_{\rm pk}$ and $L_{\rm pk}$ from Figure 8, we will proceed by hypothesizing some functional forms for the distributions. Although there is no physical reason to expect that the values of either $L_{\rm pk}$ or $t_{\rm pk}$ are in fact drawn from any distribution with a simple functional form, determining approximate forms of the distributions of t_{pk} and L_{pk} should prove useful until more physically-motivated versions can be found, for example, for estimating the likelihood of detecting future SNe in the radio. It also allows us to compare the distributions across different types of SNe, and may also provide some insight into the physics of radio emission from SNe.

We have noted in Section 5.1 that the values of both $t_{\rm pk}$ and $L_{\rm pk}$ seem relatively uniformly scattered in logarithmic space. A normal distribution in, for example, $t_{\rm pk}$ therefore seems incompatible with the measurements, whereas a normal distribution in $\log(t_{\rm pk})$, that is, a log-normal distribution in $t_{\rm pk}$, could provide a reasonable fit.



The logarithm of the probability, Figure 12. $\log_{10}[p_{\text{tot}}(t_{\text{pk}}, L_{\text{pk}})]$ of the measurements as a function of $t_{\rm pk}$ and $L_{\rm pk}$ over all our SNe. The maximum value of $p_{\rm tot}$ is 0.0039, which is approximately $1/(N_{\rm SNe} = 294)$. For a small number of SNe (e.g., SN 1993J, Figure 8 left), the values of t_{pk} and L_{pk} are well determined, and therefore some particular pair of values has $p_i \simeq 1$, thus contributing $1/N_{\rm SNe}$ to $p_{\rm tot}$. For other SNe, allowed values of $p_{\rm tot}$ lie on curved lines, whose peak values will be lower than $1/N_{\rm SNe}$ (since p_i must sum to 1 over the whole image). Finally, for SNe for which only upper limits are available, the maximum p_i is lower still since it is spread out more or less uniformly over the bottom part of the $t_{\rm pk}$ - $L_{\rm pk}$ plane. Although the probability for any particular t_{pk} and $L_{\rm pk}$ in this region is low, the integral of $p_{\rm tot}$ below, say, $L_{\rm pk} = 10^{25} \text{ erg s}^{-1} \text{ Hz}^{-1}$ is substantial, so that the probability of $L_{\rm pk} < 10^{25} \text{ erg s}^{-1} \text{ Hz}^{-1}$ is not negligible. Again, the region with $p_{\text{tot}} = 0$ at the top left above the dashed line is excluded because it would require strongly superluminal expansion.

We will therefore mostly work with the logarithms of $t_{\rm pk}$ and $L_{\rm pk}$, which we denote by $\log_{10}(t_{\rm pk}) = \log_{10}(t_{\rm pk}/d)$ and $\log_{10}(L_{\rm pk}) = \log_{10}(L_{\rm pk}/[{\rm erg \ s^{-1} \ Hz^{-1}}])$.

Since there is no strong correlation between the more probable values of $t_{\rm pk}$ and $L_{\rm pk}$ in Figure 12, we consider only separate distributions for $t_{\rm pk}$ and $L_{\rm pk}$, so the hypothesized joint probability for $t_{\rm pk}$ and $L_{\rm pk}$ can be obtained by multiplying their respective hypothesized probability distributions. This product would be the anticipated radio luminosity-risetime function for corecollapse SNe.

5.3. Finding the Most Likely Distribution Function for $L_{\rm pk}$ and $t_{\rm pk}$

We try therefore the following three forms for the distribution functions for $\log_{10}(t_{\rm pk})$ and $\log_{10}(L_{\rm pk})$:

1. A uniform distribution in $\log_{10}(x)$ where x is either $t_{\rm pk}$ or $L_{\rm pk}$. This distribution has two free parameters, namely the low and high limits, $x_{\rm low}$ and $x_{\rm high}$. (With well determined values of $t_{\rm pk}$ and $L_{\rm pk}$ the highest probability would be achieved by placing these limits at just below the smallest and just above the highest observed values. However, given that our measurements do not uniquely determine $t_{\rm pk}$ or $L_{\rm pk}$ in the majority of cases, the boundaries are flexible, and we determine the values of the limits that give the highest probability.)

2. A lognormal distribution in x, which is a normal distribution in $\log_{10}(x)$. This has also two free parameters, the mean, μ , and the standard deviation, σ , so the probability, $p(x) = \frac{1}{\sigma\sqrt{(2\pi)}}e^{-0.5(\frac{x-\mu}{\sigma})^2}$.

3. A power-law distribution, where $p(x) = Kx^q$ if $x > x_{\min}$ and p = 0 otherwise. This distribution also has two free parameters, namely q and x_{\min} . Given that Figures 10 and 11 suggest that both very small and large values of $t_{\rm pk}$ are unlikely, we consider the power-law distribution only for $L_{\rm pk}$, where the many lower limits means small values of $L_{\rm pk}$ could be likely.

In all cases we normalize the distributions over the ranges $-1 < \log_{10}(t_{\rm pk}) < 10^{4.5}$ (0.1 d to 86 yr) and $22 < \log_{10}(L_{\rm pk}) < 30.3 (10^{22} \text{ to } 2 \times 10^{30} \text{ erg s}^{-1} \text{ Hz}^{-1}).$

For each SN, we then multiply the likelihood function for $t_{\rm pk}$ and $L_{\rm pk}$ (Figure 8) by the hypothesized joint distribution of $t_{\rm pk}$ and $L_{\rm pk}$. The integral of this product over all possible values of $t_{\rm pk}$ and $L_{\rm pk}$ then gives the likelihood of the measurements for this SN for this particular hypothesized $t_{\rm pk}$, $L_{\rm pk}$ distribution. The likelihood of the measurements for all SNe given the hypothesized distributions of $t_{\rm pk}$ and $L_{\rm pk}$ is then the product of the likelihoods for the individual SNe.

Our first goal is to determine which functional form, i.e., lognormal, uniform, or power-law, is most appropriate for $t_{\rm pk}$ and $L_{\rm pk}$. Since our sample is almost certainly notably incomplete at larger distances, we use here only those SNe at D < 100 Mpc, where our sample is more complete, retaining 262 SNe from our total of 294.

We evaluate in a brute-force fashion the likelihood for each possible value of the four free parameters over the two distributions (two in $t_{\rm pk}$ and two in $L_{\rm pk}$; for example μ and σ in the case of a lognormal distribution). We find that the highest likelihood occurs for lognormal distributions in both $t_{\rm pk}$ and $L_{\rm pk}$. The maximumlikelihood estimate of the lognormal distribution function for $\log_{10}(t_{\rm pk})$ has mean, $\mu = 1.7$ and standard deviation $\sigma = 0.9$, while that for $L_{\rm pk}$ has $\mu = 25.5, \sigma = 1.5$. We give the values of the maximum likelihoods for other combinations of distribution functions relative to that for the best-fitting case of lognormal distributions in both $t_{\rm pk}$ and $L_{\rm pk}$, along with the associated parameter estimates in Table 3. A lognormal distribution in both $t_{\rm pk}$ and $L_{\rm pk}$ results in a significantly higher likelihood than any other combination of the three functions (lognormal, power-law, uniform) that we tried.

5.4. The Lognormal Distributions for Different SN types

Thus guided towards the use of lognormal distributions, we proceed to determine the distributions of $t_{\rm pk}$ and $L_{\rm pk}$ for various groups of SNe, to study whether different kinds of SNe are characterized by different distributions of $t_{\rm pk}$ and $L_{\rm pk}$. In addition to the maximum likelihood estimates of the means and standard deviations of the lognormal distributions, we also obtain the p = 68% points, being the points where the overall likelihood is 68% of that associated with the best-fit values. We give our results in Table 4.

Are different Types of SNe characterized by different distributions of $t_{\rm pk}$ and $L_{\rm pk}$? We have already seen from Figure 2 that Type I b/c SNe tend to have higher $L_{\rm pk}$ and shorter $t_{\rm pk}$. We split our set of SNe by Type as discussed in our introduction, and fit the distributions of $t_{\rm pk}$ and $L_{\rm pk}$ separately for the different Types. We use the following three main classes: Type I b/c, Type IIn, and the remainder of the Type II's. We also examine the subset of Type I b/c SNe that are broad-lined Type Ic (Ic-BL) and the Type IIb subset of the Type II SNe. The results are given in Table 4, and we plot the distributions in Figure 13.

Because of the completeness considerations mentioned earlier, we again consider only subsamples of SNe at D < 100 Mpc, with the exception of the rare BL subclass, where we include all examples regardless of D. Note that the first line of Table 4 represents the same fit as the first line of Table 3.

We find that the 110 Type I b/c SNe are characterized by values of $t_{\rm pk} \sim 3 \times$ lower and values of $L_{\rm pk} \sim 1.3 \times$ higher than are the 106 Type II SNe. The range of $L_{\rm pk}$ values is higher for Type I SNe (σ of $\log_{10}(L_{\rm pk})=1.7$) than for Type II's (σ of $\log_{10}(L_{\rm pk})=1.3$).

Type I b/c are over-represented in our sample, they form 42% of our sample at D < 100 Mpc, while they represent only 26% of all the SNe in the Lick Observatory Supernova Search (LOSS; Smith et al. 2011) and 19% of a complete nearby sample of 175 SNe from LOSS (Li et al. 2011). The reason for the over-representation is that Type I b/c's were more actively observed because of the potential association with GRBs.



Figure 13. The distributions of different Types of SNe in the $t_{\rm pk}$, $L_{\rm pk}$ plane. The ellipses show the p = 68% contour, i.e., the contour containing 68% of the SNe, of the relevant most-likely lognormal distributions for the various kinds of SNe (see Table 4). We illustrate the distributions for different samples of SNe as: All (dashed, black, n = 262), Type I b/c (red, n = 110), all Type II (except Type IIn; blue, n = 106), Type IIn (green, n = 41), Type IIb (mustard, n = 19), and Type Ic-BL SNe (magenta, n = 27). We use the samples at D < 100 Mpc, except for the rare Type Ic BL where we take all examples regardless of D. Note that the Type IIb and Ic BL distributions are rather uncertain because of the small number of SNe.

We ask whether the presence of the Type II SN 1987A, which was clearly unusual in a number of respects, and which, because of its low radio luminosity, could be detected because of its nearness, biases our derived distributions of $t_{\rm pk}$ and $L_{\rm pk}$? We redid the fit for the 105 Type II SNe excluding SN 1987A, and found that the best-fit distribution of $t_{\rm pk}$ and $L_{\rm pk}$ (Table 4) changed only slightly, so we can conclude that our derived distributions are not overly sensitive to the presence of the unusual SN 1987A.

We examined the 41 Type IIn SNe (at D < 100 Mpc) for which we have measurements. Type IIn SNe are associated with particularly strong radio emission. They are characterized by $\sim 31 \times$ longer $t_{\rm pk}$, and $\sim 17 \times$ higher values of $L_{\rm pk}$ than the remainder of the Type II population. We find that Type IIn SNe are also over-represented in our sample, they are 16% of our sample at D < 100 Mpc, while they represent only 9% of the core collapse SNe in the whole LOSS sample (Smith et al. 2011) and 5% of the nearby complete LOSS subsample (Li et al. 2011). The reason for the over-representation is that IIn's were probably more actively observed in the radio because of their strong association with radio emission.

Our Type II sample contained 19 SNe of Type IIb (none at D > 100 Mpc). Although this number is

Table 3. Distribution Functions for t_{pk} and L_{pk}

Distrib	pution functions a	Maximum l	ikelihood
$t_{ m pk}$	$L_{\rm pk}$	$\Delta \log_{10} p^b$	best-fit parameters
lognormal	lognormal	0	$\log_{10} t_{\rm pk} : \mu = 1.7, \sigma = 0.9; \ \log_{10} L_{\rm pk} : \mu = 25.5, \sigma = 1.5$
lognormal	power-law	-4.34	$\log_{10} t_{\rm pk}$: $\mu = 1.7, \sigma = 0.9; \log_{10} L_{\rm pk}$: min = 23.9, exponent = -1.24
uniform	lognormal	-5.11	$\log_{10} t_{\rm pk} : \min - 0.3, \max = 3.5; \log_{10} L_{\rm pk} : \mu = 25.6, \sigma = 1.5$
lognormal	log-uniform	-5.63	$\log_{10} t_{\rm pk} : \mu = 1.8, \sigma = 0.9; \ \log_{10} L_{\rm pk} : \min = 22.0, \max = 29.1$

^{*a*} The functional form of the distribution functions. Lognormal is a Normal (Gaussian) distribution in $\log_{10}(x)$, characterized by the mean, μ , and standard deviation, σ . Log-uniform is a uniform distribution in $\log_{10}(x)$. $t_{\rm pk}$ is in days, and $L_{\rm pk}$ is in erg s⁻¹ Hz⁻¹.

 b We give the log₁₀ maximum likelihood values relative to that for the best-fitting case where both distribution functions were lognormal.

too low to permit a very reliable determination of the $t_{\rm pk}$ and $L_{\rm pk}$ distributions, we did find some interesting trends. Type IIb's were much more likely to be detected than other types of SNe, with 79% being detected. The Type IIb's have values of $t_{\rm pk}$ in between those of Type I b/c and Type II, but closer to those of Type II. They have a high mean value of $L_{\rm pk}$, $36 \times$ higher than that of the remainder of the Type II's (excluding IIn's), and about $2 \times$ higher even than that of Type IIn's. The spread in the values of $L_{\rm pk}$ is considerably smaller than for other Types, σ in $\log_{10}(L_{\rm pk})$ being only 0.5.

Finally, we examined Type Ic SNe classified as broadlined, BL, which are the type associated with gammaray bursts. BL SNe are relatively rare, and only 6 were detected within D < 100 Mpc, so we take all 27 BL SNe in our database, regardless of D. They are characterized by a relatively short rise-time, with a mean $\log_{10}(t_{\rm pk})$ of only 1.2 ($\sigma = 0.6$), and a fairly high $\log_{10}(L_{\rm pk})$, with a mean of 26.7 with $\sigma = 1.7$, with the mean $\log_{10}(L_{\rm pk})$ being $\sim 20 \times$ higher than that for all Type I b/c's. However, since there were only 13 detected BL SNe in our sample, the distribution of t_{pk} and L_{pk} must be regarded as rather uncertain. We note that two unusually nearby BL SNe, SN 2002ap (Berger et al. 2002; Soderberg et al. 2006b) and SN 2014ad (Marongiu et al. 2019), were observed over a wide range of times and had $L_{8.4\text{GHz}} \lesssim 10^{25.5} \text{ erg s}^{-1} \text{ Hz}^{-1}$. Since our sample is probably biased in favour of radio-bright examples, it seems likely that $\sim 10\%$ of BL SNe have radio luminosities $< 10^{25.5}$ erg s⁻¹ Hz⁻¹, unless they have very short $t_{\rm pk}$ less than a few days.

6. MASS-LOSS RATES

Massive stars lose a significant fraction of their mass before exploding as SNe. This mass-loss is still poorly understood. An exploding SN provides a probe of this mass-loss, since the medium into which the SN shock expands is the circumstellar medium (CSM) which consists of the star's wind during the period before it exploded. The radio emission from the SN is due to the interaction of the SN ejecta with the CSM, and its brightness depends in part on the CSM density, which is a function of the mass-loss rate, \dot{M} , of the progenitor. Although the flux-density measurements provide useful direct constraints on \dot{M} of only a small fraction of well-observed SNe, we can use the distributions of $t_{\rm pk}$ and $L_{\rm pk}$ obtained in Section 5.4 to constrain the distribution of \dot{M} over our sample of SNe.

The SN shock is expected to both amplify the magnetic field and accelerate some fraction of the electrons to relativistic energies. The amount of synchrotron radio emission depends on the energy in the magnetic field as well as that in the relativistic electrons. In the absence of any absorption, the amount of synchrotron radiation can be estimated by assuming that constant fractions of the post-shock thermal energy density are transferred to magnetic fields and relativistic electrons (see, e.g., Chevalier 1982; Chevalier & Fransson 2006). The spectral luminosity, L_{ν} , at a given time will therefore depend on the CSM density at the corresponding shock radius. L_{ν} will also depend on the square of the shock speed and the volume of the emitting region. Although the shock speed and radius are measured using VLBI for some SNe (e.g., SN 1993J, Bartel et al. 2002; SN 2011dh de Witt et al. 2016; for a review see Bietenholz 2014) they are not measured for the great majority of SNe.

The post-shock energy density, at time, t when the shock has radius r, will be $\propto \rho_{\text{CSM}}(r)v^2(t)$, or in the case of a steady wind, $\propto \dot{M}v^2(t)r(t)^{-2}$. If there is equipartition between the relativistic electrons and the magnetic field, then a measurement of the spectral luminosity, L_{ν} , can be used to estimate \dot{M} , provided that a number of things are known or, in our case, can be assumed.

Table 4. Lognormal Distributions of t_{pk} and L_{pk}

Set of SNe	$N_{\rm SNe}$	$\log_{10}(p)$	$\log_{10}(p)$ Distribution of $\log_{10}(p)$		Distribution of log	$L_{10}(L_{\rm pk})$
		per measurement a	$_{\mu}b$	σ^{C}	$_{\mu}b$	σ^{C}
All $(D < 100 \text{ Mpc})$	262	-1.09	1.7 (1.6, 1.8)	0.9	25.5 (25.2, 25.7)	1.5
All $(D < 50 \text{ Mpc})$	189	-1.15	1.6 (1.5, 1.7)	0.8	25.5 (25.2, 25.7)	1.5
Type I b/c	110	-1.00	$1.1 \ (1.1, \ 1.3)$	0.5	25.4(24.8, 25.7)	1.7
Type II	106	-1.05	1.6 (1.4, 1.9)	1.0	$25.3 \ (25.0, \ 25.6)$	1.3
Type II w/o SN 1987 A	105	-1.05	1.7 (1.5, 2.0)	1.0	25.4 (25.1, 25.7)	1.2
IIn	41	-1.20	$3.1 \ (2.8, \ 4.1)$	0.7	$26.5\ (25.9,\ 27.0)$	1.1
IIb	19	-1.55	$1.5\ (1.3\ 1.7)$	0.6	$26.8 \ (26.7, \ 27.0)$	0.5
$_{\rm Broad-lined(BL)}d$	27	-1.00	$1.2 \ (0.9, \ 1.4)$	0.6	26.7 (25.9, 27.2)	1.7

^a The average \log_{10} of the probability per measurement if $t_{\rm pk}$ and $L_{\rm pk}$ are distributed with the most probable log-Gaussian distribution. This is more comparable over different numbers of SNe than the probability for all the measurements, which is expected to be lower the larger the number of measurements.

^b The mean, μ , of the normal distributions in $\log_{10}(t_{\rm pk})$ and $\log_{10}(L_{\rm pk})$, i.e. the lognormal distributions in $t_{\rm pk}$ and $L_{\rm pk}$, with the p = 68% confidence range in parenthesis following.

 c The standard deviation, $\sigma,$ corresponding to the mean values μ in the preceding column.

^d Due to the rarity of broad-lined (BL) SNe, we relax our restriction on D to include D > 100 Mpc for these SNe.

The first thing we need to assume is the wind speed of the progenitor, v_{wind} . L_{ν} actually depends on the density, which is proportional to \dot{M}/v_{wind} , rather than depending directly on \dot{M} . Type I b/c SNe generally have Wolf-Rayet progenitors, with fast, low-density winds, with $v_{\text{wind}} \sim 1000 \text{ km s}^{-1}$. Type II SNe, on the other hand, have supergiant progenitors, which generally have slow, dense winds with $v_{\text{wind}} \sim 10 \text{ km s}^{-1}$. In calculating \dot{M} , we will assume $v_{\text{wind}} \sim 1000 \text{ km s}^{-1}$ for the Type I b/c's, and $\sim 10 \text{ km s}^{-1}$ for the Type II's.

The next thing that we need to assume is the efficiency of the conversion of thermal energy to both magnetic field and relativistic particle energies. These efficiencies are usually expressed as the ratio between the energy density of the magnetic field and the relativistic particles to the post-shock thermal energy density, and we will denote the two ratios with ϵ_B and ϵ_e , respectively. Although the values are not well known, it is often assumed that $\epsilon_B \simeq \epsilon_e$ (equipartition), and that both are ~ 0.1 . We will here also assume $\epsilon_B = \epsilon_e = 0.1$, and we note that our values of \dot{M} must remain somewhat speculative, but we hope nonetheless instructive. We discuss the uncertainty in deriving \dot{M} from radio lightcurves further in Sec. 7.4 below.

Finally, the volume of the emitting region and the speed of the shock also needs to be known or assumed. In the case of Type I b/c SNe, the absorption producing the rising part of the lightcurve is most often synchrotron self-absorption (SSA). In this case the absorption is internal to the emitting region, and $t_{\rm pk}$ and $L_{\rm pk}$ allow an estimate of the radius at time $t_{\rm pk}$ (as noted

already in Section 4.2). If we assume the emitting region to be a spherical shell with outer radius 26% larger than the inner one, then the filling factor is 0.5, which is considered typical. We will assume f = 0.5. With these assumptions, Chevalier & Fransson (2006) and Soderberg et al. (2012) find that

$$\dot{M} = 1.1 \times 10^{-7} \left(\frac{0.1}{\epsilon_B}\right) \left(\frac{\epsilon_e}{\epsilon_B}\right)^{-8/19} \cdot \left(\frac{L_{\rm pk}}{10^{26} \, {\rm erg \, s^{-1} \, Hz^{-1}}}\right)^{-4/19} \left(\frac{t_{\rm pk}}{\rm d}\right)^2 \, {\rm M}_{\odot} \, {\rm yr}^{-1}$$
(1)

where we have recast the equation given in Soderberg et al. (2012) for our nominal frequency of 8.4 GHz, and taken $v_{\rm wind} = 1000 \text{ km s}^{-1}$.

This equation, however, is only applicable if the spectral energy distribution (SED) is dominated by SSA. As can be seen in Figs. 3 and 10, Type I b/c SNe show a wide range of lightcurve behaviors. In particular, for ones which are slow-rising and faint, the rise cannot be reproduced by SSA without assuming expansion velocities too low to be believable. In those cases, therefore, there is likely significant FFA absorption. In the presence of FFA, the radius and velocity implicit in the above calculation are only lower limits. Most Type I b/c SNe show expansion velocities of > 30,000 km s⁻¹ (Chevalier 2007). For any Type I b/c SN where $t_{\rm pk}$ and $L_{\rm pk}$ imply $v_{\rm SSA} < 20,000 \ {\rm km \ s^{-1}}$, the assumption of an SSA-dominated lightcurve is problematic, and eq. 1 therefore not applicable, and the speed of the shock volume of the emitting region must be estimated in some fashion other than from SSA.

Eq. 1 is also not applicable for Type II SNe, where the absorption is generally dominated by FFA. However, Type II SNe seem to be characterized by a relatively narrow range in expansion velocity: de Jaeger et al. (2019) found that for the 51 Type II SNe from the Berkeley sample, the standard deviation of expansion velocity measured from the H α line at t = 10 d was only 19%, suggesting a fairly narrow range of velocities ⁴.

Since the spread in the velocity for Type II SNe is not large, we assume a single representative value for all Type II SNe. Weiler et al. (2002) give expressions in this case, which are based on assuming a self-similar evolution of the SN, with $r \propto t^m$, where *m* is called the deceleration parameter, as well as assuming a single characteristic initial expansion velocity for all SNe of a particular Type (I or II). Although the latter assumption is demonstrably poor for rapidly-expanding Type I b/c SNe, which have speeds ranging up to *c*, it is probably reasonable for the slower Type I b/c's, i.e., those with $v_{\rm SSA} < 20,000 \text{ km s}^{-1}$, as well as for Type II's, which generally do not show high expansion velocities.

Although Weiler et al. (2002) take m = 1 for Type II SNe, the observations do not necessarily bear this out, with some Type II SNe showing substantial deceleration, for example, m = 0.781 for SN 1993J between ages of ~1.5 and ~5 yr (Bartel et al. 2002) and m = 0.69 observed for SN 1986J (Bietenholz et al. 2010a). Overall, the Type II SNe show a similar range of values of m as do the few Type I b/c's for which we have reliable estimates of m, and therefore we assume m = 0.8 for both Type I and Type II SNe. Following Chevalier (1982), we find in this case that

$$L_{\nu} \propto (1-m)(\dot{M}/v_{\text{wind}})^{(p-7+12m)/4} \cdot m^{(5+p)/2} t^{-(p+5-6m)/2} \nu^{-(p-1)/2}.$$
 (2)

where p is the energy index of the relativistic electron population. Representative values of p for Type I b/c and II SNe are 3 and 2.4, respectively (Weiler et al. 2002). Setting m = 0.8, this equation simplifies to

$$\text{Type I b/c}: L_{\nu} \propto (\dot{M}/v_{\text{wind}})^{1.4} t^{-1.6} \nu^{-1} \qquad (3)$$

and

Γype II :
$$L_{\nu} \propto (\dot{M}/v_{\text{wind}})^{1.25} t^{-1.3} \nu^{-0.7}$$
 (4)

Now we need to determine the constant of proportionality in eqs. 3 and 4, to use them to obtain $\dot{M}/v_{\rm wind}$. For Type I b/c SNe, we determine the constant by requiring that \dot{M} have the same value as that calculated using eq. 1 for a representative value of $L_{\rm pk} =$ 2×10^{26} erg s⁻¹ Hz⁻¹, and our rounded mean value of $t_{\rm pk}$ for Type I b/c SNe of 20 d (Table 4), which corresponds to $v_{\rm SSA} = 20,000$ km s⁻¹. We obtain

Type I b/c :
$$\dot{M} = 7.2 \times 10^{-7} \times \left(\frac{L_{\rm pk}}{10^{26} \text{ erg s}^{-1} \text{ Hz}^{-1}}\right)^{0.71} \left(\frac{t_{\rm pk}}{1\text{ d}}\right)^{1.14} \text{ M}_{\odot} \text{ yr}^{-1}$$
(5)

for $v_{\rm wind} = 1000 \text{ km s}^{-1}$.

For Type II SNe, we determine the constant of proportionality by using the mean values of $\log_{10}(\dot{M})$ determined from the absorption for four well observed SNe (SN 1970G, SN 1979C, SN 1980K, SN 1981K)⁵ given in Weiler et al. (2002). We obtain

Type II :
$$\dot{M} = 1.1 \times 10^{-7} \times \left(\frac{L_{\rm pk}}{10^{26} \text{ erg s}^{-1} \text{ Hz}^{-1}}\right)^{0.80} \left(\frac{t_{\rm pk}}{1\text{ d}}\right)^{1.04} \text{ M}_{\odot} \text{ yr}^{-1}$$
(6)

for $v_{\rm wind} = 10 \text{ km s}^{-1}$.

Given the distributions of $t_{\rm pk}$ and $L_{\rm pk}$ we obtained in Table 4, we can now calculate the corresponding distribution of \dot{M} for Type I b/c and Type II SNe. Since we found that a lognormal distribution was appropriate for $t_{\rm pk}$ and $L_{\rm pk}$, we determine the distribution of $\log_{10}(\dot{M})$.

We find that the mean of $\log_{10}(\dot{M} \text{ in } M_{\odot} \text{ yr}^{-1})$ for Type I b/c SNe is -5.6 ± 1.1 , assuming $v_{\text{wind}} = 1000 \text{ km s}^{-1}$. For Type II SNe, (excluding IIn), $\log_{10}(\dot{M}) = -6.8 \pm 1.4$, assuming $v_{\text{wind}} = 10 \text{ km s}^{-1}$. The progenitors and wind velocity of the Type IIn's are not well known and their \dot{M} rates are likely strongly time-variable, and therefore equation 6 will be poorly calibrated for them, so we do not extend this analysis to the Type IIn SNe.

7. DISCUSSION

Our large compilation of 1475 radio measurements of 294 SNe shows that the radio lightcurves of SNe are extremely varied. With our simple characterization of the lightcurves with only two parameters, $t_{\rm pk}$ (rise-time) and $L_{\rm pk}$ (peak spectral luminosity), we find that both $t_{\rm pk}$ and $L_{\rm pk}$ can vary over large ranges, at least 3 and 5 orders of magnitude, respectively (Figure 10). We

⁴ Expansion velocities from radio are expected to be somewhat higher than those from optical, e.g., from H α , because the former usually relate to the forward shock and the latter to expanding areas interior to it. However, the optical and radio velocities are expected to be well correlated, so a narrow range in H α suggests a correspondingly narrow range in forward shock velocities. See discussion in Bartel et al. (2007).

⁵ We omit SN 1982aa from this calculation, because no optical spectrum was ever obtained and its Type is therefore uncertain (see Section 7.3 below).

showed that a lognormal distribution was appropriate for both t_{pk} and L_{pk} .

We find that the normal distribution of $\log_{10}(t_{\rm pk})$ has a mean of 1.7, corresponding to 50 d, with σ (standard deviation) of 0.9 (line 2 in Table 4, using only SNe at D < 100 Mpc). Both quite short risetimes of 7 d and quite long ones of > 1 yr are within the range of $\pm 1\sigma$ and thus not uncommon.

We find that many SNe must be fairly faint in the radio. Indeed, to date, only about 31% of the SNe at D < 100 Mpc that have been observed in the radio were detected at all. The results published so far tend to be biased in favour of the detections or towards higher radio luminosities. If we include the many non-detections, we find that the most probable distribution of $\log_{10}(L_{\rm pk})$ in erg s⁻¹ Hz⁻¹ has mean of 25.5, corresponding to 3×10^{25} erg s⁻¹ Hz⁻¹, with $\sigma = 1.5$.

This distribution has a significantly lower mean $\log_{10}(L_{\rm pk})$, as well as a wider range, than the mean of 27.3, corresponding to 2×10^{27} erg s⁻¹ Hz⁻¹, with $\sigma = 1.25$ given by Lien et al. (2011), which was based on only 20 *detected* SNe. In fact, if we repeat the calculation from Table 4 for only those SNe with at least 3 detections, we find that the mean $\log_{10}(L_{\rm pk}) = 27.1$, close to that found by Lien et al. (2011). The inclusion of the many limits is crucial for obtaining the distribution of *all* radio SNe, not just the well-studied radio-bright ones.

We find that more than half of all SNe will have peak luminosities $< 10^{26}$ erg s⁻¹ Hz⁻¹ (at 4 to 8 GHz), corresponding to ~1 mJy at 10 Mpc Although SN 1987A is at the faint end of the distribution with $L_{\rm pk} \lesssim$ 10^{24} erg s⁻¹ Hz⁻¹, we expect ~11% of all SNe, or ~6% of Type II SNe, will be comparably faint in the radio.

7.1. Differences Between SNe of Type I b/c and II

It has long been accepted that Type Ib/c SNe tend to have more rapidly-evolving radio lightcurves, characterized by shorter values of $t_{\rm pk}$, than do Type II's. However, until the present work, this has only been asserted on the basis of relatively small numbers of SNe (e.g., Weiler et al. 2002, 2010). While we find the assertion to be true, with the values of $\log_{10}(t_{\rm pk})$ being characterized by a mean of 1.1 (13 d) for Type I b/c SNe, in comparison to 1.6 (40 d) for Type II SNe excluding Type IIn's, the caveat that must be stated here is the standard deviations in $\log_{10}(t_{\rm pk})$ were large for both Types, being 0.5 for I b/c and 1.0 for Type II. Therefore, as can also be seen in Figure 2, there is considerable overlap, with some Type I b/c SNe having very slow rise times up to several years, while some Type II SNe have short rise times of < 1 month, and SN 1987A has one of < 2 d.

We further find that the Type I b/c and Type II SNe reach a similar range of $L_{\rm pk}$ values. For our sample of SNe at D < 100 Mpc, the mean value of $\log_{10}(L_{\rm pk}$ in erg s⁻¹ Hz⁻¹) for Type I b/c's was 25.4, while that for Type II's (excluding IIn) was marginally lower at 25.3. The standard deviations for $\log_{10}(L_{\rm pk})$ were large, being 1.6 for Type I b/c's, and 1.3 for Type II's, so there is very significant overlap in the distribution of $\log_{10}(L_{\rm pk})$ (see also Figure 2). Some Type I b/c SNe have quite low values of $\log_{10}(L_{\rm pk}) \leq 25.5$, while many Type II SNe have high values of $\log_{10}(L_{\rm pk}) > 25$. We note, however, that for Type I b/c's, the standard deviation of the $\log_{10}(L_{\rm pk})$ distribution is notably higher than it is for Type II's, so both extreme high and low values of $L_{\rm pk}$ are more likely for Type I b/c's.

While it had been suggested on the basis of only four examples that Type I b/c SNe could be approximate radio standard candles (Weiler et al. 1998), our data (Figure 3) make clear that this is very much not the case, with the variation in $L_{\rm pk}$ extending over several orders of magnitude.

Our best-fit distributions of t_{pk} and L_{pk} are illustrated in Figure 13. The 41 SNe of Type IIn (at D < 100 Mpc) have higher and later radio peaks than the remainder of the Type II's, with mean values of $\log_{10}(t_{\rm pk})$ and $\log_{10}(L_{\rm pk})$ being 3.1 (corresponding to 3.5 yr) and 26.5, respectively, but the standard deviations in $\log_{10}(t_{\rm pk})$ and $\log_{10}(L_{\rm pk})$ are large, being 0.7 and 1.1, respectively, thus overlapping with the other SN Types. We note that Stockdale et al. (2007) suggested a much higher $\log_{10}(L_{\rm pk})$ of 28 for Type IIn's, but again this result was biased by not including non-detections. There is a possibility that Type IIn SNe have similar radio lightcurves to other Type II's initially, i.e., with risetimes on the order of $t_{\rm pk} = 40$ d, and relatively low values of $L_{\rm pk}$, but are characterized by luminous late-time radio emission, since there are relatively few observations of IIn's at earlier times (Figure 5).

We note again that the values of $L_{\rm pk}$ of the few SNe that have many measurements are notably higher than the mean (except for SN 1987A), being around $L_{\rm pk} \gtrsim 10^{27}$ erg s⁻¹ Hz⁻¹. The reason is that the radio SNe that have attracted the most attention are the most luminous examples, but our many upper limits show that the majority of SNe are in fact relatively faint.

7.2. The Synchrotron-Self-Absorption Expansion Velocity

As mentioned in Section 4.2, if SSA (synchrotron selfabsorption) is the dominant absorption mechanism, the emitting volume, and thus the radius, can be deduced from the frequency at which the SED peaks. Equivalently, for some particular frequency, ν , $L_{\rm pk}$ allows calculation of the source volume or radius at the time $t_{\rm pk}$. We call this radius $r_{\rm SSA}$, and the corresponding velocity $v_{\rm SSA} = r_{\rm SSA}/t_{\rm pk}$. Both $r_{\rm SSA}$ and $v_{\rm SSA}$ are just calculated from ν , $t_{\rm pk}$, and $L_{\rm pk}$, regardless of whether SSA is in fact the dominant absorption mechanism. Only if SSA *is* dominant do $r_{\rm SSA}$ and $v_{\rm SSA}$ correspond to the physical radius and speed.

For each of our SNe the measurements provide some constraint on $v_{\rm SSA}$, to the degree to which the measurements constrain $t_{\rm pk}$ and $L_{\rm pk}$. In Figure 8, we showed the likelihood of various values of $t_{\rm pk}$ and $L_{\rm pk}$ given our measurements for three example SNe. The lines of constant $v_{\rm SSA}$ are parallel to the dotted line in the top left corner showing $v_{\rm SSA} = 2c$ in the $t_{\rm pk}$ - $L_{\rm pk}$ plane. Integrating along lines of constant $v_{\rm SSA}$, we can therefore determine the probability of particular values of $v_{\rm SSA}$ given our measurements.

Referring again to the three example SNe shown in Figure 8, for some SNe, such as SN 1993J, $t_{\rm pk}$, $L_{\rm pk}$ and thus $v_{\rm SSA}$ are well determined, and only a single value of $v_{\rm SSA}$ is allowed by the measurements, while for others such as SN 2017gax, we have only very weak constraints on $v_{\rm SSA}$, and virtually any value of $v_{\rm SSA}$ can be accommodated by the (single) measurement. If we normalize the probability for each SN, over the range of $v_{\rm SSA} = 1 \text{ km s}^{-1}$ to 2c, we can determine a probability distribution of $v_{\rm SSA}$ over our collection of SNe by summing over all our SNe, giving each equal weight.

We show this distribution in Figure 14, showing separately the distributions for Type I b/c, Type II, and Type IIn SNe. We note that the probability we show is that of particular values of $v_{\rm SSA}$ given all our observations. A non-zero probability for some value of $v_{\rm SSA}$ means that value is allowed by the observations for some fraction of our SNe, but does not require that there exist any SN characterized by that value of $v_{\rm SSA}$. This is particularly true of the very low values of $v_{\rm SSA} < 100 \text{ km s}^{-1}$, which are allowed by the measurements for a significant number of our SNe, but which likely do not occur in any real SNe. Nonetheless, in the absence of concrete measurements of $v_{\rm SSA}$, (or better, the actual shock speed) for a large number of SNe, Fig 14 will give some insight into what values of $v_{\rm SSA}$ are allowed by the currently existing measurements.

Recall also that $v_{\rm SSA}$ is only a lower limit to the shock velocity (see Section 4.2): if free-free absorption dominates and the peak in the SED is not due to SSA, then both $r_{\rm SSA}$ and $v_{\rm SSA}$ are only lower limits to the physical r and v. In fact, given that the shock speeds observed in SNe are almost always larger than a few thousand km s⁻¹, much of the portion of Figure 14 be-



Figure 14. The normalized likelihood of different values of the SSA velocity, $v_{\rm SSA}$, over different samples of SNe. $v_{\rm SSA}$ is the velocity calculated from $t_{\rm pk}$ and $L_{\rm pk}$ assuming that the dominant absorption mechanism is synchrotron selfabsorption. The likelihood is that of all our measurements arising from SNe characterized by a particular value of $v_{\rm SSA}$. Values of $v_{\rm SSA}$ which may never occur in any actual SN can have non-zero likelihood if they are not disallowed by our flux-density measurements. Note that $v_{\rm SSA}$ is a *lower* limit to the shock speed. We weight each SN equally. We show $p(v_{\rm SSA})$ for SNe of Type I b/c SNe (red), II (blue), and IIn (green).

low 10^4 km s⁻¹ is likely due to cases where in fact FFA dominates, and the shock speed is larger than v_{SSA} .

Comparing the distributions of $v_{\rm SSA}$ for different types of SNe, we find the following: For Type I b/c SNe, high values of $v_{\rm SSA} > 20,000$ km s⁻¹ are the most probable, with values up to and even exceeding c occurring.⁶ It is generally accepted that Type I b/c SNe tend to have higher speeds than Type II's (see, e.g., Chevalier 1998, 2007), but this has only been concluded on the basis of much smaller numbers of SNe in the past, and we can now confirm this pattern on the basis of a much larger sample. Note also that some fraction of Type I b/c can have low $v_{\rm SSA} < 10,000$ km s⁻¹, so even in the case of Type I b/c SNe, $v_{\rm SSA}$ can be significantly lower than the shock speed.

The most probable values of $v_{\rm SSA}$ for Type II SNe are ~3000 km s⁻¹. Since this is lower than the expected shock speeds, we conclude that for the majority of Type II SNe, FFA is the dominant absorption mechanism rather than SSA, and $v_{\rm SSA}$ therefore is lower than the shock speed. Again our conclusion is in agree-

⁶ As mentioned earlier, $v_{\rm SSA} > c$ was seen in SN 2003dh, and the superluminal apparent expansion was confirmed directly by VLBI observations (e.g., Pihlström et al. 2007).

ment with statements made earlier (e.g., Chevalier 1998, 2007), but which had in the past been made only on the basis of a far smaller sample of SNe. Type IIn SNe are characterized by even lower values of $v_{\rm SSA}$.

7.3. Identifying the SN Type on the Basis of the Radio Lightcurve

Can the radio emission be used to determine the Type of an SN, for example in cases where there is no optical detection? Because of the large overlap in the lightcurves of different SN types, the radio lightcurve for any particular SN generally does not reliably indicate the SN type.

There are two exceptions where the radio lightcurve nevertheless can give a fairly reliable indication of the SN Type. First, a 8.4-GHz spectral luminosity $L_{8.4\text{GHz}} > 10^{28} \text{ erg s}^{-1} \text{ Hz}^{-1}$ in the first month seems to occur only for Type I b/c SNe. Such high, early luminosities therefore strongly suggest a Type I b/c SN. In general, very high radio luminosities of $L_{8.4\text{GHz}} > 10^{28.5} \text{ erg s}^{-1} \text{ Hz}^{-1}$ seem to occur predominantly for Type I b/c SNe regardless of age. Second, a very high luminosity at late times with $L_{\rm pk} > 10^{27} \text{ erg s}^{-1} \text{ Hz}^{-1}$ at $t_{\rm pk} > 1000 \text{ d}$, strongly suggests a Type IIn supernova.

Third, a high value of $v_{\rm SSA} > 30,000 \text{ km s}^{-1}$ (see Section 7.2) suggests that the SN is much more likely to be of Type I b/c, while values of $v_{\rm SSA} < 10,000 \text{ km s}^{-1}$ are more likely in Type II SNe.

In our database, there are five SNe of which no optical spectrum was obtained, and of which therefore the SN Type (I or II) is unknown: SNe 1982aa, 2000ft, 2008iz, and 2010P and Spirits 16tn. Can the Type be determined from radio observations alone on the basis of our distributions of $t_{\rm pk}$ and $L_{\rm pk}$? We show the radio lightcurves of these SNe in comparison to the remainder of the SNe in our sample in Figure 15.

SN 1982aa in NGC 6052 was detected in the radio and reached a very late and high peak (Yin 1994). Although the explosion date is uncertain, the values of $t_{\rm pk}$ and $L_{\rm pk}$ are fairly well determined at ~ 10^{2.5} d and ~ 10^{29.0} erg s⁻¹ Hz⁻¹, respectively. These values of $t_{\rm pk}$ and $L_{\rm pk}$ strongly suggest a Type IIn (Figures 5, 13), although the SN is exceptional regardless of Type.

SN 2000ft in NGC 7469 was detected only after the radio peak (Alberdi et al. 2006), and the explosion time is again uncertain. Although it was optically detected (Colina et al. 2007), no spectrum was obtained. The values of $t_{\rm pk}$ and $L_{\rm pk}$ are fairly well determined at ~ $10^{2.0}$ d and ~ $10^{28.1}$ erg s⁻¹ Hz⁻¹, respectively. It is quite luminous compared to the majority of SNe, but the lightcurve and the values of $t_{\rm pk}$ and $L_{\rm pk}$ are equally

SN 2008iz was detected in the radio in M82, and was never detected optically despite the close distance (3.8 Mpc), presumably because of very strong optical extinction. It has a very unusual radio lightcurve (Marchili et al. 2010; Brunthaler et al. 2010). The values of $t_{\rm pk}$ and $L_{\rm pk}$ are fairly well determined at ~ $10^{1.8}$ d and ~ $10^{27.3}$ erg s⁻¹ Hz⁻¹, respectively. It showed both an unusually slow rise and a relatively shallow decay, and seems to be showing a late-time rise after $t \simeq 1000$ d. Although a Type II has been suggested, the radio lightcurve is equally compatible with either Type I b/c or II, although $L_{\rm pk}$ was higher than the average for either Type. Its SN Type therefore also remains unknown.

SN 2010P, in Arp 299, was discovered in the infrared and subsequently detected in the radio (Kankare et al. 2014; Romero-Cañizales et al. 2014, and references therein). Infrared observations and an optical spectrum were obtained by Kankare et al. (2014). The spectrum had relatively low signal-to-noise ratio due to the high extinction, and was compatible with an SN of either Type Ib or a IIb. In this case the peak of the radio lightcurve is not well determined, and the first measurement occurred only at t = 523 d, so a wide range of t_{pk} and L_{pk} are compatible with the measurements. The likely values of $\log_{10}(t_{\rm pk})$ are between 1.2 and 2.5 and those of $\log_{10}(L_{pk})$ between 27 and 29, with the higher values of $L_{\rm pk}$ occurring in conjunction with earlier values of $t_{\rm pk}$. While Kankare et al. (2014) suggest that the radio evolution precludes a Type Ib, we find (see Figure 15) that, when compared to our broad sample. SN 2010P's radio evolution is not inconsistent with that seen in some Type I b/c's. It is, however, more luminous than the mean of any of our SN Types. Although the optical spectrum rules out a normal Type II, whether SN 2010P was of Type Ib or IIb must remain uncertain.

Spirits 16tn was a heavily obscured SN, detected in the infrared, for which spectroscopic classification was not possible (Jencson et al. 2018). Fairly low limits of $L_{6\rm GHz} \lesssim 10^{24.3}$ erg s⁻¹ Hz⁻¹ were placed on the radio luminosity (Jencson et al. 2018). However, as can be seen from Figs. 13 and 15, such low values can occur for either Type I b/c or Type II SNe, therefore its SN Type must also remain uncertain.

7.4. Determining Mass Loss Rates from Radio Emission

From the distribution of t_{pk} and L_{pk} , a distribution of mass-loss rates, \dot{M} , for the progenitors can be estimated. For both SN Type I b/c and Type II the mean



Figure 15. The radio lightcurves or limits of five SNe which were not spectroscopically identified, compared to those of the spectroscopically confirmed SNe of Types I b/c (red) and II (blue) and IIn (green) in our sample. SNe 1982aa, 2000ft, 2008iz, 2010P and Spirits 16tn are highlighted in violet, light green, black, magenta and maroon, respectively. Spirits 16tn was not detected, but two fairly low limits were obtained, shown as larger maroon triangles between t = 100 and 300 d. SN 1982aa is most likely of Type IIn. For SNe 2000ft, 2008iz and 2010P, and Spirits 16tn, the radio lightcurves do not permit a conclusive identification of the SN Type.

values of M are low compared to the published values for well-studied SNe. This is because the well-studied SNe tend to be substantially brighter than the average, and thus tend to have denser CSM to produce the stronger radio emission than the average. We found that the average value of $\log_{10}(M)$ for a Type I b/c SNe was $-5.6 \pm 1.1 \text{ (in } M_{\odot} \text{ yr}^{-1}; \text{ assuming } v_{\text{wind}} = 1000 \text{ km s}^{-1}),$ while the equivalent value for Type II's (excluding IIn's) was lower at -6.8 ± 1.4 (assuming $v_{\text{wind}} = 10 \text{ km s}^{-1}$). However, we caution against over-interpreting this difference, since our values of M rely on a number of assumptions (see Section 6), and there may be systematic biases dependent on the type of the SN since the massloss rates are calculated differently for Type I b/c's than for Type II's. Also, as with t_{pk} and L_{pk} , the range of values of M is quite large, with the standard deviations of $\log_{10}(M)$ over our sample being 1.1 and 1.4, respectively, so here also there is considerable overlap between Type I b/c's and Type II's.

To indicate the uncertainty in estimating the mass loss rate we give as an example SN 1993J, whose 8.4-GHz lightcurve we showed in Figure 1. Despite being one of the most intensely studied SNe with extensive, multi-frequency radio lightcurves as well as X-ray data, estimating \dot{M} seems to be far from straightforward, and various authors have reported a considerable range of $\log_{10}(\dot{M})$ for its progenitor⁷. In $\log_{10}(\dot{M}/M_{\odot} \text{ yr}^{-1})$, for $v_{\text{wind}} = 10 \text{ km s}^{-1}$, Weiler et al. (2007) reports values between -6.3 and -5.2, Björnsson (2015) reports -5.0, Weiler et al. (2002) report -4.6, and Fransson & Björnsson (1998) report -4.3. We found that the distribution of $\log_{10}(\dot{M})$ for all Type II SNe excluding Type IIn's had a mean of -6.8 and standard deviation of 1.4. The cited values for SN 1993J, are higher than the mean of the distribution, although not outlandish, with, for example, ~4% of SNe having \dot{M} higher than even the highest of the values reported for SN 1993J ($\log_{10}(\dot{M}) = -4.3$). Since SN 1993J was exceptionally radio luminous, it is not surprising that it has a denser CSM, and thus that its progenitor had a higher \dot{M} than the population average.

In the standard self-similar model of an SN, the rise of the radio lightcurve to L_{pk} is relatively rapid, regardless of whether FFA or SSA is the dominant absorption mechanism, and is followed by a slower decay. For external FFA by a uniform wind medium, with density $\propto r^{-2}$, the optical depth, τ , is expected to decay with time as $\tau \propto t^{-\delta}$ with $\delta \simeq 2.5$ (see, e.g. Weiler et al. 2002). Inspection of Figure 2 shows that such steeply rising lightcurves are not common. We used $\delta = 1$, which is more representative, for our lightcurve model. To check the dependence of our results on the assumed $\delta = 1$, we tried a model with $\delta = 2.5$ but obtained a much lower likelihood than with $\delta = 1$. We note that $\delta = 0.9$ produces a marginally higher likelihood than $\delta = 1$, but the difference is small, and our results should not be compromised by our use of $\delta = 1$.

If the rising part of the lightcurve (in a self-similar model) were due to SSA within the emitting region, rather than FFA, then the rising part would be expected to be a power-law, $L_{\nu} \propto t^a$, rather than exponential as seen with FFA. In the case of pure SSA, values of $a \sim 2.1$ are expected (Chevalier 1998). This expected value again is much steeper than the majority of the observed lightcurves in Figure 2. We fitted our data with SSA-like model lightcurves with a power-law, rather than an exponential rise. We found that a fit with a = 2.1 produces much lower likelihood than our standard model (exponential rise with $\delta = 1$). Smaller values of a produced higher likelihoods, although over the whole sample, an exponential rise with $\tau \propto t^{-1}$ pro-

⁷ We note that part of the variation in \dot{M} derives from different assumptions about the poorly-known values of the efficiencies ϵ_B and ϵ_e (Sec. 6). However, some of the values of \dot{M} , e.g., those of Weiler et al. (2007), are derived from absorption only, and independent of any assumed values of ϵ_B and ϵ_e , so the variation in derived values of \dot{M} cannot be ascribed entirely to the use of different values of ϵ_B and ϵ_e .

duced a somewhat higher likelihood than a power-law rise for any value of a.

Weiler et al. (2002) also found that absorption by a uniform wind medium cannot fit the rising portions of many SN radio lightcurves. They appeal to geometrical effects from a clumpy absorbing medium to flatten the rise and produce a better fit to the rising part of the lightcurves. Björnsson & Keshavarzi (2017) model the effect of an inhomogeneous synchrotron-emitting region in SSA-dominated SNe, and find the effect is to flatten the part of the SED below the peak, which would also tend to make the rise in the lightcurve less steep.

From our much larger collection of measurements we can conclude that the slowly rising lightcurves are a fairly general phenomenon and that therefore some form of geometrical effect, such as clumpiness in the CSM or inhomogeneity in the synchrotron-emitting region, are common.

7.5. Structure of a Comprehensive Radio SN Observing Program

Our sample is of necessity heterogeneous and incomplete, with only a fraction of SNe being observed at all in the radio, and even if observed, often with very sparse sampling in time. While a census of southern SNe is one of the goals of the Variables and Slow Transients survey (VAST; Murphy et al. 2013), to be conducted with the Australian Square Kilometre Array Pathfinder at 1.4 GHz, as well as of the ThunderKAT transients programme underway at MeerKAT (Fender et al. 2016), a systematic multi-frequency program of observing SNe in the radio would be desirable to obtain a clearer picture of the radio SN luminosity-risetime function. What would such a program entail?

We found the mean $\log_{10}(L_{\rm pk})$ for our sample of all Types of core-collapse SNe was 25.5 (in erg s⁻¹ Hz⁻¹) with a standard deviation of 1.6 (Table 4). This mean $\log_{10}(L_{\rm pk})$ corresponds to a flux density of 30 μ Jy at 30 Mpc.

To achieve a 5σ detection of 30 µJy requires ~30 min at 10 GHz or ~3 h at 1.5 GHz with the VLA, ~4 h at 10 GHz or ~7 h at 2 GHz with ATCA, and ~2 h at 1.3 GHz with MeerKAT. Note that this is only the mean $\log_{10}(L_{\rm pk})$, so at this sensitivity level ~50% of SNe would remain undetected. On the Transient Name Server⁸, the rate of classified SNe with D < 30 Mpc is ~16 yr⁻¹ over the whole sky. Observations to fully sample the luminosity distribution of radio SNe will therefore be challenging with current instrumentation. Given the wide range of $t_{\rm pk}$ that we have found, with the 1- σ range being from 7 d to 1 yr, an observing program with at least 7 logarithmically-spaced observations of each SN starting after about one week and extending to at least t = 1 yr would be required to get reasonably complete sampling and provide accurate constraints on $t_{\rm pk}$ and $L_{\rm pk}$. Obviously such a program will miss the ~17% of SNe with $t_{\rm pk} < 1$ week but observations on a shorter timescale would be hard to schedule.

A systematic program to provide more robust statistics with a more complete sample than we have been able to do with the existing ad-hoc sample would therefore be a challenging and long-term project with current instrumentation, but would certainly be an important project with the Square Kilometre Array, whose sensitivity will greatly surpass that of current instruments (Perez-Torres et al. 2015a). Notwithstanding the difficulty of obtaining a complete sample with current instrumentation, it is still well worthwhile to observe nearby or unusual SNe on a case-by-case basis, and we encourage observers to publish non-detections.

8. CONCLUSION

We examined a large number of radio flux density measurements for 294 SNe at between 5 and 10 GHz. We parameterize the radio lightcurves by a simple model consisting of an optically-thick rise over time $t_{\rm pk}$ from the explosion, to a maximum value of the spectral luminosity, $L_{\rm pk}$, followed by a power-law decay with $L_{\nu} \propto t^{-1.5}$. We concentrate here only on the part of the lightcurve near the initial peak, and disregard any late-time rises in flux density, such as observed in, e.g., SN 1987A.

We find that both $t_{\rm pk}$ and $L_{\rm pk}$ vary over large ranges. In the case of $t_{\rm pk}$, some SNe (such as SN 1987A) had $t_{\rm pk}$ of a couple of days or even less, while others (such as SN 1986J) do not reach the peak until $t_{\rm pk} \gtrsim 1000$ d.

The range in $L_{\rm pk}$ is even larger: SN 1987A reached $L_{\rm pk}$ of only $\sim 10^{23.6}$ erg s⁻¹ Hz⁻¹, while that of SN 1998bw was $\sim 10^{29}$ erg s⁻¹ Hz⁻¹.

1. We find that, over our sample of SNe, lognormal distributions of t_{pk} and L_{pk} provide a reasonable fit to the measurements, including the many upper limits.

2. Many SNe in our sample have low values of $L_{\rm pk}$. At 8.4 GHz, 50% of all SNe have $L_{\rm pk} < 10^{25.5}$ erg s⁻¹ Hz⁻¹ or flux densities $< 30 \ \mu$ Jy at D = 30 Mpc.

3. The median value of $L_{\rm pk}$ is ~30 times lower than that obtained if one does not consider the many upper limits in addition to the detections.

4. For Type I b/c SNe at D < 100 Mpc, the mean value and standard deviation of $t_{\rm pk}$ were $10^{1.1\pm0.5}$ d, and those

 $^{^8}$ https://wis-tns.weizmann.ac.il; we looked at the SNe listed between 2020 Jan 1 and June 30, of which 8 had D<30 Mpc.

of $L_{\rm pk}$ were $10^{25.4\pm1.7}$ erg s⁻¹ Hz⁻¹. Type I b/c SNe are characterized by more rapid rises than are Type II's, but they reach similar values of $L_{\rm pk}$.

5. For Type II SNe, at D < 100 Mpc and excluding Type IIn's, the mean value and standard deviation of $t_{\rm pk}$ were $10^{1.6\pm1.0}$ d, and those of $L_{\rm pk}$ were $10^{25.3\pm1.3}$ erg s⁻¹ Hz⁻¹.

6. Type IIn SNe are characterized by long risetimes, $10^{3.1\pm0.7}$ d but high values of $L_{\rm pk}$ of $10^{26.5\pm1.1}$ erg s⁻¹ Hz⁻¹.

7. Type IIb SNe seem to be characterized by considerably higher $L_{\rm pk}$ than the remainder of the Type II's of $10^{26.8\pm0.5}$ erg s⁻¹ Hz⁻¹, and also a narrower range of $L_{\rm pk}$ than other Types. However, our sample contained only 19 Type IIb SNe, so this distribution is somewhat uncertain.

8. In general, given the wide distributions, the values of $t_{\rm pk}$ and $L_{\rm pk}$ for any particular SN do not reliably indicate whether the SN is of Type I b/c or II.

9. The exception to item 8. above is that $L_{\nu} > 10^{28}$ erg s⁻¹ Hz⁻¹ in the first month strongly suggests a Type I b/c SN.

10. From the distribution of $t_{\rm pk}$ and $L_{\rm pk}$ values we estimated also the distribution of mass-loss rates, \dot{M} . We found that for Type I b/c SNe, $\dot{M} = 10^{-5.6\pm1.1} \,\mathrm{M_{\odot} yr^{-1}}$, while for Type II SNe excluding Type IIn's, $\dot{M} = 10^{-6.8\pm1.4} \,\mathrm{M_{\odot} yr^{-1}}$, for assumed $v_{\rm wind}$ of 1000 km s⁻¹ and 10 km s⁻¹, respectively. We caution, however, that the determination of \dot{M} from $t_{\rm pk}$ and $L_{\rm pk}$ is very imprecise, and possibly subject to biases that could be dependent on the SN Type.

11. We find that the rising part of the lightcurves is in most cases too shallow to be described either by synchrotron self-absorption (SSA) or free-free absorption in a uniform medium. This relative flatness suggests that geometrical effects, such as a clumpy CSM or non-spherically symmetric structure in the ejecta or the CSM, are likely common among SNe.

ACKNOWLEDGEMENTS

This research was supported by both the National Research Foundation of South Africa and the National Sciences and the Engineering Research Council of Canada. In addition to the authors, Mekuanint Hailemariam and Nceba Mhlalo reduced some archival VLA data, which has contributed to this paper. The first author thanks Richard Grumitt for helpful discussions, and we thank the anonymous referee for his or her helpful suggestions. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. The Australia Telescope Compact Array (ATCA) is part of the Australia Telescope National Facility, which is funded by the Australian Government for operation as a National Facility managed by CSIRO. We acknowledge the Gomeroi people as the traditional owners of the ATCA Observatory site. (e-)MERLIN is a National Facility operated by the University of Manchester at Jodrell Bank Observatory on behalf of the Science and Technology Facilities Council of the UK. We have made use of NASA's Astrophysics Data System Abstract Service.

Software: AIPS (Associated Universities 1999), CASA (International Consortium Of Scientists 2011), e-Merlin Pipeline (Argo 2014)

REFERENCES

- Alberdi, A., Colina, L., Torrelles, J. M., et al. 2006, ApJ, 638, 938
- Argo, M. 2007, in From Planets to Dark Energy: the Modern Radio Universe, 82
- Argo, M. 2014, e-MERLIN data reduction pipeline, ascl:1407.017
- Argo, M., Torres, M. P., Beswick, R., & Wrigley, N. 2017a, The Astronomer's Telegram, 10421
- —. 2017b, The Astronomer's Telegram, 10472
- Argo, M. K., Beswick, R. J., Muxlow, T. W. B., & Pedlar, A. 2007, The Astronomer's Telegram, 1084, 1
- Argo, M. K., Beswick, R. J., Muxlow, T. W. B., et al. 2008, The Astronomer's Telegram, 1469, 1

- Argo, M. K., Romero-Canizales, C., Beswick, R., & Prieto, J. L. 2016, The Astronomer's Telegram, 9147, 1
- Associated Universities, I. 1999, AIPS: Astronomical Image Processing System, ascl:9911.003
- Bannister, K., Corsi, A., & Sand, D. 2017, The Astronomer's Telegram, 10660, 1
- Bartel, N., & Bietenholz, M. F. 2008, ApJ, 682, 1065
- Bartel, N., Bietenholz, M. F., Rupen, M. P., & Dwarkadas, V. V. 2007, ApJ, 668, 924
- Bartel, N., Bietenholz, M. F., Rupen, M. P., et al. 2002, ApJ, 581, 404
- Bauer, F. E., Dwarkadas, V. V., Brandt, W. N., et al. 2008, ApJ, 688, 1210

- Berger, E., Kulkarni, S. R., & Chevalier, R. A. 2002, ApJL, 577, L5
- Berger, E., Kulkarni, S. R., Frail, D. A., & Soderberg, A. M. 2003, ApJ, 599, 408
- Beswick, R. J., Fenech, D., Thrall, H., et al. 2005, IAUC, 8572, 1
- Beswick, R. J., Muxlow, T. W. B., Argo, M. K., & Pedlar, A. 2004, IAUC, 8332, 2
- Bietenholz, M. 2014, in 12th European VLBI Network Symposium and Users Meeting (2014), published by SISSA, Trieste, ed. A. Tarchi, M. Giroletti, & L. Feretti, 51
- Bietenholz, M., & Bartel, N. 2007a, The Astronomer's Telegram, 1254, 1
- —. 2008a, The Astronomer's Telegram, 1525, 1
- —. 2008b, The Astronomer's Telegram, 1657, 1
- —. 2014, The Astronomer's Telegram, 6429, 1
- Bietenholz, M. F., & Bartel, N. 2005, ApJL, 625, L99
- —. 2007b, ApJL, 665, L47
- —. 2017a, ApJ, 839, 10
- —. 2017b, ApJ, 851, 7
- Bietenholz, M. F., Bartel, N., & Rupen, M. P. 2002, ApJ, 581, 1132
- —. 2010a, ApJ, 712, 1057
- Bietenholz, M. F., De Colle, F., Granot, J., Bartel, N., & Soderberg, A. M. 2014, MNRAS, 440, 821
- Bietenholz, M. F., Kamble, A., Margutti, R., Milisavljevic, D., & Soderberg, A. 2018, MNRAS, 475, 1756
- Bietenholz, M. F., Soderberg, A. M., & Bartel, N. 2009, ApJL, 694, L6
- Bietenholz, M. F., Soderberg, A. M., Bartel, N., et al. 2010b, ApJ, 725, 4
- Björnsson, C. I. 2015, ApJ, 813, 43
- Björnsson, C. I., & Keshavarzi, S. T. 2017, ApJ, 841, 12
- Bostroem, K. A., Valenti, S., Horesh, A., et al. 2019, MNRAS, 485, 5120
- Brunthaler, A., Marti-Vidal, I., Menten, K. M., et al. 2010, in 10th European VLBI Network Symposium and EVN Users Meeting: VLBI and the New Generation of Radio Arrays, Vol. 10, 55
- Bufano, F., Pignata, G., Bersten, M., et al. 2014, MNRAS, 439, 1807
- Cano, Z., Wang, S.-Q., Dai, Z.-G., & Wu, X.-F. 2017, Advances in Astronomy, 2017, 8929054
- Cendes, Y., Gaensler, B. M., Ng, C. Y., et al. 2018, ApJ, 867, 65
- Chakraborti, S., Ray, A., Smith, R., et al. 2013, ApJ, 774, 30
- Chakraborti, S., Soderberg, A., Chomiuk, L., et al. 2015, ApJ, 805, 187

- Chandra, P., Chevalier, R. A., Chugai, N., et al. 2012, ApJ, 755, 110
- Chandra, P., Chevalier, R. A., Chugai, N., Fransson, C., & Soderberg, A. M. 2015, ApJ, 810, 32
- Chandra, P., Nayana, A. J., Björnsson, C. I., et al. 2019, ApJ, 877, 79
- Chandra, P., Ray, A., & Bhatnagar, S. 2002, IAUC, 7982, 2
- Chandra, P., & Soderberg, A. 2007a, The Astronomer's Telegram, 1182, 1
- —. 2007b, The Astronomer's Telegram, 1271, 1
- —. 2008a, The Astronomer's Telegram, 1869, 1
- —. 2008b, The Astronomer's Telegram, 1366, 1
- —. 2008c, The Astronomer's Telegram, 1359, 1
- —. 2008d, The Astronomer's Telegram, 1409, 1
- —. 2008e, The Astronomer's Telegram, 1410, 1
- —. 2009a, The Astronomer's Telegram, 1891, 1
- —. 2009b, The Astronomer's Telegram, 2070, 1
- —. 2009c, The Astronomer's Telegram, 2335, 1
- —. 2009d, The Astronomer's Telegram, 2351, 1
- —. 2009e, The Astronomer's Telegram, 2358, 1
- Chandra, P., Soderberg, A., Chevalier, R., Fransson, C., & Chugai, N. 2010, The Astronomer's Telegram, 2532, 1
- Chandra, P., Stockdale, C. J., Chevalier, R. A., et al. 2009, ApJ, 690, 1839
- Chevalier, R. A. 1982, ApJ, 259, 302
- -. 1998, ApJ, 499, 810
- Chevalier, R. A. 2007, in Revista Mexicana de Astronomia y Astrofisica, vol. 27, Vol. 30, Revista Mexicana de Astronomia y Astrofisica Conference Series, 41–48
- Chevalier, R. A., & Fransson, C. 2006, ApJ, 651, 381
- Colina, L., Díaz-Santos, T., Alonso-Herrero, A., et al. 2007, A&A, 467, 559
- Corsi, A., Ho, A., & Kulkarni, S. 2018, The Astronomer's Telegram, 11295, 1
- Corsi, A., Ofek, E. O., Frail, D. A., et al. 2011, ApJ, 741, 76
- Corsi, A., Ofek, E. O., Gal-Yam, A., et al. 2012, ApJL, 747, L5
- de Jaeger, T., Zheng, W., Stahl, B. E., et al. 2019, MNRAS, 490, 2799
- de Witt, A., Bietenholz, M. F., Kamble, A., et al. 2016, MNRAS, 455, 511
- Dobie, D., Ravi, V., Ho, A., Kasliwal, M., & Murphy, T. 2018a, The Astronomer's Telegram, 11795
- Dobie, D., Ravi, V., Ho, A., Kasliwal, M., & Murphy, T. 2018b, The Astronomer's Telegram, 11818
- Dobie, D., Ravi, V., Ho, A., Kasliwal, M., & Murphy, T. 2018c, The Astronomer's Telegram, 11862
- Drout, M. R., Soderberg, A. M., Mazzali, P. A., et al. 2013, ApJ, 774, 58

- Drout, M. R., Milisavljevic, D., Parrent, J., et al. 2016, ApJ, 821, 57
- Dwarkadas, V. V., Romero-Cañizales, C., Reddy, R., & Bauer, F. E. 2016, MNRAS, 462, 1101
- Fender, R., Woudt, P. A., Corbel, S., et al. 2016, in MeerKAT Science: On the Pathway to the SKA, 13
- Fransson, C., & Björnsson, C.-I. 1998, ApJ, 509, 861
- Hancock, P., & Horesh, A. 2016, The Astronomer's Telegram, 8504, 1
- Ho, A. Y. Q., Goldstein, D. A., Schulze, S., et al. 2019, ApJ, 887, 169
- Ho, A. Y. Q., Kulkarni, S. R., Perley, D. A., et al. 2020a, ApJ, 902, 86
- Ho, A. Y. Q., Corsi, A., Cenko, S. B., et al. 2020b, ApJ, 893, 132
- Horesh, A., Cao, Y., Mooley, K., & Carpenter, J. 2013a, The Astronomer's Telegram, 5198, 1
- Horesh, A., Kasliwal, M., Carpenter, J., et al. 2011, The Astronomer's Telegram, 3512, 1
- Horesh, A., Stockdale, C., Fox, D. B., et al. 2013b, MNRAS, 436, 1258
- Horesh, A., Kulkarni, S. R., Corsi, A., et al. 2013c, ApJ, 778, 63
- Horesh, A., Sfaradi, I., Ergon, M., et al. 2020, ApJ, 903, 132
- International Consortium Of Scientists. 2011, CASA:
- Common Astronomy Software Applications, ascl:1107.013 Jacobson-Galán, W. V., Margutti, R., Kilpatrick, C. D.,
- et al. 2020, ApJ, 898, 166 Jencson, J. E., Kasliwal, M. M., Adams, S. M., et al. 2018, ApJ, 863, 20
- Kamble, A., Margutti, R., Milisavljevic, D., Soderberg, A., & Parrent, J. 2015, The Astronomer's Telegram, 7845, 1
- Kamble, A., & Soderberg, A. 2013, The Astronomer's Telegram, 5154, 1
- Kamble, A., Soderberg, A., Margutti, R., Parrent, J., & Milisavljevic, D. 2014a, The Astronomer's Telegram, 6724
- Kamble, A., Soderberg, A. M., Chomiuk, L., et al. 2014b, ApJ, 797, 2
- Kamble, A., Margutti, R., Soderberg, A. M., et al. 2016a, ApJ, 818, 111
- Kamble, A., Margutti, R., Alexander, K., et al. 2016b, The Astronomer's Telegram, 8911
- Kankare, E., Fraser, M., Ryder, S., et al. 2014, A&A, 572, A75
- Kasliwal, M., Frail, D., Quimby, R., et al. 2010a, The Astronomer's Telegram, 3090, 1
- Kasliwal, M., Kulkarni, S., Arcavi, I., et al. 2010b, The Astronomer's Telegram, 2864, 1
- Kasliwal, M. M., Frail, D., Quimby, R., et al. 2010c, The Astronomer's Telegram, 3093, 1

- Kelley, M. T., Stockdale, C. J., Sramek, R. A., et al. 2006, Central Bureau Electronic Telegrams, 495, 1
- Kimani, N., Sendlinger, K., Brunthaler, A., et al. 2016, A&A, 593, A18
- Krauss, M. I., Soderberg, A. M., Chomiuk, L., et al. 2012, ApJL, 750, L40
- Kulkarni, S. R., Frail, D. A., Wieringa, M. H., et al. 1998, Nature, 395, 663
- Kundu, E., & Ryder, S. 2019, The Astronomer's Telegram, 13040, 1
- Kundu, E., Ryder, S. D., Filipovic, M. D., et al. 2020a, The Astronomer's Telegram, 13477, 1
- -. 2020b, The Astronomer's Telegram, 13805, 1
- Lacey, C. K., Van Dyk, S. D., Weiler, K. W., et al. 1999, IAUC, 7336, 2
- Lacey, C. K., Weiler, K. W., Sramek, R. A., & van Dyk, S. D. 1998, IAUC, 7068, 2
- Li, W., Leaman, J., Chornock, R., et al. 2011, MNRAS, 412, 1441
- Lien, A., Chakraborty, N., Fields, B. D., & Kemball, A. 2011, ApJ, 740, 23
- Lundqvist, P., Kundu, E., Pérez-Torres, M. A., et al. 2020, ApJ, 890, 159
- Marchili, N., Martí-Vidal, I., Brunthaler, A., et al. 2010, A&A, 509, A47
- Margutti, R., Soderberg, A. M., Wieringa, M. H., et al. 2013, ApJ, 778, 18
- Margutti, R., Milisavljevic, D., Soderberg, A. M., et al. 2014, ApJ, 780, 21
- Margutti, R., Kamble, A., Milisavljevic, D., et al. 2017, ApJ, 835, 140
- Margutti, R., Metzger, B. D., Chornock, R., et al. 2019, ApJ, 872, 18
- Marongiu, M., Guidorzi, C., Margutti, R., et al. 2019, ApJ, 879, 89
- Martí-Vidal, I., Marcaide, J. M., Alberdi, A., et al. 2011, A&A, 526, A142
- —. 2007, A&A, 470, 1071
- Milisavljevic, D., Margutti, R., Soderberg, A. M., et al. 2013, ApJ, 767, 71
- Milisavljevic, D., Margutti, R., Kamble, A., et al. 2015, ApJ, 815, 120
- Milisavljevic, D., Patnaude, D. J., Raymond, J. C., et al. 2017, ApJ, 846, 50
- Montes, M. J., Van Dyk, S. D., Weiler, K. W., Sramek,
 R. A., & Panagia, N. 1997, ApJL, 482, L61
 —. 1998, ApJ, 506, 874
- Montes, M. J., Weiler, K. W., Van Dyk, S. D., et al. 2000, ApJ, 532, 1124

- Mould, J. R., Huchra, J. P., Freedman, W. L., et al. 2000, ApJ, 529, 786
- Murphy, T., Chatterjee, S., Kaplan, D. L., et al. 2013, PASA, 30, 6
- Nayana, A. J., Chandra, P., & Ray, A. K. 2018, ApJ, 863, 163
- Palliyaguru, N. T., Corsi, A., Frail, D. A., et al. 2019, ApJ, 872, 201
- Panagia, N., Sramek, R. A., & Weiler, K. W. 1986, ApJL, 300, L55
- Perez-Torres, M., Alberdi, A., Beswick, R. J., et al. 2015a, in Advancing Astrophysics with the Square Kilometre Array (AASKA14), 60
- Perez-Torres, M., Argo, M., Martí-Vidal, I., et al. 2015b, The Astronomer's Telegram, 8452, 1
- Pérez-Torres, M. A., Alberdi, A., Colina, L., et al. 2009, MNRAS, 399, 1641
- Pihlström, Y. M., Taylor, G. B., Granot, J., & Doeleman, S. 2007, ApJ, 664, 411
- Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020, A&A, 641, A6
- Romero-Cañizales, C., Herrero-Illana, R., Pérez-Torres, M. A., et al. 2014, MNRAS, 440, 1067
- Roming, P. W. A., Pritchard, T. A., Brown, P. J., et al. 2009, ApJL, 704, L118
- Roy, R., Kumar, B., Maund, J. R., et al. 2013, MNRAS, 434, 2032
- Ryder, S., Kundu, E., Marnoch, L., Chomiuk, L., & Sarbadhicary, S. 2019a, The Astronomer's Telegram, 12820, 1
- Ryder, S., Covarrubias, R., Amy, S., et al. 2010a, Central Bureau Electronic Telegrams, 2242, 1
- Ryder, S. D., Amy, S. W., Stockdale, C. J., et al. 2010b, The Astronomer's Telegram, 2450, 1
- —. 2011a, The Astronomer's Telegram, 3370, 1
- Ryder, S. D., Kool, E., Stockdale, C. J., & Kotak, R. 2015, The Astronomer's Telegram, 7762, 1
- Ryder, S. D., Kool, E. C., Filipovic, M., et al. 2019b, The Astronomer's Telegram, 12373
- Ryder, S. D., Kool, E. C., Stockdale, C. J., & Kotak, R. 2016a, The Astronomer's Telegram, 8836, 1
- Ryder, S. D., Kool, E. C., Stockdale, C. J., et al. 2016b, The Astronomer's Telegram, 9475
- —. 2017, The Astronomer's Telegram, 10147, 1
- Ryder, S. D., Kool, E. C., Stockdale, C. J., Romero-Canizales, C., & Kotak, R. 2016c, The Astronomer's Telegram, 8815, 1
- Ryder, S. D., Kundu, E., Filipovic, M. D., et al. 2020, The Astronomer's Telegram, 13642, 1

- Ryder, S. D., Kundu, E., Velovic, V., et al. 2019c, The Astronomer's Telegram, 13136, 1
- Ryder, S. D., Sadler, E. M., Subrahmanyan, R., et al. 2004, MNRAS, 349, 1093
- Ryder, S. D., Stockdale, C. J., Immler, S., et al. 2011b, The Astronomer's Telegram, 3764, 1
- Salas, P., Bauer, F. E., Stockdale, C., & Prieto, J. L. 2013, MNRAS, 428, 1207
- Sanders, N. E., Soderberg, A. M., Valenti, S., et al. 2012, ApJ, 756, 184

Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in Astronomical Society of the Pacific Conference Series, Vol. 77, Astronomical Data Analysis Software and Systems IV, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes, 433

- Schinzel, F. K., Taylor, G. B., Stockdale, C. J., Granot, J., & Ramirez-Ruiz, E. 2009, ApJ, 691, 1380
- Shivvers, I., Zheng, W., Van Dyk, S. D., et al. 2017, MNRAS, 471, 4381
- Smith, N., Li, W., Filippenko, A. V., & Chornock, R. 2011, MNRAS, 412, 1522
- Smith, N., Cenko, S. B., Butler, N., et al. 2012, MNRAS, 420, 1135
- Smith, N., Kilpatrick, C. D., Mauerhan, J. C., et al. 2017, MNRAS, 466, 3021
- Soderberg, A. 2007, PhD thesis, California Institute of Technology
- —. 2008, The Astronomer's Telegram, 1811, 1
- Soderberg, A., & Chandra, P. 2008, The Astronomer's Telegram, 1470, 1
- Soderberg, A. M., Brunthaler, A., Nakar, E., Chevalier, R. A., & Bietenholz, M. F. 2010, ApJ, 725, 922
- Soderberg, A. M., Chevalier, R. A., Kulkarni, S. R., & Frail, D. A. 2006a, ApJ, 651, 1005
- Soderberg, A. M., Kulkarni, S. R., Berger, E., et al. 2005, ApJ, 621, 908
- Soderberg, A. M., Nakar, E., Berger, E., & Kulkarni, S. R. 2006b, ApJ, 638, 930
- Soderberg, A. M., Kulkarni, S. R., Nakar, E., et al. 2006c, Nature, 442, 1014
- Soderberg, A. M., Berger, E., Page, K. L., et al. 2008, Nature, 453, 469
- Soderberg, A. M., Margutti, R., Zauderer, B. A., et al. 2012, ApJ, 752, 78
- Sramek, R. A., Panagia, N., & Weiler, K. W. 1984, ApJL, 285, L59
- Stockdale, C. J., Kelley, M., van Dyk, S. D., et al. 2005, IAUC, 8603, 2

- Stockdale, C. J., Kelley, M. T., Weiler, K. W., et al. 2007, in American Institute of Physics Conference Series, Vol. 937, Supernova 1987A: 20 Years After: Supernovae and Gamma-Ray Bursters, ed. S. Immler, K. Weiler, & R. McCray, 264–268
- Stockdale, C. J., Sramek, R. A., Van Dyk, S. D., Weiler, K. W., & Panagia, N. 2003, IAUC, 8153, 2
- Stockdale, C. J., Van Dyk, S. D., Sramek, R. A., et al. 2004, IAUC, 8282, 2
- Stockdale, C. J., Weiler, K. W., Immler, S., et al. 2008a, The Astronomer's Telegram, 1484, 1
- -. 2008b, The Astronomer's Telegram, 1477, 1
- —. 2008c, The Astronomer's Telegram, 1439, 1
- —. 2008d, The Astronomer's Telegram, 1883, 1
- Stockdale, C. J., Weiler, K. W., Soderberg, A., et al. 2008e, The Astronomer's Telegram, 1452, 1
- Stockdale, C. J., Weiler, K. W., Immler, S., et al. 2009a, The Astronomer's Telegram, 1912, 1
- 2009b, The Astronomer's Telegram, 2016, 1
- —. 2009c, The Astronomer's Telegram, 1915, 1
- Stockdale, C. J., Heim, M. S., Weiler, K. W., et al. 2009d, The Astronomer's Telegram, 1925, 1
- Stockdale, C. J., Rentz, B., Vandrevala, C. M., et al. 2009e, IAUC, 9056, 1
- Stockdale, C. J., Heim, M. S., Vandrevala, C. M., et al. 2009f, Central Bureau Electronic Telegrams, 1714, 1
- Stritzinger, M., Philips, M. M., Morrell, N., Salgado, F., & Folatelli, G. 2009, Central Bureau Electronic Telegrams, 1751, 1
- Terreran, G., Margutti, R., Bersier, D., et al. 2019, ApJ, 883, 147
- Turtle, A. J., Campbell-Wilson, D., Bunton, J. D., et al. 1987, Nature, 327, 38
- van der Horst, A. J., Kamble, A. P., Paragi, Z., et al. 2010, The Astronomer's Telegram, 2612, 1
- —. 2011, ApJ, 726, 99
- van Dyk, S. D., Lacey, C. K., Sramek, R. A., & Weiler, K. W. 1999, IAUC, 7322, 2
- van Dyk, S. D., Montes, M. J., Weiler, K. W., Sramek, R. A., & Panagia, N. 1998, AJ, 115, 1103
- van Dyk, S. D., Sramek, R. A., Montes, M. J., Weiler, K. W., & Panagia, N. 1996a, IAUC, 6528, 1
- van Dyk, S. D., Sramek, R. A., Weiler, K. W., Montes, M. J., & Panagia, N. 1996b, IAUC, 6378, 2

- van Dyk, S. D., Sramek, R. A., Weiler, K. W., et al. 1996c, IAUC, 6375, 1
- van Dyk, S. D., Sramek, R. A., Weiler, K. W., & Panagia, N. 1993a, ApJ, 409, 162
- van Dyk, S. D., Weiler, K. W., Sramek, R. A., & Panagia, N. 1993b, ApJL, 419, L69
- van Dyk, S. D., Weiler, K. W., Sramek, R. A., et al. 1996d, AJ, 111, 1271
- Weiler, K. W., Panagia, N., Montes, M. J., & Sramek, R. A. 2002, ARA&A, 40, 387
- Weiler, K. W., Panagia, N., & Sramek, R. A. 1990, ApJ, 364, 611
- Weiler, K. W., Panagia, N., Sramek, R. A., et al. 1989, ApJ, 336, 421
- -. 2010, Mem. Soc. Astron. Italiana, 81, 374
- Weiler, K. W., Panagia, N., Stockdale, C., et al. 2011, ApJ, 740, 79
- Weiler, K. W., Sramek, R. A., Panagia, N., van der Hulst, J. M., & Salvati, M. 1986, ApJ, 301, 790
- Weiler, K. W., van Dyk, S. D., Discenna, J. L., Panagia, N., & Sramek, R. A. 1991, ApJ, 380, 161
- Weiler, K. W., van Dyk, S. D., Montes, M. J., Panagia, N., & Sramek, R. A. 1998, ApJ, 500, 51
- Weiler, K. W., van Dyk, S. D., Pringle, J. E., & Panagia, N. 1992, ApJ, 399, 672
- Weiler, K. W., Williams, C. L., Panagia, N., et al. 2007, ApJ, 671, 1959
- Wellons, S., Soderberg, A. M., & Chevalier, R. A. 2012, ApJ, 752, 17
- Wieringa, M. H., Kulkarni, S. R., & Frail, D. A. 1999, A&AS, 138, 467
- Williams, C. L., Panagia, N., Van Dyk, S. D., et al. 2002, ApJ, 581, 396
- Wilson, W. E., Ferris, R. H., Axtens, P., et al. 2011, MNRAS, 416, 832
- Woosley, S. E., & Bloom, J. S. 2006, ARA&A, 44, 507
- Yadav, N., Ray, A., Chakraborti, S., et al. 2014, ApJ, 782, 30
- Yaron, O., Perley, D. A., Gal-Yam, A., et al. 2017, Nature Physics, 13, 510
- Yin, Q. F. 1994, ApJ, 420, 152
- Zanardo, G., Staveley-Smith, L., Ball, L., et al. 2010, ApJ, 710, 1515