

Central Lancashire Online Knowledge (CLOK)

Title	A comparison of the pharmacodynamic effects of intravenous ketamine-xylazine with alfaxalone in mute swans (Cygnus olor) presenting at a wildlife veterinary hospital
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/39037/
DOI	https://doi.org/10.1016/j.vaa.2021.03.014
Date	2021
Citation	Baldrey, Vicki, Stanford, Michael and Bacon, Heather (2021) A comparison of the pharmacodynamic effects of intravenous ketamine-xylazine with alfaxalone in mute swans (Cygnus olor) presenting at a wildlife veterinary hospital. <i>Veterinary Anaesthesia and Analgesia</i> . ISSN 1467-2987
Creators	Baldrey, Vicki, Stanford, Michael and Bacon, Heather

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.vaa.2021.03.014>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Table 1. Scoring system used to assess the quality of anaesthetic induction, maintenance and recovery in Mute Swans following the intravenous administration of either 10 mg kg⁻¹ alfaxalone or 0.28 mg kg⁻¹ xylazine/12.5 mg kg⁻¹ ketamine and subsequent administration of isoflurane in oxygen.

Score	Quality	Anaesthesia	Recovery
5	Excellent	Smooth induction, good muscle relaxation, physiological variables stable	Smooth, calm recovery, standing at first attempt
4	Very good	Smooth induction, some muscle tension during procedure but physiological variables stable	Smooth, calm recovery, mild ataxia when attempting to stand
3	Good	Minor agitation on induction, muscle tension during procedure, physiological variables stable but higher concentration isoflurane required	Some wing movements or ataxia but no manual restraint required
2	Poor	Agitated induction, muscle tension during procedure, transient apnoea post induction, transient manual ventilation required	Moderate wing flapping, disorientation, manual restraint required
1	Very poor	Agitated induction, poor muscle relaxation, prolonged apnoea following induction, manual ventilation required	Violent wing flapping, rolling or ataxia, manual restraint required