

Central Lancashire Online Knowledge (CLOK)

Title	A comparison of the pharmacodynamic effects of intravenous ketamine-xylazine with alfaxalone in mute swans (Cygnus olor) presenting at a wildlife veterinary hospital
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/39037/
DOI	https://doi.org/10.1016/j.vaa.2021.03.014
Date	2021
Citation	Baldrey, Vicki, Stanford, Michael and Bacon, Heather (2021) A comparison of the pharmacodynamic effects of intravenous ketamine-xylazine with alfaxalone in mute swans (Cygnus olor) presenting at a wildlife veterinary hospital. <i>Veterinary Anaesthesia and Analgesia</i> . ISSN 1467-2987
Creators	Baldrey, Vicki, Stanford, Michael and Bacon, Heather

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.vaa.2021.03.014>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Table 3 Presenting condition or indication for anaesthesia of swans included in this study.

Birds were anaesthetized with intravenous alfaxalone (Group A, $n = 27$) or ketamine/xylazine (Group KX, $n = 31$). For drug doses see Table 2 legend.

	Group A ($n = 27$)	Group KX ($n = 31$)
Air sac endoscopy	22	21
Radiography	1	4
Suture wound or debride abscess	3	3
Remove severely damaged feathers	1	2
Remove external fixator	0	1