

Central Lancashire Online Knowledge (CLOK)

Title	Should we be giving dietary advice to prevent periodontal disease? The effect of a low-carbohydrate diet in reducing periodontal inflammation
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/40069/
DOI	https://doi.org/10.1038/s41407-021-0783-9
Date	2021
Citation	Littlemore, Bevereley and Duerden, Sarah (2021) Should we be giving dietary advice to prevent periodontal disease? The effect of a low-carbohydrate diet in reducing periodontal inflammation. BDJ Team.
Creators	Littlemore, Bevereley and Duerden, Sarah

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1038/s41407-021-0783-9>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

***‘Should we be giving dietary advice to prevent periodontal disease?
The effect of a low-carbohydrate diet in reducing periodontal
inflammation’***



BEV LITTLEMORE

MSC, PGDIP, PGCERT TLHE, BSc (HONS), DIP DHDT, FHEA

SENIOR LECTURER AND COURSE LEAD DENTAL THERAPY

UNIVERSITY OF CENTRAL LANCASHIRE



SARAH DUERDEN

MSC, PGCERT TLHE, BSc (HONS), DIP DHDT, FHEA

SENIOR LECTURER AND YEAR LEAD DENTAL THERAPY

UNIVERSITY OF CENTRAL LANCASHIRE

INTRODUCTION

Dietary advice in the dental setting predominantly focuses on the prevention of dental caries with very little thought given to periodontal disease, indeed the common risk factor approach proposed by Sheiham and Watt (2000) links diet to dental decay, obesity, cancers and cardiovascular disease – it does not associate diet with periodontal disease¹. However, periodontal disease is the 6th most prevalent disease affecting approximately 20-50% of the global population ². At least 1 in 10 adults worldwide suffer from periodontal disease, making it more prevalent than cardiovascular disease^{3 4 5}. Severe periodontitis has a standardised prevalence of 11.2% worldwide⁶ and is a significant public health concern⁷. Recent studies have questioned the role of dietary intake in the reduction of periodontal inflammation, it is

plausible that future dietary advice may include specific advice to reduce such periodontal inflammation.

PERIODONTAL DISEASE AND INFLAMMATION

Periodontal diseases are mediated as a result of an inflammatory response to microorganisms within the dental biofilm⁸. The bacteria within dental biofilm release biologically active components, such as lipopolysaccharides, chemotactic peptides, and organic acids⁹. Detection of microbial pathogen-associated molecular patterns (PAMPs) by the host immune pattern recognition receptors (PRRs) induces the immune response and the production of pro-inflammatory cytokines including tumour necrosis factor alpha (TNF- α), interleukin- (IL-) 6, IL-8, IL-12, interferon- γ (IFN- γ) and acute-phase proteins such as C-reactive protein (CRP)^{10 11 12 13}. Activation of the complement system produces complement peptides, anaphylatoxins C3a, C4a and C5a that attract host immune cells monocytes, lymphocytes and neutrophils¹⁴.

Not only is complement is a central component of the innate immune system, it is an important mediator of the adaptive immune responses¹⁵. The complement cascade can be activated through three pathways: classical, lectin, and the alternative pathways. Periodontal diseases primarily activate the alternative pathway via bacterial polysaccharides, including aggregated IgA and lipopolysaccharide, through properdin (factor P) to cleave C3. Factors B and D, along with C3b convert C5 into C5a and C5b and the cascade continues until completion^{8 16}.

This signals the transition from the innate to the acquired immune response. The presence of pro-inflammatory cytokines results in the recruitment of host immune cells and the infiltration of neutrophils, natural killer (NK) cells, and lymphocytes to recognise antigens on dendritic cells¹⁷. T-lymphocytes, B-lymphocytes, macrophages and plasma cells predominate, collagenolytic activity increases, blood flow decreases⁸. T-lymphocytes express CD8+ (cytotoxic) and CD4+ (helper T-cell) cells that generate pro-inflammatory cytokines IL-1, IL-4, IL-10, TNF- α , IFN- γ and transforming growth factor β (TGF- β)¹¹. Additionally, CD4+ lymphocytes produce the cytokine RANK-L, that signals the stimulation of reactive oxygen species (ROS) in osteoclasts^{18 19}. The presence of ROS is increasingly implicated in the connective tissue degradation, particularly the alveolar bone^{18 20}.

The presentation of antigens to the dendritic cells attracts polymorphonuclear leukocytes (PMNs) to the site; degranulation of PMNs releases proteases and other lytic enzymes, defensins and ROS that help in degrading the pathogens during phagocytosis²¹. However, whilst this aids in the elimination of the microbial burden, it also has the effect of connective tissue fibres destruction, resulting in an additional local damage.

The excessive inflammatory response (both innate and acquired) mediates the connective tissue damage and alveolar bone loss²². Furthermore, the tissue breakdown products provide a nutrient source for the pathogenic bacteria thus inducing an unfortunate self-sustainable feed-forward cycle of escalating dysbiosis and disease progression^{23 13 24}. Consequently, although the initial inflammatory response in periodontal inflammation is of microbial origin, the subsequent tissue destruction is predominantly host mediated¹³.

LOW-CARBOHYDRATE DIETS

A Western lifestyle, high in carbohydrates with a high ratio of Omega-6 to Omega-3 fatty acids can promote inflammatory processes²⁵. Diets containing glucose-rich refined carbohydrates cause exaggerated postprandial surges in glucose and triglycerides²⁶; ROS is generated at a rate that exceeds that of antioxidant defences, resulting in oxidative stress^{27 28}. This postprandial dysmetabolism is associated with the initiation of inflammation^{26 28}.

The National Diet and Nutrition Survey²⁹ report a daily average intake of carbohydrates of 198g for women and 252g for men, representing approximately 47-48% of food energy intake. The parameters of a low-carbohydrate diet range between 100-150g of carbohydrates daily and due to claims of weight loss success, have gained substantial popularity in recent years^{30 31}. As carbohydrates work directly to raise the blood glucose concentration, a low-carbohydrate diet is often recommended in the management of type 2 diabetes³². Similarly, the palaeolithic or 'stone age' diet attempts to emulate the diet of our ancestors and restricts food intake to lean meats, fish, insects, eggs, fruit, berries, vegetables, and nuts while limiting grains and legumes. Notably, dairy products and refined carbohydrates are excluded from the diet³³. It has been observed that adopting a palaeolithic diet can not only promote weight loss but is also associated with reduced systemic inflammation^{34 35}.

Several studies have investigated the role of nutrition in the prevention and management of periodontal inflammation^{36 37}. It is known that both micro and macronutrients modulate pro-inflammatory or anti-inflammatory cascades, influencing inflammatory status²⁶. Much of this research is focused around micronutrients^{38 39 40 41}. It is considered that such inflammatory processes can be resolved with the aid of Omega-3 fatty acids⁴². Additionally, dietary antioxidants are crucial for maintaining redox equilibrium to reduce oxidative stress⁴³; several studies have demonstrated a positive impact of antioxidants Vitamin C and D on periodontal inflammation⁴⁴. Few studies have assessed the effects of macronutrients⁴⁴, even less investigating the effects of a low-carbohydrate diet on periodontal inflammation³⁸.

This review critically evaluates the empirical evidence to establish whether reducing carbohydrate intake can reduce periodontal inflammation in patients with periodontal disease.

METHOD

A systematic review was conducted using PICO methods to align the inclusion and exclusion criteria with the research question to facilitate development of the research strategy. Concepts and synonyms were used to identify key words to be used when searching databases. After searching the Cochrane Register for existing systematics reviews on the research topic, bibliographic databases Medline, Dentistry and Oral Sciences Source, CINAHL Complete and ProQuest were searched. Investigation of the grey literature and citation chaining completed the searches.

INCLUSION AND EXCLUSION CRITERIA

The inclusion and exclusion criteria were defined using the PICO model⁴⁵ to objectively identify those papers relevant to the research question⁴⁶. The exclusion criteria include factors or characteristics that may be confounders for the outcome parameter (see Table 1).

SEARCH RESULTS


An initial 354 papers were identified and following de-duplication, 342 abstracts were screened for relevance with 331 papers excluded based on inclusion and exclusion criteria. Full text papers were retrieved for 11 studies; a further 7 papers were excluded at this stage. A final 4 papers were included in the review: 3 RCTs and 1 case series.

The process by which the papers were identified can be seen in the PRISMA flow diagram (Fig. 2).

Table 1. Inclusion and Exclusion Criteria

PICO	Inclusion	Exclusion
Population	<p>It is expected that papers in this review will include the following:</p> <ul style="list-style-type: none"> Subjects with gingivitis/gingival inflammation – gingival erythema, oedema, $\geq 10\%$ gingival bleeding <p>AND/OR</p> <ul style="list-style-type: none"> Periodontitis – pocket probing depths $\geq 4\text{mm}$, evidence of clinical attachment loss, bleeding on probing $\geq 10\%$ 	<p>It is expected that papers in this review will exclude the following:</p> <ul style="list-style-type: none"> Subjects with systemic disease that may affect the periodontium e.g. diabetes Pregnancy/lactating mothers Subjects with a carbohydrate or insulin related disease e.g. diabetes Medications that may affect the periodontium (calcium channel blockers, cyclosporine, phenytoin) Medications that might influence bleeding and /or gingival inflammation including non-steroidal anti-inflammatories, anticoagulants and cortisone Subjects who have undergone treatment of periodontitis within the preceding 12 months Smokers or previously smoked within the last 10 years Intake of any other nutrient supplements Subjects regularly exceeding recommended daily limits for alcohol ingestion
Intervention/s	<ul style="list-style-type: none"> Diet conditions with a main criterion of a maximum processed carbohydrate intake of 25%/130g TDE <p>OR</p> <ul style="list-style-type: none"> Dietary intake consisting solely of the limited carbohydrate regimes akin to those consumed in the stone age or Palaeolithic time periods. 	<ul style="list-style-type: none"> Papers evaluating the effect of micronutrients on periodontal health Evaluating the effects of nutrients on dental caries Evaluating the effects of low-carbohydrate diet/nutrition of plaque bacteria
Comparator/s	<ul style="list-style-type: none"> Normal or Western diet high in carbohydrates 	<ul style="list-style-type: none"> Comparators of different population groups or interventions
Outcome/s	<ul style="list-style-type: none"> Improved periodontal clinical parameters: <ul style="list-style-type: none"> Reduced gingival inflammation (GI) Reduced bleeding on probing (BoP) Reduced pocket probing depth (PPD) Reduced plaque index (PI) Periodontal inflamed surface area (PISA) 	<ul style="list-style-type: none"> Dental caries only (dmf-t/DMF-t) Serological parameters only Microbiota only

PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources



From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ 2021;[372:n71](https://doi.org/10.1136/bmj.n71). doi: 10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>

Assessment of Methodological Quality

Identified studies were reviewed, data extracted and critically appraised for their methodological quality using the Effective Public Health Practice Project (EPHPP) Tool⁴⁷. This tool has the advantage of continuity and consistency as it is applicable across a variety of intervention study designs and not specific to any one study design⁴⁸. The overall outcome of the quality assessment for each study is rated as either strong, moderate, or weak; the results are summarised in Table 2. Critical analysis of the methodological quality of the included papers enabled consideration of impact of the quality assessment on the review findings and recommendations.

Table 2. Summary of Quality Assessment using EPHPP Tool

Reference	Selection Bias	Study Design	Confounders	Blinding	Data Collection Methods	Withdrawals/ Dropouts	Global Rating
Baumgartner et al., 2009 ⁴⁹	Moderate	Moderate	Moderate	Weak	Strong	Moderate	Moderate
Woelber et al., 2016 ⁵⁰	Moderate	Strong	Strong	Weak	Strong	Strong	Moderate
El Makaky et al., 2019 ⁵¹	Moderate	Strong	Strong	Moderate	Strong	Moderate	Strong
Woelber et al., 2019 ⁵²	Moderate	Strong	Strong	Moderate	Strong	Moderate	Strong

After assessment of the individual quality components, two of the RCTs^{51 52} had a strong global rating, characterised as having no weak ratings in any component. The paper by Baumgartner et al., (2009)⁴⁹ and the RCT by Woelber et al., (2016)⁵⁰ were appraised as being of moderate quality due to both being considered weak when assessing the blinding component of the quality assessment⁴⁷. As no papers received a global rating of weak, no paper was excluded

on this basis. However, the various limitations observed across the papers will need to be considered when interpreting the results.

DATA EXTRACTION AND STUDY CHARACTERISTICS OF INCLUDED STUDIES

The data extraction form was adapted from the Cochrane Effective Practice and Organisation of Care (EPOC) data extraction tool for randomised and non-randomised trials⁵³. Two reviewers extracted and evaluated relevant data to enable a comparison of methodology and outcomes to facilitate the assessment of the effectiveness of the regimes⁵⁴.

A summary of findings, including pertinent characteristics of participants, oral hygiene, and dietary regimes along with the outcomes have been summarised in Tables 3, 4, 5 and 6.

The considerable heterogeneity in terms of interventions, populations and study duration precluded statistical synthesis in the form of meta-analysis, consequently, data extracted from the studies was synthesised to form a narrative in which to determine the range of clinical outcomes reported⁵⁵.

Table 3: Study Characteristics: General Information, Population Characteristics, Interventions and Outcomes

Author/s	Baumgartner S, Imfeld T, Schicht O, Rath C, Persson RE, Persson GR.	Woelber, J.P., Bremer, K., K. Vach, K., König, D., Hellwig, E., Ratka-Krüger, P., Al-Ahmad, A., Tennert, C.	El Makaky, Y., Beltagy, T., El Makakey, A.	Woelber JP, Gärtner M, Breuninger L, Anderson A, König D, Hellwig E, Al-Ahmad A, Vach K, Dötsch A, Ratka-Krüger P, Tennert C.
Year of Publication	2009	2016	2019	2019
Study Duration	4 weeks	8 weeks	4 weeks	8 weeks
Country of Study	Switzerland	Germany	Egypt	Germany
Study Design	Longitudinal case series	Pilot Randomised Controlled Trial	Randomised Controlled Trial	Randomised Controlled Trial
Study Aims/Objectives	To assess the oral microbiota and clinical data in subjects without access to traditional oral hygiene methods and who ate a diet available to humans during the Stone Age.	To evaluate an oral health optimized diet low in carbohydrates, and rich in Omega 3-fatty acids, vitamins C and D, antioxidants and rich in fibre.	To assess the effects of an anti-inflammatory diet on gingival health and serological parameters in child participants.	To investigate the influence of an anti-inflammatory diet on clinical, serological and subgingival microbiome parameters.
Ethical approval obtained	Yes, ethical approval obtained from the University of Zurich.	Yes, ethical approval was obtained from the University of Freiburg Ethics committee and the study was registered with the German Clinical Trials Register.	Yes, ethical approval obtained from the Research Ethics Committee, Faculty of Dentistry; Kafrelsheikh University.	Yes, ethical approval was obtained by the University of Freiburg Ethics committee and the study was registered with the German Clinical Trials Register.
Study Quality (EPHPP)	Moderate	Moderate	Strong	Strong
No. of Participants Recruited	10	16	40 (Intervention Group n = 20 Control Group n = 20)	38
Withdrawal/Dropouts	None	1 intervention group participant	None	6 withdrawn due to unsuitable dietary regime 2 control group participants dropped out due to medical reasons
No. of participants after withdrawals/dropouts	10	15 (Intervention Group n = 10 Control Group n = 5)	40 (Intervention Group n = 20 Control Group n = 20)	30 (Intervention Group n = 15 Control Group n = 15)
Patient demographics: <ul style="list-style-type: none"> Gender Ratio Age 	5 males: 5 females 7 adults (18-46yrs) 3 children (8-12yrs)	Intervention Group: 4 males: 6 females Mean age 34.4 ± 14.1 years, ranging from 23 to 70 years Control Group: 2 males: 3 females Mean age 34.0 ± 16.5 years, ranging from 24 to 63 years	Intervention Group: 9 boys: 11 girls Mean age 11.90 ± 1.410 years, ranging from 10 to 14 years Control Group: 8 boys: 12 girls Mean age 11.75±1.410 years, ranging from 10 to 14 years	Intervention Group (n=15): 6 males: 9 females Mean age 27.2 ±4.7 years Control Group (n=15): 7 males: 8 females Mean age 33.7 ±13.1 years

Author/s	Baumgartner S, Imfeld T, Schicht O, Rath C, Persson RE, Persson GR.	Woelber, J.P., Bremer, K., K. Vach, K., König, D., Hellwig, E., Ratka-Krüger, P., Al-Ahmad, A., Tennert, C.	El Makaky, Y., Beltagy, T., El Makakey, A.	<ul style="list-style-type: none"> Woelber JP, Gärtner M, Breuninger L, Anderson A, König D, Hellwig E, Al-Ahmad A, Vach K, Dötsch A, Ratka-Krüger P, Tennert C.
Follow-up Time	After 4 weeks	Weekly intervals at weeks 5, 6, 7, 8	After 4 weeks	Weekly intervals at weeks 5, 6, 7, 8
Oral Hygiene Regime Intervention Group	None allowed other than the use of twigs and other natural products foraged.	At the start of the first week all participants were instructed to stop all interdental hygiene procedures for the next eight weeks.	Normal toothbrushing but no interdental cleaning	Participants were instructed not to perform any interdental hygiene throughout the study period.
Oral Hygiene Regime Control Group	N/A	At the start of the first week all participants were instructed to stop all interdental hygiene procedures for the next eight weeks.	Normal toothbrushing but no interdental cleaning	Participants were instructed not to perform any interdental hygiene throughout the study period.
Dietary Regime Control Group	N/A	Continue with usual diet mainly based on carbohydrates	No change in usual dietary habits	The control group was instructed not to profoundly change their diet for the next 6 weeks
Dietary Regime Intervention Group	Restricted Stone Age diet consisting of: <ul style="list-style-type: none"> basic supply of whole grains of barley, wheat, spelt salt, herbs, honey, milk, meat from domestic animals (goats and hens), berries, edible plants fish 	Reduction of the intake of carbohydrates <130 g/d: See Table 5: Dietary Regime Dietary recommendations were delivered verbally in an information brochure. This was done after the 2 nd baseline measurements were taken. 2 further weeks were given to allow participants to adjust to the new dietary regime. Participants were required to encouraged to follow the new det for the remaining 4 weeks, completing a daily food diary.	Low-carbohydrates diet <130 g/ day: See Table 5: Dietary Regime Plus: <ul style="list-style-type: none"> almonds and walnuts gluten-free whole grains olive oil and soy-based foods herbs and spices Detailed verbal data about dietary protocol was given to each patient and participant's parents/ caregivers.	At baseline, both groups had to continue their Western diet for 2 weeks. After this, the test group had to change to an anti-inflammatory diet (AID) protocol for 4 weeks after two transitional weeks. Reduction of the intake of starches <130 g/day: See Table 5: Dietary Regime Participants received detailed verbal introduction into the AID protocol for 30 min. <ul style="list-style-type: none"> All participants were instructed to fill out a 24 hr-dietary diary for 1 week at the second, fifth and eighth week
Statistical analysis	<ul style="list-style-type: none"> Paired t-test to assess changes in the clinical indices PD, BOP, GI, and PI over time. Significance declared at the P <0.001 level. 	<ul style="list-style-type: none"> Mixed linear regression analysis used to test for differences between groups. Multiple testing was corrected using the <u>Scheffe</u> method 	<ul style="list-style-type: none"> Paired t-test to detect general changes within groups. Mean and standard deviation for simple descriptive analysis (age and sex). Student t-test used to assess changes between groups. P-value 0.05 was considered as a level of significance. 	<ul style="list-style-type: none"> Linear regression model used to analyse changes between groups. Linear mixed model used to compare changes between groups over time, Bonferroni used to correct for multiple testing. All analyses regarding the clinical (and serological) data were calculated with STATA 14.2.
Conclusions	Diet restriction, coupled with abstinence from oral hygiene, did not result in increased gingival inflammation; decreases in BOP and PDs were observed.	A low-carbohydrate diet that is rich in Omega-3 fatty acids, vitamins C and D, antioxidants and fibre can significantly reduce periodontal inflammation.	A low-carbohydrate, anti-inflammatory diet was able to significantly gingival inflammation.	The evaluated anti-inflammatory diet was able to significantly reduce gingival inflammation in a clinically relevant range.

Table 4: Dietary Regime Restricted Carbohydrate Diets < 130g

<p>Dietary Regime: Restricted Carbohydrates <130g</p> <ul style="list-style-type: none"> • El Makaky et al., 2019 • Woelber et al., 2016 • Woelber et al., 2019 	<p>Macronutrients:</p> <ul style="list-style-type: none"> • Reduction in the amount of processed (fibre free) carbohydrates (disaccharides) such as sweetened beverages and meals, flour containing foods, <u>rice</u> and potatoes • Reduction in the amount of omega-6 fatty acids (such as margarine, safflower oil, corn oil, sunflower oil, sesame oil) • Decrease the amount of industrial animal proteins (such as processed meat products and industrial dairy) as far as possible • Reduction in the quantity of trans-fatty acids • Daily intake of supplements of omega-3 fatty acids (such as two spoons of flaxseed oil, a portion of the sea fish, etc.) • Daily intake of vegetables such as tomatoes, broccoli, carrots, spinach, sweet potatoes, beets, cabbage, and beans • Daily intake of fruits such as orange, cherries, blueberries, strawberries, cantaloupe, watermelon, avocados, and kiwifruit <p>Micronutrients:</p> <ul style="list-style-type: none"> • Daily intake of vitamin C (from fruit/vegetables such as oranges, kiwi, bell peppers) • Daily intake of vitamin D • Daily intake of nitrates such as beet/spinach • Daily intake of fibre (such as legumes, fruit, vegetables, bran) • Daily intake of antioxidants (such as green tea, one pinch of Curcuma, coffee without milk, berries, and ginger)
--	---

Table 5. Results: Primary Outcomes

Key: Group A: Control Group Group B: Intervention Group

	Gingival Index						Bleeding on Probing (%)					
	Group A Baseline	Group A End of Study	Level of Significance	Group B Baseline	Group B End of Study	Level of Significance	Group A Baseline	Group A End of Study	Level of Significance	Group B Baseline	Group B End of Study	Level of Significance
Baumgartner et al., 2009	N/A	N/A	N/A	0.38	0.43	Not stat. significant	N/A	N/A	N/A	34.8%	12.6%	p < 0.001
Woelber et al., 2016	1.01 ± 0.14	1.22±0.17	p < 0.001	1.10 ±0.51	0.54±0.30	p < 0.001	46.46% ±15.61	64.06% ± 11.27	p=0.012	53.57% ±18.65	24.17% ±11.57	p=0.012
El Makaky et al., 2019	1.445 ± 0.4084	1.660 ± 0.3775	p=0.000	1.450 ± 0.4335	1.145 ± 0.4443	p=0.000	N/A	N/A	N/A	N/A	N/A	N/A
Woelber et al., 2019	0.92 ±0.25	0.74 ± 0.18	p < 0.05	1.03 ± 0.21	0.61 ± 0.29	p < 0.05	28.39% ± 13.32	27.09% ± 10.03	p=0.864	30.35% ± 11.07	23.55% ± 13.61	p=0.864

Table 6. Results: Secondary Outcomes

Key: Group A: Control Group Group B: Intervention Group

	Plaque Index						Pocket Probing Depth (mm)						PISA (mm) ²					
	Group A Baseline	Group A End of Study	Level of Signif.	Group B Baseline	Group B End of Study	Level of Signif.	Group A Baseline	Group A End of Study	Level of Signif.	Group B Baseline	Group B End of Study	Level of Signif.	Group A Baseline	Group A End of Study	Level of Signif.	Group B Baseline	Group B End of Study	Level of Signif.
Baumgartner et al., 2009	N/A	N/A	N/A	0.68	1.47	p<0.001	N/A	N/A	N/A	2.23 ± 0.6	2.09 ± 0.63	P <0.001	N/A	N/A	N/A	N/A	N/A	N/A
Woelber et al., 2016	0.75 ± 0.63	0.97 ± 0.70	p = 0.084	1.10 ± 0.51	0.84 ± 0.47	p = 0.084	2.31 ± 0.43	2.52 ± 0.40	Not given	2.19 ± 0.34	2.11 ± 0.35	Not given	662.24 ± 420.05	963.24 ± 373.78	p<0.01	638.88 ± 305.41	284.83 ± 174.14	p<0.001
El Makaky et al., 2019	1.540 ± 0.7051	1.655 ± 0.6589	p= 0.001	1.550 ± 0.7585	1.560 ± 0.7618	p=0.733	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Woelber et al., 2019	0.58 ± 0.14	0.48 ± 0.12	p=1.0	0.58 ± 0.12	0.48 ± 0.13	p=1.00	1.82 ± 0.24	2.00 ± 0.14	p=0.018	1.85 ± 0.27	1.84 ± 0.17	p= 0.018	270.50 ± 140.97	286.00 ± 114.02	p=0.6	315.27 ± 148.68	252.37 ± 151.78	p=0.6

REVIEW FINDINGS

Clinical Parameters

The included studies were appropriate to address the aims of this review. There were disparities in terms of participants and study design, particularly between the case series⁴⁹ and the three RCTs^{50 51 52}. Despite this, in relation to the research question, there were comparable similarities observed in the primary and secondary outcomes measured.

All four papers observed reductions in periodontal inflammation, as measured by reduced gingival inflammation, bleeding on probing and reduced pocket probing depths. The three RCTs observed significant reductions in GI and BoP^{50 51 52}, only the case series observed an increase in GI from baseline to the end of the study however, it was of no statistical significance⁴⁹. Similar reductions were observed for bleeding on probing with Baumgartner et al., (2009)⁴⁹ and Woelber et al., (2016)⁵⁰ reporting significant reductions. BoP also reduced significantly in the intervention group in the RCT by Woelber et al., (2019)⁵², however when compared to the control group, the difference was not significant. These results, in conjunction with the increases seen in the control groups are indicative of the effectiveness of the intervention as both gingival index and bleeding on probing are accurate indicators of gingival health⁷. Likewise, probing depth is an important clinical parameter in the assessment of the clinical status of periodontal tissues⁵⁶. The three papers that evaluated the effect of the dietary regime on PPD all observed significant reductions within the intervention groups^{49 50 52}.

A relatively new index, PISA is considered beneficial for quantifying periodontal inflammation⁵⁷ as it integrates multiple periodontal indices, such as BoP, PPD, and PI into a single numerical index. The two RCTs that carried out this index observed reduced inflammation in the intervention groups compared to the control groups^{50 52}. Although the reduction observed in the more recent RCT⁵² was not deemed statistically significant, the results align with the findings of the other periodontal indices.

Age Differences

The participants in one RCT⁵¹ were children between the ages of 10-14-years; there are inherent challenges in assessing periodontal indices in children with a mixed dentition due to the presence of false pocketing as the permanent teeth are only partially erupted⁶⁷, additionally, possibly due to hormonal changes, the prevalence of gingivitis peaks during puberty⁵⁸. It is acknowledged in the case series⁴⁹ that such challenges may impact the results however, they observed no trends of differences to suggest any such impact that would differentiate the children from the adult participants within their study.

Oral Hygiene Regimes and Plaque

Interproximal cleaning is important in maintaining interproximal gingival health⁷ and the consequence of having no access to oral hygiene aids and being prohibited from carrying out interdental cleaning is reflected by the increase in supragingival plaque scores observed in two of the studies^{49 51}. Interestingly, the increases in plaque levels observed were not accompanied by a corresponding increase in the severity of gingival inflammation which would normally be expected⁷. Indeed, it was concluded that even in the presence persistent plaque levels, an anti-inflammatory diet significantly decreased gingival inflammation⁵⁰.

Dietary Regimes

When assessing compliance with dietary regimes, two studies^{50 52} assessed dietary habits using self-reported diaries. Using diaries is not without challenges; they rely on subjects to accurately recall and record the relevant data and have questionable validity and reliability⁵⁹.

Determining compliance was not necessary in the simulated Stone-Age case series⁴⁹, although the authors acknowledge that it was difficult to assess precise dietary intake because analysis of carbohydrate consumption and other macro/micronutrients was not possible. However, this simulated palaeolithic environment likely precludes subjects exceeding carbohydrate limits imposed in the other studies included in this review.

Carbohydrates and Inflammation

Diets high in refined carbohydrates can cause an exaggerated postprandial surge in glucose and triglycerides²⁶. This postprandial hyperglycaemia results in increased production of ROS and release of inflammatory cytokines⁶⁰. It has already been established that dietary antioxidants maintain redox equilibrium, reducing oxidative stress⁴³ and not only do the dietary requirements in the three RCTs restrict carbohydrates to less than 130g daily, they also incorporate elements that are anti-inflammatory; as such, the reduced clinical inflammation observed may be related to these components. Both studies by Woelber et al., (2016; 2019)⁵⁰ ⁵² acknowledge it may be difficult to attribute the improved clinical parameters observed in the intervention group to any one specific component of the diet, although regression analysis did show significant association between reduced clinical parameters and carbohydrate-reduction in the 2016 trial⁵⁰.

Whilst the case series⁴⁹ did not specifically refer to this within their study, the dietary regime the participants were subjected to included berries high in antioxidants⁶¹. Although honey isn't a refined carbohydrate, it is a pure sugar and as such one might have expected this to be reflected in the results observed, however the reduced periodontal inflammation observed by the authors is possibly due to the anti-inflammatory and anti-bacterial properties of honey⁶². Indeed, the authors concluded it was likely the combination of a restriction of refined carbohydrates and the supplemental intake of antioxidants that reduced the observed periodontal inflammation and that consequently, dietary advice for patients with gingivitis and periodontitis may be very important.

Additional outcomes

One RCT⁵² observed significant weight loss in the intervention group likely due to the reduced total energy intake associated with the low-carbohydrate regime. As evidence suggests low calorie intake is associated with improved periodontal health⁶³, it is possible this had an adjunctive effect on the reduced inflammation observed in this study. Additionally, they found no significant differences between or within groups in relation to the subgingival microbiome. The significance of this being that the clinical results observed were likely a result of an altered immunological response due the dietary regime and not resulting from an altered microbial composition of the subgingival biofilm. This supports the findings observed

in the case series⁴⁹, which despite reporting an increase in subgingival bacterial counts found no pathogenic bacteria associated with periodontitis.

CONCLUSION AND RECOMMENDATIONS

Despite the heterogeneity and variable methodological quality of the studies, all four studies reported similar outcomes in that despite reduced oral hygiene measures, the restricted carbohydrate intake reduced periodontal inflammation.

However, all papers included anti-inflammatory components to the dietary regimes, consequently how much of this decrease in periodontal inflammation can be attributed to the reduction in carbohydrates alone and how much was as a result of additional anti-inflammatory components in the diets is difficult to ascertain.

Nevertheless, the results provided by these studies highlight the potential of combining dietary advice relating to anti-inflammatory low-carbohydrate diets with appropriate oral health regimes.

The implications of the findings of this review may have a wider impact on patients' general health and on healthcare; if periodontal inflammation can be reduced through the adoption of a low-carbohydrate diet, then systemic inflammation and the risk for other inflammatory conditions may also be reduced. This supports previous research reporting improved systemic inflammation as a result of low-carbohydrate dietary regimes^{34 35}. Although the most recent RCT⁵² observed no significant difference in serological markers between control and intervention groups, another⁵¹ reported significant reductions in serological inflammatory parameters TNF- α and IL-6.

As periodontal disease has been linked to several non-communicable diseases, reducing the incidence and prevalence of periodontal disease may subsequently decrease the incidence and prevalence of such conditions as cardiovascular disease. Not only will this reduce the overall burden of disease and the associated financial impact of such, but more importantly, improve the quality of life of patients.

Unfortunately, the paucity of high-quality research currently available suggests despite the positive results of this review, it is far from enough to inform policy change; it may however

generate enough interest to influence further research in this arena. On considering a change in recommended treatment modalities in the management and prevention of periodontal disease, it seems that the next logical step would be a well-structured, sufficiently powered RCT investigating the effects of a low-carbohydrate diet, possibly as an adjunctive treatment to NSPT with a comparator group receiving NSPT alone.

From the limited evidence currently available, it is plausible that in addition to periodontal therapy, in the management of periodontal diseases dental professionals should consider assessing dietary habits with particular reference to intake of refined carbohydrates and an anti-inflammatory diet, providing dietary advice to reduce periodontal inflammation. However, further high quality longitudinal RCTs are needed in this topic area to strengthen the evidence base.

REFERENCES

1. Sheiham, A., & Watt, R.G. (2000) 'The Common Risk Factor Approach: a rational basis for promoting oral health'. *Community Dentistry and Oral Epidemiology*, 28, pp. 399-406.
2. Nazir, M.A. (2017) 'Prevalence of periodontal disease, its association with systemic diseases and prevention', *International journal of health sciences*, 11 (2), pp. 72–80.
3. Ower, P. (2015) 'The diseased root surface in periodontitis' in Eaton, K.A. and Ower, P. (eds.) *Practical Periodontics*. Edinburgh: Elsevier Health Sciences, pp. 259-269.
4. Kassebaum, N.J., Smith, A.G.C., Bernabé, E., Fleming, T.D., Reynolds, A.E., Vos, T. & Oral Health Collaborators (2017) 'Global, regional, and national prevalence, incidence, and disability adjusted life years for oral conditions for 195 countries, 1990–2015: A systematic analysis for the global burden of diseases, injuries, and risk factors', *Journal of Dental Research*, 96 (4), pp. 380–387.
5. Janakiram, C., & Dye, B.A. (2020) 'A public health approach for prevention of periodontal disease', *Periodontol 2000*, 84 (1), pp. 202-214.
6. Kassebaum, N.J., Bernabé, E., Dahiya, M., Bhandari, B., Murray, C.J., Marcenes, W. (2014) 'Global burden of severe periodontitis in 1990-2010: a systematic review and

- meta-regression', *J Dent Res*, 93 (11): 1045-53. doi: 10.1177/0022034514552491. Epub 2014 Sep 26. PMID: 25261053; PMCID: PMC4293771.
7. Chapple, I.L.C., Van der Weijden, F., Doerfer, C., Herrera, D., Shapira, L., Polak, D., Madianos, P., Louropoulou, A., Machtei, E., Donos, N., Greenwell, H., Van Winkelhoff, A.J., Eren Kuru, B., Arweiler, N., Teughels, W., Aimetti, M., Molina, A., Montero, E., Graziani, F. (2015) 'Primary prevention of periodontitis: managing gingivitis', *J Clin Periodontol*, 42 (16), pp. S71–S76.
 8. Cekici, A., Kantarci, A., Hasturk, H., Van Dyke, T.E. (2014) 'Inflammatory and immune pathways in the pathogenesis of periodontal disease', *Periodontology 2000*, 64 (1), pp. 57-80.
 9. Lamont, R.J., Koo, H., Hajishengallis, G. (2018) 'The oral microbiota: dynamic communities and host interactions', *Nature reviews. Microbiology*, 16 (12), pp. 745–759.
 10. Scannapieco, F.A. (2004) 'Periodontal inflammation: from gingivitis to systemic disease?', *Compend Contin Educ Dent*, 25 (7), pp. 16-25.
 11. Graves, D. (2008) 'Cytokines that promote periodontal tissue destruction', *J Periodontol*, 79 (8), pp. 1585–1591.
 12. Chapple, I.L.C. (2009) 'Periodontal diagnosis and treatment - where does the future lie?', *Periodontol 2000*, 51, pp. 9-24.
 13. Khan, S.A., Kong, E.F., Meiller, T.F., Jabra-Rizk, M.A. (2015) 'Periodontal Diseases: Bug Induced, Host Promoted', *PLoS Pathog*, 11 (7), e1004952. Available at: <https://doi.org/10.1371/journal.ppat.1004952>.
 14. Merle, N.S., Noe, R., Halbwachs-Mecarelli, L., Fremeaux-Bacchi, V., Roumenina, L.T. (2015) 'Complement System Part II: Role in Immunity', *Front Immunol*, 26; 6: p. 257. doi: 10.3389/fimmu.2015.00257. PMID: 26074922; PMCID: PMC4443744.
 15. Degn, S.E., & Thiel, S. (2013) 'Humoral Pattern Recognition and the Complement System', *Scand J Immunol*, 78 (2), pp. 181-193.
 16. Hajishengallis, G., Kajikawa, T., Hajishengallis, E., Maekawa, T., Reis, E.S., Mastellos, D.C., Yancopoulou, D., Hasturk, H., Lambris, J.D. (2019) 'Complement-Dependent Mechanisms and Interventions in Periodontal Disease', *Frontiers in immunology*, 10, 406. Available at: <https://doi.org/10.3389/fimmu.2019.00406>.

17. Benakanakere, M., Kinane, D.F. (2012) 'Innate cellular responses to the periodontal biofilm', *Frontiers of Oral Biology*, 15, pp. 41–55.
18. Ha, H., Kwak, H.B., Lee, S.W., Jin, H.M., Kim, H.M., Kim, H.H., Lee, Z.H. (2004) 'Reactive oxygen species mediate RANK signaling in osteoclasts', *Exp Cell Res*, 10, 301 (2), pp.119-27.
19. Vernal, R., Dutzan, N., Hernández, M. (2006) 'High expression levels of receptor activator of nuclear factor-kappa B ligand associated with human chronic periodontitis are mainly secreted by CD4⁺ T lymphocytes', *Journal of Periodontology*, 77 (10), pp. 1772–1780.
20. Huang, X., Xie, M., Xie, Y. (2020) 'The roles of osteocytes in alveolar bone destruction in periodontitis', *J Transl Med*, 18, 479. Available at: <https://doi.org/10.1186/s12967-020-02664-7>.
21. Khurshid, Z., Naseem, M., Sheikh, Z., Najeeb, S., Shahab, S., Zafar, M.S. (2015) 'Oral antimicrobial peptides: Types and role in the oral cavity', *Saudi Pharm. J*, 24 (5), pp. 515-524.
22. Hajishengallis, G., & Korostoff, J.M. (2017) 'Revisiting the Page & Schroeder model: the good, the bad and the unknowns in the periodontal host response 40 years later', *Periodontol 2000*, 75 (1), pp. 116-151.
23. Hajishengallis, E., & Hajishengallis, G. (2014) 'Neutrophil Homeostasis and Periodontal Health in Children and Adults', *Journal of Dental Research*, 93 (3), pp. 231-237.
24. Van Dyke, T.E., Bartold, P.M., Reynolds, E.C. (2020) 'The Nexus Between Periodontal Inflammation and Dysbiosis', *Frontiers in immunology*, 11, p. 511. Available at: <https://doi.org/10.3389/fimmu.2020.00511>.
25. Bosma-den Boer, M.M., Van Wetten, M.L., Pruimboom, L. (2012) 'Chronic Inflammatory Diseases Are Stimulated by Current Lifestyle: How Diet, Stress Levels and Medication Prevent Our Body from Recovering', *Nutr. Metab*, 9, 32.
26. O'Keefe, J., & Bell, D. (2007) 'Postprandial hyperglycemia/hyperlipidemia (postprandial dysmetabolism) is a cardiovascular risk factor', *Am J Cardiol*, 100 (5), pp. 899-904.
27. Weissman, A., Lowenstein, L., Peleg, A., Thaler, I., Zimmer, E.Z. (2006) 'Power spectral analysis of heart rate variability during the 100g oral glucose tolerance test in pregnant women', *Diabetes Care*, 29 (3), pp. 571-574.

28. O'Keefe, J.H., Gheewala, N.M., O'Keefe, J.O. (2008) 'Dietary strategies for improving post-prandial glucose, lipids, inflammation and cardiovascular health', *J Am Coll Cardiol*, 51 (3), pp. 249-255.
29. Public Health England (2016) 'The National Diet and Nutrition Survey'. Available at: <https://www.gov.uk/government/collections/national-diet-and-nutrition-survey>.
30. Naude, C.E., Schoonees, A., Senekal, M., Young, T., Garner, P., Volmink, J. (2014) 'Low-carbohydrate versus isoenergetic balanced diets for reducing weight and cardiovascular risk: a systematic review and meta-analysis', *PLoS One*, 9, e100652.
31. Abbasi, J. (2018) 'Interest in the Ketogenic Diet Grows for Weight Loss and Type 2 Diabetes', *JAMA*, 319 (3), pp. 215–217.
32. Schofield, G.M., Henderson, G.D., & Thornley, S. (2016). Very low-carbohydrate diets in the management of diabetes revisited. *The New Zealand medical journal*, 129 1432, 67-74 .
33. Leech, B., McEwen, B., Sekyere, E.O. (2020) 'Diet, Digestive Health, and Autoimmunity: The Foundations to an Autoimmune Disease Food Pyramid—Part 2', *Alternative and Complementary Therapies*, 26 (4), pp. 158-167.
34. McEwen, B.J. (2018) 'The Palaeolithic diet and cardiometabolic syndrome: Can an ancient diet be the way of the future?', *Adv Integrat Med*, 5, pp. 38–40.
35. de Menezes, E.V.A., Sampaio, H.A.D.C., Carioca, A.A.F. (2019) 'Influence of Paleolithic diet on anthropometric markers in chronic diseases: systematic review and meta-analysis', *Nutr J*, 18, 41. Available at: <https://doi.org/10.1186/s12937-019-0457-z>.
36. Enwonwu, C.O., & Ritchie, C.S. (2007) 'Nutrition and inflammatory markers', *J Am Dent Assoc*, 138 (1), pp. 70-73.
37. Hujoel, P.P., & Lingström, P. (2017) 'Nutrition, dental caries and periodontal disease: a narrative review', *J Clin Periodontol*, 4 (S18), pp. 79-84.
38. Van der Velden, U., Kuzmanova, D., Chapple, I.L.C. (2011) 'Micronutritional approaches to periodontal therapy', *J Clin Periodontol*, 38 (11), pp. 142–58.
39. Luo, P.P, Xu, H.S., Chen, Y.W., Wu, S.P. (2018) 'Periodontal disease severity is associated with micronutrient intake', *Aust Dent J*, 63 (2), pp. 193-201.
40. Dommisch, H., Kuzmanova, D., Jönsson, D., Grant, M., Chapple, I. (2018) 'Effect of micronutrient malnutrition on periodontal disease and periodontal therapy', *Periodontol 2000*, 78 (1), pp. 129-153.

41. Cagetti, M.G., Wolf, T.G., Tennert, C., Camoni, N., Lingström, P., Campus, G. (2020) 'The Role of Vitamins in Oral Health. A Systematic Review and Meta-Analysis', *International journal of environmental research and public health*, 17 (3), p. 938.
42. Calder, P.C. (2013) 'Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology?', *British journal of clinical pharmacology*, 75 (3), pp. 645–662.
43. Pizzino, G., Irrera, N., Cucinotta, M., Pallio, G., Mannino, F., Arcoraci, V., Squadrito, F., Altavilla, D., Bitto, A. (2017) 'Oxidative Stress: Harms and Benefits for Human Health. Oxidative medicine and cellular longevity'. Available at: <https://doi.org/10.1155/2017/8416763> [Accessed 12/01/2021].
44. Najeeb, S., Zafar, M.S., Khurshid, Z., Zohaib, S., Almas, K. (2016) 'The Role of Nutrition in Periodontal Health: An Update', *Nutrients*, 8 (9), p. 530.
45. da Costa Santos, C.M., de Mattos Pimenta, C.A., Nobre, M.R. (2007) 'The PICO strategy for the research question construction and evidence search', *Rev Lat Am Enfermagem*, 15 (3), pp. 508-511.
46. Garg R. (2016) 'Methodology for research', *Indian journal of anaesthesia*, 60 (9), pp. 640–645.
47. Effective Public Health Practice Project, EPHPP (2009) Quality Assessment Tool for Quantitative Studies. Available at: <http://www.ephpp.ca/tools.html>.
48. Armijo-Olivo, S., Stiles, C.R., Hagen, N.A., Biondo, P.D., Cummings, G.G. (2012) 'Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public Health Practice Project Quality Assessment Tool: methodological research', *Journal of Evaluation in Clinical Practice*, 18 (1), pp. 12-18.
49. Baumgartner, S., Imfeld, T., Schicht, O., Rath, C., Persson, R.E., Persson, G.R. (2009) 'The impact of the stone age diet on gingival conditions in the absence of oral hygiene', *J Periodontol*, 80 (5), pp. 759-768.
50. Woelber, J.P., Bremer, K., Vach, K., König, D., Hellwig, E., Ratka-Krüger, P., Al-Ahmad, A., Tennert, C. (2016) 'An oral health optimized diet can reduce gingival and periodontal inflammation in humans - a randomized controlled pilot study', *BMC Oral Health*, 17 (1), 28 DOI 10.1186/s12903-016-0257-1. Erratum in: *BMC Oral Health*. 2016 Oct 6, 16(1), 109.

51. El Makaky, Y., Beltagy, T., El Makakey, A. (2019) 'The Effects of an Anti-inflammatory Diet on Gingival Health in Children (Randomized controlled trial)', *Egyptian Dental Journal*, 65 (3), pp. 1995-2002.
52. Woelber, J.P., Gärtner, M., Breuninger, L., Anderson, A., König, D., Hellwig, E., Al - Ahmad, A., Vach, K., Dötsch, A., Ratka - Krüger, P., Tennert, C. (2019) 'The influence of an anti - inflammatory diet on gingivitis. A randomized controlled trial' , *Journal of Clinical Periodontology*, 46 (4), pp. 481-490.
53. Cochrane Effective Practice and Organisation of Care [EPOC] (2017) 'Data collection form. EPOC Resources for review authors'. Available at: <https://epoc.cochrane.org/resources/epoc-resources-review-authors>.
54. Boland, A., Cherry, M.G. & Dickson R. (2017) 'Doing a systematic review: a student's guide', 2nd Edition. London: Sage Publishing.
55. Li, T., Higgins, J.P.T., Deeks, J.J. (editors). (2021) 'Chapter 5: Collecting data', In: Higgins, J.P.T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M.J., Welch, V.A. (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.2 (updated February 2021). Cochrane. Available at: www.training.cochrane.org/handbook.
56. Meseli, S., Kuru, B., Kuru, L. (2017) 'Relationships between initial probing depth and changes in the clinical parameters following non-surgical periodontal treatment in chronic periodontitis', *Journal of Istanbul University Faculty of Dentistry*, 51 (3), pp. 11-17.
57. Park, S.Y., Ahn, S., Lee, J.T., Yun, P.Y., Lee, Y.J., Lee, J.Y., Song, Y.W., Chang, Y.S., Lee, H.J. (2017) 'Periodontal inflamed surface area as a novel numerical variable describing periodontal conditions', *J Periodontal Implant Sci*, 47 (5), pp. 328-338.
58. British Society of Periodontology (2012) 'Guidelines for periodontal screening and management of children and adolescents under 18 years of age'. Available at: https://www.bsperio.org.uk/assets/downloads/bsp_bspd-perio-guidelines-for-the-under-18s-2012.pdf. [Accessed 19/05/2021].
59. Nicolson, P.J.A., Hinman, R.S., Wrigley, T.V., Stratford, P.W., Bennell, K.L. (2018) 'Self-reported Home Exercise Adherence: A Validity and Reliability Study Using Concealed Accelerometers', *Journal of Orthopaedic & Sports Physical Therapy*, 48 (12), pp. 943-950.

60. Buyken, A.E., Flood, V., Empson, M., Roach, E., Barclay, A.W., Brand-Miller, J., Mitchell, P. (2010) 'Carbohydrate nutrition and inflammatory disease mortality in older adults', *The American Journal of Clinical Nutrition*, 92 (3), pp. 634-643.
61. Carlsen, M.H., Halvorsen, B.L., Holte, K., Bøhn, S.K., Dragland, S., Sampson, L., Willey, C., Senoo, H., Umezono, Y., Sanada, C., Barikmo, I., Berhe, N., Willett, W.C., Phillips, K.M., Jacobs, D.R., Jr, Blomhoff, R. (2010) 'The total antioxidant content of more than 3100 foods, beverages, spices, herbs and supplements used worldwide', *Nutrition Journal*; 9 (3). Available at: <https://doi.org/10.1186/1475-2891-9-3>.
62. Vallianou, N. (2014) 'Honey and its Anti-Inflammatory, Anti-Bacterial and Anti-Oxidant Properties', *General Medicine: Open Access*. Available at: <https://www.researchgate.net/publication/269552034>.
63. Park, H.S., Nam, H.S., Seo, H.S., Hwang, S.J. (2015) 'Change of periodontal inflammatory indicators through a 4 - week weight control intervention including caloric restriction and exercise training in young Koreans: A pilot study' , *BMC Oral Health*, 15, p. 109.

