

Central Lancashire Online Knowledge (CLOK)

Title	First and second trimester ultrasound in pregnancy: A systematic review and metasynthesis of the views and experiences of pregnant women, partners, and health workers
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/40175/
DOI	https://doi.org/10.1371/journal.pone.0261096
Date	2021
Citation	Moncrieff, Gill, Finlayson, Kenneth William, Cordey, Sarah Elizabeth, McCrimmon, Rebekah, Harris, Catherine orcid iconORCID: 0000-0001-7763-830X, Barreix, Maria, Tunçalp, Özge and Downe, Soo (2021) First and second trimester ultrasound in pregnancy: A systematic review and metasynthesis of the views and experiences of pregnant women, partners, and health workers. PLOS ONE, 16 (12). e0261096.
Creators	Moncrieff, Gill, Finlayson, Kenneth William, Cordey, Sarah Elizabeth, McCrimmon, Rebekah, Harris, Catherine, Barreix, Maria, Tunçalp, Özge and Downe, Soo

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1371/journal.pone.0261096>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>


RESEARCH ARTICLE

First and second trimester ultrasound in pregnancy: A systematic review and metasynthesis of the views and experiences of pregnant women, partners, and health workers

Gill Moncrieff^{1*}, Kenneth Finlayson¹, Sarah Cordey¹, Rebekah McCrimmon², Catherine Harris³, Maria Barreix⁴, Özge Tunçalp⁴, Soo Downe¹

1 Research in Childbirth and Health Group, THRIVE Centre, University of Central Lancashire, Preston, United Kingdom, **2** School of Health and Community Studies, University of Central Lancashire, Preston, United Kingdom, **3** Applied Health Research Hub, University of Central Lancashire, Preston, United Kingdom, **4** UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction, Department of Sexual and Reproductive Health and Research, World Health Organization, Geneva, Switzerland

* gmoncrieff1@uclan.ac.uk



OPEN ACCESS

Citation: Moncrieff G, Finlayson K, Cordey S, McCrimmon R, Harris C, Barreix M, et al. (2021) First and second trimester ultrasound in pregnancy: A systematic review and metasynthesis of the views and experiences of pregnant women, partners, and health workers. PLoS ONE 16(12): e0261096. <https://doi.org/10.1371/journal.pone.0261096>

Editor: Carla Betina Andreucci Polido, Universidade Estadual de Campinas, BRAZIL

Received: September 17, 2021

Accepted: November 22, 2021

Published: December 14, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0261096>

Copyright: © 2021 Moncrieff et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background

The World Health Organization (WHO) recommends one ultrasound scan before 24 weeks gestation as part of routine antenatal care (WHO 2016). We explored influences on provision and uptake through views and experiences of pregnant women, partners, and health workers.

Methods

We undertook a systematic review (PROSPERO CRD42021230926). We derived summaries of findings and overarching themes using metasynthesis methods. We searched MEDLINE, CINAHL, PsycINFO, SocIndex, LILACS, and AIM (Nov 25th 2020) for qualitative studies reporting views and experiences of routine ultrasound provision to 24 weeks gestation, with no language or date restriction. After quality assessment, data were logged and analysed in Excel. We assessed confidence in the findings using Grade-CERQual.

Findings

From 7076 hits, we included 80 papers (1994–2020, 23 countries, 16 LICs/MICs, over 1500 participants). We identified 17 review findings, (moderate or high confidence: 14/17), and four themes: *sociocultural influences and expectations; the power of visual technology; joy and devastation: consequences of ultrasound findings; the significance of relationship in the ultrasound encounter*. Providing or receiving ultrasound was positive for most, reportedly increasing parental-fetal engagement. However, abnormal findings were often shocking.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: The work was commissioned to the University of Central Lancashire by the UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), a cosponsored program executed by the World Health Organization (WHO). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Some reported changing future reproductive decisions after equivocal results, even when the eventual diagnosis was positive. Attitudes and behaviours of sonographers influenced service user experience. Ultrasound providers expressed concern about making mistakes, recognising their need for education, training, and adequate time with women. Ultrasound sex determination influenced female feticide in some contexts, in others, termination was not socially acceptable. Overuse was noted to reduce clinical antenatal skills as well as the use and uptake of other forms of antenatal care. These factors influenced utility and equity of ultrasound in some settings.

Conclusion

Though antenatal ultrasound was largely seen as positive, long-term adverse psychological and reproductive consequences were reported for some. Gender inequity may be reinforced by female feticide following ultrasound in some contexts. Provider attitudes and behaviours, time to engage fully with service users, social norms, access to follow up, and the potential for overuse all need to be considered.

Introduction

Antenatal ultrasound is a routine and established component of antenatal care within high-income countries [1]. In low- and middle-income countries ultrasound scanning in pregnancy is more recent [2]. In many of these settings, provision is not universal [3], and it is often restricted to high level and/or private facilities, limiting access for many [2, 4]. In 2016, the World Health Organization first recommended ultrasound as a routine aspect of antenatal care [5]. This recommendation was for one ultrasound scan before 24 weeks gestation, to estimate gestational age, improve detection of fetal anomalies and multiple pregnancies, reduce induction of labour for post-term pregnancy, and improve a woman's pregnancy experience. Part of the rationale for the establishment of this recommendation within guidelines was to better regulate the use of antenatal ultrasound, and to increase equitable access for pregnant women in low- and middle-income settings.

For many expectant parents, antenatal ultrasound provides a positive experience [6]. Health workers value its use for gestational age estimation, multiple pregnancy identification and assessment of physiological or potentially pathological fetal growth [1]. Identification of fetal anomalies is also an intrinsic part of ultrasound examination in early pregnancy [1]. As imaging has become more sophisticated, there has been increasing potential to identify markers of uncertain significance [7]. This can bring many benefits, but it has also resulted in concerns relating to overdiagnosis as well as the psychological risks for women, birthing people, and partners when the implications of these markers are not clear [8, 9]. Some have expressed eugenic concerns, as ultrasound-identified fetal abnormalities force parents to decide between giving birth to a child with disabilities, or termination [10], while in some social, cultural and religious contexts, termination is not an option [11]. In some social settings, ultrasound sex determination is associated with female feticide [12], and possibly sex distribution skew [13], raising moral, ethical, and gender equity issues.

Because of the rapid technical improvements in first and second trimester ultrasound, and the spread in routine use, the WHO recommended updating of their early ultrasound recommendation. This qualitative systematic review was carried out to inform the update, enabling

the consideration of values and preferences, and acceptability, feasibility, and equity implications, and the opportunity to share insights into successful implementation and service provision. These considerations are integral to implementation of antenatal ultrasound where it is not yet a routine component of antenatal care, as well as the improvement of existing services.

We undertook a rapid scoping search of the existing literature but did not identify any previous systematic reviews of experiences of first and second trimester ultrasound that were suitable to inform WHO guidelines on this subject. There is one previous systematic review on experiences of antenatal ultrasound, but this was published in 2002. It did not include the perspectives of health workers, or studies from low- or middle-income countries [6].

To inform guidelines and practice in the area of first and second trimester ultrasound we aimed to examine the following questions, for maternity service users (including birth companions), health workers, policy makers and funders in all settings:

- a. What views, beliefs, concerns and experiences have been reported in relation to routine ultrasound examination in pregnancy?
- b. What are the influencing factors associated with appropriate or inappropriate use of routine antenatal ultrasound scanning?

Methods

Search strategy and selection criteria

We undertook a systematic review using thematic synthesis to develop our review findings and analytic themes [14]. The study protocol is registered on PROSPERO (CRD42021230926).

Searches. We undertook searches in Medline (Ovid), CINAHL, PsycINFO, and SocIndex (via EBSCO), and LILACS and AIM (via Global Index Medicus) on Nov 25th and 26th 2020, with no language or date restrictions. Additional relevant papers were identified through searching reference lists and citation searches of included studies. A log was used to record inclusion/exclusion at each stage of selection. One member of the review team (CH) undertook the searches, and de-duplication of results using both automated and manual methods in EndNote.

Inclusion criteria. Our protocol specified searches for qualitative, survey, and mixed-methods studies. For this paper, we report on findings from qualitative studies. We included papers addressing routine use of ultrasound during antenatal care, including to detect fetal viability, gestational age, fetal growth, fetal abnormality, multiple pregnancy, and any other routine application, where this was a standard part of the routine ultrasound offer for the population in the country(ies) where the study was set.

Included participants were pregnant or postnatal women, families of such women, and related community members, antenatal health workers, managers, funders, or policy makers involved in the receipt, provision, management or funding of routine antenatal ultrasound scanning.

We included all settings (low-, high- and middle-income), and all types of health care design and provision (including public, private and mixed models of provision), and localities (hospital facilities, birth centres, or local communities).

Exclusion criteria. We excluded papers if ultrasound was undertaken for specific indications, for example following IVF procedures, or after women's reports of reduced fetal movements.

We excluded controlled studies, cohort studies, and epidemiological studies.

Screening. Initial screening by title and abstract was refined through blind screening 100 records in two teams to ensure agreement in the screening process. Uncertainties were discussed amongst the review team, and a further 100 hits were then screened until sufficient

agreement was reached. For full text screening, batches of ten records were screened in each team until sufficient agreement was reached, after which three members of the review team (GM, SC, RM) screened the remaining records independently.

Data extraction and analysis

Studies assessed as eligible for inclusion were quality assessed [15]. Quality assessment was undertaken by GM, SC, RM and KF. SD independently assessed 10% of studies to calibrate the assessments of the teams. Very low-quality studies were logged for transparency but were not included in the analysis.

The authors name, the date, characteristics, and setting of included papers, and the key findings, were logged on the study-specific Excel file. Translation of non-English studies was carried out using Google translate.

Analytic procedure. We initially derived review findings and overarching themes using a thematic synthesis approach [14]. We started by logging themes and findings highlighted by the authors, or, where these were not clear, reviewer generated findings from the quote material and author narratives (GM, SC, RM). As each subsequent paper was coded, themes were generated (GM, KF, SD) and entered iteratively onto a separate worksheet of the study Excel file, resulting in an initial thematic framework. The findings continued to develop as the data from each paper were added. This included looking for what was similar between papers and for what contradicted ('disconfirmed') the review findings. All authors involved in the primary analysis (GM, KF, SD), consciously looked for data that would contradict our prior beliefs and views.

Confidence in each finding was assessed using GRADE-CERQual [16]. Review findings were graded using a classification system ranging from 'high' to 'moderate' to 'low' to 'very low' confidence. Following CERQual assessment the review findings were grouped into higher order analytic themes and the final framework was agreed by consensus amongst the authors.

Analysis of subgroups or subsets. Findings were logged by country income status (HIC vs LMIC), and by trimester of scan (first, second, or both). Interpretation of the findings and themes includes these subgroups where they can be clearly differentiated in the data.

Reflexive statement. Based on our collective and individual experiences (as midwives, academics, service users, and researchers), we anticipated that the findings of our review would reveal that women and their partners generally look forward to ultrasound but may be unprepared for it to reveal abnormalities; that health workers like to use it as it gives them a sense of certainty in diagnosis; and that policy makers and funders see it as a useful source of revenue and/or of attracting women to use facilities. We maintained awareness of these prior beliefs and their potential impact on our analysis to ensure we were not over-interpreting data that supported our prior beliefs, or over-looking disconfirming data.

Results

Of the 7076 records generated by our search, 181 studies met the initial inclusion criteria to be included in our synthesis. 4656 records were excluded at the initial abstract screening stage, primarily because they were unrelated to the focus of this review. Full text screening excluded 574 studies, primarily because they did not focus on perceptions/experiences of routine ultrasound. Of the 181 studies initially identified as being eligible for inclusion, 80 were qualitative and 98 were quantitative or mixed methods studies. Due to the large number of qualitative papers identified, the decision was made to focus on the qualitative studies, and to analyse the qualitative/mixed methods studies separately. Eighty qualitative papers were therefore included before quality screening, and three more were identified from reference lists of the

included papers. Following quality appraisal, 3 studies were rated D and excluded. [Fig 1](#) outlines the screening and selection process.

Of the 80 studies included in our review, eight were rated A, 52 B, and 20 were rated C. They were published between 1994 and 2020 and were from 23 different countries, with 16 studies from LICs/MICs. They represent the views of over 1500 participants. The majority of papers reported the views of women or women and their partners; 19 reported provider perspectives; seven reported the views of both. There were no eligible studies that included the views of funders or policy makers. Study characteristics and quality appraisal grades are presented in [Table 1](#).

Findings

Our analysis generated 17 review findings, synthesised into four over-arching analytic themes. Three findings represent the views of women and their partners only, three represent the views of healthcare professionals only, and 11 describe findings from both groups. Most were graded moderate or high confidence. The Summary of Findings and CERQual assessment are provided in [Table 2](#).

Sociocultural influences and expectations. For many women, ultrasound was seen as an integral part of pregnancy and an opportunity not to be missed [17–26]. It offered parents the chance to ‘meet’ their baby and receive an image of the scan that they could share with friends and family [21, 25, 27, 28]. Fathers’ attendance was seen as a demonstration of their commitment to their family and to facilitate involvement with the pregnancy [19, 25, 28, 29–34]. For health workers however, these views sometimes conflicted with their role in providing a medical assessment and potential diagnosis [35–37]. It also sometimes conflicted with parent’s autonomy in terms of whether attending ultrasound was seen as a choice, or a decision to be made [17, 18, 21, 22, 38–42]. Some felt that they had not been offered an actual choice due to the routine nature of ultrasound in antenatal care, whilst others felt they should follow the authoritative advice of health professionals to ensure wellbeing of their baby [43–47]. In some contexts, healthcare professionals actively directed women towards ultrasound with the belief that this would inevitably result in better outcomes, and women were seen as irresponsible if they declined the offer of a scan [39, 44, 48–52].


‘Yes I’m sure it is (optional) but I think everybody else does it . . . well maybe not . . . but anyway I wouldn’t miss it.’ (Sweden) [25]

‘I don’t know if it is good or bad. They provide it for us so we use it.’ (Australia) [46]

‘The ones that choose not to are far more informed than the ones that choose to—because you have to go against the system.’ (Australia) [50]

In some low-income settings, access to ultrasound was limited due to lack of staff and other resources, as well as the costs incurred for women and the distance they would have to travel to attend appointments [53–56]. Some midwives in these contexts expressed the desire for training in the use of ultrasound, so that they could make decisions when other staff were not available [55, 57]. There were varying beliefs in relation to the safety of ultrasound as well as the diagnosis that could be made through its use [19, 34, 41, 49, 52, 58]. In some contexts, social and religious beliefs influenced the utility of a diagnosis if the only solution to a finding of fetal abnormality was termination [44, 59–61].

‘She [pregnant woman] didn’t go for ultrasound even though she was told to do so, she refused because of the cost.’ (Tanzania) [54]



* Reasons for exclusion: not experiences of ultrasound; not antenatal ultrasound; does not fit method criteria; unable to obtain full text; indistinguishable ultrasound data; limited ultrasound data

Fig 1. Screening and selection process.

<https://doi.org/10.1371/journal.pone.0261096.g001>

Table 1. Characteristics and quality rating of included studies.

First author	Date	Country	Resource setting	Participants	Sample size	Scan trimester	Study design/methods	Quality rating
Ahman ¹⁷	2010	Sweden	HIC	Women	11	Second	Naturalistic inquiry, in-depth interviews	A-
Baillie ¹⁸	2000	UK	HIC	Women	24	Second	Interpretative phenomenological analysis, semi-structured interviews	B+
Bashour ¹⁹	2005	Syria	LMIC	Women	30	General	Qualitative, semi-structured interviews	C
Carolan ²⁰	2009	Canada	HIC	Women	10	Second and third	Constructivist grounded theory	B-
Ekelin ²¹	2004	Sweden	HIC	Women (22) and partners (22)	44	Second	Grounded theory, interviews	B+
Larsson ²²	2010	Sweden	HIC	Women (5) and partners (4)	9	Second	Grounded theory, interviews	B+
Lou ²³	2017	Denmark	HIC	Women	20	First	Ethnography, semi-structured interviews	B+
Mitchel ²⁴	2004	Canada	HIC	Women	42	Second	Qualitative, semi-structured interviews	B-
Molander ²⁵	2010	Sweden	HIC	Women	10	First and second	Qualitative descriptive, interviews	B+
Thorpe ²⁶	1993	UK	HIC	Women	42	General	Qualitative, interviews	C-
Barr ²⁷	2013	UK	HIC	Women (17), partners (5), health workers (22)	44	First	Qualitative, focus groups	B-
Walsh ²⁸	2014	USA	HIC	Partners	22	Second	Qualitative, observation and semi-structured interviews	A-
Ahman ²⁹	2012	Sweden	HIC	Fathers	17	Second	Naturalistic inquiry, in-depth interviews	B+
Dheensa ³⁰	2013	UK	HIC	Women (6) and partners (12)	18	General	Grounded theory, semi structured	B
Dheensa ³¹	2015	UK	HIC	Fathers	12	General	Grounded theory, in-depth, semi-structured interviews	B-
Draper ³³	2002	UK	HIC	Fathers	18	Second	Ethnography, interviews	C
Pereira Silva Cardoso ³³	2018	Brazil	UMIC	Women	11	Second and third	Qualitative descriptive, semi-structured interviews	B+
Williams ³⁴	2005	UK	HIC	Women	15	First	Qualitative, semi-structured in-depth interviews	B
Edvardsson ³⁵	2018	Norway	HIC	Obstetricians	20	General	Qualitative, in-depth interviews	B
Hadicre ³⁶	2020	UK	HIC	Sonographers	14	General	Qualitative, semi-structured interviews	B-
Schwennesen ³⁷	2012	Denmark	HIC	Sonographers	7	First	Ethnography, semi-structured interviews	C
Ahman ³⁸	2015	Sweden	HIC	Obstetricians	11	General	Qualitative, interviews	A-
Ahman ³⁹	2019	Norway	HIC	Midwives	24	General	Qualitative, focus groups and interviews	B
Edvardsson ⁴⁰	2016	Sweden	HIC	Midwives	25	General	Exploratory qualitative, focus groups	B-
Firth ⁴¹	2011	Tanzania	LIC	Women	25	General	Descriptive, semi-structured and structured interviews	C
Ockleford ⁴²	2003	UK	HIC	Women	41	Second	Qualitative, semi-structured interviews	B-
Georges ⁴³	1996	Greece	HIC	Women (26) and health workers (16)	42	General	Ethnography, interviews, and observation	C-
Harris ⁴⁴	2008	UK	HIC	Women	34	General	Qualitative, interviews	C+
Jones ⁴⁵	2020	Kenya	LMIC	Women	50	First and second	Qualitative, in-depth semi-structured interviews	B
Liamputtong ⁴⁶	2002	Australia	HIC	Women	67	General	Ethnography, in-depth interviews	B+
Øyen ⁴⁷	2016	Norway	HIC	Women	8	General	Phenomenology, interviews	B

(Continued)

Table 1. (Continued)

First author	Date	Country	Resource setting	Participants	Sample size	Scan trimester	Study design/methods	Quality rating
Gammeltoft ⁴⁸	2007	Vietnam	LMIC	Women (116) and health workers (23)	139	General	Mixed methods, interviews, and observation	C
Gammeltoft ⁴⁹	2007	Vietnam	LMIC	Women	32	General	Phenomenology, in depth interviews	C
Edvardsson ⁵⁰	2015	Australia	HIC	Midwives	37	General	Qualitative, focus groups	C+
Sandelowski ⁵¹	1994	USA	HIC	Women	62	General	Qualitative, interviews	C-
Tsianakas ⁵²	2002	Australia	HIC	Women	15	General	Qualitative, in-depth interviews	B+
Ahman ⁵³	2016	Tanzania	LIC	Physicians	16	General	Qualitative, interviews	B+
Ahman ⁵⁴	2018	Tanzania	LIC	Midwives	31	General	Qualitative, focus groups	B+
Holmlund ⁵⁵	2017	Rwanda	LIC	Midwives	23	General	Qualitative, focus groups	B+
Scott ⁵⁶	2020	India	LMIC	Health workers	30	General	Qualitative, in-depth interviews	A-
Vesel ⁵⁷	2019	Kenya	LIC	Health workers	32	General	Qualitative, In-depth interviews and focus group discussions	B
Temam ⁵⁸	2011	USA	HIC	Women	25	General	Ethnography, interviews	B+
Gitsels ⁵⁹	2015	Holland	HIC	Women	12	General	Qualitative, interviews	C
Lewando-Hundt ⁶⁰	2001	Israel	HIC	Women (16) and health workers (20)	36	Second	Qualitative, in-depth interviews	C
Rice ⁶¹	1999	Australia	HIC	Women	30	Second	Qualitative, interviews and observation	C-
Gottfreosdottir ⁶²	2009	Iceland	HIC	Women (10) and partners (10)	20	First	Qualitative, semi-structured interviews	B-
Ledward ⁶³	2017	UK	HIC	Women	6	Second and third	Grounded theory, semi-structured interviews	C+
Doering ⁶⁴	2015	New Zealand	HIC	Women	13	Second	Qualitative descriptive, interviews	B-
Kristjansdottir ⁶⁵	2014	Iceland	HIC	Women	14	First	Phenomenology, semi-structured interviews	B+
Hawthorne ⁶⁶	2009	Australia	HIC	Women	20	First	Hermeneutic phenomenology, semi-structured interviews	B-
Larsson ⁶⁷	2009	Sweden	HIC	Women (5) and partners (4)	9	Second	Grounded theory, interviews	B+
Ekelin ⁶⁸	2016	Sweden	HIC	Women (10) and partners (6)	16	Second	Qualitative, interviews	B-
Gomes ⁶⁹	2007	Brazil	UMIC	Women	3	General	Qualitative, questionnaire and interviews	C-
Mabuke ⁷⁰	2011	Uganda	LIC	Women (50) and health workers (30)	80	General	Qualitative exploratory, semi-structured interviews	C
Bhagat ⁷¹	2012	India	MIC	Women (26) and girls (16)	42	General	Ethnography, focus groups	C
Ranji ⁷²	2012	Sweden	HIC	Women (9) and partners (9)	18	Second	Qualitative exploratory, in-depth interviews	B-
Denny ⁷³	2014	UK	HIC	Women	7	Second and third	Qualitative, semi-structured interviews	B-
Gomes ⁷⁴	2007	Brazil	UMIC	Women	3	General	Collective case study, semi-structured interviews	C
Edvardsson ⁷⁵	2014	Australia	HIC	Obstetricians	14	General	Qualitative, semi-structured interviews	A-
Edvardsson ⁷⁶	2015	Vietnam	LMIC	Obstetricians	17	General	Qualitative, semi-structured interviews	B+
Edvardsson ⁷⁷	2016	Rwanda	LIC	Physicians	19	General	Exploratory qualitative, semi-structured interviews	B
Dykes ⁷⁸	2001	Sweden	HIC	Women	12	Second	Grounded theory, in-depth interviews	B
Gagnon ⁷⁹	2020	Canada		Women	25	General	Qualitative, interviews	A
Walsh ⁸⁰	2020	USA	HIC	Women (22), partners (20), sonographers (7)	49	Second	Qualitative, observation	B

(Continued)

Table 1. (Continued)

First author	Date	Country	Resource setting	Participants	Sample size	Scan trimester	Study design/methods	Quality rating
Stephenson ⁸¹	2017	Australia	HIC	Health workers	27	First and second	Qualitative, interviews	B
Edvardsson ⁸²	2015	Australia	HIC	Obstetricians	14	General	Qualitative, interviews	B+
Holmlund ⁸³	2020	Vietnam	LMIC	Midwives	25	General	Qualitative, focus groups	A-
Stephenson ⁸⁴	2016	Australia	HIC	Women	26	First and second	Qualitative, semi-structured interviews	B
Gottfreisdottir ⁸⁵	2009	Iceland	HIC	Women (10) and partners (10)	20	First	Qualitative, semi-structured interviews	A-
Oscarsson ⁸⁶	2015	Sweden	HIC	Women	10	Second	Grounded theory, semi-structured interviews	B+
Asplin ⁸⁷	2012	Sweden	HIC	Women	27	Second	Exploratory descriptive, semi-structured interviews	B
Cristofalo ⁸⁸	2006	USA	HIC	Women	34	Second	Mixed methods, interviews	B
Van der Zalm ⁸⁹	2006	USA	HIC	Women	13	General	Qualitative, interviews	B
Sommerseth ⁹⁰	2010	Norway	HIC	Women	22	Second	Phenomenology, semi-structured interviews	B
Williams ⁹¹	2002	UK	HIC	Health workers	32	First	Qualitative, focus groups	B
Hammond ⁹²	2020	UK/ Netherlands	HIC	Women (15) and partners (1)	16	Second	Qualitative, semi-structured interviews	B
Denney-Koelsch ⁹³	2015	USA	HIC	Women (16) and partners (14)	30	General	Qualitative, naturalistic interviews	C+
Gammeltoft ⁹⁴	2007	Vietnam	LMIC	Women (30) and health workers (23)	53	General	Ethnography	B-
Jansson ⁹⁵	2010	Sweden	HIC	Nurses (4) and midwives (9)	13	Second	Qualitative, semi-structured interviews	B+
Reiso ⁹⁶	2020	Norway	HIC	Midwives	13	General	Qualitative, semi-structured interviews	B

<https://doi.org/10.1371/journal.pone.0261096.t001>

'We perceive that it is not out our job, but our wish as midwives is to be able to perform ultrasound so that we can play a role in the mother's care and make decisions without necessarily waiting for the availability of the doctor.' (Rwanda) [55]

'In our society it would be too late to do anything about that because the woman is not allowed, according to our religion, to have an abortion. Hence there is no point in doing tests during pregnancy. It's only a waste of time, money and effort.' (Israel) [60]

For some, beliefs about what was important to know during pregnancy, the value placed on ultrasound, and the impact of a diagnosis, appeared to be influenced by the vicarious experiences of friends, family and community members [17, 19, 28, 62, 63]. Information about the provision and nature of the ultrasound assessment appeared to also be mediated through community members in some cases, rather than healthcare professionals [29, 64, 65]. This extended to support after the scan which was often provided by friends and family [66–68].

'I needed help to sort out all my feelings and questions, my husband was a great support to me, but I would have liked to talk to my midwife.' (Iceland) [65]

Finding out the fetal sex was important for respondents in a range of contexts, in terms of imaging their future baby, and practical planning [28, 45, 48, 68–70]. However, in some circumstances, this knowledge had negative consequences [30, 71]. As reported by both health workers and community members, this was particularly (but not only) apparent in cultures

Table 2. Summary of Findings and CERQual assessment.

Overarching theme	Review finding	Findings	CERQual assessment			
			Methodological limitations	Adequacy of data	Coherence	Relevance
Sociocultural influences & expectations	Prenatal ultrasound as a clinical assessment or a social event	Quoties				
	Parents: Ultrasound is generally viewed as an integral part of pregnancy, to look forward to and not to be missed. It offers couples the chance to meet and bond with their baby and to share the news of their pregnancy with others. Fathers view attendance at the scan as part of their pregnancy, and a demonstration of their commitment to their partner and child. Attendance is also felt to be necessary for fathers to support their partner if complications are detected. For some couples, it offers a way to actively facilitate partner involvement in the pregnancy. Health workers: Providers sometimes found it difficult to reconcile their role as a clinician working in an environment assessing risk, with the expectations of parents who viewed the scan as an exciting event where they would see and ultimately share an image of their child for the first time.	Parents <i>No, the thought hadn't crossed my mind... I think that it's part of the pregnancy in some way to have an ultrasound.</i> (Ekelin 2004, Sweden) <i>It gives me a sense of security. With the first look at the fetus, even my husband would directly feel a sense of parenthood. He will be encouraged and you can feel that he has changed into a responsible person. Men should be involved in women's matters. They should not stop removed from them.</i> (Bashour 2005, Syria) Health workers: <i>The majority of them don't really come with any great belief that it's about anything other than tell me what gender it is and that I'm going to get lots of nice pictures of my baby.</i> (Hedder 2020, England)	Minor concerns about the methodological limitations of 5/23 studies contributing to the review finding, mainly around data collection and analysis phases	Few or minor concerns about adequacy of data as the finding is supported by rich data from a number of studies	Few or minor concerns around coherence as the data is consistent and supported by information from women and health workers	Few or minor concerns about relevance as the finding relates directly to the review question in HIC contexts
	Impact of routine ultrasound screening on women's autonomy and decision making	Parents <i>There's no reason to say no... we just read through it and we said, 'no harm', see, I think we're not experts in this baby thing anyway, so it's like whatever they offer, we would just take it, as long as it's not harming me or the baby, that's fine.</i> (Williams 2005, England) <i>The only thing is that looking back now I can't remember been given the opportunity to decide what I want. I was just recommended to me just being told that it was the normal procedure to have the test, to have the ultrasound. I can't remember been given the choice as such.</i> (Tsanakas 2002, Australia) Health workers: <i>Now it's almost like if you do not accept the offer (of an ultrasound examination), then you are a bad mother almost. (...) Are you irresponsible then? I don't know.</i> (Ahman 2019, Norway) <i>Do women get a choice though? Usually it's you get the slip to go and get your ultrasound.</i> (Edvardsson 2015b, Australia)	Minor concerns about the methodological limitations of 4/20 studies contributing to the review finding, mainly around data collection and analysis phases	Minor concerns around adequacy of data as the finding is supported by relatively rich data from a variety of settings and contexts	Minor concerns around coherence as the finding is framed around women's autonomy and incorporates both societal and professional influences with the latter more prevalent in LMICs	Few or minor concerns about relevance as the finding relates directly to the review question
The personal and social consequences of fetal gender identity	Parents: Finding out the sex of their child is important to parents across contexts. This can be to enable planning based on gender expectations, and knowledge of fetal sex appears to aid bonding for some. However, in some contexts, the desire to know fetal sex is driven by cultural and family preference for a male baby. Carrying a fetus identified as being of an undesirable sex can be a heavy burden for some, with severe consequences. Women in some cultural contexts report that ultrasound can result in female feticide. Some service users report that health workers may not disclose fetal sex if they are aware of the potential for culturally and socially influenced preferences and consequences. Health workers: In most settings and contexts the disclosure of gender identity following a scan was acknowledged to have significant consequences, both positive and negative. In some contexts, providers were aware of a preference for male babies and the potential for selective abortion. In some settings there is a policy of non-disclosure of gender identity to address this concern.	Parents <i>My baby was a boy and I was so happy. You know, having a boy is so important in Afghanistan. I wanted to have a boy and so did my mother.</i> (Ranjit 2012, Sweden) <i>... via USC people can know about sex of the baby and can get the girl child aborted.</i> (Bhagat 2012, India) Health workers: <i>There is this stigma between girls and boys, in some communities they want to know if it's a boy or a girl so that they may be able to either prevent the pregnancy from going on.</i> (Ahman 2016, Tanzania) <i>What often happens in ultrasound is you tell the woman the sex of the child she did not want. I think that this can be disturbing psychologically for the mother and, in the end, it can be harmful to the child.</i> (Holmlund 2017, Rwanda)	Minor concerns about the methodological limitations of 6/19 studies contributing to the review finding, mainly around data collection and analysis phases	Few or minor concerns around adequacy of data as the finding highlights the importance of scans in identifying the gender of the fetus (for parents) with the caveat that gender preference may have tragic implications in some contexts	Minor concerns around coherence as the finding highlights the importance of scans in identifying the gender of the fetus (for parents) with the caveat that gender preference may have tragic implications in some contexts	Few or very minor concerns about relevance as the finding relates directly to the review question

(Continued)

Table 2. (Continued)

Expectations about the nature and impact of antenatal ultrasound influences uptake	Findings		CERQual assessment					
	Parents	Parents, 11 studies: Bashour 2005, Syria (C); Firth 2011, Tanzania (C); Gamsjost 2007b, Vietnam (C); Griseis 2015, Holland (C); Kristiansdottir 2014, Iceland (B+); Lewandowski 2001, Israel (C)+; Rice 1999, Australia (C)+; Temam 2011, USA (B+); Tiansaka 2002, Australia (B+); Williams 2005, England (B)+	Moderate concerns about the methodological limitations of 7/17 studies supporting the review finding, largely related to data collection procedures and analysis of data	Minor concerns around adequacy of data as the finding is supported by relatively rich data from women and health workers in variety of different settings and contexts	Minor concerns around coherence as the finding incorporates a range of beliefs and understandings about ultrasound	Few or minor concerns about relevance as the finding relates directly to the review question	Moderate	Finding downgraded because of concerns about the methodological quality of some of the contributing studies
Parents: Fears around potential for harming the unborn baby; stories of misdiagnosis and false alarms, and religious and social beliefs with respect to the morals and timing of pregnancy termination for fetal abnormality may influence whether ultrasound is acceptable or believed to be necessary for women and their partner. Some women may misunderstand the nature of the diagnosis that can be made through ultrasound. Health workers: In some contexts, there were societal misunderstandings about how ultrasound scanning could harm the baby. Some felt the power of the technology was overestimated in society in general.	Parents <i>I went quite late in my pregnancy. I wanted to make sure that I was beyond two months. You see, I've heard ultrasound is not good in the first months of the pregnancy.</i> (Bashour 2005, Syria) <i>This is why I panic, because where do you draw the line, because people get things wrong with them at different severities don't they? And I think if you've got a baby, you love it—whatever it's got wrong with it, you still love it and protect it, don't you? I know people who would say, 'no, that baby's got something wrong with it, I'm not having it', ... but that's why the test would be hard for me, because I wouldn't be straightforward, 'oh, if there's something wrong, I'm not having it.'</i> (Williams 2005, England) Health workers <i>I think... most members of the public think an ultrasound is a more powerful tool than it is.</i> (Edvardsson 2014, Australia)	Parents, 11 studies: Bashour 2005, Syria (C); Firth 2011, Tanzania (C); Gamsjost 2007b, Vietnam (C); Griseis 2015, Holland (C); Kristiansdottir 2014, Iceland (B+); Lewandowski 2001, Israel (C)+; Rice 1999, Australia (C)+; Temam 2011, USA (B+); Tiansaka 2002, Australia (B+); Williams 2005, England (B)+ Health workers, 6 studies: Ahman, 2018, Tanzania (B); Edvardsson, 2014, Australia (A+); Edvardsson 2015b; Australia (C+); Holmlund 2014, Australia (A+); Holmlund Vietnam, 2020 (A+); Vesel 2019, Kenya (B)	Moderate concerns about the methodological limitations of 7/17 studies supporting the review finding, largely related to data collection procedures and analysis of data	Minor concerns around adequacy of data as the finding is supported by relatively rich data from women and health workers in variety of different settings and contexts	Minor concerns around coherence as the finding incorporates a range of beliefs and understandings about ultrasound	Few or minor concerns about relevance as the finding relates directly to the review question	Moderate	Finding downgraded because of concerns about the methodological quality of some of the contributing studies
	Parents <i>I know that other doctors show everything to the woman on the scan, but I kept going to that doctor anyway because he was competent. He is famous!</i> (Bashour 2005, Syria) <i>I've got a friend whose daughter is 15 with Down syndrome and it's not the worst.</i> (Ledward 2017, England) <i>I needed help to sort out all my feelings and questions, my husband was a great support to me, but I would have liked to talk to my midwife.</i> (Kristiansdottir 2014, Iceland)	Parents, 13 studies: Ahman 2010, Sweden (A)+; Ahman 2012, Sweden (B)+; Bashour 2005, Syria (C); Döring 2015, NZ (B+); Ekelin 2004, Sweden (B)+; Firth 2011, Tanzania (C); Gottfredsdottir 2009a, Iceland (B)+; Hammond 2020, England/Netherlands 2020, (B)+; Hawthorne 2009, Australia (B)+; Larsson 2009, Sweden (B)+; Ledward 2017, England (C)+; Vesel 2019, Kenya (B); Ocasarsson 2015, Sweden (B)+; Wadh 2014, USA (A)+	Minor concerns about the methodological limitations of 5/12 studies contributing to the review finding, mainly around data collection and analysis phases	Moderate concerns about adequacy of data due to relative inadequacy in how rich the data supporting this finding is	No or very minor concerns relating to coherence	Moderate concerns about relevance, as the majority of studies supporting this finding are from high income settings	Moderate	Minor concerns around adequacy of data as the finding is supported by relatively rich data from women and health workers in variety of different settings and contexts
	Coping with limited resources Health workers: In a variety of LMIC contexts providers recognized that limited resources affected their ability to provide an equitable service. In some contexts, women were only offered a scan in the event of a complication whilst in others a lack of equipment and trained staff restricted access. Health workers reported that, even where equipment and trained professionals were available, a lack of personal resources limited uptake by women. In some settings, midwives expressed the desire for training due to lack of availability of trained staff.	People from remote areas know about it [ultrasound] and would want to use it, but they cannot easily come here because they live far away from the hospital. (Holmlund 2017, Rwanda) <i>I wish in the future we will get training for us physicians, even nurses, for the basics, at least the basics to know a few things about how to do, how to examine the woman by using ultrasound, or how to determine different things which are important. It would be easier even if you are at night and you have one ultrasound machine here we can use.</i> (Ahman 2016, Tanzania)	Health workers, 6 studies: Ahman, 2018, Tanzania (B); Ahman 2016, Tanzania (B+); Holmlund 2017, Rwanda (B+); Scott, 2020, India (A+); Edvardsson 2016; Rwanda (B+); Vesel 2019, Kenya (B)	Few or very minor concerns about the methodological limitations of the studies contributing to the review finding	Moderate concerns about adequacy of data as the finding is supported by relatively thin data from a limited number of studies	Minor concerns about coherence as the finding reflects a variety of resource constraints as well as attempts to overcome resource limitations (by training other health workers to conduct scans)	Moderate concerns about relevance as the finding is supported by few data from MICs	Moderate
Overarching theme	Findings		CERQual assessment					
	Review finding	Quotes	Supporting studies	Adequacy of data	Coherence	Relevance	CERQual assessment	Comments

(Continued)

Table 2. (Continued)

The power of visual technology	Findings	CERQual assessment		
Parents: For many women, ultrasound is trusted as safe and is a valued technology that provides reassurance and a sense of security that their baby is developing normally. Health workers: Across a broad spectrum of settings and contexts healthcare providers viewed ultrasound as an essential component of pregnancy care, especially in complicated pregnancies. It is seen as a trusted intervention that optimises pregnancy outcomes and provides pleasure and reassurance to women and their partners.	Parents <i>'It's important for me to know if there is life inside, if everything looks fine.'</i> (Oyen 2016, Norway) <i>'I feel comfortable. The scan makes me feel psychologically eased. There is no point in going to the doctor if the scan is not available... It is my duty to go every month and follow up the situation of the fetus.'</i> (Bashour 2005, Syria) Health workers <i>'The stream [of requests] for early ultrasound is absolutely huge. People want to look as soon as they are pregnant. That is not how it was before.'</i> (Ahman 2019, Norway) <i>'Initially, I can say it came as an extra tool without really knowing why I have to do this. But, through getting used to the tools and doing it regularly, I came to get used to it and think right now I can say it is something we feel we cannot do without.'</i> (Vesd 2019, Kenya) <i>'I think it's been a big game changer in obstetric care and modern obstetrics is ingrained with ultrasound.'</i> (Edvardsson 2014, Australia)	Parents, 18 studies: Bashour 2005, Syria (C); Dheensa 2015, England (B-); Dheensa 2013, England (B-); Doering 2015, NZ (B-); Ekelin 2016, Sweden (B+)⁺; Firth 2011, Tanzania (C); Georges 1996, Greece (C); Gomes 2007b, Brazil (C); Gottfredsdottir 2009b, Iceland (B-); Harris 2008, Australia (C-); Hawthorne 2009, Australia (B-); Jones 2020, Kenya (B)⁺; Liangputing 2002, Australia (B-); Lou 2017, Denmark (B+); Oyen 2016, Norway (B); Rice 1999, Australia (C-); Tsanakas 2002, Australia (B+); Walsh 2014, USA (A+)⁺ Health workers, 14 Studies: Ahman, 2015, Sweden (A-); Ahman 2016, Tanzania (B+); Ahman 2018, Tanzania (B); Ahman 2019, Norway (B); Edvardsson 2014, Australia (A-); Edvardsson 2015, Vietnam (B+); Edvardsson 2015(b) Australia (C+); Edvardsson 2016, Australia (B-); Edvardsson 2016(b), Rwanda (B); Edvardsson 2018, Norway (B); Gamdlohoff & Nguyen 2007(a), Vietnam (C); Holmlund 2017, Rwanda (B+); Holmlund 2020, Vietnam (A-); Vesel 2019, Kenya (B)	Few or minor concerns about the methodological limitations of 6/29 studies contributing to the review finding, mainly around data collection and analysis phases Few or minor concerns around adequacy of data as the finding is supported by rich data from a number of studies Few or minor concerns about coherence as the finding is relatively consistent across all settings Few or minor concerns about relevance as the finding relates directly to the review question	High
Parents: In some contexts, the sense of contact provided by ultrasound, as well as the need to ensure ongoing normality, drives a need for frequent scanning. Ultrasound can also be prioritised over other forms of clinical assessment. For some women and their partners, there may be an unjustified expectation of the ability of ultrasound to detect and resolve complications. Health workers: In many settings, providers noted increased demand for antenatal scans and, in some contexts, the potential for overuse. Some providers in these contexts highlighted the unregulated nature of the scanning business, along with associated safety concerns. Health workers also noted the potential for diminishing clinical skills even in public care settings due to the overuse of ultrasound. In some clinical settings providers described examples of potentially serious undiagnosed conditions as women replaced formal antenatal appointments with scan appointments.	Parents: <i>'That's why I had scans constantly. To see how it was developing.'</i> (Gammliehoff 2007b, Vietnam) <i>'The first time I saw the baby, I was crazy with happiness. It was a contact with the child. Every time I went to the doctor, I wanted to see the child again.'</i> (Georges 1996, Greece) Health workers <i>'You don't know what to do and so you put on the probe and sometimes a few too many ultrasounds are done without any indication.'</i> (Ahman 2016, Sweden) <i>'Private clinics just conduct ultrasound scans for patients, but no examinations or tests. Therefore, they don't know in what state their patients are... In our department, some patients had surgery and died because they were in serious states of multi-organ dysfunction.'</i> (Holmlund 2020, Vietnam)	Parents 9 studies: Bashour 2005, Syria (C); Denny 2014, England (C+)⁺; Doering 2015, NZ (B-); Gammliehoff 2007a, Vietnam (C); Gammliehoff 2007b, Vietnam (C); Georges 1996, Greece (C); Ockelford 2003, England (B-)⁺; Oscarson 2015, Sweden (B+)⁺; Tsanakas 2002, Australia (B+) Health workers, 10 Studies: Ahman 2016, Tanzania (B+); Ahman, 2018 Tanzania (B); Edvardsson 2014, Australia; Edvardsson 2015, Vietnam (B+); Edvardsson 2015b, Australia; Edvardsson, 2016(b), Rwanda (C+); Edvardsson, 2019, Norway (B-); Holmlund Rwanda, 2017 (B+); Holmlund Vietnam, 2020 (A-); Vesel 2019, Kenya (B)	Minor concerns about the methodological limitations of 5/18 studies contributing to the review finding, mainly around data collection and analysis phases Minor concerns around coherence as the issues are highlighted by both health workers and service users. Minor concerns about relevance as the finding relates directly to the review question	High
Parents: For many women, visualisation of their baby through ultrasound offers objective confirmation of pregnancy and the existence of their child. The ultrasound scan provides a significant moment for couples to connect with their child, and to begin to visualise their future together, as a family. This opportunity may be particularly pertinent for fathers. Some parents begin to envisage their child's potential characteristics and personality through the scan image. Health workers: A number of providers felt that the visual representation of the fetus on a screen conferred identity as a person and facilitated parental bonding.	Parents <i>'And it was really fun to see this little baby on the screen and see it moving around, and that was really good. So, despite my initial anxiety it was good, it was a good experience, and I'd actually say to others it was a really nice thing to have done.'</i> (Harris 2008, Australia) <i>'And it became so very alive and I felt very close to the baby. Yes it felt like a fine moment, it was a very philosophic, emotional moment... It felt very good.'</i> (Ekelin 2004, Sweden) <i>'So it feels fun that we could get to know her personality already there.'</i> (Ekelin 2004, 2004 Sweden) Health workers <i>'All we needed was for you to talk to her! Now everything I read her to do she's doing. I think you got lucky with this one.'</i> (Walsh 2020, USA) <i>'You dealt with the unknown, when very few pregnant women had an ultrasound. Today I notice it more, but I myself have some trouble seeing the fetus as a fetus, I realize I want to think of it as a child.'</i> (Ahman 2019, Norway)	Parents, 20 papers: Denny 2014, England (C+)⁺; Dheensa 2013, England (B-); Draper 2002, England (C)⁺; Dykes 2001, Sweden (B)⁺; Ekelin 2004, Sweden (B+)⁺; Ekelin 2016, Sweden (B+)⁺; Firth 2011, Tanzania (C); Gagnon 2020, Canada (A)⁺; Georges 1996, Greece (C); Gomes 2007b, Brazil (C); Harris 2008, England (C+); Hawthorne 2009, Australia (B-); Lou 2017, Denmark (B+); Oyen 2016, Norway (B); Rasiq 2012, Sweden (B+)⁺; Rice 1999, Australia (C-); Stephenson 2016, Australia (B)⁺; Tsanakas 2002, Australia (B+); Walsh 2020, USA (B)⁺; Walsh 2014, USA (A-)⁺ Health workers 8 Studies: Ahman, 2015, Sweden (A-); Ahman 2016, Tanzania (B+); Ahman 2018, Tanzania (B); Ahman 2019, Norway (B); Edvardsson 2015, Vietnam (B-); Edvardsson 2015(c) Australia (B+); Edvardsson 2016, Australia (B-); Edvardsson 2018, Norway (B)	Few or minor concerns about the methodological limitations of 5/25 studies contributing to the review finding, rich data from a large number of studies Few or minor concerns around coherence as the finding is relatively consistent across all settings Minor concerns about relevance as the finding relates directly to the review question	High
Ultrasound legitimises the pregnancy and frames the fetus as a person Parents: Once mothers have seen the image of their baby on the screen, the potential consequences of the ultrasound, and decisions that may have to be made, become more complex. Health workers: Providers reported conflict between the welfare of the mother and that of the fetus. Many health workers commented that mothers would go to any lengths, often to their own detriment, to potentially improve the wellbeing of or treat their baby. Health workers frequently discussed the challenges of dealing with complex ethical and moral issues in the course of their work. These issues were often framed around the difficulties of prioritizing one life over another or the capacity to set aside personal beliefs when women and/or their partners held a different opinion. For some providers the scan image confirmed identity both as a person and a patient.	Parents <i>'Obviously I know they can't do these tests without showing you the scan, but it's easy to sit at home and say, "right, if they say this, we will obviously terminate the pregnancy", but when you see that baby on the screen, you don't care what it's got wrong with it, you just see that it's there and you know it's inside you... it must be a horrible decision once you've actually seen this is the baby inside you, to suddenly say, "no, I don't want to carry on with it". I think that must be quite a heartbreaking decision to make.'</i> (Williams 2005, England) Health workers <i>'We are accustomed to putting the mother's health first and foremost but that is sort of a balancing act'</i> (Ahman 2019, Norway) <i>'When she [the pregnant woman] understands that you are going to do something to help her baby, she does not refuse, she bears with it.'</i> (Holmlund 2017, Rwanda)	Parents, 5 studies: Baillie 2000, England; Ekelin 2016, Sweden (B+)⁺; Williams 2005, England (B-)⁺ Health workers, 13 Studies: Ahman, 2015, Sweden (A-); Ahman 2016, Tanzania (B+); Edvardsson 2014, Australia (A-); Edvardsson 2015, Vietnam (B+); Edvardsson 2015(b) Australia (C+); Edvardsson 2015(c) (B+); Edvardsson 2016, Australia (B-); Edvardsson 2018, Norway (B); Gamdlohoff & Nguyen 2007(c), Vietnam (B-)⁺; Holmlund 2017, Rwanda (B+); Holmlund 2020, Vietnam (A-); Stephenson 2017, Australia (B); Williams 2002, UK (C)⁺	Minor concerns about the methodological limitations of 2/17 studies contributing to the review finding, rich narrative from a number of studies Few or minor concerns as the finding title incorporates the range of moral and ethical dilemmas identified by health workers and service users Moderate concerns about the relevance as the finding relates directly to the review question, but there are few studies from the perspective of service users and they are all from two HICs	Moderate

(Continued)

Table 2. (Continued)

Findings		CERQual assessment							
Health workers	Findings	Supporting studies	Methodological limitations	Adequacy of data	Coherence	Relevance	CERQual assessment	Comments	
Health workers: Providers were sometimes left with feelings of uncertainty if the image was perceived to be ambiguous and feelings of anxiety if they thought they may have missed something. In some HIC settings these anxieties were enhanced by the potential for cesarean (or even litigation) in the event of an abnormality going undetected.	<i>'It's important to have knowledge about what you can see [with ultrasound], and to use the tool (...) so that you do the right things [during an ultrasound examination]. It can have consequences both ways, in terms of not detecting what is there and seeing things that aren't really there.'</i> (Ahman 2019, Norway) <i>'This would be down side of the area of medicine we're in is that the tension or the pressure is on, not to miss anything. That has increased dramatically.'</i> (Edvardsson 2014, Australia)	Parents, 9 studies: Ahman, 2015, Sweden (A-)*; Ahman 2018, Norway (B); Edvardsson 2014, Australia (A-)*; Edvardsson 2015, Australia (B+)*; Hadlere, 2020, UK (B-)*; Resio 2020, Norway (B); Scott 2020, India (A-)*; Stephenson, 2017, Australia (B); Williams 2002, UK (C)	Few or minor concerns about the methodological limitations of 1/9 studies contributing to the review finding	Minor concerns around adequacy of data as the finding is supported by relatively rich data from health workers	Moderate concerns about the finding encompasses the competencies, responsibilities and uncertainties of professional practice only (+) study from India)	Moderate concerns about relevance as the finding relates directly to the review question, though largely confined to HICs	Low	Finding downgraded because of limited data from LMICs and a lack of coherence across studies	
Overarching theme	Review finding	Quotes	Supporting studies	Methodological limitations	Adequacy of data	Coherence	Relevance	CERQual assessment	Comments
Key and evaluation: consequences of ultrasound findings	Adverse clinical findings are shocking and unexpected	Parents <i>'We were full of expectation and thought it would be very exciting to see our baby.'</i> (Larsson 2010, Sweden) <i>'It was a shock like this, because what we expect is that I will be everything perfect.'</i> (Gomes 2007a, Brazil) <i>'I realized that they actually weren't really looking as that they could give me a nice day it was actually been really looking as they were measuring all the bits and pieces.'</i> (Barr 2013, England) Health workers <i>'A lot of them think that they're going there, yay we get to see the baby and we get to find out the sex. They don't actually understand the significance of the scan and what we're looking for. And then if something does turn up, then they're completely devastated because nobody told me we were looking for that, I thought we were going to find out the sex of the baby.'</i> (Edvardsson 2015b, Australia)	Parents, 20 studies: Ahman 2010, Sweden (A)*; Ahman 2012, Sweden (B+)*; Asplin 2012, Sweden (B)*; Baillie 2012, England (B +)*; Barr 2013, England (B-)*; Carolan 2009, Canada (C+)*; ; Cristofalo 2006, USA (B)*; ; Denny 2014, England (C+)*; ; Denny-Koeth 2015, USA (B-)*; Draper 2002, England (C)*; Games 2007a, Brazil (C+); Hawthorne 2009, Australia (B-)*; Kristjansson 2014, Iceland (B+)*; Larsson 2009, Sweden (B+)*; Larsson 2010, Sweden (B+)*; ; Mitchell 2010, Canada (B+)*; ; Ocarsson 2015, Sweden (B+)*; ; Sommerseth 2010, Norway (B)*; Stephenson 2016, Australia (B)*; ; van der Zim 2006, USA (B) 2018, Norway (B); Barr 2013, UK (B-)*; Edvardsson 2016, Australia (B-)*; Edvardsson 2015(b) Australia (B); Edvardsson 2016(b) Rwanda (B); Hadlere, 2020, UK (B-)*; Williams 2002, UK (C)*	Few or minor concerns about the methodological limitations of 4/27 studies contributing to the review finding	Few or minor concerns about adequacy of data as the finding is supported by rich narrative data from a large number of studies	Few or minor concerns about coherence as the finding title incorporates both the adverse findings and the issue of lack of preparedness for them	Moderate concerns about relevance as the finding relates directly to the review question, but there are very little data from LMICs	Moderate	Finding downgraded because of limited data from LMICs
	A diagnosis of potential abnormality has both short- and long-term psychological consequences (even if the fetus is later found to be healthy)	Parents <i>'I can't ignore that there was actually something they saw there. Maybe it's an irrational fear but... I don't know how much I... you know with bonding, accepting the pregnancy.'</i> (Ahman 2010) <i>'There is another uncertainty in its place. That things may go wrong with a subsequent pregnancy and whether a new syndrome will arise or not during the next pregnancy.'</i> (Hammond 2020, England/Netherlands) <i>'I just want to put it out of my mind [the scans and worry]... I suppose I was a bit annoyed about it, but they get a all worried for nothing.'</i> (Carolan 2009) <i>'I don't want any more children. And maybe it's because of the stress I've had because of the uncertainties. I just never want to experience it again.'</i> (Hammond 2020, England/Netherlands) Health workers <i>'Those people are angry and bitter later when the child is normal and the pregnancy's been overshadowed by a lot of distress because of comments at the time of the ultrasound but it... a lot of stuff is grey and it's hard to get it right but there are definitely patients for whom we make the pregnancy more stressful.'</i> (Edvardsson 2014)	Parents, 16 studies: Ahman 2010, Sweden (A)*; Ahman 2012, Sweden (B+)*; Asplin 2012, Sweden (B)*; Baillie 2012, England (B +)*; ; Carolan 2009, Canada (C+)*; ; Cristofalo 2006, USA (B)*; ; Denny 2014, England (C+)*; ; ; Gottfredsdottir 2009b, Iceland (A-)*; ; Hammond 2020, England/Netherlands 2020, (B)*; ; Hawthorne 2009, Australia (B+)*; ; Kristjansson 2014, Iceland (B+)*; Larsson 2009, Sweden (B+)*; ; Larsson 2010, Sweden (B+)*; ; Mitchell 2010, Canada (B+)*; ; Ocarsson 2015, Sweden (B+)*; ; van der Zim 2006, USA (B) 2006, USA (B) Health workers, 7 studies: Ahman, 2015, Sweden (A-)*; Ahman 2018, Tanzania (B); Ahman 2019, Norway (B); Edvardsson 2018, Norway(B); Gundloft & Nguyen 2007(c) Vietnam (B-)*; Höhnland 2017, Rwanda (B+); Schwennsen 2012, Denmark (C)*	Few or very minor concerns about the methodological limitations of 5/23 studies contributing to the review finding	Minor concerns about adequacy of data as the finding is supported by rich narrative data from a variety of settings and contexts	Few or minor concerns about coherence as the finding title incorporates the impact of uncertainty of potentially abnormal diagnosis in the context of a range of actual outcomes	Moderate concerns about relevance, especially for service users, as there is relatively little data from LMICs	Moderate	Finding downgraded because of concerns about relevance in LMICs
	For pregnant women and their families, ultrasound in pregnancy enables planning and preparation when abnormalities are detected	<i>'If you are in such situation as this you want to know...'</i> (Firth 2011, Tanzania) <i>'It would make it possible to be prepared before the baby was born... if there is something to be done immediately or if any extra medication is needed... just when the baby is born.'</i> (Ekelin 2016, Sweden) <i>'I haven't told my son about the pregnancy yet, because I don't want to go to the scan and find out there is something wrong and have to terminate and then have to tell him... It's going to be hard enough me dealing with terminating this baby, let alone a 7 year old coping too.'</i> (Hawthorne 2009, Australia)	Parents, 10 studies: Ekelin 2004, Sweden (B+)*; Ekelin 2016, Sweden (B+)*; ; Firth 2011, Tanzania, Gottfredsdottir 2009b (B+)*; Hawthorne 2009 (B-)*; Larsson 2010, Sweden (B+)*; ; Mölander 2010, Sweden (B+)*; ; Ocarsson 2015, Sweden (B+)*; ; Tsimakas 2002, Australia (B+); Williams 2005 (B-)*	No or very minor concerns about the methodological limitations of the studies supporting this finding	Moderate concerns about adequacy due to lack of depth to the data, and relatively few studies supporting this finding	Few or minor concerns	Moderate concerns about relevance as this finding includes data from only one LMIC and the 5/10 of the HIC studies are from one country	Moderate	Finding downgraded due to lack of studies from LMICs and relatively few HIC countries represented, as well as lack of richness in the included data
	A positive, exciting, significant moment in pregnancy (the scan is 'normal')	<i>'It's a unique moment, there's nothing equal.'</i> (Gomes 2007b, Brazil) <i>'The scan, it stands out a mile'; 'a defining moment so far'</i> (Draper 2002, England) <i>'I was relieved. Not that I had been going round thinking about it before rather than that I had been in the back of my head. By the way I reacted I could feel that I was relieved anyway.'</i> (Ekelin 2004, Sweden)	Parents, 12 studies: Bashour 2005, Syria (C); Draper 2002, England (C)*; ; Dikes 2001, Sweden (B+)*; Ekelin 2004, Sweden (B+)*; ; Ekelin 2016, Sweden (B+)*; ; Firth 2011, Tanzania (C); Gammehoff 2007b, Vietnam (C); Gomes 2007b, Brazil (C); Jones 2020 Nairobi (B)* ; ; Oyen 2016, Norway (B); Stephenson 2016, Australia (B)* ; ; Williams 2005, England (B-)*	Moderate concerns about the methodological limitations of 4/11 of the studies contributing to the review finding, largely related to data collection procedures and analysis of data	Minor concerns about adequacy as the finding is supported by rich data and 12 studies	No or very minor concerns about coherence	Minor concerns about relevance as although the supporting studies are from a wide variety of setting, they include data from women only	Moderate	Finding downgraded because of concerns about methodological limitations

(Continued)

Table 2. (Continued)

The significance of relationship in the ultrasound encounter	Findings		CERQual assessment				
	Impact of staff attitudes, behaviours and communication skills on women and families		Few or very minor concerns about the methodological limitations of 2/23 studies contributing to the review finding	Few or minor concerns around adequacy of data as the finding is supported by rich data from a large number of studies	Minor concerns around coherence as both women and health workers identified relevant provider attributes	Moderate concerns about relevance as the finding is largely supported by data from HICs and more than half of the contributing studies come from Sweden and 6 from the USA	Moderate concerns about relevance because of concerns in particular the lack of data from LMICs and the prevalence of data from 2 countries
	Parents: Women and their partners want health workers to welcome them, to acknowledge the importance and unique nature of the situation for them as parents, and to provide relevant information. However, some women and couples experience a lack of any meaningful interaction and information during their ultrasound scan, leaving them excluded from their experience, and uncertain about the results. In some contexts, women will not ask health providers proactively for the information they need. Women were highly sensitive to non-verbal cues from providers during the scan. Long silences and being excluded from conversations, or not being able to view the ultrasound screen was anxiety-provoking for many. In contrast, being welcomed to and engaged in the scanning episode, and being provided with coherent and timely information during and after the scan, had a positive impact on the experience, even if it then resulted in an adverse diagnosis. Health workers: Providers often referred to communication as a significant but challenging aspect of their role. They highlighted the need for more time during a consultation to establish a rapport with parents with differing expectations, and to offer empathic and compassionate care when needed. This included professional, non-directive conversation, and facilitating parental engagement with the fetus whilst simultaneously looking for anomalies.	Parents: <i>‘Maybe that the sonographing midwife would ask a little about ... what expectations we had and ... if we had seen ultrasound imaging before, ... how we experienced that and ... what we hoped for and ... pause the imaging and ... well that she would ask if we were worried about something ... a little more time.’</i> (Molander 2010, Sweden) <i>‘I was like the expecting like a, ‘Hey how are you doing? Are you pretty excited about it? Like asking me, ‘How you feeling about this? I’d probably feel more welcome.’</i> (Walsh 2014, USA) <i>‘I can’t recall her looking at me, maybe just glancing, just looking at my tummy basically, I can’t recall her ever looking at me and saying, ‘Would you like to see your baby now?’ or ‘Would you like me to explain some things to you?’</i> (van der Zim 2006, USA) <i>‘With a little more empathy’ clinicians can ‘better guide people in dealing with uncertainty.’</i> (Hammond 2020, England/Netherlands) Health workers: <i>‘Even after all these years, there’s still times when you get a reaction or a question that you weren’t expecting and you stumble over your words and it’s almost like being back in the first couple of times you did it, again it’s totally thrown you.’</i> (Hardacre 2020, England)	Parents, 17 studies: Asplin 2012, Sweden (B)⁺; Baillie 2000, England (B)⁺; Badoir 2005, Syria (C); Cristofalo 2006, USA (B)⁺; Denney-Koelch 2015, USA (B); Ekelin 2004, Sweden (B)⁺; Gottfredsdottir 2009a, Iceland (B)⁻; Hammond 2020, England/Netherlands 2020, (B)⁺; Jones 2020, Kenya (B)⁺; Larsson 2009, Sweden (B)⁺; Larsson 2010, Sweden (B)⁺; Molander 2010, Sweden (B)⁺; Ranji 2012, Sweden (B)⁺; Sanddowski 1994, USA; van der Zim 2006, USA (B); Walsh 2014, USA (A-)⁺; Walsh 2020, USA (B)⁺ Health workers, 7 studies: Alhman 2015, Sweden (A-); Barr 2013, UK (B-); Hardacre 2020, UK (B-); Jansson 2010, Sweden (B)⁺; Reiso 2020, Norway (B); Stephenson 2017, Australia (B); Williams 2002, UK (C)[*]				
	A challenging role, with a need for training and emotional support Health workers: Providers expressed satisfaction in their ability to guide prospective parents through an ultrasound assessment and offer support during difficult conversations. However, they also discussed the difficulty of maintaining a professional and supportive persona whilst also dealing with their own emotions. A few also talked about their sense of responsibility, the relative isolation of their role and the importance of peer support in their personal and professional development. Particular challenges were a lack of time to form relationships and communicate results, a lack of adequate training in the communication of abnormal results and the need for a more holistic approach to their engagement with service users.	<i>‘I know that I am providing good care to women at a terrible time in their lives. And whether the outcome is good or bad, they know that what could have been done, reasonably was done, that it was done by people who cared about them and knew what they were talking about. And that’s very rewarding.’</i> (Edvardsson 2014, Australia) <i>‘If you didn’t deliver the happy-clappy scan to them ... they would complain. Irrespective if you’ve had ... you know ... you’d told the previous patients some really bad news, you weren’t allowed to be just a little bit down or you had to put on the show for the next patient.’</i> (Hardacre 2020, England)	Health workers, 12 Studies: Alhman 2015, Sweden (A-); Alhman 2018, Tanzania (B); Alhman 2019, Norway (B); Edvardsson 2015(c) Australia (B+); Edvardsson 2016(b); Rwanda (B); Hardacre 2020, UK (B-); Holmlund 2017, Rwanda (B+); Gamellott & Nguyen 2007(c) Vietnam (B-); Reiso 2020, Norway (B); Stephenson 2020, Australia (B); Vest 2019, Kenya (B); Williams 2002, UK (C)[*]				

* = first trimester ultrasound scan; ** = second trimester scan

<https://doi.org/10.1371/journal.pone.0261096.t002>

where there is a preference for male babies. In these contexts, the disclosure of female fetal sex through ultrasound could result in feticide [19, 53, 71, 72]. To avoid this potential outcome there was a policy of non-disclosure relating to fetal sex to avoid this outcome [19, 53, 54, 71, 72].

'USG is done to know the sex of the child and then abortion is done if its female child.' (India) [71]

There is this stigma between girls and boys, in some communities they want to know if it's a boy or a girl so that they may be able to either prevent the pregnancy from going on.' (Tanzania) [53]

'... via USG people can know about sex of the baby and can get the girl child aborted.' (India) [71]

The power of visual technology. For most respondents, ultrasound was seen as central to antenatal care. Women generally trusted it as a valued technology that could provide confirmation of their pregnancy and reassurance of fetal wellbeing [19, 28, 30, 43, 64, 66, 73]. For providers, it was an important tool, particularly for the detection and management of complications [39, 43, 53, 57, 74–76]. However, some respondents reported that a reliance on ultrasound results in the potential for overuse, and consequent neglect of other forms of antenatal care [19, 53, 74]. Some participants felt compelled towards ultrasound to visualise their baby and for reassurance [19, 31, 43, 47, 61]. For some women and healthcare professionals, ultrasound held greater value than other forms of antenatal assessment. The overuse of ultrasound was felt to result in reduced clinical skills and the potential to miss complications that were not picked up through this form of assessment [38, 43, 48, 55, 74].

'The scan is very necessary; there is no point in visiting the doctor without seeing the fetus and knowing how well it is doing. You would not benefit at all!' (Syria) [19]

'Initially, I can say it came as an extra tool without really knowing why I have to do this. But, through getting used to the tools and doing it regularly, I came to get used to it and think right now I can say it is something we feel like we cannot do without.' (Kenya) [57]

'I think that in Vietnam nowadays, obstetric ultrasound is the most important investigation to monitor the pregnancy. Some other investigations like blood test, urine test also have importance but they cannot be compared to the obstetric ultrasound.' (Vietnam) [75]

'I think it's a very useful tool, I think we're getting to the situation where many people can do nothing without an ultrasound, so those clinical skills have gone to a large extent.' (Australia) [74]

For many women and healthcare professionals, the power of the ultrasound image was significant [32, 32, 43, 50, 54, 66, 73, 75–77]. Some women appeared to lack trust that they were pregnant until they were able to visualise the image of their baby [21, 25, 44, 52, 68, 72, 78]. The capacity for visualisation was particularly valued by fathers and other parents [21, 28, 61, 78]. The scan image offered the chance to visualise the future together as family. For some, it represented an opportunity to construct their child's future personality and characteristics [32, 21, 73, 79]. However, this sense of connection also complicated decisions around termination of pregnancy [18, 30, 34, 68].

‘Before I found out I was pregnant I’d always said if I knew I was having a handicapped baby, I’d have a termination, but then when I went for the very first scan and saw the baby moving about and saw his heart beating, I thought afterwards I don’t know whether I could do it now, because he’s alive, it’s a person.’ (England) [18]

Some providers were concerned that the clarity of the ultrasound image meant that all complications should be visible and identified [39]. Some feared the potential for consequences for both the mother, and for their professional security, if abnormalities were missed [36, 39, 76]. In some LMIC contexts, concerns were also expressed about the lack of appropriate training and the potential for this to result in missed complications or misdiagnosis [38, 39, 76, 80]. Some respondents described professional and moral dilemmas around prioritising either mother or fetus in their clinical assessments [35, 40, 81, 82], as well ethical concerns when parents made decisions that did not fit with personal or professional beliefs [55, 80, 82]. Some also expressed concern that women would go to any lengths to protect the wellbeing of their baby, even when this was to their own detriment [38, 75, 81].

‘No special training on ultrasound, that’s the limitation, that’s why you can sometimes miss some complications if I find something I am not understanding.’ (Rwanda) [76]

‘I have never met an expectant mother who has hesitated to expose herself to something that might be harmful to her health as long as it benefits the fetus.’ (Sweden) [38]

Both joy and devastation; consequences of ultrasound findings. The scan appointment was a source of great excitement, joy and relief for many couples, providing a chance to bond with their baby, whilst also instilling a sense of responsibility, particularly amongst fathers and other co-parents [19, 21, 32, 41, 45, 68, 77, 83]. For some, it also offered the potential for choice and the opportunity to plan when complications were detected [22, 68, 84, 85]. However, for many, the identification of abnormalities was completely unexpected [17, 18, 20, 24, 65, 69, 73, 80, 86–88]. Some reported deep shock and distress on hearing this news [17, 65, 67, 69, 73, 86–89]. Both service users and healthcare professionals reflected on how this shock could be compounded by couples’ expectations that the scan appointment is a happy event that would provide confirmation of wellbeing [24, 36, 65, 83]. The difficulty in getting the balance right in preparing couples for potential consequences of the scan was also discussed by healthcare professionals. Some felt that they lacked time to do this, amongst all the other issues to be discussed in an appointment, and they struggled to get the balance between discussing risk and maintaining a sense of normality prior to the scan [27, 37, 90].

... it’s making sure that they know enough but not frightening them or making them feel very negative about the pregnancy ... not put too much emphasis on the possibility of problems.’ (England) [27]

‘We were so naive. We thought we were going to see the baby and get a nice photo.’ (Canada) [24]

“It was a shock like this, because what we expect is that it will be everything perfect” (Brazil) [69]

‘You come to find out the sex of the baby and have the bomb dropped on you.’ (USA) [87]

Uncertain findings that could, but may not, indicate abnormality, were particularly difficult for many couples, resulting in feelings of having lost their pregnancy, and a shift to a new

tentative, risky state [18, 20, 29, 91]. Some women reported detaching themselves from their pregnancy and/or baby while also experiencing constant worry in relation to their baby's well-being [17, 18, 22]. This state persisted into the long term for some, even after a follow-up diagnosis that all was well [18, 20, 91]. In some cases, this concern persisted even into infancy, with, at the extreme, the decision not to pursue previously planned future pregnancies [18, 20, 91]. Some health professionals were acutely aware of the impact of uncertain findings on parents, resulting in dilemmas around whether these should be disclosed [36, 74, 81]. Parents were also conflicted about the benefits versus the harms of disclosing these findings [17, 29]. Some expressed regret in retrospect about the negative impact on their pregnancy [20, 87, 65, 67].

'Because of this I wouldn't have a third child. . . I'm not putting myself through this stress again ever, and I would have gone on to have a third one. We're stopping at two.' (England) [18]

'The more you see sometimes the more uncertain things get. And you can ruin a pregnancy quite a bit like that. So I'm not sure whether it's always good.' (Australia) [74]

The significance of relationship in the ultrasound encounter. Women and partners expressed a desire for scan providers to recognise the unique nature of the scan experience for them, to make them feel welcome, and to provide information and the opportunity to ask questions [21, 22, 25, 76, 65, 88]. Their actual experiences ranged from health workers being cold, disinterested, and lacking time to provide information, to those who were warm and engaging, and actively fostered questions and interest in the scan [18, 19, 22, 72, 80, 92]. In some contexts, women reported that they were unable to ask questions and that their experience was completely in the hands of the healthcare professional [19, 92]. Some women and their partners reported being completely excluded from their scan experience, unable to see the image of their baby, and left in silence to guess through body language what might be happening [18, 19, 22, 87].

'He was staring for a long time at the screen. You see he is very good. He keeps looking [she waves as if she is reading from a book], and he keeps explaining. He told me about the [amniotic fluid]. My previous doctor was different. She does the scan very quickly and tells you: 'Hey stand up. . . you have nothing' and that's all. I tell you, I felt the difference between those two doctors.' (Syria) [19]

For some health workers supporting women through difficult findings was a rewarding aspect of their role; but they expressed the desire for more training in the communication of abnormal results, as well as more professional support to confirm findings [36, 37, 93–95]. A lack of time to form relationships and properly communicate results meant that some providers felt the need to distance themselves, in order to protect their own emotions and to enable them to perform consecutive scans within a limited time period [36, 90, 95].

'It's the responsibility of being alone in such a small place, I'm the only one looking. . . I miss a colleague, so I could say "Could you take a look with me, let's discuss this together.' (Norway) [95]

'You've got to protect yourself, you've got to . . . not harden your heart, but you do have to protect yourself and not get too emotionally involved, because otherwise you wouldn't survive very long in our job.' (England) [36]

Discussion

In 2019, the WHO maternal and perinatal health steering group prioritised updating their early ultrasound scan recommendations [5]. This systematic review informs the subsequent recommendations and will inform living guideline updates of this recommendation [96]. The potential drivers for appropriate or inappropriate use of ultrasound were captured in the four study themes.

In line with other studies [6], the experience of providing or receiving ultrasound was generally seen as positive in our analysis [21, 25, 34, 38, 39, 41, 97], generating high demand for scans [19, 39, 43, 49, 50, 55, 64, 74], but the consequences of adverse findings was sometimes devastating [18, 20, 50, 65, 67, 73, 74, 87]. Importantly, in this review, we found that even when an initial concern was later ruled out, there were very significant long-term adverse consequences for some service users [17, 18, 20, 67, 91]. Respondents also reported overuse, with implications for the provision of other antenatal assessments and potential loss of clinical skills [19, 38, 48, 53, 55, 74, 82]. This reinforces previously published survey data from a range of settings [98–100].

Provider attitudes and behaviours were influential in the service user experience [18, 19, 21, 22, 72, 86, 88], as were local social norms [18, 21, 25, 34, 41, 52, 58, 60, 61] and access to follow up investigations and support [21, 22, 67, 86, 87]. Providers reported concerns around missing important features of the scan [38, 39, 75, 96], and a lack of sufficient time and training to appropriately carry out ultrasound assessments [36, 38, 76, 90, 95].

Previous survey research has found mixed evidence about the impact of ultrasound screening on maternal anxiety [101]. Our data suggest possible drivers for the varying perceptions of ultrasound screening. The power of the visual in making the fetus ‘real’ is evident in our analysis [21, 23, 28, 32, 35, 43, 44, 50, 73], reinforcing the validity of concepts of what has been termed the ‘*tentative pregnancy*’, in which women put their sense of being pregnant on hold until they have visual evidence of the fetus, and of its wellbeing [102]. Our data show that visual markers with unknown provenance or meaning can be unsettling for health workers as well as for service users [17, 18, 20, 38, 50, 74, 81]. The value of diagnosing abnormality was less clear in contexts where termination was not an option [58, 60, 61]. The critical, ethical and equity issue of female feticide reported in some settings underpins growing concerns about sex selection, linked to a much lower female-male sex ratio than would be expected in some countries [13, 68, 69, 103].

Our findings raise questions about the utility of ultrasound in pregnancy as a screening tool in settings where the implications of features on the scans are not always understood by practitioners or service users [100, 104–107], and/or if there are no effective follow up, treatment, or solution to some ultrasound findings [108–110]. They raise concerns about the use of ultrasound as a deliberate ‘draw’ to bring women into antenatal care, if the consequence is overuse by undertrained staff, without time to undertake the scan effectively, including provision of tailored information and psychosocial support where needed; and without effective, affordable, equitable referral pathways.

The strengths of this review include the comprehensive search that was not restricted by language or date, and the inclusion of 80 qualitative studies covering countries from most regions of the world. Fourteen of the 17 review findings were assessed as high or moderate confidence evidence using the GRADE CERQual approach [16]. We have included the experiences and perspectives of women and their partners, as well as health workers, from low-, middle- and high-income countries. Limitations include that we were unable to distinguish between first and second trimester ultrasound in our findings, as the findings were not clearly separated, or they were similar in both trimesters. We were also unable to include the views of

policy makers or funders, as our search did not retrieve any eligible studies that included this perspective. Furthermore, many of the findings relate to identification and diagnosis of abnormality, rather than to assessment of gestational age, fetal growth, or multiple pregnancy. The majority of studies in our review are from high-income countries, which was anticipated, but the inclusion of more studies from low-income settings may have provided further implications for the use of ultrasound services in this context. Thirteen of the included studies were from the CROss-Country Ultrasound Study (CROCUS). However, these studies explored the views of both providers and service users, from a number of different low-, middle-, and high-income countries.

This review offers a critical insight into how countries can introduce and maintain optimal routine antenatal ultrasound services. The findings reinforce the psychological and emotional benefits of such services from the point of view of most women and their partners, and the clinical benefits as perceived by service providers. However, there are implications for implementation in settings where antenatal ultrasound is not yet a routine component of antenatal care, and improvements that can be made in other settings where use of this technology is already established. In all settings, and particularly those with restricted resources, adequate education and training in both the use of obstetric ultrasound and in positive interactions with service users is essential, as well the allowance of sufficient time to undertake the scan effectively and with attention to the needs of the parents. Mitigation against overuse is important, to ensure that the use of ultrasound is appropriately balanced with the provision of expert clinical antenatal care. The potential for, and consequences of fetal sex disclosure must also be considered, especially in contexts where there is sociocultural bias towards male sex. Improvements can be made in all settings to ensure that women and their partners make autonomous informed decisions relating to the uptake of antenatal ultrasound; that they are adequately involved during the scanning procedure; and that information relating to the results is provided in a timely and supportive manner.

Future research should consider the ways in which ultrasound might be implemented to ensure equity of access, follow up, and longer term social and psychological support where this is needed, so that the positive aspects are maintained, while limiting the potential for overuse and for adverse impacts. There is a need to determine what is necessary and optimal to disclose with regard to markers of unclear significance and to consider how couples can be optimally supported through uncertain findings, and through to future reproductive decision making. Consideration should be given to the whole maternity and health care system into which ultrasound is introduced. Research into the use of portable ultrasound may be relevant for all settings, but particularly within LMICs, where this may be a requirement for rural and remote provision of ultrasound. This would require the ability to produce scan images of sufficient quality, as well as consideration of the findings of this review.

Supporting information

S1 Checklist. PRISMA 2020 checklist.
(PDF)

S1 Table. Search strategy.
(PDF)

Author Contributions

Conceptualization: Maria Barreix, Özge Tunçalp, Soo Downe.

Data curation: Gill Moncrieff.

Formal analysis: Gill Moncrieff, Kenneth Finlayson, Soo Downe.

Funding acquisition: Soo Downe.

Investigation: Gill Moncrieff, Kenneth Finlayson, Sarah Cordey, Rebekah McCrimmon, Catherine Harris.

Methodology: Gill Moncrieff, Kenneth Finlayson, Soo Downe.

Project administration: Maria Barreix, Özge Tunçalp, Soo Downe.

Supervision: Kenneth Finlayson, Soo Downe.

Validation: Gill Moncrieff, Kenneth Finlayson, Soo Downe.

Visualization: Gill Moncrieff, Kenneth Finlayson.

Writing – original draft: Gill Moncrieff, Soo Downe.

Writing – review & editing: Gill Moncrieff, Kenneth Finlayson, Sarah Cordey, Rebekah McCrimmon, Catherine Harris, Özge Tunçalp, Soo Downe.

References

1. Whitworth M, Bricker L, Mullan C. Ultrasound for fetal assessment in early pregnancy. *Cochrane Database Syst Rev*. 2015: CD007058. <https://doi.org/10.1002/14651858.CD007058.pub3> PMID: 26171896
2. Stewart KA, Navarro SM, Kambala S, et al. Trends in ultrasound use in low and middle income countries: A systematic review. *Int J MCH AIDS*. 2020; 9:103–120. <https://doi.org/10.21106/ijma.294> PMID: 32123634
3. Goley SM, Sakula-Barry S, Adofo-Ansong N, et al. Investigating the use of ultrasonography for the antenatal diagnosis of structural congenital anomalies in low-income and middle-income countries: a systematic review. *BMJ Paediatr Open*. 2020; 4: e000684. <https://doi.org/10.1136/bmjpo-2020-000684> PMID: 32864479
4. Hitimana R, Lindholm L, Krantz G, Nzayirambaho M, Pulkki-Brännström A-M. Cost of antenatal care for the health sector and for households in Rwanda. *BMC Health Serv Res*. 2018; 18. <https://doi.org/10.1186/s12913-018-3013-1> PMID: 29631583
5. World Health Organisation. WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience; 2016.
6. Garcia J, Bricker L, Henderson J, et al. Women's views of pregnancy ultrasound: a systematic review. *Birth*. 2002; 29: 225–250. <https://doi.org/10.1046/j.1523-536x.2002.00198.x> PMID: 12431263
7. Ebrashy A, Kurjak A, Adra A, et al. Controversial ultrasound findings in mid trimester pregnancy. Evidence based approach. *J Perinat Med*. 2016; 44: 131–137. <https://doi.org/10.1515/jpm-2015-0223> PMID: 26506099
8. Kaplan R, Adams S. Incidental fetal ultrasound findings: Interpretation and management. *J Midwifery Womens Health*. 2018; 63: 323–329. <https://doi.org/10.1111/jmwh.12754> PMID: 29763964
9. Prentice G, Welsh A, Howat A, Ross D, Henry A. Isolated 'soft signs' of fetal choroid plexus cysts or echogenic intracardiac focus—consequences of their continued reporting. *Australas J Ultrasound Med*. 2019; 22: 253–264. <https://doi.org/10.1002/ajum.12180> PMID: 34760567
10. Garel M, Gosme-Seguret S, Kaminski M, Cuttini M. Ethical decision-making in prenatal diagnosis and termination of pregnancy: a qualitative survey among physicians and midwives. *Prenat Diagn*. 2002; 22: 811–817. <https://doi.org/10.1002/pd.427> PMID: 12224077
11. Bakst S, Romano-Zelekha O, Ostrovsky J, Shohat T. Determinants associated with making prenatal screening decisions in a national study. *J Obstet Gynaecol*. 2019; 39: 41–48. <https://doi.org/10.1080/01443615.2018.1463977> PMID: 30244627
12. Ghosh R, Sharma AK. Missing female fetus: a micro level investigation of sex determination in a peri-urban area of Northern India. *Health Care Women Int*. 2012; 33: 1020–1034. <https://doi.org/10.1080/07399332.2011.610537> PMID: 23066964
13. Pham BN, Hall W, Hill PS. Indirect evidence of the contribution of prenatal sex selection practices to the high sex ratio at birth in Vietnam. *J Popul Res*. 2011; 28: 293–299.

14. Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol*. 2008; 8: 45. <https://doi.org/10.1186/1471-2288-8-45> PMID: 18616818
15. Walsh D, Downe S. Appraising the quality of qualitative research. *Midwifery*. 2006; 22: 108–119. <https://doi.org/10.1016/j.midw.2005.05.004> PMID: 16243416
16. Lewin S, Booth A, Glenton C, et al. Applying GRADE-CERQual to qualitative evidence synthesis findings: introduction to the series. *Implement Sci*. 2018; 13. <https://doi.org/10.1186/s13012-017-0688-3> PMID: 29384079
17. Ahman A, Runestam K, Sarkadi A. Did I really want to know this? Pregnant women's reaction to detection of a soft marker during ultrasound screening. *Patient Educ Couns*. 2010; 81: 87–93. <https://doi.org/10.1016/j.pec.2009.12.011> PMID: 20083367
18. Baillie C, Smith J, Hewison J, Mason G. Ultrasound screening for chromosomal abnormality: Women's reactions to false positive results. *Br J Health Psychol*. 2000; 5: 377–394.
19. Bashour H, Hafez R, Abdulsalam A. Syrian women's perceptions and experiences of ultrasound screening in pregnancy: Implications for antenatal policy. *Reprod Health Matters*. 2005; 13: 147–154. [https://doi.org/10.1016/s0968-8080\(05\)25164-9](https://doi.org/10.1016/s0968-8080(05)25164-9) PMID: 16035608
20. Carolan M, Hodnett E. Discovery of soft markers on fetal ultrasound: maternal implications. *Midwifery*. 2009; 25: 654–664. <https://doi.org/10.1016/j.midw.2007.11.002> PMID: 18314233
21. Ekelin M, Crang-Svalenius E, Dykes A-K. A qualitative study of mothers' and fathers' experiences of routine ultrasound examination in Sweden. *Midwifery*. 2004; 20: 335–344. <https://doi.org/10.1016/j.midw.2004.02.001> PMID: 15571882
22. Larsson A-K, Svalenius EC, Lundqvist A, Dykes A-K. Parents' experiences of an abnormal ultrasound examination—vacillating between emotional confusion and sense of reality. *Reprod Health*. 2010; 7: 10. <https://doi.org/10.1186/1742-4755-7-10> PMID: 20546610
23. Lou S, Frumer M, Schlütter MM, Petersen OB, Vogel I, Nielsen CP. Experiences and expectations in the first trimester of pregnancy: a qualitative study. *Health Expect*. 2017; 20: 1320–1329. <https://doi.org/10.1111/hex.12572> PMID: 28521069
24. Mitchell LM. Women's experiences of unexpected ultrasound findings. *J Midwifery Womens Health*. 2004; 49: 228–234. <https://doi.org/10.1016/j.jmwh.2003.11.004> PMID: 15134676
25. Molander E, Alehagen S, Berterö CM. Routine ultrasound examination during pregnancy: a world of possibilities. *Midwifery*. 2010; 26: 18–26. <https://doi.org/10.1016/j.midw.2008.04.008> PMID: 18571818
26. Thorpe K, Harker L, Pike A, Marlow N. Women's views of ultrasonography. A comparison of women's experiences of antenatal ultrasound screening with cerebral ultrasound of their newborn infant. *Soc Sci Med*. 1993; 36: 311–315. [https://doi.org/10.1016/0277-9536\(93\)90015-v](https://doi.org/10.1016/0277-9536(93)90015-v) PMID: 8426975
27. Barr O, Skirton H. Informed decision making regarding antenatal screening for fetal abnormality in the United Kingdom: a qualitative study of parents and professionals: Informed consent and antenatal screening. *Nurs Health Sci*. 2013; 15: 318–325. <https://doi.org/10.1111/nhs.12034> PMID: 23347127
28. Walsh T, Tolman R, Davis R, Palladino C, Romero V, Singh V. Moving Up the "Magic Moment": Fathers' Experience of Prenatal Ultrasound. *Fathering: A Journal of Theory, Research, and Practice about Men as Fathers*. 2014; 12: 18–37.
29. Åhman A, Lindgren P, Sarkadi A. Facts first, then reaction—expectant fathers' experiences of an ultrasound screening identifying soft markers. *Midwifery*. 2012; 28: e667–75. <https://doi.org/10.1016/j.midw.2011.07.008> PMID: 21920645
30. Dheensa S, Williams R, Metcalfe A. Shattered Schemata and Fragmented Identities: Men's Experiences of Antenatal Genetic Screening in Great Britain. *J Fam Issues*. 2013; 34: 1081–1103.
31. Dheensa S, Metcalfe PA, Williams R. What do men want from antenatal screening? Findings from an interview study in England. *Midwifery*. 2015; 31: 208–214. <https://doi.org/10.1016/j.midw.2014.08.011> PMID: 25242107
32. Draper J. 'It was a real good show': the ultrasound scan, fathers and the power of visual knowledge. *Sociol Health Illn*. 2002; 24: 771–795.
33. Pereira Silva Cardoso VE, Da Silva Junior AJ, Bonatti AF, Soares dos Santos GW, Nascimento Ribeiro TA. The partner's involvement in the prenatal routine through the pregnant women perspective / A participação do parceiro na rotina pré-natal sob a perspectiva da mulher gestante. *Rev Pesqui Cuid é Fundam*. 2018; 10: 856–862.
34. Williams C, Sandall J, Lewando-Hundt G, Heyman B, Spencer K, Grellier R. Women as moral pioneers? Experiences of first trimester antenatal screening. *Soc Sci Med*. 2005; 61: 1983–1992. <https://doi.org/10.1016/j.socscimed.2005.04.004> PMID: 15899542

35. Edvardsson K, Åhman A, Fagerli TA, et al. Norwegian obstetricians' experiences of the use of ultrasound in pregnancy management. A qualitative study. *Sex Reprod Healthc*. 2018; 15: 69–76. <https://doi.org/10.1016/j.srhc.2017.12.001> PMID: 29389504
36. Hardicre NK, Arezina J, McGuinness A, Johnson J. Managing the unmanageable: A qualitative study exploring sonographer experiences of and training in unexpected and difficult news delivery. *Radiography*. 2021; 27: 369–376. <https://doi.org/10.1016/j.radi.2020.09.015> PMID: 33077330
37. Schwennesen N, Koch L. Representing and intervening: 'doing' good care in first trimester prenatal knowledge production and decision-making: 'Doing' good care in prenatal decision-making. *Sociol Health Illn*. 2012; 34: 283–298. <https://doi.org/10.1111/j.1467-9566.2011.01414.x> PMID: 22257243
38. Åhman A, Persson M, Edvardsson K et al. Two sides of the same coin—an interview study of Swedish obstetricians' experiences using ultrasound in pregnancy management. *BMC Pregnancy Childbirth*. 2015; 15. <https://doi.org/10.1186/s12884-015-0743-5> PMID: 26589489
39. Åhman A, Edvardsson K, Fagerli TA, et al. A much valued tool that also brings ethical dilemmas—a qualitative study of Norwegian midwives' experiences and views on the role of obstetric ultrasound. *BMC Pregnancy Childbirth*. 2019; 19: 33. <https://doi.org/10.1186/s12884-019-2178-x> PMID: 30651083
40. Edvardsson K, Lalos A, Åhman A, Small R, Graner, MD, PhD S, Mogren I. Increasing possibilities—Increasing dilemmas: A qualitative study of Swedish midwives' experiences of ultrasound use in pregnancy. *Midwifery*. 2016; 42: 46–53. <https://doi.org/10.1016/j.midw.2016.09.009> PMID: 27788415
41. Firth ER, Mlay P, Walker R, Sill PR. Pregnant women's beliefs, expectations and experiences of antenatal ultrasound in Northern Tanzania. *Afr J Reprod Health*. 2011; 15: 91–107. PMID: 22590896
42. Ockleford E, Berryman J, Hsu R. Do women understand prenatal screening for fetal abnormality? *Br J Midwifery*. 2003; 11: 445–449
43. Georges E. Fetal ultrasound imaging and the production of authoritative knowledge in Greece. *Med Anthropol Q*. 1996; 10: 157–175. <https://doi.org/10.1525/maq.1996.10.2.02a00040> PMID: 8744082
44. Harris G, Connor L, Bisits A, Higginbotham N. "Seeing the Baby": Pleasures and Dilemmas of Ultrasound Technologies for Primiparous Australian Women. *Med Anthropol Q*. 2004; 18: 23–47. <https://doi.org/10.1525/maq.2004.18.1.23> PMID: 15098426
45. Jones RM, Vesel L, Kimenju G et al. Implementation of the INTERGROWTH-21st gestational dating and fetal and newborn growth standards in Nairobi, Kenya. *Glob Health Action*. 2020; 13: 1770967. <https://doi.org/10.1080/16549716.2020.1770967> PMID: 32544027
46. Liamputtong P, Watson L. The voices and concerns about prenatal testing of Cambodian, Lao and Vietnamese women in Australia. *Midwifery*. 2002; 18: 304–313. <https://doi.org/10.1054/midw.2002.0331> PMID: 12473445
47. Øyen L, Aune I. Viewing the unborn child—pregnant women's expectations, attitudes and experiences regarding fetal ultrasound examination. *Sex Reprod Healthc*. 2016; 7: 8–13. <https://doi.org/10.1016/j.srhc.2015.10.003> PMID: 26826039
48. Gammeltoft T, Nguyen HTT. The commodification of obstetric ultrasound scanning in Hanoi, Viet Nam. *Reprod Health Matters*. 2007; 15: 163–171.
49. Gammeltoft T, Nguyen HTT. Fetal conditions and fatal decisions: ethical dilemmas in ultrasound screening in Vietnam. *Soc Sci Med*. 2007; 64: 2248–2259. <https://doi.org/10.1016/j.socscimed.2007.02.015> PMID: 17403563
50. Edvardsson K, Mogren I, Lalos A, Persson M, Small R. A routine tool with far-reaching influence: Australian midwives' views on the use of ultrasound during pregnancy. *BMC Pregnancy Childbirth*. 2015; 15. <https://doi.org/10.1186/s12884-015-0632-y> PMID: 26311437
51. Sandelowski M. Separate, but less unequal: Fetal ultrasonography and the transformation of expectant mother/fatherhood. *Gend Soc*. 1994; 8: 230–245.
52. Tsianakas V, Liamputtong P. Prenatal testing: the perceptions and experiences of Muslim women in Australia. *J Reprod Infant Psychol*. 2002; 20: 7–24. <https://doi.org/10.1080/02646830220106767> PMID: 12784816
53. Åhman A, Kidanto H, Ngarina M, Edvardsson K, Small R, Mogren I. 'Essential but not always available when needed'—an interview study of physicians' experiences and views regarding use of obstetric ultrasound in Tanzania. *Glob Health Action*. 2016; 9: 31062. <https://doi.org/10.3402/gha.v9.31062> PMID: 27452066
54. Åhman A, Edvardsson K, Lesio Kidanto H, Ngarina M, Small R, Mogren I. 'Without ultrasound you can't reach the best decision'—Midwives' experiences and views of the role of ultrasound in maternity care in Dar Es Salaam, Tanzania. *Sexual & Reproductive Healthcare*. 2018; 15: 28–34. <https://doi.org/10.1016/j.srhc.2017.11.007> PMID: 29389498

55. Holmlund S, Ntaganira J, Edvardsson K et al. Improved maternity care if midwives learn to perform ultrasound: a qualitative study of Rwandan midwives' experiences and views of obstetric ultrasound. *Glob Health Action*. 2017; 10: 1350451. <https://doi.org/10.1080/16549716.2017.1350451> PMID: 28764602
56. Scott K, Gupta S, Williams E, Arthur M, Somayajulu U, Noguchi L. "I can guess the month . . . but beyond that, I can't tell" an exploratory qualitative study of health care provider perspectives on gestational age estimation in Rajasthan, India. *BMC Pregnancy Childbirth*. 2020; 20: 529. <https://doi.org/10.1186/s12884-020-03201-6> PMID: 32917163
57. Vesel L, Nimako K, Jones RM, et al. Implementing the INTERGROWTH-21st gestational dating and fetal and newborn growth standards in peri-urban Nairobi, Kenya: Provider experiences, uptake and clinical decision-making. *PLoS One*. 2019; 14: e0213388. <https://doi.org/10.1371/journal.pone.0213388> PMID: 30849125
58. Teman E, Ivry T, Bernhardt BA. Pregnancy as a proclamation of faith: Ultra-Orthodox Jewish women navigating the uncertainty of pregnancy and prenatal diagnosis. *Am J Med Genet A*. 2011; 15: 69–80. <https://doi.org/10.1002/ajmg.a.33774> PMID: 21204212
59. Gitsels-van der Wal JT, Martin L, Manniën J, Verhoeven P, Hutton EK, Reinders HS. A qualitative study on how Muslim women of Moroccan descent approach antenatal anomaly screening. *Midwifery*. 2015; 31: e43–9. <https://doi.org/10.1016/j.midw.2014.12.007> PMID: 25592301
60. Lewando-Hundt G, Shoham-Vardi I, Beckerleg S, Belmaker I, Kassem F, Jaafar AA. Knowledge, action and resistance: the selective use of pre-natal screening among Bedouin women of the Negev, Israel. *Soc Sci Med*. 2001; 52: 561–569. [https://doi.org/10.1016/s0277-9536\(00\)00160-x](https://doi.org/10.1016/s0277-9536(00)00160-x) PMID: 11206653
61. Rice PL, Naksook C. Pregnancy and technology: Thai women's perceptions and experience of prenatal testing. *Health Care Women Int*. 1999; 20: 259–278. <https://doi.org/10.1080/073993399245755> PMID: 10633624
62. Gottfredsdóttir H, Björnsdóttir K, Sandall J. How do prospective parents who decline prenatal screening account for their decision? A qualitative study. *Soc Sci Med*. 2009; 69: 274–277. <https://doi.org/10.1016/j.socscimed.2009.05.004> PMID: 19481846
63. Ledward. Pregnant women's experiences of screening for fetal abnormalities according to NICE guidelines: How should midwives communicate information? *Evidence Based. Midwifery*. 2017; 15: 112–119.
64. Doering K, Patterson J, Griffiths C. Experience of the New Zealand maternity care system by a group of Japanese women in one centre. *N Z. Coll Midwives*. 2015; 51: 5–10.
65. Kristjansdóttir H, Gottfredsdóttir H. Making sense of the situation: women's reflection of positive fetal screening 11–21 months after giving birth. *Midwifery*. 2014; 30: 643–649. <https://doi.org/10.1016/j.midw.2013.10.024> PMID: 24332751
66. Hawthorne F, Ahern K. "Holding our breath": the experiences of women contemplating nuchal translucency screening. *App Nurs Res*. 2009; 22: 236–242. <https://doi.org/10.1016/j.apnr.2008.02.007> PMID: 19875037
67. Larsson A-K, Crang-Svalenius E, Dykes A-K. Information for better or for worse: interviews with parents when their foetus was found to have choroid plexus cysts at a routine second trimester ultrasound. *J Psychosom Obstet Gynaecol*. 2009; 30: 48–57. <https://doi.org/10.1080/01674820802621775> PMID: 19308783
68. Ekelin M, Persson L, Välimäki A, Crang Svalenius E. To know or not to know—parents' attitudes to and preferences for prenatal diagnosis. *J Reprod Infant Psychol*. 2016; 34: 356–369.
69. Gomes AG, Piccinini CA. Impressões e sentimentos das gestantes sobre a ultra-sonografia e suas implicações para a relação materno-fetal no contexto de anormalidade fetal. *PSICO*. 2007; 38: 65–76.
70. Mubuke AG. An exploratory study of the views of Ugandan women and health practitioners on the use of sonography to establish fetal sex. *Pan Afr Med J*. 2011; 9:36. <https://doi.org/10.4314/pamj.v9i1.71214> PMID: 22355436
71. Bhagat N, Laskar AR, Sharma N. Women's perception about sex selection in an urban slum in Delhi. *J Reprod Infant Psychol*. 2012; 30: 92–104.
72. Ranji A, Dykes A-K, Ny P. Routine ultrasound investigations in the second trimester of pregnancy: the experiences of immigrant parents in Sweden. *J Reprod Infant Psychol*. 2012; 30: 312–325.
73. Denny E, Quinlan-Jones E, Bibila S, Kilby M. The experience of pregnant women with a diagnosis of fetal lower urinary tract obstruction (LUTO). *Midwifery*. 2014; 30: 636–642. <https://doi.org/10.1016/j.midw.2013.10.023> PMID: 24290423

74. Edvardsson K, Small R, Persson M, Lalos A, Mogren I. 'Ultrasound is an invaluable third eye, but it can't see everything': a qualitative study with obstetricians in Australia. *BMC Pregnancy Childbirth*. 2014; 14: 363. <https://doi.org/10.1186/1471-2393-14-363> PMID: 25336335
75. Edvardsson K, Graner S, Thi LP, et al. 'Women think pregnancy management means obstetric ultrasound': Vietnamese obstetricians' views on the use of ultrasound during pregnancy. *Glob Health Action*. 2015; 8: 28405. <https://doi.org/10.3402/gha.v8.28405> PMID: 26519131
76. Edvardsson K, Ntaganira J, Åhman A, Sengoma JPS, Small R, Mogren I. Physicians' experiences and views on the role of obstetric ultrasound in rural and urban Rwanda: a qualitative study. *Trop Med Int Health*. 2016; 21: 895–906. <https://doi.org/10.1111/tmi.12718> PMID: 27125579
77. Dykes K, Stjernqvist K. The importance of ultrasound to first-time mothers' thoughts about their unborn child. *J Reprod Infant Psychol*. 2001; 19: 95–104.
78. Gagnon R. L'enfant à naître: dispositifs de contrôle dès que la vie paraît. *Enfances, Familles, Generations*. 2020
79. Walsh TB. Your baby is so happy, active, uncooperative: How prenatal care providers contribute to parents' mental representations of the baby. *Midwifery*. 2020; 83: 102630. <https://doi.org/10.1016/j.midw.2020.102630> PMID: 32006801
80. Stephenson N, Mills C, McLeod K. "Simply providing information": Negotiating the ethical dilemmas of obstetric ultrasound, prenatal testing and selective termination of pregnancy. *Fem Psychol*. 2017; 27: 72–91.
81. Edvardsson K, Small R, Lalos A, Persson M, Mogren I. Ultrasound's 'window on the womb' brings ethical challenges for balancing maternal and fetal health interests: obstetricians' experiences in Australia. *BMC Med Ethics*. 2015; 16. <https://doi.org/10.1186/s12910-015-0023-y> PMID: 25953252
82. Holmlund S, Lan PT, Edvardsson K, et al. Sub-optimal use of ultrasound examinations may result in underperformance of Vietnamese maternity care—A qualitative study of midwives' experiences and views. *Sex Reprod Healthc*. 2020; 24: 100508. <https://doi.org/10.1016/j.srhc.2020.100508> PMID: 32278314
83. Stephenson N, McLeod K, Mills C. Ambiguous encounters, uncertain fetuses: Women's experiences of obstetric ultrasound. *Fem Rev*. 2016; 113: 17–33.
84. Gottfredsdóttir H, Sandall J, Björnsdóttir K. 'This is just what you do when you are pregnant': a qualitative study of prospective parents in Iceland who accept nuchal translucency screening. *Midwifery*. 2009; 25: 711–720. <https://doi.org/10.1016/j.midw.2007.12.004> PMID: 18295382
85. Oscarsson M, Gottvall T, Swahnberg K. When fetal hydronephrosis is suspected antenatally—a qualitative study. *BMC Pregnancy Childbirth*. 2015; 15. <https://doi.org/10.1186/s12884-015-0791-x> PMID: 26694546
86. Asplin N, Wessel H, Marions L, Georgsson Öhman S. Pregnant women's experiences, needs, and preferences regarding information about malformations detected by ultrasound scan. *Sex Reprod Healthc*. 2012; 3: 73–78. <https://doi.org/10.1016/j.srhc.2011.12.002> PMID: 22578754
87. Cristofalo EA, Dipietro JA, Costigan KA, Nelson P, Crino J. Women's response to fetal choroid plexus cysts detected by prenatal ultrasound. *J Perinatol*. 2006; 26: 215–223. <https://doi.org/10.1038/sj.jp.7211489> PMID: 16554849
88. Van der Zalm JE, Byrne PJ. Seeing baby: women's experience of prenatal ultrasound examination and unexpected fetal diagnosis. *J Perinatol*. 2006; 26: 403–408. <https://doi.org/10.1038/sj.jp.7211540> PMID: 16801957
89. Sommereth E, Sundby J. Women's experiences when ultrasound examinations give unexpected findings in the second trimester. *Women Birth*. 2010; 23: 111–116. <https://doi.org/10.1016/j.wombi.2010.01.001> PMID: 20153992
90. Williams C, Alderson P, Farsides B. Dilemmas encountered by health practitioners offering nuchal translucency screening: a qualitative case study. *Prenat Diagn*. 2002; 22: 216–220. <https://doi.org/10.1002/pd.289> PMID: 11920897
91. Hammond J, Klapwijk JE, Hill M, et al. Parental experiences of uncertainty following an abnormal fetal anomaly scan: Insights using Han's taxonomy of uncertainty. *J Genet Couns*. 2021; 30: 198–210. <https://doi.org/10.1002/jgc4.1311> PMID: 32638447
92. Denney-Koelsch EM, Côté-Arsenault D, Lemcke-Berno E. Parents' experiences with ultrasound during pregnancy with a lethal fetal diagnosis. *Glob Qual Nurs Res*. 2015; 2. <https://doi.org/10.1177/2333393615587888> PMID: 26878030
93. Gammeltoft T. Sonography and sociality: obstetrical ultrasound imaging in urban Vietnam. *Med Anthropol Q*. 2007; 21: 133–153. <https://doi.org/10.1525/maq.2007.21.2.133> PMID: 17601081

94. Jansson C, Adolfsson A. A Swedish study of midwives' and nurses' experiences when women are diagnosed with a missed miscarriage during a routine ultrasound scan. *Sex Reprod Healthc.* 2010; 1: 67–72. <https://doi.org/10.1016/j.srhc.2010.01.002> PMID: 21122599
95. Reiso M, Langli B, Sommerseth E, Johannessen A. A qualitative study of the work experiences of midwives performing obstetric ultrasound in Norway. *BMC Pregnancy Childbirth.* 2020; 20: 641. <https://doi.org/10.1186/s12884-020-03333-9> PMID: 33087091
96. Vogel JP, Dowswell T, Lewin S, et al. Developing and applying a 'living guidelines' approach to WHO recommendations on maternal and perinatal health. *BMJ Glob Health.* 2019; 4: e001683 <https://doi.org/10.1136/bmjgh-2019-001683> PMID: 31478014
97. Gomes AG, Piccinini CA. Impressões e sentimentos de gestantes em relação à ultra-sonografia obstétrica no contexto de normalidade fetal. *Psicologia: Reflexão e Crítica.* 2007; 20: 179–187.
98. Holmlund S, Lan P, Edvardsson K et al. Health professionals' experiences and views on obstetric ultrasound in Vietnam: a regional, cross-sectional study. *BMJ Open.* 2019; 9: e031761. <https://doi.org/10.1136/bmjopen-2019-031761> PMID: 31548354
99. Ker C-R, Kuo Y-L, Chan T-F. A survey of current use, dilemma and outlook of antenatal ultrasonography in Taiwan. *Taiwan J Obstet Gynecol.* 2019; 58: 820–826. <https://doi.org/10.1016/j.tjog.2019.09.017> PMID: 31759535
100. Ranji A, Dykes A-K. Ultrasound screening during pregnancy in Iran: womens' expectations, experiences and number of scans. *Midwifery.* 2012; 28: 24–29. <https://doi.org/10.1016/j.midw.2010.10.005> PMID: 21041007
101. Lou S, Mikkelsen L, Hvidman L, Petersen OB, Nielsen CP. Does screening for Down's syndrome cause anxiety in pregnant women? A systematic review. *Acta Obstet Gynecol Scand.* 2015; 94: 15–27. <https://doi.org/10.1111/aogs.12482> PMID: 25146713
102. Rothman BK. *The Tentative Pregnancy: Prenatal Diagnosis and the Future of Motherhood.* Penguin Books; 1987.
103. Roberts LR, Montgomery SB. India's distorted sex ratio: Dire consequences for girls. *J Christ Nurs.* 2016; 33: E7–E15. <https://doi.org/10.1097/CNJ.0000000000000244> PMID: 28286369
104. Gourounti K, Lykeridou K, Daskalakis G, Glentis S, Sandall J, Antsaklis A. Women's perception of information and experiences of nuchal translucency screening in Greece. *Fetal Diagn Ther.* 2008; 24: 86–91. <https://doi.org/10.1159/000142133> PMID: 18648204
105. Kinnevey C, Kawooya M, Tumwesigye T, Douglas D, Sams S. Addressing obstetrical challenges at 12 rural Ugandan health facilities: Findings from an international ultrasound and skills development training for midwives in Uganda. *Int J MCH AIDS.* 2016; 5: 46–52. <https://doi.org/10.21106/ijma.106> PMID: 28058192
106. Matiang'i M, Joosse K, Ngunju P, et al. Barriers and enablers that influence utilization of ultrasound screening services among antenatal women in kajiado and Kisii counties Kenya. *Open J Clin Diagn.* 2021; 11: 1–17.
107. Ugwu A, Osungbade E, Erundu F. Maternal perspectives of prenatal sonogram in a north-eastern population in Nigeria. *Libyan J Med.* 2009; 4: 140–142. <https://doi.org/10.4176/090424> PMID: 21483536
108. Franklin HL, Mirza W, Swanson DL, et al. Factors influencing referrals for ultrasound-diagnosed complications during prenatal care in five low and middle income countries. *Reprod Health.* 2018; 15: 204. <https://doi.org/10.1186/s12978-018-0647-8> PMID: 30541560
109. Stefanovic V. Role of obstetric ultrasound in reducing maternal and neonatal mortality in developing countries: From facts to acts. *J Ultrasound Obstet Gynecol.* 2020; 14: 43–49.
110. Swanson DL, Franklin HL, Swanson JO, et al. Including ultrasound scans in antenatal care in low-resource settings: Considering the complementarity of obstetric ultrasound screening and maternity waiting homes in strengthening referral systems in low-resource, rural settings. *Semin Perinatol.* 2019; 43: 273–281. <https://doi.org/10.1053/j.semperi.2019.03.017> PMID: 30979599