

Central Lancashire Online Knowledge (CLOK)

Title	Cross-sectional and longitudinal relationships between cardiorespiratory fitness and health-related quality of life in primary school children in England: the mediating role of psychological correlates of physical activity
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/43418/
DOI	https://doi.org/10.1177/17579139221118771
Date	2022
Citation	Solera-Sanchez, Alba, Christian, Danielle, Beltran-Valls, Maria Reyes, Adelantado- Renaua,, Mireira, Martin-Smith, Rhona, MacDonald, Mahiri, Tyler, Richard and Fairclough, Stuart (2022) Cross-sectional and longitudinal relationships between cardiorespiratory fitness and health-related quality of life in primary school children in England: the mediating role of psychological correlates of physical activity. Perspectives in Public Health.
Creators	Solera-Sanchez, Alba, Christian, Danielle, Beltran-Valls, Maria Reyes, Adelantado- Renaua,, Mireira, Martin-Smith, Rhona, MacDonald, Mahiri, Tyler, Richard and Fairclough, Stuart

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1177/17579139221118771>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Cross-sectional and longitudinal relationships between cardiorespiratory fitness and health-related quality of life in primary school children in England: the mediating role of psychological correlates of physical activity

Alba Solera-Sanchez^a, Danielle L. Christian^b, Maria Reyes Beltran-Valls^a, Mireia Adelantado-Renau^a, Rhona Martin-Smith^c, Mahiri J. MacDonald^c, Richard Tyler^c, Stuart J. Fairclough^c

^a LIFE Research Group, Department of Education, University Jaume I, CP 12071 Castellon, Spain.

^b Faculty of Health and Wellbeing, University of Central Lancashire, Preston, Lancashire, PR1 2HE, UK.

^c Movement Behaviours, Health and Wellbeing Research Group, Department of Sport and Physical Activity, Edge Hill University, Ormskirk, Lancashire L39 4QP, UK.

Corresponding author: Maria Reyes Beltran Valls, Av. de Vicent Sos Baynat, s/n, PC 12071 Castellon, Spain. Phone: 00 34 964 72 96 82, Fax: 00 34 964 72 92 64, email: vallsm@uji.es.

Acknowledgements: We are grateful for the organisational support from West Lancashire Sport Partnership, and thank the children and teachers for their engagement in the project. Written informed consent and assent were required to participate in the project in accordance with the project approvals granted by the University Research Ethics Committee (#SPA-REC-2015-182).

Funding: The study was funded by West Lancashire Borough Council and Lancashire County Council.

Author's contributions: SJF was involved in the planning and development of the project, data analysis and manuscript write-up; DLC was involved in the planning and development of the project and manuscript write-up; ASS was involved in the literature search and screening, data analysis, and manuscript write-up; MRBV and MAR contributed to data analysis, and manuscript write-up; RMS, MJM and RT were in the manuscript write-up. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests: All authors declare that they have no competing interests.

Abstract

Purpose. The aims were (i) to analyse the cross-sectional and longitudinal associations between children's cardiorespiratory fitness (CRF) and health-related quality of life (HRQoL), and (ii) to examine whether these associations were mediated by physical activity self-efficacy and physical activity enjoyment.

Methods. This study involved 383 children (10.0 ± 0.5 years) recruited from 20 primary schools in northwest England. Data were collected on two occasions 12 weeks apart. The number of laps completed in the 20m Shuttle Run Test was used as the CRF indicator. HRQoL was assessed using the KIDSCREEN-10 questionnaire. Physical activity self-efficacy and enjoyment were assessed with the social-cognitive and Physical Activity Enjoyment Scale questionnaires, respectively. Linear mixed models with random intercepts (schools) assessed associations between CRF and HRQoL cross-sectionally, and longitudinally. Boot-strapped mediation procedures were performed, and indirect effects (IE) with 95% confidence intervals (CI) not including zero considered as statistically significant. Analyses were adjusted for sex, time of the year, socioeconomic status, waist-to-height ratio, maturation and physical activity.

Results. CRF was cross-sectionally associated with HRQoL ($\beta=0.09$; 95%CI=0.02, 0.16, $p=0.015$). In the longitudinal analysis, CRF at baseline was associated with HRQoL at 12 weeks after additionally controlling for baseline HRQoL ($\beta=0.08$; 95%CI=0.002, $p=0.15$, $p=0.045$). Cross-sectionally, physical activity self-efficacy and enjoyment acted individually as mediators in the relationship between CRF and HRQoL (IE=0.069; 95%CI:0.038; $p=0.105$ and IE=0.045; 95%CI:0.016; $p=0.080$, respectively). In the longitudinal analysis physical activity self-efficacy showed a significant mediating effect (IE=0.025; 95%CI=0.004; $p=0.054$).

Conclusions. Our findings highlight the influence of CRF on children's psychological correlates of physical activity and their overall HRQoL.

Key words: Health, physical fitness, quality of life, youth.

1. Introduction

Health-related quality of life (HRQoL) is a multidimensional concept which reflects an individual's own perception of their physical, mental, social health, and functionality.¹ HRQoL has been highlighted as an important health indicator² since perceived well-being and functionality are considered important components of health surveillance.³ Indeed, investigating HRQoL has been nowadays considered relevant due to its relationship with both self-reported chronic diseases (e.g., diabetes, breast cancer, arthritis, and hypertension) and their risk factors (e.g., body mass index, physical inactivity, sleep patterns, diet quality, and smoking status).⁴ Measuring HRQoL can help to determine the burden of preventable disease, injuries, and disabilities, and can provide valuable new insights into the relationships between HRQoL and risk factors.⁵ Thus, over the past twenty-five years, HRQoL has become an important outcome in healthy children, being commonly examined by professionals, such as clinicians, caregivers, educators, or public health authorities due to a collective interest towards the subjective perception and evaluation of an individual's own life.⁶⁻⁸ Given the importance of HRQoL, identifying factors that may contribute to improving children's HRQoL is a public health priority.

Among possible factors influencing children's HRQoL, previous cross-sectional studies have revealed significant positive associations with cardiorespiratory fitness (CRF). CRF represents a measure of the body's ability to deliver and use oxygen to support muscular activity during physical activity⁹ and is considered an important health marker.¹⁰ Previous research suggests that CRF may be a potentially useful strategy to enhance children's HRQoL, however evidence of this relationship is limited to cross-sectional studies.¹¹⁻¹³ For example, Andersen et al.'s study of 1129 schoolchildren aged 10 years, showed that CRF was positively associated with overall HRQoL.¹¹ Another study including 415 children aged between 8 and 9 years reported a positive weak correlation between CRF and HRQoL in boys, but not in girls.¹² Moreover, the study of Redondo-Tébar et al., which involved 1413 younger children, aged 4 to 7 years, concluded that children with higher CRF levels had greater HRQoL.¹³ While informative, these studies cannot explain the dynamic processes that could occur over time, neither provide a long-term perspective of the influence that CRF might have on HRQoL, which could contribute to understanding the determinants of children's health outcomes.¹⁴ Thus, to strengthen the current evidence base, investigation into the longitudinal associations between CRF and HRQoL is warranted.¹⁵

CRF has been considered a physiological component that has been reported to influence psychological correlates of physical activity.^{16,17} This is possibly due to the impact that sufficient levels of CRF have on brain functioning (e.g., serotonin), self-worth, life satisfaction¹⁶ and the reward system.¹⁷ Indeed, previous literature reported that children with higher levels of CRF had stronger psychological correlates of physical activity, such as physical activity self-efficacy¹⁸ and physical activity enjoyment¹⁶ compared to low CRF peers. Thus, CRF seems to be an important

attribute positively influencing psychological correlates. On the other hand, two previous studies which implemented new school playground activities, reported positive associations between children's physical activity enjoyment and HRQoL.^{20,21} Taken together, it is plausible that the positive association between CRF and HRQL in children is explained through the influence that CRF exerts on the psychological correlates.

Based on this previous research, there is a need for future studies to investigate variables influencing HRQoL, specifically focusing on CRF as a variable that could impact HRQoL as well as other physical activity correlates,²² which could act as possible underlying mechanisms in that association. This will be of interest for health authorities seeking to improve children's overall HRQoL through the implementation of educational interventions at schools and the design of public health strategies. Therefore, the aims of this research were (i) to analyse the cross-sectional and longitudinal association between children's CRF and HRQoL, and (ii) to examine whether these associations were mediated by physical activity self-efficacy and physical activity enjoyment separately, as key psychological correlates of physical activity.

2. Material and methods

2.1 Study design

This observational study used baseline and follow-up data from the *Active West Lancs* primary school physical activity and wellbeing programme. The aim of this programme was to evaluate the impact of a combined educational and exercise programme designed to promote and enhance children's physical activity behaviours and knowledge, fitness, and wellbeing. The programme aligned to the UK government's Childhood Obesity Strategy recommendation for children to engage in 30 minutes of physical activity during the school day.²³ The programme was delivered in four clusters of five schools over four consecutive 12-week phases between 2018 and 2019. As no significant pre-post changes were observed in CRF and HRQoL outcomes over the 12-weeks, for this study the baseline data were treated as cross-sectional, and the combined 12-week follow-up longitudinal data were treated as longitudinal.

2.2 Participants

The 20 schools were situated in West Lancashire, northwest England. All year 5 children (age 9-10 years) in the schools were informed about the project and received an information pack to share with their parents/carers. Written informed consent and assent were required from parents/carers and children respectively, before children could participate in the project in accordance with the project approvals granted by the University Research Ethics Committee (#SPA-REC-2015-182). Children were included if they provided the required informed parental

consent, assent, and medical screening forms, which indicated an absence of any medical conditions or disabilities preventing participation in the data collection and/or regular physical education lessons. The analytical sample consisted of 383 children (44.4% girls) at baseline and 272 children (43.4% girls) at 12-week follow-up. The participants' drop-out at follow-up was primarily due to absence from school on data collection days. This study used participants' valid data for CRF at baseline and HRQoL at baseline and at 12-weeks follow-up.

2.3 Active West Lancs Programme

The Active West Lancs programme consisted of classroom-based healthy lifestyle education lessons based on the 'Dr Feelwell' concept developed by MerseyCare National Health Service Foundation Trust (<https://www.merseycare.nhs.uk/>), and structured 'Born to Move' physical activity lessons (<https://www.lesmills.com/borntomove/>). Both were taught once per week for 45-60 minutes by physical activity specialists from an organisation which delivers physical education, physical activity, health, and wellbeing sessions in West Lancashire primary schools. The lessons complemented the regular curriculum and did not replace mandatory subjects that cover physical activity, health, and wellbeing concepts (e.g., physical education). The data reported in the present study are from the 20 schools involved in the four phases of the programme (January-April, April-July, September-December 2018, and January-April 2019) (supplementary figure 1).

2.4 Measures

2.2.4.1 Cardiorespiratory fitness

The 20-m multistage shuttle run test (20mSRT)²⁴ was conducted to provide an estimate of CRF. This test has been used extensively with participants of a similar age to those in the current study.²⁵ Prior research showed its validity (corrected mean r at the population level [95% CI]: $r_p = 0.78$ [0.72-0.85]) and reliability (intra-class correlation coefficients ranging from 0.78 to 0.93) in children.²⁶ Participants were encouraged to run for as long as possible until exhaustion or until they had reached their maximal effort. Otherwise, the test ended if the participant failed to reach within 2m of the marked line on two consecutive occasions. The 20mSRT was administered by the research team on a flat, clean surface indoors (e.g., sports/assembly hall) or outdoors (e.g., school playground) depending on available facilities and was completed in groups of up to 10 children. The total number of completed laps (shuttles) was used as a proxy indicator of CRF.

2.2.4.2 Health-related quality of life

The KIDSCREEN-10 Index questionnaire was used as a measure of global HRQoL³. KIDSCREEN-10 is a 10-item questionnaire, which asks participants how they felt in the last week. Items reflect the factors of physical well-being, psychological well-being, autonomy,

parent relations, peers and social support, and school environment, which are derived from the 27-item version of KIDSCREEN and are presented using a 1-5 Likert scale (i.e., 1 = “nothing” and 5 = “very much”).¹ Cronbach’s alphas are 0.82 and test–retest reliability was also generally satisfactory with internal consistent coefficients (ICCs) ranging from 0.61 to 0.70.²⁷ The Cronbach’s alpha for internal consistency of this questionnaire was 0.73 and 0.71 for the cross-sectional and longitudinal samples, respectively. Raw scores were converted to T-scores using the methodology described in the KIDSCREEN administration manual.³ The questionnaire was completed in classrooms following instructions from the research team and in the presence of the class teachers.

2.2.4.3 Socioeconomic status

Neighbourhood-level socioeconomic status (SES) was calculated for each child using the 2019 Indices of Multiple Deprivation (IMD).²⁸ The IMD is a UK government-produced deprivation measure for England comprising income, employment, health, education, housing, environment, and crime.²⁸ IMD rank scores were generated from parent-reported home postcodes using the National Statistics Postcode Directory database. Every neighbourhood in England is ranked from one (most deprived area) to 32,844 (least deprived area).²⁸

2.2.4.4. Anthropometric variables

Height was measured using a portable stadiometer (Leicester Height Measure, Seca, Birmingham, UK), and body mass was measured using calibrated scales (813 model, Seca). Body mass index (BMI) was calculated for each participant, BMI z-scores were assigned,²⁹ and International Obesity Task Force BMI cut-points applied to classify the participants as underweight, normal weight or overweight/obese.³⁰ Waist circumference was measured, using an anthropometric tape measure from the minimal waist site to the nearest millimetre, with participants in the standing position and at the end of expiration. Waist-to-height ratio (WHtR) was calculated as a measure of central obesity.³¹ Age at peak height velocity (APHV) was used as a proxy somatic measure of biological maturation. This method is based on anthropometric variables to predict APHV, which is a commonly used indicator of biological maturity.³² The method employs validated sex-specific regression equations which include participants’ chronological age and height.³² All the measurements were undertaken by trained researchers. To ensure accurate and standardised measurements all researchers firstly completed a six-hour training and supervised practice session using the assessment protocols. In addition, in order to avoid interindividual variability each researcher was responsible for administering the same measures during baseline and follow up assessment periods.

2.2.4.5. Moderate-to-vigorous physical activity

Self-reported moderate-to-vigorous physical activity (MVPA) data were collected using the Youth Activity Profile (YAP) English version.³³ The YAP is a 15-item questionnaire comprised of three sections (school-day MVPA, out-of-school MVPA, and sedentary behaviour), with five questions per section. Participants are asked to recall their MVPA and sedentary behaviour over the past 7 days during context-specific time segments (e.g., active travel to and from school, break time, etc.). The out-of-school segment refers to activity levels before school, immediately after school, evening, and at weekends. All questions were structured using a 5-point Likert scale (e.g., for active travel to school, a score of 1 indicated 0 days per week of active travel, whereas a score of 5 indicated 4–5 days per week). For this study, only data from the school-day and out-of-school MVPA questions were used. For each child, mean values for school-day and out-of-school MVPA were calculated and averaged resulting in a score for overall MVPA (1=low, 5=high). The YAP was completed in classrooms following instructions from the research team and in the presence of the class teachers.

2.2.4.6. Psychological correlates of physical activity: self-efficacy and enjoyment

Self-efficacy was measured using a valid and reliable questionnaire which contained 8 items related to the child's ability to be physically active.³⁴ The items were rated on a 5-point Likert scale ranging from 1 (very easy / disagree a lot) to 5 (very difficult / agree a lot). The Cronbach's alpha for the internal consistency of the cross-sectional sample was 0.77 and 0.78 for the longitudinal sample. Enjoyment was assessed through the Physical Activity Enjoyment Scale (PACES) for children.³⁵ A 5-point Likert-type scale (1 = "disagree a lot" to 5 = "agree a lot") is used to answer 16 statements. The average of the answers assigned to the 16 items is the final score. The Cronbach's alpha for the internal consistency of the sample was 0.87 and 0.88 for the cross-sectional and longitudinal samples, respectively.

2.5 Statistical analyses

Preliminary analyses involved checking all variables for normality using normal probability plots and Kolmogorov–Smirnov tests. The data assumed a normal distribution and descriptive statistics were calculated for all continuous measures using means (SD) and percentages for categorical variables. As exploratory analyses did not show a significant interaction of sex and CRF in relation to HRQoL ($p > 0.05$), the main analyses were performed with the total mixed-sex sample.

For study aim (i), mixed linear models examined the cross-sectional association between CRF and HRQoL with adjustment for sex, time of year, SES, WHtR, APHV, and MVPA; and the longitudinal association between CRF at baseline and HRQoL 12-weeks later adjusted for sex, time of year, SES, WHtR, APHV, MVPA, and HRQoL at baseline. Schools were included as random intercepts for aim (i) analysis. For study aim (ii), mediation analyses were conducted to assess the mediating role of each psychological correlate of physical activity (i.e., physical

activity self-efficacy and physical activity enjoyment) on the association between CRF and HRQoL with adjustment for sex, time of year, SES, WHtR, APHV, MVPA and schools. Cross-sectional mediation analyses were performed with CRF as the independent variable, HRQoL as the dependent variable and physical activity self-efficacy, and physical activity enjoyment individually introduced as mediator variables, with adjustment for the covariates. Further, longitudinal mediation analyses were performed with CRF at baseline as the independent variable, HRQoL at 12-week follow-up as the dependent variable and each psychological correlate individually introduced as mediator variables, with adjustment for the same covariates, with the addition of HRQoL at baseline. Effect sizes (Cohen's d) were calculated for both cross-sectional and longitudinal linear mixed models as suggested by Brysbaert and Stevens³⁶ and defined as: small (<0.2), medium ($0.2-0.5$), and large ($0.5-0.8$). For the mediation analyses effect sizes, R^2 was used to calculate f^2 ranges, carried out as proposed by Cohen³⁷ and defined as small (<0.02), medium ($0.02-0.15$), and large ($0.15-0.35$). The PROCESS SPSS Macro version 2.16.3, model 4, with 5000 bias-corrected boot-strap samples and 95% confidence intervals (CIs) was used for these analyses³⁸. Mediation was assessed by the indirect effect of CRF (independent variable) on HRQoL (dependent variable) through (i) self-efficacy, and (ii) enjoyment (mediators). Indirect effects ($a*b$ paths) with confidence intervals not including zero were considered significant. Mediation percentage (P_M) indicates how much of the association between CRF and HRQoL was explained by the mediator variables.³⁸ We performed post-hoc power statistical analyses to examine the impact of the changes from 383 to 272 on the results presented. Statistical significance was set at $p<.05$ for all analyses which were performed using IBM SPSS Statistics version 23 (IBM, Armonk, NY).

3. Results

Participants' baseline characteristics are presented in Table 1. The percentage of children at baseline and 12-week follow-up in the normal weight group was 78.1% and 82.7%, respectively. The drop-out from overweight and obese participants from baseline to follow up was 21.9% and 17.3%, respectively. On average, for the 20mSRT test, children performed 32.5 shuttles at baseline and 36.3 at 12-week follow-up. Mean HRQoL scores were 50.4 at baseline and 50.3 at 12-week follow-up. The psychological correlates of physical activity showed the same values at baseline and at 12-week follow-up.

Table 1. Characteristics of the participants at baseline and at 12-weeks follow-up. 263

	Baseline (n = 383)	Follow-up (n = 264)
Variable	Mean (SD) or frequency (%)	
Sex		265
Boys	213 (55.6%)	154 (56.6%) 266
Girls	170 (44.4%)	118 (43.4%) 267
Age (y)	10.0 (0.5)	10.2 (0.4) 268
SES (IMD rank)	15902.2 (10201.5)	16513.47 (9911.7) 269
WHtR	0.5 (0.1)	0.5 (0.1) 270
APHV (years)	-2.5 (0.7)	-2.3 (0.7) 271
Height (cm)	139.8 (6.4)	140.8 (6.4) 272
Mass (kg)	35.4 (7.8)	35.9 (7.6) 273
BMI (kg·m ⁻²)	18.0 (3.0)	18.0 (2.9) 274
Weight status		275
Under Weight	24 (6.2%)	17 (6.3%) 276
Normal Weight	275 (71.8%)	203 (74.6%) 277
Overweight/Obese	84 (21.9%)	52 (19.1%) 278
Waist circumference (cm)	65.3 (8.0)	65.7 (8.8) 279
YAP MVPA score	3.4 (0.8)	3.7 (0.8) 280
CRF (shuttles)	32.5 (16.0)	36.3 (17.1) 281
HRQoL	50.4 (9.7)	50.3 (10.0) 282
Physical activity self-efficacy	3.6 (0.7)	3.6 (0.8) 283
Physical activity enjoyment	4.3 (0.6)	4.3 (0.7) 284
Data are presented as mean (\pm SD) or frequencies (percentages). Differences between baseline and follow-up were examined by paired <i>t</i> -test ($p < 0.05$). SD: standard deviation; %: percentage; SES: socioeconomic status; IMD: indices of multiple deprivation; WHtR: waist to height ratio; APHV: peak height velocity; BMI: body mass index; YAP: youth activity profile; MVPA: moderate to vigorous physical activity; CRF: cardiorespiratory fitness; HRQoL: health-related quality of life. 285		

284 The results of the linear mixed model showing the cross-sectional associations between CRF and
285 HRQoL are presented in Table 2. A positive association was observed between CRF and HRQoL
286 ($p=0.015$) after adjusting for sex, time of year, SES, WHtR, APHV, and MVPA. A medium effect
287 size ($d = 0.26$) was found for this model. The unadjusted cross-sectional linear mixed model
288 showing the association between CRF and HRQoL is presented in supplementary table 1. Table
289 3 presents the linear mixed model outcome analysing the longitudinal associations between CRF
290 and HRQoL. The analysis revealed a positive association between CRF at baseline and HRQoL
291 at 12-week follow-up ($p=0.045$) after adjusting for covariates including HRQoL at baseline. A
292 medium effect size ($d = 0.35$) was found for this model. The unadjusted longitudinal linear mixed

model showing the association between CRF at baseline and HRQoL at 12-week follow-up is presented in supplementary table 2.

Table 2. Cross-sectional associations between cardiorespiratory fitness and HRQoL (n= 383).

	Model 1		
	β	95% CI	<i>p</i>
Intercept	57.67	44.43 – 70.90	<0.001
Sex	2.80	-0.54 – 6.14	0.100
Project phase	-0.71	-1.75 – 0.32	0.158
SES	5.31	-5.41 – 0.00	0.324
WHtR	-18.24	-39.04 – 2.56	0.086
APHV	-0.09	-2.45 – 2.27	0.941
YAP MVPA	0.16	-1.09 – 1.40	0.806
Cardiorespiratory fitness	0.09	0.02 – 0.16	0.015

Model 1: adjusted for the fixed effects of sex, time of the year, socioeconomic status, waist to height ratio, peak height velocity and moderate to vigorous physical activity. Clustering for analysis was schools. Health related quality of life was measured using KIDSCREEN-10. Data are presented as standardized regression coefficient (β) and 95% confidence interval (CI). Statistically significant values are in bold. HRQoL: health-related quality of life; SES: socioeconomic status; WHtR: waist to height ratio; APHV: peak height velocity; YAP: youth activity profile; MVPA: moderate to vigorous physical activity.

Table 3. Longitudinal associations between cardiorespiratory fitness at baseline and HRQoL at 12-weeks follow-up (n= 272).

	Model 2		
	β	95% CI	<i>p</i> -value
Intercept	21.46	6.7 – 36.22	0.005
Baseline HRQoL	0.52	0.42 – 0.63	<0.001
Sex	3.41	0.04 – 6.78	0.47
SES	1.53	-8.52 – 0.00	0.764
Project phase	-0.63	-1.91 – 0.65	0.337
WHtR	0.50	-21.83 – 22.82	0.965
APHV	-0.58	-2.82 – 1.67	0.615
YAP MVPA	-0.05	-1.28 – 1.18	0.940
Cardiorespiratory fitness	0.08	0.02 – 0.15	0.045

Model 2: adjusted for the fixed effects of sex, time of the year, socioeconomic status, waist to height ratio, peak height velocity, moderate to vigorous physical activity and baseline HRQoL. Clustering for analysis was schools. Health related quality of life was measured using KIDSCREEN-10.

Data are presented as standardized regression coefficient (β) and 95% confidence interval (CI). Statistically significant values are in bold. HRQoL: health-related quality of life; SES:

socioeconomic status; WHtR: waist to height ratio; APHV: peak height velocity; YAP: youth activity profile; MVPA: moderate to vigorous physical activity.

Figure 1 shows the adjusted cross-sectional mediating effect of self-efficacy and enjoyment in the association between CRF and HRQoL. There was a significant indirect effect (path $a*b$) between CRF and HRQoL when each psychological correlate of physical activity was individually included in the analyses. CRF was positively associated with both single psychological correlates (a path; all $p<0.001$), which were also positively associated with HRQoL (b path; all $p<0.001$). However, in each model the direct effect between CRF and HRQoL was not significant (c' path; all $p>0.05$). The outcome of these cross-sectional mediation analyses suggested that CRF could indirectly influence HRQoL through its effects on children's physical activity self-efficacy ($P_M=82.7\%$) and enjoyment ($P_M=54.1\%$). Mediation analyses effect sizes were medium to large, with R^2 ranging from 0.07 to 0.19 for enjoyment and from 0.09 to 0.22 for self-efficacy. For the cross-sectional mediation models the post-hoc power of the regressions included ranged from 99.7% to 100%.

The results of the adjusted longitudinal mediating effects of both physical activity self-efficacy and enjoyment on the association between CRF at baseline and HRQoL at 12-week follow-up are shown in Figure 2. There was a significant indirect effect of baseline self-efficacy in the longitudinal association between baseline CRF and HRQoL at 12-week follow-up (path $a*b$), but not for enjoyment. Moreover, baseline CRF was significantly associated with baseline self-efficacy (a path; $p<0.01$), whereas a non-significant association was found with baseline enjoyment (a path; $p>0.05$). Baseline self-efficacy and enjoyment were positively associated with HRQoL at 12-week follow-up (b path; all $p<0.01$). Finally, the direct effect between baseline CRF and HRQoL at 12-week follow-up was non-significant (c' path; all $p>0.05$). The results of the longitudinal mediation analyses suggested that baseline CRF could indirectly influence HRQoL 12 weeks later through its effects on children's baseline physical activity self-efficacy ($P_M=29.4\%$) and enjoyment ($P_M=15.8\%$), separately. R^2 ranged from 0.10 to 0.20 for enjoyment, and from 0.09 to 0.25 for self-efficacy (i.e., medium to large effects). For the longitudinal mediation models the post-hoc power of the regressions included was 100%.




Figure 1. Psychological correlates of physical activity (i.e., self-efficacy and enjoyment) mediation models of the cross-sectional relationship between CRF and HRQoL, adjusted for sex, time of the year, socioeconomic status, waist to height ratio, peak height velocity, moderate to vigorous physical activity and schools ($n = 383$). Results are showed as unstandardized regression coefficients; p -value. IE = indirect effect [lower and upper levels for 95% confidence interval of the indirect effect between CRF and HRQoL]. P_M : percentage of mediation; CRF: cardiorespiratory fitness; HRQoL: health-related quality of life.




Figure 2. Psychological correlates of physical activity (i.e., self-efficacy and enjoyment) of the longitudinal relationship between CRF at baseline and HRQoL at 12-week follow-up, adjusted for sex, time of the year, socioeconomic status, waist to height ratio, peak height velocity, moderate to vigorous physical activity, schools, and HRQoL at baseline (n = 272). Results are showed as unstandardized regression coefficients; *p*-value. IE = indirect effect [lower and upper levels for 95% confidence interval of the indirect effect between CRF at baseline and HRQoL at 12-week follow-up]. P_M: percentage of mediation; CRF: cardiorespiratory fitness; HRQoL: health-related quality of life.

4. Discussion

The results of our cross-sectional and longitudinal analyses showed that among northwest England primary school children (i) CRF was positively associated with HRQoL and (ii) this association was mediated by self-efficacy and enjoyment as psychological correlates of physical activity. Our findings expand prior knowledge about the association between CRF and HRQoL in children and reveal for the first time potential underlying mechanisms involved in the association between CRF and HRQoL, highlighting the significant roles of single physical activity correlates such as self-efficacy and enjoyment.

Our cross-sectional results showed a positive association between CRF and HRQoL. Similar findings were found in previous studies,^{11–13,39} which reported that children with higher levels of CRF had better HRQoL. For instance, a study in Norwegian 10-year-olds revealed that CRF had a small to medium effect size (R^2 ranging from 0.17 to 0.5) in its positive association with all HRQoL domains (i.e., physical and psychological well-being, autonomy and parents, social support and school environment).¹¹ Also, Redondo et al. reported a small (all $R^2 < 0.5$) positive association between CRF and HRQoL among children aged 4 to 7 years.¹² Regarding longitudinal associations between CRF and HRQoL, our novel findings revealed a positive association between CRF at baseline and children's HRQoL at 12-week follow-up after adjusting for confounders. This outcome is partially supported by previous longitudinal research in different age groups by confirming the individual positive small association of CRF at baseline on HRQoL over a 2-year period follow-up ($R^2 < 0.5$).⁴⁰ These findings might be somewhat explained by the positive influence that CRF has on both physical and mental dimensions of health in children² over time,^{41,42} which may positively impact children's HRQoL. We hypothesise that the similarity between our study's effect sizes and the ones of previous evidence might be due to the several dimensions of HRQoL which could not be fully influenced by CRF.

Since mediation analysis assumes that the independent variable influences the mediator, our cross-sectional and longitudinal results suggest that CRF at baseline influenced the psychological

variables, which, in turn, may affect HRQoL at baseline and 12-weeks later. With respect to path *a*, our findings could be partially supported by a previous cross-sectional study which reported that children with higher CRF levels had higher physical self-efficacy and physical activity enjoyment than their peers with low CRF.¹⁸ Regarding our longitudinal results, we were not able to make comparisons since no evidence relating CRF and physical activity self-efficacy over time was found. We hypothesise that it is plausible that CRF influenced physical activity self-efficacy and enjoyment through motor skill development/proficiency and sport experiences. Children's CRF levels are associated with increased motor competence,⁴³ positive sport and physical activity experiences,^{44,45} which in turn may affect several domains of their HRQoL. However, there is paucity of evidence in this area and further research is warranted. With respect to path *b*, our findings are in line with previous cross-sectional²¹ and interventional²⁰ studies which reported a positive association of children's physical activity enjoyment and their HRQoL. However, no previous studies have examined the cross-sectional and longitudinal associations between children's physical activity self-efficacy and their HRQoL. The association found in our study between both psychological correlates and HRQoL may be related to the mental domain of the construct, predisposing children to higher scores of psychological well-being.^{21,46}

The results obtained in the present study through mediation analyses, a powerful statistical technique that can be used to clarify the process underlying the relationship between two variables,³⁸ add support for the psychological correlates of physical activity being an intermediate step on the causal pathway between CRF and children's HRQoL. Thus, our findings are consistent with the idea that the promotion of children's physical activity self-efficacy and physical activity enjoyment may be of importance to improve their HRQoL.

Our mediation results are partially supported by only one previous cross-sectional study. This involved overweight adolescents, and showed the mediating role that motivational variables (i.e., self-determined motivation) have in the association between CRF and HRQoL.⁴⁷ However, the mediating roles of physical activity self-efficacy and enjoyment in children have not been previously investigated. Yet, based on prior cross-sectional research in other populations framed by self-determined motivation, being more physically fit leads to the need for more autonomy and competence during physical activity practice and, therefore, the development of more autonomous forms of self-regulations, which might benefit persistence and mental well-being^{47,48} with a positive impact on their HRQoL.

Given the need of further research on correlates of physical activity due to its influence on behavioural change¹⁷ and the temporal trends in physical fitness reporting a global declining tendency over the years,⁴⁹ our data may have significant implications for HRQoL improvement. Indeed, maintaining children's HRQoL is important for current health, as well as, has transferable value for future societal health. Our findings are of interest to educators and policy makers, to

raise the importance of CRF for improving children's psychological correlates of physical activity and their HRQoL.

Strengths of this study include the homogeneous age-matched and relatively large sample of children. The multilevel analyses accounted for school-level variance and adjusted for important fixed effects confounders. Furthermore, the mediation models added significant novelty to provide improved insights into the CRF-HRQoL relationships. There are also limitations which warrant consideration. The findings obtained from the cross-sectional elements of the study preclude claims of causal inferences and directionality between CRF and HRQoL, whereas there is more confidence about causality in those from the longitudinal aspects which controlled for baseline HRQoL and confounders. The 12-weeks duration of the follow-up is short which limits the significance of the longitudinal results. Moreover, the sample was drawn from one geographical region of northwest England, therefore the results may not be generalizable to populations elsewhere. MVPA was assessed using a self-report instrument which is open to recall and social desirability biases; however, the YAP is a validated method that was administered in the same way at both time points, thus limiting variation in responses between baseline and follow-up. Lastly, we acknowledge that more accurate estimates of CRF could have been obtained using a laboratory-based physiological direct measure. However, such measures were not feasible within our study, and the 20mSRT is the most widely used field-based test of CRF in children, which demonstrates criterion validity against gas-analyzed peakVO₂, and has strong ecological validity and feasibility in school settings.

5. Conclusion

The results of the current study showed that CRF was cross-sectionally and longitudinally associated with HRQoL in primary school children in England. Furthermore, self-efficacy and enjoyment as psychological correlates of physical activity act separately as mediators in the positive association between CRF and HRQoL. Therefore, we contribute to the comprehension of the relationship between these key factors, suggesting that both optimal CRF levels and better psychological correlates of physical activity are important for children's HRQoL. Our findings should be considered when designing education and public health interventions and strategies aiming to improve HRQoL during childhood.

References

1. Ravens-Sieberer U, Gosch A, Rajmil L, et al. KIDSCREEN-52 quality-of-life measure for children and adolescents. *Expert Rev Pharmacoeconomics Outcomes Res* 2005; 5: 353–364.
2. Ortega FB, Ruiz JR, Castillo MJ, et al. Physical fitness in childhood and adolescence: A powerful marker of health. *Int J Obes* 2008; 32: 1–11.
3. Ravens-Sieberer U, Gosh A, Erhart M, et al. *The KIDSCREEN questionnaires: Quality of life questionnaires for children and adolescents; Handbook*. 2006.
4. Prevention C for DC and. *Measuring Healthy Days: Population Assessment of Health-Related Quality of Life*. Atlanta, Georgia, 2000.
5. Lin X-J, Lin I-M, Fan S-Y. Methodological issues in measuring health-related quality of life. *Tzu Chi Med J* 2013; 25: 8–12.
6. Meyer M, Oberhoffer R, Hock J, et al. Health-related quality of life in children and adolescents: Current normative data, determinants and reliability on proxy-report. *J Paediatr Child Health* 2016; 52: 628–631.
7. Solans M, Pane S, Estrada MD, et al. Health-Related Quality of Life Measurement in Children and Adolescents: A Systematic Review of Generic and Disease-Specific Instruments. *Value Heal* 2008; 11: 742–764.
8. Ravens-Sieberer U, Erhart M, Wille N, et al. Health-related quality of life in children and adolescents in Germany: results of the BELLA study. *Eur Child Adolesc Psychiatry* 2008; 17 Suppl 1: 148–156.
9. Armstrong N, Tomkinson GR, Ekelund U. Aerobic fitness and its relationship to sport, exercise training and habitual physical activity during Youth. *Br J Sports Med* 2011; 45: 849–858.
10. Ruiz JR, Castro-Piñero J, Artero EG, et al. Predictive validity of health-related fitness in youth: A systematic review. *Br J Sports Med* 2009; 43: 909–923.
11. Andersen JR, Natvig GK, Aadland E, et al. Associations between health-related quality of life, cardiorespiratory fitness, muscle strength, physical activity and waist circumference in 10-year-old children: the ASK study. *Qual Life Res* 2017; 26: 3421–3428.
12. Saucedo-Araujo RG, Huertas-Delgado FJ, Villa-González E, Ávila-García M, Gálvez-

- 487 Fernández P TP. Is children's health-related quality of life associated with physical fitness
488 and mode of commuting? PREVIENE Project. *Perspect Public Health* 2021; 141: 102–
489 110.
- 490 13. Redondo-Tébar A, Ruíz-Hermosa A, Martínez-Vizcaíno V, et al. Associations between
491 health-related quality of life and physical fitness in 4–7-year-old Spanish children: the
492 MOVIKIDS study. *Qual Life Res* 2019; 28: 1751–1759.
- 493 14. Morrow V, Crivello G. International Journal of Social Research Methodology What is the
494 value of qualitative longitudinal research with children and young people for international
495 development? *Int J Soc Res Methodol* 2016; 18: 267–280.
- 496 15. Gálvez Casas A, Rodríguez García PL, García-Cantó E, et al. Capacidad aeróbica y
497 calidad de vida en escolares de 8 a 12 años. *Clínica e Investig en Arterioscler* 2015; 27:
498 239–245.
- 499 16. Raghuveer G, Hartz J, Lubans DR, et al. Cardiorespiratory Fitness in Youth: An Important
500 Marker of Health: A Scientific Statement from the American Heart Association.
501 *Circulation* 2020; 142: E101–E118.
- 502 17. Bauman AE, Reis RS, Sallis JF, et al. Correlates of physical activity: why are some people
503 physically active and others not? *Lancet* 2012; 380: 258–271.
- 504 18. Vitali F, Robazza C, Bortoli L, et al. Enhancing fitness, enjoyment, and physical self-
505 efficacy in primary school children: A DEDIPAC naturalistic study. *PeerJ* 2019; 7: e6436.
- 506 19. Fu Y, Gao Z, Hannon JC, et al. Effect of the SPARK program on physical activity,
507 cardiorespiratory endurance, and motivation in middle-school students. *J Phys Act Heal*
508 2016; 13: 534–542.
- 509 20. Hyndman BP, Benson AC, Ullah S, et al. Evaluating the effects of the Lunchtime
510 Enjoyment Activity and Play (LEAP) school playground intervention on children's quality
511 of life, enjoyment and participation in physical activity. *BMC Public Health* 2014; 14:
512 164.
- 513 21. Hyndman B, Benson AC, Lester L, et al. Is there a relationship between primary school
514 children's enjoyment of recess physical activities and health-related quality of life? A
515 cross-sectional exploratory study. *Heal Promot J Aust* 2017; 28: 37–43.
- 516 22. Rhodes RE, Janssen I, Bredin SSD, et al. Physical activity: Health impact, prevalence,
517 correlates and interventions. *Psychol Health* 2017; 32: 942–975.
- 518 23. Government UK. Childhood obesity: a plan for action.

- 519 24. Léger LA, Mercier D, Gadoury C, et al. The multistage 20 metre shuttle run test for aerobic
520 fitness. *J Sports Sci* 1988; 6: 93–101.
- 521 25. Tomkinson GR, Carver KD, Atkinson F, et al. European normative values for physical
522 fitness in children and adolescents aged 9-17 years: Results from 2 779 165 Eurofit
523 performances representing 30 countries. *Br J Sports Med* 2018; 52: 1445–1456.
- 524 26. Tomkinson GR, Lang JJ, Blanchard J, et al. The 20-m Shuttle Run: Assessment and
525 Interpretation of Data in Relation to Youth Aerobic Fitness and Health. *Pediatr Exerc Sci*
526 2019; 31: 152–163.
- 527 27. Ravens-Sieberer U, Herdman M, Devine J, et al. The European KIDSCREEN approach to
528 measure quality of life and well-being in children: Development, current application, and
529 future advances. *Qual Life Res* 2014; 23: 791–803.
- 530 28. Department D for C and LD. English indices of deprivation - Statistical Release. 2015.
531 DOI: London: DCLG.
- 532 29. Cole TJ, Freeman J V., Preece MA. Body mass index reference curves for the UK, 1990.
533 *Arch Dis Child* 1995; 73: 25–29.
- 534 30. Cole TJ, Bellizzi MC, Flegal KM, et al. Establishing a standard definition for child
535 overweight and obesity worldwide: International survey. *Br Med J* 2000; 320: 1240–1243.
- 536 31. Mokha JS, Srinivasan SR, DasMahapatra P, et al. Utility of waist-to-height ratio in
537 assessing the status of central obesity and related cardiometabolic risk profile among
538 normal weight and overweight/obese children: The Bogalusa Heart Study. *BMC Pediatr*
539 2010; 10: 73.
- 540 32. Moore SA, McKay HA, Macdonald H, et al. Enhancing a Somatic Maturity Prediction
541 Model. *Med Sci Sports Exerc* 2015; 47: 1755–1764.
- 542 33. Fairclough SJ, Christian DL, Saint-Maurice PF, et al. Calibration and validation of the
543 youth activity profile as a physical activity and sedentary behaviour surveillance tool for
544 english youth. *Int J Environ Res Public Health* 2019; 16: 3711.
- 545 34. Motl RW, Dishman RK, Trost SG, et al. Factorial validity and Invariance of
546 Questionnaires Measuring Social-Cognitive Determinants of Physical Activity among
547 Adolescent Girls. *Prev Med (Baltim)* 2000; 31: 584–594.
- 548 35. Moore JB, Yin Z, Hanes J, et al. Measuring Enjoyment of Physical Activity in Children:
549 Validation of the Physical Activity Enjoyment Scale. *J Appl Sport Psychol* 2009; 21: 116–
550 129.

36. Brysbaert M, Stevens M. Power Analysis and Effect Size in Mixed Effects Models: A Tutorial. *J Cogn* 2018; 1: 9.
37. Cohen J. A power primer. *Psychol Bull* 1992; 112: 155–159.
38. AF H. *Introduction to Mediation, Moderation, and Conditional Process Analysis: A Regression-Based Approach*. New York, NY: Guilford Press, 2013.
39. Morales PF, Sánchez-López M, Moya-Martínez P, et al. Health-related quality of life, obesity, and fitness in schoolchildren: The Cuenca study. *Qual Life Res* 2013; 22: 1515–1523.
40. Evaristo OS, Moreira C, Lopes L, et al. Cardiorespiratory fitness and health-related quality of life in adolescents: A longitudinal analysis from the LabMed Physical Activity Study. *Am J Hum Biol* 2019; 31: e23304.
41. Castro-Piñero J, Perez-Bey A, Segura-Jiménez V, et al. Cardiorespiratory Fitness Cutoff Points for Early Detection of Present and Future Cardiovascular Risk in Children: A 2-Year Follow-up Study. *Mayo Clin Proc* 2017; 92: 1753–1762.
42. Kandola AA, Osborn DPJ, Stubbs B, et al. Individual and combined associations between cardiorespiratory fitness and grip strength with common mental disorders: a prospective cohort study in the UK Biobank. *BMC Med* 2020; 18: 1–11.
43. Lang JJ, Chaput JP, Longmuir PE, et al. Cardiorespiratory fitness is associated with physical literacy in a large sample of Canadian children aged 8 to 12 years 11 Medical and Health Sciences 1117 Public Health and Health Services. *BMC Public Health* 2018; 18: 1–13.
44. Eberline A, Judge LW, Walsh A, et al. Relationship of Enjoyment, Perceived Competence, and Cardiorespiratory Fitness to Physical Activity Levels of Elementary School Children. *Phys Educ* 2018; 75: 394–413.
45. Welk GJ. The youth physical activity promotion model: A conceptual bridge between theory and practice. *Quest* 1999; 51: 5–23.
46. Bandura A. Health promotion from the perspective of social cognitive theory. *Psychol Health* 2007; 13: 623–649.
47. Riiser K, Ommundsen Y, Småstuen MC, et al. The relationship between fitness and health-related quality of life and the mediating role of self-determined motivation in overweight adolescents. *Scand J Public Health* 2014; 42: 766–772.
48. Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic motivation,

social development, and well-being. *Am Psychol* 2000; 55: 68–78.

49. Tomkinson GR, Kaster T, Dooley FL, et al. Temporal Trends in the Standing Broad Jump Performance of 10,940,801 Children and Adolescents Between 1960 and 2017. *Sport Med* 2021; 51: 531–548.

Supplementary material

Supplementary table 1. Cross-sectional associations between cardiorespiratory fitness and HRQoL (n= 383).


Model 1			
	β	95% CI	<i>p</i>
Intercept	57.30	44.94 – 49.66	<0.001
Cardiorespiratory fitness	0.09	0.04 – 0.16	0.002

Model 1: Unadjusted. Health related quality of life was measured using Kidscreen-10. Data are presented as standardized regression coefficient (β) and 95% confidence interval (CI). Statistically significant values are in bold.

Supplementary table 2. Cross-sectional associations between cardiorespiratory fitness and HRQoL (n= 383).

Model 1			
	β	95% CI	<i>p</i>
Intercept	47.75	44.06 – 49.43	<0.001
Cardiorespiratory fitness	0.11	0.03 – 0.18	0.004

Model 1: Unadjusted. Health related quality of life was measured using Kidscreen-10. Data are presented as standardized regression coefficient (β) and 95% confidence interval (CI). Statistically significant values are in bold.



Supplementary figure 1. Active West Lincs Programme phases of data collection.