



## Article

# Understanding Post-Crisis Trauma Recovery for the Past Decade in Uniformed Services: A Narrative Review

de silva, Raneesha, Ireland, Jane Louise, Birch, Philip, Ireland, Carol Ann, Lewis, Michael, Dissanayaka, Ravindra and Atapattu, Methma

Available at <http://clock.uclan.ac.uk/44195/>

*de silva, Raneesha, Ireland, Jane Louise ORCID: 0000-0002-5117-5930, Birch, Philip, Ireland, Carol Ann ORCID: 0000-0001-7310-2903, Lewis, Michael ORCID: 0000-0001-5567-3569, Dissanayaka, Ravindra and Atapattu, Methma (2022) Understanding Post-Crisis Trauma Recovery for the Past Decade in Uniformed Services: A Narrative Review. Mental Health Review Journal .*

It is advisable to refer to the publisher's version if you intend to cite from the work.

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.



**Understanding Post-Crisis Trauma Recovery for the Past Decade in Uniformed Services: A Narrative Review**

Journal:	<i>Mental Health Review Journal</i>
Manuscript ID	MHRJ-11-2021-0081.R2
Manuscript Type:	Research Paper
Keywords:	trauma recovery, crisis respondents, post-disaster management, military, police

SCHOLARONE™  
Manuscripts

## Abstract

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 This narrative review explores mental health difficulties, including risk and protective  
3 factors, which may impact on symptom severity post exposure to crisis situations (war,  
4 terrorism, or natural disasters), among first responders from uniformed services. Peer-  
5 reviewed journal articles published in English between January 2012 to March 2022 were  
6 searched in ProQuest, Wiley, Google Scholar, and PubMed databases. Twelve articles were  
7 obtained from an original screening of 94,058 articles. Full article texts were screened for  
8 content and quality by two reviewers, with high agreeability. Post-traumatic Stress Disorder  
9 (PTSD) and depression were the most common diagnoses. Risk factors identified were (1)  
10 *Pre-deployment factors* of overweight, low cognitive ability and social support, existing  
11 emotional difficulties, negative childhood experiences, and stressful life events; (2) *During*  
12 *crisis situations factors* of higher frequency and subjective severity of combat, increased rates  
13 of combat stress reaction (CSR), high levels of concerns for life and family, more stressful  
14 mission position, threat of death/severe injury, and high rate of killing the enemy, and (3)  
15 *Post-deployment factors*, such as low social support and physical health, lack of coping  
16 mechanisms and use of avoidance strategies, and social stigma. Protective factors increasing  
17 resilience and lessening symptom severity were reported as pre-deployment cognitive ability,  
18 high social support, stable physical health, effective coping, post-traumatic growth, and high  
19 levels of perceived adequacy in pre-deployment preparation and training. In addition to main  
20 findings, data about author(s), publication type, population, age, type of crisis, and evaluation  
21 measures were extracted. Key findings and related theories, gaps in literature, and  
22 recommendations are discussed.

23 KEYWORDS: trauma recovery; crisis respondents; post-disaster management; military;  
24 police.  
25

## Understanding Post-Crisis Trauma Recovery for the Past Decade in Uniformed Services: A Narrative Review

Independent of the type of crisis (i.e., combat and/or natural disasters) and location (Western or Eastern countries), the uniformed services are regularly required to contribute their services as frontline crisis respondents. Constant exposure to stressful crisis situations may manifest as post-crisis psychological symptoms and difficulties such as Post-Traumatic Stress Disorder (PTSD), depression, and/or anxiety (Bowler et al., 2016; Ikin et al., 2020). Due to decreased income or unemployment, substance abuse and/or other personal factors (e.g., young age and ethnicity; Bowler et al., 2016; Elbogen et al., 2012; MacManus et al., 2019), these difficulties may be aggravated. In contrast, some factors, such as social support, stable living conditions, pre-deployment preparedness, and peer support (Elbogen et al., 2012; Kline et al., 2013) may reduce the intensity of crisis experiences being developed into severe mental health difficulties. However, at present, research lacks a collated overview of these findings, as aforementioned factors are reported from single and separate studies. In particular, there is an evident lack of research regarding protective factors of uniformed personnel, which may alleviate their mental health difficulties. Therefore, the present narrative review is a timely necessity.

While several theories could contribute to this research domain, *Pluralistic Trauma Theory* (PTT: Balaev, 2018), *Self-Determination Theory* (SDT: Ryan & Deci, 2000), *Effort-Reward Imbalance model* (ERI: Siegrist, 1996), and *Job Demand-Control-Support model* (JDACS: Johnson, and Hall, 1988) represent prominent theoretical frameworks to be considered. PTT (Balaev, 2018) suggests that trauma is not only about past experiences, but also about relations among experience, language, and knowledge. This approach conceptualises trauma as an event that changes perception and identity but awakes new

1 knowledge about one's self and the external world. This theory relies heavily on external  
2 stressors; trauma can happen in specific bodies, times, cultures, and places, resulting in a  
3 specific representation of trauma for each person. According to this theory, the effect of  
4 trauma is an interplay between external (e.g., cultural factors) and internal factors (e.g.,  
5 personality traits), which creates a link between singular vs. collective experiences of trauma.  
6 This subjective perception is important in determining, "what is viewed as salient, how it is  
7 interpreted and encoded at the time of the event, what is most important as autobiographical  
8 functions, and what is socially acceptable to speak/what must remain hidden and  
9 unacknowledged" (Balaev, 2018). In contrast to more traditional models of trauma (Freud,  
10 1895; Caruth, 1996), which attributes more to unobserved neurobiological functions, PTT  
11 suggests that what remains hidden and unacknowledged is due to the influence of cultural  
12 values (Balaev, 2018). An earlier complementary theory that is also worth noting is Self-  
13 Determination Theory (SDT; Ryan & Deci, 2000). Emphasis here is placed on the notion that  
14 human functionality depends on the social context; the interplay between intrinsic and  
15 extrinsic motivational factors determines an individual's response to different social  
16 situations and further influences social, cognitive and personality development. It is  
17 suggested here that exposure to a traumatic event could serve as a social event that impacts  
18 on individual functionality, impacting on development, although this has not been applied to  
19 trauma per se but rather to symptoms that could be secondary to this, such as depression and  
20 anxiety. However, what it does note is how optimal levels of development comprise of  
21 competence (confidence to perform), relatedness (need to perform), and autonomy  
22 (independence to perform), and that once these are met, functionality is raised (Legault,  
23 2017). How exposure to traumatic events impacts on these areas is not yet known.

24         Alongside these more general models of understanding human reactions are those  
25 specific to the work environment, since focus here is on trauma exposure as a result of

1 exposure through work. The Effort-Reward Imbalance model (ERI: Siegrist, 1996), for  
2 example, captured the negative effects of occupational stress and work conditions in  
3 developed/rapidly-developing countries, due to an imbalance between high cost spent and  
4 low gain received in return. This repeated imbalance could arguably frustrate related circuits  
5 of the brain's reward system and increase the risk for developing stress-related disorders,  
6 such as depression (Siegrist, 2016). This suggests that occupations where there is a high cost  
7 but a low gain could be particularly vulnerable to this. Arguably working in an occupation  
8 such as the uniformed services, where there is a potential high cost to employees, raises the  
9 risk for stress-related disorders, of which trauma could be a core example. Alongside this is  
10 the need to capture the Job Demand-Control-Support (JDCS) model (Johnson & Hall, 1988),  
11 which outlines how employees are primarily stressed when there is a high demand (due to  
12 work overload, time pressure, and role conflict), low control (lack of autonomy), and  
13 inadequate social support. This builds on what is referred to as the 'strain' hypothesis and has  
14 parallels to the ERI 'high risk' components. The competing 'buffer' hypothesis views 'social  
15 support' as a protective factor, which moderates the negative impact of high demand—low  
16 control (Van der Doef & Maes, 1999). This raises the importance of capturing social support  
17 as a variable of potential significant interest in understanding the impacts of exposure to  
18 trauma within occupational settings.

19 As yet, however, research into the factors that could serve as risk and/or protective  
20 factors are not clearly indicated in terms of post-crisis recovery. The *types* of impact are  
21 researched to a greater degree but not the factors that could serve as 'strain' and/or 'buffers'  
22 to the demands that crisis responders are exposed to. Clearly, available theory can suggest  
23 possible factors of importance, but the empirical basis remains less clear. In consideration of  
24 this, the narrative review presented here aims to identify and assess the existing research  
25 literature, evidenced in the last decade, to capture the following aims: 1.) To understand what

1 mental health difficulties may manifest in a crisis responder, from uniformed services, when  
2 exposed to crisis situations; 2.) To capture the risk factors aggravating mental health  
3 difficulties; and 3.) To identify what factors protect crisis respondents from mental health  
4 difficulties.

## 5 **Method**

6 This narrative review was conducted in accordance with recommended guidelines of  
7 the Synthesis Without Meta-analysis Reporting Items (SWiM; Campbell et al., 2020)  
8 checklist (e.g. Grouping studies, describing methods, identify key study characteristics).

### 9 **Data Sources and Search**

10 Databases such as ProQuest, Wiley, Google Scholar, and PubMed were searched for  
11 peer-reviewed full text journal articles published in English between January 2012 and March  
12 2022. This specific timeline is therefore able to provide an overview of findings for the  
13 period from the first emerging publications using the DSM-5 to the current date. The  
14 keywords used for the search strategy are: “Crisis respondents\*” OR “Post-disaster  
15 management\*” AND “Mental health difficulties\*” OR “Mental health symptoms\*” OR  
16 “\*Trauma” OR “Trauma recovery\*” AND “Risk factors\*” OR “Protective factors\*” AND  
17 “War\*” OR “Terrorism\*” OR “Natural disasters\*” OR “Emergency\*” OR “Rescue\*” OR  
18 “Violence\*” AND “Military\*” OR “Police\*”. Articles were excluded if they were duplicates,  
19 not relevant, not a primary research study, had weak methodology (i.e., high risk of sampling  
20 bias with convenience and snowball sampling, low sampling size, imprecise outcome  
21 measures, and limited generalisability due to over/under emphasis on multiple demographic  
22 factors), and if the publication language was not English (See Figure 1).

## **Data Extraction and Quality Appraisal**

Titles and abstracts of the articles were initially evaluated by Reviewer 1 (RS). Thereafter, articles eligible for full-text screening were assessed by two reviewers (RS and MA) who reached high agreement (78%) on the final decision. Each paper was independently assessed for its quality (i.e., the overall appraisal is presented as, “Include”, “Exclude” or “Seek further info”) using the Critical Appraisal Checklist for Systematic Reviews and Research Syntheses (Joanna Briggs Institute; 2017) by two reviewers (RS and MA), who reported high agreement (82%). Data were extracted under following domains: author(s), publication type, population, age, and type of crisis (Table 1), in addition to key findings, which were tabulated as mental health difficulties, and risk and protective factors (Table 2).

## **Results**

### **Literature Search**

The initial search resulted in 94,058 articles. After 564 duplicates were removed, 93,494 article titles and abstracts were screened as either “Relevant” or “Not relevant”. Thereafter, 93,309 articles were excluded in total because they were: “Not relevant” ( $n=91,344$ ), “Not a primary research study” ( $n=1,948$ ), and “Publication language is not English” ( $n=17$ ). This resulted in 185 articles to be screened for full text; out of which, 175 articles were excluded (“Not relevant”;  $n=136$ , “Not a primary research study”;  $n=28$ , and “Weak methodology”;  $n=11$ ). Additionally, 32 articles were screened as a result of hand searching full-text references. Only two articles were assessed [from the hand-searched lists] for eligibility, with 30 articles excluded because they were, “Not relevant” ( $n=21$ ), “Not a primary research study” ( $n=7$ ), or presented with “Weak methodology” ( $n=2$ ). This results in 12 articles meeting the inclusion criteria for the narrative review.



1  
2  
3 1  
4  
5  
6 2 <Insert Figure 1 here>  
7  
8 3  
910 4 **Study Characteristics**

11  
12 5 The majority of the active duty or retired crisis respondents ( $n=37,267$ ) were exposed  
13  
14 6 to combat during deployment to Iraq and/or Afghanistan. Other occupation roles included  
15  
16 7 first responders from the police, fire department, and coastguards. All studies were from  
17  
18 8 Western countries, such as the United States of America, Sweden, and Denmark. The mean  
19  
20 9 age ranged between 24 years and 64 years. Validated measures were utilised in each study to  
21  
22 10 assess mental health difficulties and risk/protective factors. Detailed information is collated  
23  
24 11 and presented as Table 1.  
25  
26  
27  
28  
29  
30  
31 12 <Insert Table 1 here>  
32  
33  
34

35 15 **Main Findings**

36  
37 16 This narrative review identified core themes and sub themes through the Thematic Narrative  
38  
39 17 Analysis, described below. Main findings are presented in Table 2.  
40  
41  
42

43 18 ***Pre-deployment Factors***

44  
45 19 Only one study (Sørensen et al., 2016) had assessed the presence of mental health  
46  
47 20 difficulties prior to deployment. When Danish soldiers ( $n=42$ ) were screened for PTSD, six  
48  
49 21 weeks prior deployment, an inverse relationship (0.97) was revealed between PTSD (PCL  $\geq$   
50  
51 22 44; moderate to severe) and cognitive abilities.  
52  
53  
54

55 23 The impact of pre-deployment cognitive abilities on the risk of developing PTSD  
56  
57 24 symptoms was analysed in this study (Sørensen et al., 2016). The resilient categories  
58  
59 25 exhibited low distress levels across six assessments before, during, and after deployment.  
60

1  
2  
3 1 However, they did not have higher mean pre-deployment cognitive ability scores compared  
4  
5 2 with the mild distress trajectory, which exhibited moderate symptoms that decreased after  
6  
7 3 deployment. PTSD symptoms measured as a PCL score revealed an inverse association with  
8  
9 4 pre-deployment cognitive ability (Sørensen et al., 2016). This association remained  
10  
11 5 significant when adjusted for perceived war zone stress, pre-deployment symptoms levels,  
12  
13 6 pre-deployment traumatic life events, and education. Further affirming these findings, an  
14  
15 7 association was revealed between lower pre-deployment cognitive ability scores and the  
16  
17 8 relieved-worsening trajectory (i.e., a subgroup with PTSD symptoms). This subgroup  
18  
19 9 reported the highest level of pre-deployment emotional problems and the lowest level of  
20  
21 10 perceived social support (Sørensen et al., 2016).

### 11 ***During-deployment Factors***

12 As the only study that had assessed the veterans during deployment, Sørensen et al.,  
13 (2016) reported an association between the higher mean score of pre-deployment cognitive  
14 ability and lower reports of PTSD symptoms — participants from the most resilient subgroup  
15 (i.e., exhibiting low levels of distress) stated the highest level of danger/injury score during  
16 the mission, a higher proportion of soldiers being wounded or injured during the mission, and  
17 a higher proportion of having killed an enemy.

### 18 ***Post-deployment Factors***

19 The following sub themes were identified for post-deployment factors: mental health  
20 difficulties, and risk and protective factors for difficulties.

#### 21 Mental Health Difficulties

22 The following mental health symptoms were noted in uniformed personnel exposed to  
23 crisis situations.

1  
2  
3           ***PTSD and Depression.*** According to Chapman et al. (2014), between deployed and  
4  
5           2 non-deployed combat medics, medics who have been deployed were found to seek out  
6  
7           3 assistance with their mental health than their peers. The rate of depression was high in  
8  
9           4 deployed medics than those who have never been deployed. However, no significant  
10  
11           5 difference was found at three to 12 months post-deployment with controlled demographic  
12  
13           6 variables (Chapman et al. 2014). Hierarchical regression reveals that PTSD was still present  
14  
15           7 in combat medics, three months post-deployment. However, it was not significant in  
16  
17           8 comparison with the peer group. Amongst a group of combat medics, those who screened  
18  
19           9 positive for a mental health problem reported higher concerns regarding stigma and barriers  
20  
21           10 to care, where stigma was perceived as a more prominent issue than barriers of care.  
22  
23           11 Additionally, those who required the highest level of assistance perceived more barriers to  
24  
25           12 mental health care and stigma for seeking treatment (Chapman et al. 2014).

26  
27           13 Sørensen et al., (2016) conducted a similar study with a special emphasis on  
28  
29           14 symptoms of PTSD. Results revealed its presence in both studies as a post-combat mental  
30  
31           15 health difficulty. These findings were further supported by Shea et al., (2013) in which it was  
32  
33           16 reported that 12.6% of the veterans fulfilled the full criteria for PTSD within first six months  
34  
35           17 of post deployment. Adding to aforementioned research findings, Shea et al., (2017)  
36  
37           18 presented that veterans who endorsed danger to themselves, or exposure to death/serious  
38  
39           19 injury of others, revealed increased levels of depression, anxiety, avoidance, re-experiencing,  
40  
41           20 and hyperarousal symptoms; whilst killing someone else had not significantly associated with  
42  
43           21 any symptom measures or guilt.

44  
45           22 Albeit previous findings, Presseau et al., (2019), investigated the prevalence of trauma  
46  
47           23 types, prevalence of PTSD within these trauma types, and the association among trauma  
48  
49           24 types, mental health, and behavioural outcomes in an epidemiological sample of service  
50  
51           25 members. It was revealed that Life Threat to Self (LTS; 51%) and Life Threat to Others

1 (LTO; 31%) were the most endorsed events. Moral Injury – Self (MIS) and Aftermath of  
2 Violence (AV) were the least endorsed events. Nearly 40% of the participants endorsed  
3 multiple events. These complex traumas were not associated with more severe symptoms of  
4 mental health difficulties (Presseau et al., 2019). There were also no differences in the PTSD  
5 prevalence by the trauma type. Only 10% of those exposed to Criterion A events (i.e.,  
6 “exposure to actual or threatened death, serious injury, or sexual violence”) were diagnosed  
7 of PTSD. Although the majority of participants who endorsed Criterion A events reported  
8 high use of alcohol, anxiety, depression, suicide ideation, PTSD symptom severity, and  
9 combat exposure, 90% of them were not diagnosed with PTSD. Aftermath of Violence (AV)  
10 was positively correlated with the number of PTSD symptoms (Presseau et al., 2019). LTO  
11 and LTS did not report arousal or anxiety, but LTS was related to depression. Participants  
12 who endorsed LTS and AV events in Criterion A trauma types reported severe suicidal  
13 ideation and their deployment experience to be more distressing (Presseau et al., 2019).

14 ***Aggression and Increased Psychiatric Symptoms.*** Worthen et al., (2021) reported an  
15 association between post-deployment [perceived] difficulties in controlling violent  
16 behaviours, and deployment related traumas and PTSD symptom severity. With each  
17 additional deployment, violent behaviours, post-deployment traumas, and PTSD symptom  
18 severity worsened. This prevalence was not impacted by the use of alcohol, age, gender,  
19 and/or marital status. However, findings should be interpreted cautiously as only 3% of the  
20 respondents reported [perceived] difficulties in controlling violent behaviours.

21 In addition Worthen et al., (2021) only assessed violent behaviours in a general  
22 manner, whereas Litz et al., (2018) researched about the frequency of different trauma types  
23 in a population of service members seeking treatment for PTSD. They note how “moral  
24 injury – self” and “traumatic loss trauma” types reported more severe re-experiencing, guilt,  
25 and sadness, compared to the “life threat – self” type. Compared to the “life threat – self”

1  
2  
3 1 type, the “violence trauma” type reported more peri- and post traumatic sadness. The “Moral  
4  
5 2 injury – others” type reported greater betrayal/humiliation and more frequent aggressive  
6  
7 3 behaviours. Participants reported ‘non-threat’ events to be a more distressing war zone  
8  
9 4 experience than ‘threat’ events. ‘Life threat-self’ events were mostly endorsed by young  
10  
11 5 participants, who had a low number of service years and exposure to war stressors (Litz et al.,  
12  
13 6 2018).

14  
15  
16  
17 7 **Risk Factors Aggravating Mental Health Difficulties of First Respondents, Post**  
18  
19 8 **Exposure.**

20  
21 9 These were collapsed into personal circumstances and situational circumstances, as  
22  
23 10 follows.

24  
25  
26 11 *Personal Circumstances.* A sample of National Guards and Reserve units were  
27  
28 12 screened for PTSD, utilising the Clinician Administered PTSD Scale (CAPS). A few pre-  
29  
30 13 deployment risk factors for the development of PTSD were hypothesised: negative  
31  
32 14 temperament, prior history of stressful life events, and perceived inadequacy of training. Out  
33  
34 15 of 238 participants, 12.6% met full criteria of PTSD within the first six months post-  
35  
36 16 deployment (Shea et al., 2013). Participants with PTSD scored differently in all aspects of the  
37  
38 17 aforementioned risk factors compared to those without PTSD, except on unit support and  
39  
40 18 perceived preparation and training. Demographic factors, such as gender, ethnicity, and age  
41  
42 19 were not significant predictors of PTSD. Negative temperament and pre-deployment life  
43  
44 20 events independently predicted PTSD, while deployment-related variables, such as combat  
45  
46 21 experience, and life and family concerns, were found to be significant predictors of PTSD  
47  
48 22 (Shea et al., 2013).

49  
50  
51  
52  
53  
54 23 Arble & Arnetz (2016) identified that participants who have been exposed to stressful  
55  
56 24 experiences practiced more avoidant strategies, received low social support, and exhibited a  
57  
58 25 lack of positive coping mechanisms (i.e., substance use). In conclusion, it was discovered that  
59  
60

1 post-traumatic growth, physical health, social support, and positive coping mechanisms  
2 positively impacted the participants' well-being. Substance use and avoidant coping had,  
3 conversely, negatively impacted participants' well-being.

4 Adding to Arble & Arnetz (2016), Chapman et al., (2014) revealed that well-being of  
5 combat medics who screened positive for mental health difficulties were affected by their  
6 reluctance to seek professional intervention due to concerns about stigma associated with  
7 mental health and barriers to care. A positive correlation was revealed where higher the level  
8 of assistance needed, the perceived barriers to their care and stigmas were greater.

9 Further adding to Arble & Arnetz (2016), Blakey et al., (2022) reported that veterans  
10 diagnosed with substance use disorder (SUD) and comorbid PTSD-SUD revealed higher  
11 rates of risk in lifetime homelessness, violent behaviour, suicidal ideation and attempts, and  
12 poorer quality in vocational, financial, and social well-being. However, there was no  
13 difference in vocational and financial well-being compared to veterans diagnosed with PTSD  
14 only. In contrast to findings of Arble & Arnetz (2016) and Blakey et al., (2022), no  
15 significant association was reported by Worthen et al., (2021) for the relationship between  
16 post-deployment violent behaviour and alcohol abuse.

17 In a sample of veterans who were screened for risk factors for cardiovascular disease  
18 and/or diabetes, the "Low Treatment Need" (LTN) group screened less likely for the four  
19 main risk factors: obesity/overweight, smoking, at-risk for alcohol use, and PTSD or  
20 depression symptoms (Funderburk et al, 2014). The "Moderate Treatment Need" (MTN)  
21 group had a relatively higher probability to screen positively for smoking and at-risk alcohol  
22 use. The "High Treatment Need" (HTN) group reported the highest probability to screen for  
23 all four risks. This indicates that a high prevalence of PTSD or depression symptoms could be  
24 a risk factor for cardiovascular disease and/or diabetes among veterans (Funderburk et al,  
25 2014).

1  
2  
3 1 Affirming Funderburk et al., (2014) findings, Groer et al., (2014) reported that from  
4  
5 2 the 68% of active duty enlisted and reservist soldiers who had experienced previous combat,  
6  
7 3 half were smokers and reported depression (14%), headaches (14%), anxiety (17%), and  
8  
9 4 tinnitus (25%). When screened for PTSD, 21% of the sample reported moderate to severe on  
10  
11 5 the PCL-M scale with 11 veterans with no PTSD symptoms. It was important to note that  
12  
13 6 these 11 veterans had no combat exposure. In the overall sample, infectious illnesses, chronic  
14  
15 7 inflammation, autoimmune processes, or cardiovascular disease was indicated through high  
16  
17 8 levels of C-reactive Protein (CRP). Depression, PTSD, and CRP levels were highly  
18  
19 9 correlated with hair cortisol levels; a marker of chronic stress. However, these factors had a  
20  
21 10 low correlation with Combat Exposure Scale (CES) scores. High depression scores and  
22  
23 11 combat exposure were found to be significant contributors to high scores of PTSD. However,  
24  
25 12 hair cortisol and CRP levels could not predict high scores of PTSD when depression and  
26  
27 13 combat experience were added to the regression model.  
28  
29  
30  
31  
32

33 14 Gender differences in risk factors that may aggravate mental health difficulties in  
34  
35 15 veterans were revealed by Adams et al., (2021): the majority of female veterans reported to  
36  
37 16 be younger, unmarried, college graduates, and had less combat exposure but were more likely  
38  
39 17 to have lifetime mental health difficulties (e.g., PTSD, depression, and suicidal ideation).  
40  
41 18 They also reported low psychological resilience but were more likely to use mental health  
42  
43 19 services and psychotropic medications. No differences were reported for childhood  
44  
45 20 abuse/neglect, stressful events in the past year or lifetime traumas, social support, alcohol  
46  
47 21 misuse, and/or self-esteem, between female and male veterans. However, it should be noted  
48  
49 22 that these findings were derived by analysing a small number of female respondents (5%; n =  
50  
51 23 85) from the total sample.  
52  
53  
54  
55

56 24 ***Situational Circumstances.*** Exposure to combat is a widely regarded risk factor of  
57  
58 25 PTSD. Amongst a sample of recently deployed, returned National Guards, and reserve  
59  
60



1  
2  
3 1 members (Shea et al., 2017), over 50% of the exposure types reported were: being attacked or  
4  
5 2 ambushed, receiving small arms fire, exposure to improvised explosive devices, seeing dead  
6  
7 3 bodies/remains, and/or knowing someone seriously injured or killed. The less commonly  
8  
9 4 reported was: having killed the enemy (11%) and the killing of non-combatants (1.3%),  
10  
11 5 where “having killed” was not significantly associated with guilt and/or other symptom  
12  
13 6 measures. Hyperarousal symptoms, depression, re-experiencing, avoidance, and anxiety were  
14  
15 7 uniquely and significantly associated with “danger to oneself” and “exposure to death or  
16  
17 8 serious injury of others”. Both “danger to oneself” and “exposure to death or serious injury of  
18  
19 9 others” were significantly associated with the total Clinician-Administered PTSD Scale for  
20  
21 10 DSM–IV (CAPS-IV) score.  
22  
23  
24

25  
26 11 These findings by Shea et al., (2017) study was further supported by Litz et al.,  
27  
28 12 (2018), which revealed that among participants receiving treatment for PTSD, more severe  
29  
30 13 symptoms were reported by veterans who endorsed traumatic loss trauma, moral injury-  
31  
32 14 others, moral injury-self, and violence trauma type than those who endorsed life threat-self.  
33  
34 15 Life threat-self events were mostly reported by young participants with a short service  
35  
36 16 duration and low exposure to combat. In contrast to the other findings of Shea et al., (2017),  
37  
38 17 participants from Litz et al., (2018) reported non-threatening events (i.e., concern about  
39  
40 18 family), to be more distressing than threatening experiences in the war zone. Furthermore,  
41  
42 19 Presseau et al., (2019) reported that Life Threat to Self (LTS; 51%) and Life Threat to Other  
43  
44 20 (LTO; 31%) were more commonly endorsed by participants, even though there was no  
45  
46 21 association of these events with symptoms of mental health difficulties.  
47  
48  
49  
50  
51  
52  
53

54 23 **Protective Factors Supporting Positive Management/Emergence of Mental Health**  
55  
56 24 **Difficulties.**  
57  
58  
59  
60



1  
2  
3 1 According to Arble & Arnetz (2016), the well-being of veterans who have been  
4  
5 2 exposed to combat has been positively influenced by stable physical health, high social  
6  
7 3 support, approach to coping mechanisms, and post-traumatic growth. Adding to Arble &  
8  
9 4 Arnetz (2016), Worthen et al., (2021) revealed that post-deployment violent behaviours were  
10  
11 5 comparatively lower for Caucasian veterans who reported higher levels of education  
12  
13 6 qualifications and who received social support.  
14  
15

16  
17 7 Furthermore, Shea et al., (2013) revealed that participants with high levels of  
18  
19 8 perceived adequacy of training and low exposure to combat did not report PTSD, meaning  
20  
21 9 pre-deployment preparation and training, and low severity of exposure during deployment  
22  
23 10 have contributed as protective factors against developing mental health difficulties amongst  
24  
25 11 first respondents.  
26  
27  
28  
29  
30

31 <Insert Table 2 here>  
32  
33  
34

## 35 15 Discussion

### 36 16 Population Characteristics

37  
38  
39 17 The majority of study participants were middle-aged Caucasian and non-Hispanic  
40  
41 18 men from the United States of America, whose responses were recorded in English. The  
42  
43 19 exceptions were studies with non-English speaking participants from Denmark (Sørensen et  
44  
45 20 al., 2016) and Sweden (Arble & Arnetz, 2016). Female participants were analysed in seven  
46  
47 21 studies (Adams et al., 2021; Chapman et al., 2014; Groer et al., 2014; Litz et al., 2018; Shea  
48  
49 22 et al.; 2013; Shea et al., 2017; and Worthen et al., 2021), yet the number of female  
50  
51 23 participants were disproportionately small in comparison to men. Therefore, applicability of  
52  
53 24 the review findings primarily applies to male population. Nine studies (Adams et al., 2021;  
54  
55 25 Blakey et al., 2022; Chapman et al., 2014; Groer et al., 2014; Litz et al., 2018; Presseau et al.,  
56  
57  
58  
59  
60

1  
2  
3 1 2019; Shea et al., 2013; Shea et al., 2017; and Worthen et al., 2021) collected information on  
4  
5  
6 2 the marital status of participants, which was analysed as “social support” when considering  
7  
8 3 its impact on trauma recovery. Chapman et al., (2014) excluded veterans with severe physical  
9  
10 4 injuries requiring overnight hospitalisation, due to the high correlation between physical  
11  
12 5 injury and mental health difficulties.  
13  
14

### 6 **Most Prevalent Psychological Difficulties and Related Risk Factors**

15  
16  
17 7 PTSD and depression are the most common diagnoses, followed by anxiety (Chapman et al.,  
18  
19 8 2014; Groer et al., 2014). These mental health difficulties are prone to be aggravated by  
20  
21 9 specific risk factors, which can be broadly categorised as personal and situational. Identified  
22  
23 10 personal factors were decreased physical health (e.g., obesity), low cognitive abilities,  
24  
25 11 existing mental health difficulties, job position, threat to self, peers and family, and use of  
26  
27 12 avoidant strategies (e.g., substance abuse; Arble & Arnetz, 2016; Blakey et al., 2022; Litz et  
28  
29 13 al., 2018; Shea et al., 2013; Sørensen et al., 2016). Situational factors were identified such as  
30  
31 14 lack of social support, past stressful experiences, combat frequency and severity, and social  
32  
33 15 stigma (Arble & Arnetz, 2016; Blakey et al., 2022; Chapman et al., 2014; Worthen et al.,  
34  
35 16 2021). In contrast, post-crisis mental health difficulties could be minimised with the influence  
36  
37 17 of protective factors, such as high cognitive ability, social support, effective coping strategies,  
38  
39 18 pre-deployment preparedness, and/or robust physical health (Arble & Arnetz, 2016; Elbogen  
40  
41 19 et al., 2012; Kline et al., 2013; Sørensen et al., 2016).  
42  
43  
44  
45  
46  
47

48  
49 20 PTSD and depression symptoms in some cases were reported even six months post  
50  
51 21 deployment and sustained for 12-months post deployment (Chapman et al., 2014; Shea et al.,  
52  
53 22 2013). This implies a short onset yet lasting effects of the combat exposure on an individual’s  
54  
55 23 mental health state. The relationship between PTSD and combat exposure can be clearly  
56  
57 24 stated, independent of any confounding variables, as service personnel who have not been  
58  
59  
60

1  
2  
3 1 exposed to crisis situations, or have a short period of exposure to combat report less or no  
4  
5 2 symptoms of PTSD (Chapman et al., 2014; Groer et al., 2014).  
6  
7

8 3 Contrary to the common assumption that an individual may be more distressed by  
9  
10 4 having to kill someone due to a sense of morality, the review findings revealed that veterans  
11  
12 5 who reported a higher number of mental health symptoms expressed more concern about  
13  
14 6 their personal safety and apprehension for fellow veterans (Shea et al., 2017). A strong sense  
15  
16 7 of self and solidarity are demonstrated here, over the effects of moral injury. This arguably  
17  
18 8 aligns with both Self-Determination Theory (SDT; Ryan & Deci, 2000) and the Pluralistic  
19  
20 9 Trauma Theory (PTT; Balaev, 2018), namely that individuals are more affected by their  
21  
22 10 subjective perception of what is considered to be socially acceptable (e.g., eradicating  
23  
24 11 terrorism is essential), rather than by the objective outcome (e.g., death) of a crisis situation—  
25  
26 12 especially during militarised tasks where individual autonomy could be compromised but the  
27  
28 13 need and confidence to perform are boosted.  
29  
30  
31  
32

33 14 As discussed, several research findings have continually revealed that exposure to  
34  
35 15 crisis situations may manifest as mental health difficulties in veterans. However, in some  
36  
37 16 research where multiple types of trauma (life and/or moral threat to self and/or others) are  
38  
39 17 analysed, findings indicated no significant relationship between the exposure to crisis  
40  
41 18 situations and mental health difficulties (Presseau et al., 2019). This differential finding  
42  
43 19 cannot seemingly be attributed to the status of the sample or geographical location (i.e.,  
44  
45 20 American veterans in active duty), as other research had analysed samples of American  
46  
47 21 veterans in active duty (Groer et al., 2014; Litz et al., 2018). Therefore, further analysis of the  
48  
49 22 sample, measures, and methodology is warranted to determine the possibility of veterans not  
50  
51 23 significantly affected by exposure to crisis situations. However, this differential finding may  
52  
53 24 be somewhat explained by Litz et al., (2018), who reported that effects of life threat-self  
54  
55 25 events were mostly endorsed by participants of young age with low exposure to combat and  
56  
57  
58  
59  
60

1 service period, which is supported by the Job Demand-Control-Support (JDCS) model  
2 (Johnson & Hall, 1988), which states that individuals may be stressed by their tasks *only* if  
3 there is high demand, low control and social support. Alternatively, it is plausible, as per the  
4 Effort-Reward Imbalance model (Siegrist, 1996), that uniformed services were not affected  
5 by the exposure to crisis situations, as they found their experience to be highly rewarding  
6 (e.g., safeguarding civilians). Additionally, the PTT also states that the impact of trauma  
7 comes to fruition *as a result* of the interactions between external (i.e., situation) and internal  
8 (i.e., personality) factors. Therefore, due to their subjective perception and diminished  
9 exposure to crisis situations, the trauma response may not have manifested as mental health  
10 difficulties.

11 Key personal and situational risk factors that were discussed earlier in this section are  
12 supported by both PTT and SDT theories, which focus on the influence of internal/external  
13 stressors and intrinsic/extrinsic motivational factors, which contributes to an individual's  
14 social, cognitive and personality development, including the specific manifestation of trauma.

15 All aforementioned risk factors are domains that will aggravate mental health issues, post-  
16 deployment upon return. Addressing these factors require treatment and intervention  
17 strategies over a preventive approach.

18 In order to lessen the effect of combat exposure manifesting as mental health  
19 difficulties, the aforementioned intrinsic/extrinsic risk factors should be a focal point to be  
20 identified and assessed *pre-deployment*. This will contribute to a more effective recruitment  
21 and training within uniformed forces, and improved trauma recovery post deployment. Even  
22 though Shea et al., (2013) and Worthen et al., (2021) reported that demographic factors (e.g.,  
23 age/gender) and pre-deployment variables (e.g., negative temperament) could not predict  
24 PTSD; low cognitive ability, overweight/obesity, pre-deployment emotional problems,  
25 [perceived] low social support, negative childhood experiences, and stressful experiences

1 during a participant's lifetime were reported by other research studies as pre-deployment  
2 factors that may increase the vulnerability of a veteran to be effected by combat exposure  
3 (Sørensen et al., 2016).

4 Crisis respondents and their respective organisations have low control regarding  
5 factors during deployment that may enhance the risk to develop and worsen mental health  
6 difficulties. However, it is useful to be mindful of the intrinsic/extrinsic risk factors for post-  
7 deployment assessment, treatment, and intervention, as their importance is emphasised by  
8 PTT and SDT theories. Higher frequency and subjective severity of combat exposure,  
9 holding a more stressful position during the mission, increased rates of combat stress reaction  
10 (CSR), non-threatening events such as high levels of concerns for life and family,  
11 danger/injury during combat to self and/or others, exposure to death, and a higher kill rate of  
12 the enemy were the most prominent risk factors revealed by this narrative review (Litz et al.,  
13 2018; Shea et al., 2013; Shea et al., 2017; Sørensen et al., 2016; Shea et al., 2013). Even  
14 though Shea et al., (2017) reported that "having killed" an enemy did not contribute to mental  
15 health difficulties, this can be explained by the low number of "having killed" item endorsed  
16 by study participants as an event experienced during combat. Contrary to aforementioned  
17 research studies, Presseau et al., (2019) reported that even though Life Threat to Self (LTS)  
18 and Life Threat to Other (LTO) were more commonly endorsed by the participants, these  
19 events did not contribute to the mental health difficulties of the participants. As previously  
20 noted, these findings align with PTT (i.e., the impact of subjective perception and societal  
21 acceptance of trauma experience and response), SDT (i.e., the encouragement of need,  
22 confidence and freedom to perform) theories and the Effort-Reward Imbalance model (i.e.,  
23 high gain over cost). Additionally, more in-depth analysis of methodology and populations in  
24 study of Presseau et al (2019) is required in order to determine if it is an exception to existing  
25 research findings.

## 1 **Helpful Protective Factors in Trauma Recovery for First Respondents**

2           Despite its vital role in trauma recovery, there appears to be less reporting of factors  
3 that could lessen the severity of mental health difficulties of crisis respondents post exposure  
4 to traumatic events. This may be either due to lack of such factors or the low interest of  
5 researchers in investigating the existence and influence of protective factors. The impact of  
6 combat exposure on mental health was arguably lessened by factors such as post-traumatic  
7 growth, high social support, physical health, effective strategies of coping, and high levels of  
8 [perceived adequacy in] pre-deployment preparation and training (Arble & Arnetz., 2016;  
9 Shea et al., 2013; Worthen et al., 2021). Even though the most stable subgroup did not report  
10 a higher mean score of pre-deployment cognitive ability compared to the subgroup of  
11 participants with symptoms with moderate severity, Sørensen et al., (2016) revealed an  
12 inverse relationship between pre-deployment cognitive ability and PTSD symptom severity.

## 13 **Limitations of the Present Narrative Review**

14           The majority of articles included in the narrative review assessed English speaking  
15 middle-aged Caucasian and non-Hispanic male crisis respondents from Western countries.  
16 Therefore, this review lacks insight into mental health difficulties, and risk/protective factors  
17 post exposure to crisis situations across populations with different socioeconomic status, age,  
18 ethnicity, religion, gender, cultures, and geographical locations. Additionally, the majority of  
19 the crisis situations were combat exposure, where uniformed forces have been deployed to  
20 foreign countries. The effect of other crisis situations, such as natural disasters and the  
21 response of service personnel in attending to crisis situations in their home country, has not  
22 been explored. The findings can also not be generalised, as the search strategy was limited to  
23 English publications in the past decade.

24           Evidently, all articles except one (Sørensen et al., 2016) had focused on post-  
25 deployment phase of the soldiers. Therefore, the current research domain is greatly

1  
2  
3 1 disadvantaged due to the lack of insight during pre- and during deployment stages. As a  
4  
5 2 result, it would be difficult for practitioners to encourage prevention, in addition to treatment.  
6  
7 3 As noted, it was also difficult to determine the impact of gender, mainly due to the nature of  
8  
9 4 recruitment and deployment in uniformed forces; females have a comparatively lower  
10  
11 5 opportunity to serve as first respondents. This is a major limitation as current research is  
12  
13 6 unaware of the influential nature of gender, such as whether it serves as a risk or protective  
14  
15 7 factor against mental health difficulties, in combination with other contributing variables post  
16  
17 8 exposure to crisis situations.  
18  
19  
20

21 9 Except for Shea et al. (2017), which conducted CAPS-IV structured interviews (Blake  
22  
23 10 et al., 1995), other studies used questionnaires to gather information. This arguably limits the  
24  
25 11 responses of participants. Moreover, it is recommended that future research verify the  
26  
27 12 responses of self-report measures with collateral information obtained from independent third  
28  
29 13 parties. Future research should, overall, seek to address the gaps in the literature revealed by  
30  
31 14 this narrative review and expand inquiry beyond Western understandings. In doing so it  
32  
33 15 should adopt a wider application of method to this important area of study. Future research  
34  
35 16 could also consider application of a meta-analysis, using a wider sampling of studies as the  
36  
37 17 area continues to develop following continued embedding of DSM-5. Such analysis could  
38  
39 18 account more thoroughly for estimates of effect and heterogeneity between study results,  
40  
41 19 allowing for an enhanced objective appraisal of evidence.  
42  
43  
44  
45  
46

## 47 20 **Implications for Practice, Policy, and Future Research**

48  
49 21 As noted by the current review, there is an evident relationship between mental health  
50  
51 22 difficulties and an influence of personal and/or situational characteristics for first  
52  
53 23 respondents. Therefore, they can be assessed and treated for these identified vulnerabilities,  
54  
55 24 and trained to improve resilience prior to deployment. This pre-deployment screening process  
56  
57 25 could provide insights to organisations to implement changes to their need-based training  
58  
59  
60

1  
2  
3 1 processes and recruit individuals who are considered the best fit for deployment. However,  
4  
5 2 more inclusive research is needed to understand the impact of diverse demographic factors  
6  
7 3 across multiple geographical locations for different types of crises and deployment.  
8  
9

10 4 It is also important to assess the specific mental health difficulties and related  
11  
12 5 risk/protective factors identified by this review. This should occur during deployment in  
13  
14 6 order to enhance the well-being of first respondents in the course of active service. Such  
15  
16 7 efforts will not only contribute to deployed mission objectives overall and support optimal  
17  
18 8 individual performance, but could also minimise the risk of imminent and/or post-deployed  
19  
20 9 mental health difficulties. Despite logistical difficulties likely to arise in conducting research  
21  
22 10 during deployment, it is highly encouraged as an area future research could explore, and one  
23  
24 11 that could identify potential means preventing the aggravation of future trauma.  
25  
26  
27

28 12 Ultimately, this review provides scholarly support to researchers and practitioners in  
29  
30 13 identifying the most common mental health difficulties of first respondents, post exposure to  
31  
32 14 crisis situations. The reported personal and situational risk and/or protective factors may be  
33  
34 15 helpful in future recruitment, training, and/or trauma recovery interventions. It could also  
35  
36 16 help to direct the specific factors that should be identified and follow-up as part of  
37  
38 17 longitudinal research.  
39  
40  
41  
42  
43  
44

## 45 19 **Conclusion**

46  
47 20 This narrative review reveals that there is a clear indication of mental health  
48  
49 21 difficulties, post exposure to crisis situations among first responders, which are enhanced or  
50  
51 22 minimised by various risk and/or protective factors. Post-traumatic Stress Disorder and  
52  
53 23 depression were the most common diagnoses, followed by the impact of risk and protective  
54  
55 24 factors that could be considered for effective pre-deployment recruitment and training, and  
56  
57 25 post-deployment trauma recovery.  
58  
59  
60



## References


- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Adams, R., Hu, Y., Figley, C., Urosevich, T., Hoffman, S., Kirchner, H., Dugan, R., Boscarino, J., Withey, C. and Boscarino, J., (2021). Risk and Protective Factors Associated with Mental Health Among Female Military Veterans: Results from the Veterans' Health Study. *BMC Women's Health*, 21(1). <https://doi.org/10.1186/s12905-021-01181-z>
- Arble, E., & Arnetz, B. (2016). A Model of First-responder Coping: An Approach/Avoidance Bifurcation. *Stress and Health*, 33(3), 223-232. <https://doi.org/10.1002/smi.2692>
- Balaev, M. (2018). Trauma Studies. In D. Richter, *A Companion to Literary Theory* (pp. 360-373). Wiley-Blackwell.
- Blakey, S., Dillon, K., Wagner, H., Simpson, T., Beckham, J., Calhoun, P. and Elbogen, E., (2022). Psychosocial Well-Being Among Veterans with Posttraumatic Stress Disorder and Substance Use Disorder. *Psychological Trauma: Theory, Research, Practice, and Policy*, 14(3), pp.421-430. <https://doi.org/10.1037/tra0001018>
- Bowler, R., Kornblith, E., Li, J., Adams, S., Gocheva, V., Schwarzer, R., & Cone, J. (2016). Police Officers Who Responded to 9/11: Comorbidity of PTSD, Depression, and Anxiety 10-11 Years Later. *American Journal of Industrial Medicine*, 59(6), 425-436. <https://doi.org/10.1002/ajim.22588>
- Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, Hartmann-Boyce J, Ryan R, Shepperd S, Thomas J, Welch V, Thomson H. (2020) Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline BMJ 2020;368:l6890 <http://dx.doi.org/10.1136/bmj.l6890>
- Chapman, P., Elnitsky, C., Pitts, B., Figley, C., Thurman, R., & Unwin, B. (2014). Mental Health, Help Seeking, and Stigma and Barriers to Care Among 3- and 12-Month

- 1  
2  
3 1 Postdeployed and Never Deployed U.S. Army Combat Medics. *Military*  
4  
5 2 *Medicine*, 179(8S), 55-62. <https://doi.org/10.7205/milmed-d-12-00367>  
6  
7  
8 3 Elbogen, E., Johnson, S., Wagner, H., Newton, V., Timko, C., Vasterling, J., & Beckham, J.  
9  
10 4 (2012). Protective Factors and Risk Modification of Violence in Iraq and Afghanistan  
11  
12 5 War Veterans. *The Journal of Clinical Psychiatry*, 73(06), e767-e773.  
13  
14 6 <https://doi.org/10.4088/jcp.11m07593>  
15  
16  
17 7 Funderburk, J., Kenneson, A., & Maisto, S. (2014). Identifying Classes of Veterans with  
18  
19 8 Multiple Risk Factors. *Military Medicine*, 179(10), 1119-1126.  
20  
21 9 <https://doi.org/10.7205/milmed-d-14-00119>  
22  
23  
24 10 Groer, M., Kane, B., Williams, S., & Duffy, A. (2014). Relationship of PTSD Symptoms  
25  
26 11 with Combat Exposure, Stress, and Inflammation in American Soldiers. *Biological*  
27  
28 12 *Research for Nursing*, 17(3), 303-310. <https://doi.org/10.1177/1099800414544949>  
29  
30  
31 13 Ikin, J., McKenzie, D., Gwini, S., Kelsall, H., Creamer, M., & McFarlane, A. et al.  
32  
33 14 (2020). *Major Depression and Depressive Symptoms in Australian Gulf War Veterans*  
34  
35 15 *20 Years after the Gulf War*. <https://doi.org/10.1016/j.jad.2015.09.016>  
36  
37  
38 16 Joanna Briggs Institute. (2017). Checklist for Systematic Reviews and Research Syntheses.  
39  
40 17 [https://joannabriggs.org/ebp/critical\\_appraisal\\_tools](https://joannabriggs.org/ebp/critical_appraisal_tools)  
41  
42  
43 18 Kline, A., Ciccone, D., Weiner, M., Interian, A., Hill, L., & Falca-Dodson, M. et al. (2013).  
44  
45 19 Gender Differences in the Risk and Protective Factors Associated With PTSD: A  
46  
47 20 Prospective Study of National Guard Troops Deployed to Iraq. *Psychiatry Interpersonal*  
48  
49 21 *& Biological Processes*, 76(3), 256-72. <https://doi.org/10.1521/psyc.2013.76.3.256>  
50  
51  
52 22 Legault, L. (2017). Self-Determination Theory. *Encyclopedia of Personality and Individual*  
53  
54 23 *Differences*, 1-9. [https://doi.org/10.1007/978-3-319-28099-8\\_1162-1](https://doi.org/10.1007/978-3-319-28099-8_1162-1)  
55  
56 24 Litz, B., Contractor, A., Rhodes, C., Dondanville, K., Jordan, A., & Resick, P. et al. (2018).  
57  
58  
59  
60

- 1  
2  
3 1 Distinct Trauma Types in Military Service Members Seeking Treatment for  
4  
5 2 Posttraumatic Stress Disorder. *Journal of Traumatic Stress*, 31(2), 286-295.  
6  
7  
8 3 <https://doi.org/10.1002/jts.22276>  
9  
10 4 MacManus, D., Dickson, H., Short, R., Burdett, H., Kwan, J., & Jones, M. et al. (2019). Risk  
11  
12 5 and Protective Factors for Offending among UK Armed Forces Personnel after They  
13  
14 6 Leave Service: A Data Linkage Study. *Psychological Medicine*, 1-8.  
15  
16 7 <https://doi.org/10.1017/S0033291719003131>  
17  
18 8 Presseau, C., Litz, B., Kline, N., Elsayed, N., Maurer, D., & Kelly, K. et al. (2019). An  
19  
20 9 Epidemiological Evaluation of Trauma Types in a Cohort of Deployed Service  
21  
22 10 Members. *Psychological Trauma: Theory, Research, Practice, And Policy*, 11(8), 877-  
23  
24 11 885. <https://doi.org/10.1037/tra0000465>  
25  
26  
27 12 Shah, R., Shah, A., & Links, P. (2012). Post-traumatic Stress Disorder and Depression  
28  
29 13 Comorbidity: Severity Across Different Populations. *Neuropsychiatry*, 2(6), 521-529.  
30  
31 14 <https://doi.org/10.2217/npv.12.56>  
32  
33  
34 15 Shea, M., Reddy, M., Tyrka, A., & Sevin, E. (2013). Risk Factors for Post-deployment  
35  
36 16 Posttraumatic Stress Disorder in National Guard/Reserve Service Members. *Psychiatry*  
37  
38 17 *Research*, 210(3), 1042-1048. <https://doi.org/10.1016/j.psychres.2013.08.039>  
39  
40  
41 18 Shea, M., Presseau, C., Finley, S., Reddy, M., & Spofford, C. (2017). Different Types of  
42  
43 19 Combat Experiences and Associated Symptoms in OEF and OIF National Guard and  
44  
45 20 Reserve Veterans. *Psychological Trauma: Theory, Research, Practice, And*  
46  
47 21 *Policy*, 9(Suppl 1), 19-24. <https://doi.org/10.1037/tra0000240>  
48  
49  
50 22 Siegrist, J. (2016). Effort-Reward Imbalance Model. *Stress: Concepts, Cognition, Emotion,*  
51  
52 23 *and Behavior*, 81-86. <https://doi.org/10.1016/b978-0-12-800951-2.00009-1>  
53  
54  
55 24 Sørensen, H., Andersen, S., Karstoft, K., & Madsen, T. (2016). The Influence of Pre-  
56  
57 25 deployment Cognitive Ability on Post-traumatic Stress Disorder Symptoms and  
58  
59  
60

- 1  
2  
3 1 Trajectories: The Danish USPER Follow-up Study of Afghanistan Veterans. *Journal of*  
4  
5 2 *Affective Disorders*, 196, 148-153. <https://doi.org/10.1016/j.jad.2016.02.037>  
6  
7  
8 3 Van der Doef, M., & Maes, S. (1999). The Job Demand-Control (-Support) Model and  
9  
10 4 psychological well-being: A review of 20 years of empirical research. *Work &*  
11  
12 5 *Stress*, 13(2), 87-114. <https://doi.org/10.1080/026783799296084>  
13  
14  
15 6 Wang, H., Jin, H., Nunnink, S., Guo, W., Sun, J., & Shi, J. et al. (2011). Identification of  
16  
17 7 Post-Traumatic Stress Disorder and Risk Factors in Military First Responders 6 Months  
18  
19 8 after Wen Chuan Earthquake in China. *Journal of Affective Disorders*, 130(1-2), 213-  
20  
21 9 219. <https://doi.org/10.1016/j.jad.2010.09.026>  
22  
23  
24 10 Worthen, M., Rathod, S., Cohen, G., Sampson, L., Ursano, R., Gifford, R., Fullerton, C.,  
25  
26 11 Galea, S. and Ahern, J., (2021). Risk and Protective Factors for Difficulty Controlling  
27  
28 12 Violent Behavior in National Guard and Reserve Service Members. *Journal of*  
29  
30 13 *Interpersonal Violence*, 36(3-4), pp.1049-1067.  
31  
32 14 <https://doi.org/10.1177/0886260517737552>  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1. Extraction of Articles



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

Table 1: Study Characteristics

Author	Sample	Age (M, SD)	Type of Crisis
Adams et al., 2021	Community-based military veterans from an outpatient clinic, USA (N=1,730)	$M=59$ ( $SD=12$ )	Combat
Arble & Arnetz, 2016	First responders: coast guard, customs control, military, emergency medical services, fire department and police services, Sweden (N = 3,656)	67% in the age range was between 30 -55 years	Not available
Blakey et al., 2022	Military veterans who served on or after September 11, 2001, in Operation Iraqi Freedom (OIF) or Operation Enduring Freedom (OEF); active duty or serving in the National Guard or Reserves, USA (N=1,102; SUD only, N= 207, 18.81%; PTSD only, N = 127, 11.47%; PTSD/SUD, N = 96, 8.75%; and neither disorder, N= 672, 60.97%)	Neither ( $M=38.16$ , $SD=9.27$ ); SUD only ( $M=32.81$ , $SD=8.85$ ); PTSD only ( $M=35.70$ , $SD=8.06$ ); PTSD-SUD ( $M=32.81$ , $SD=8.20$ )	Combat
Chapman et al., 2014	Army combat medics served both as a soldier and medic, USA (N=799)	Never deployed ( $M=25.86$ , $SD=6.01$ ); 3 months post-deployment ( $M=27.97$ , $SD=6.16$ ); 12 months post-deployment ( $M=31.32$ , $SD=6.36$ )	Combat
Funderburk et al., 2014	Veterans, USA (N=28,578)	Exploratory ( $M=63.96$ , $SD=13.41$ ); Confirmatory ( $M=63.88$ , $SD=13.53$ )	Not available
Groer et al., 2014	Active-duty Army and National Guard soldiers (enlisted and reservists) in Bold Quest at Camp Atterbury, USA (N=52)	Mean age of 25 years and a range between 19 and 42 years.	Combat
Litz et al., 2018	Active-duty military service members and recently retired veterans, USA (N=999)	Life threat-self ( $M=32.06$ , $SD=7.28$ ); Life threat-others ( $M=33.67$ , $SD=6.87$ ); Aftermath of violence ( $M=34.28$ , $SD=7.97$ ); Traumatic loss ( $M=32.14$ , $SD=7.47$ ); Moral self-injury	Combat

				(M=32.68, SD=6.66); Moral injury-others (M=32.99, SD=7.33)	
Presseau et al., 2019	Active-duty military members, USA (N=789)	service	Initial sample (M=26.57, SD=6.08); post-deployment sample (M=26.57, SD=6.08); Final sample (M=27.47, SD=6.5)		Combat
Shea et al., 2013	Returnees of the National Guard and Reserve units from Iraq and Afganistan, USA (N=238)		M=33.5 (SD=9.5)		Combat
Shea et al., 2017	Returnees of the National Guard and Reserve units from Iraq and Afganistan, USA (N=238)		M=33.79 (SD=9.65)		Combat
Sørensen et al., 2016	Danish soldiers deployed to Afganistan, Denmark (T1:N=743; T6: N=429)		M=24		Combat
Worthen et al., 2021	National Guard and Reserve veterans, USA (N=1,293)		Not available		Combat



Table 2: Critical Findings

Author	Critical Findings
Adams et al., 2021	From the total sample, only 5% were female (n = 85). Compared to male veterans, female veterans were younger (<65 years; 95%), unmarried (49.4% vs 21.2%), college graduates (47% vs. 24%), had less combat exposure (2% vs. 25%), but were more likely to have lifetime PTSD (29% vs. 12%), depression (46% vs. 21%), suicidal ideation (27% vs. 11%), and lifetime use of mental health services (67% vs. 47%). Female veterans reported low psychological resilience, but more likelihood to use psychological services and psychotropic medications. No sex differences were reported for childhood abuse/neglect, stressful events in the past year or lifetime traumas, social support, alcohol misuse, and/or self-esteem.
Arble & Arnetz, 2016	This study identified that participants who have been exposed to stressful experiences practiced more avoidant strategies, received low social support, and exhibited a lack of positive coping mechanisms. It was noted that the reported lack of appropriate coping strategies, social support, and increased avoidance strategies contributed to post-traumatic growth. Exposure to stress and avoidant coping strategies have contributed to increased substance use. In conclusion, it was discovered that post-traumatic growth, physical health, social support, and positive coping mechanisms positively impacted the participants' wellbeing. Substance use and avoidance coping were found to negatively impact participants' wellbeing.
Blakey et al., 2022	Veterans screening positive for any disorder (SUD-only/PTSD-only/comorbid PTSD-SUD) reported significantly higher risk for rates of lifetime homelessness (15.30% vs 2.05%), violent behaviour (63.16% vs 8.47%), suicidal ideation (43.84% vs 6.51%), and suicide attempts (9.21% vs 0.97%), compared with veterans with neither PTSD nor SUD. Furthermore, veterans in the comorbid PTSD/SUD group reported poorer quality in vocational, financial, and social well-being than veterans with SUD only or with neither disorder, but the PTSD-SUD group did not significantly differ from the PTSD-only group on vocational and financial well-being.
Chapman et al., 2014	Between deployed and undeployed combat medics, medics who have been deployed were found to seek out assistance with their mental health than their peers. The rate of depression was high in deployed medics than those who have never been deployed. However, no significant difference was found at three to 12 months post-deployment with controlled demographic variables. Hierarchical regression reveals that PTSD was still present in combat medics three months post-deployment however, it was not significant in comparison with the peer group. Amongst a group of combat medics, those who screened positive for a mental health problem reported higher concerns regarding stigma and barriers to care, where stigma was perceived as a more prominent issue than barriers of care. Additionally, those who required the highest level of assistance perceived more barriers to mental health care and stigma for seeking treatment.
Funderburk et al., 2014	In a sample of veterans who were primary care patients who were screened for risk factors for cardiovascular disease and/or diabetes, the "Low Treatment Need" (LTN) group screened less likely for the four main risk factors: obesity/overweight, smoking, at-risk for alcohol use, and PTSD or depression symptoms. The "Moderate Treatment Need" (MTN) group had a relatively higher probability to screen positively for smoking and at-risk alcohol use. The "High Treatment Need" (HTN) group reported the highest probability to screen for all four risks. This indicates that a high prevalence of PTSD or depression symptoms could be a risk factor for cardiovascular disease and/or diabetes among veterans.
Groer et al., 2014	Amongst a population of active duty enlisted and reservist soldiers, 68% of the participants have experienced previous combat exposure. Out of the 68%, half were smokers and reported depression (14%), headaches (14%), anxiety (17%), and tinnitus (25%). When screened for PTSD, 21% of the sample reported moderate to severe on the PCL-M scale with 11 veterans with no PTSD symptoms. It was important to note that these 11 veterans have had no combat exposure. In the overall sample, infectious illnesses, chronic inflammation, autoimmune processes, or cardiovascular disease was indicated through high levels of C-reactive Protein (CRP). Depression, PTSD, and CRP levels were highly correlated with hair cortisol levels; a marker of chronic stress. However, these factors had a low correlation with Combat Exposure Scale (CES) scores. High depression scores and combat exposure were found to be significant contributors to high scores for PTSD. However, hair cortisol and CRP levels could not predict high PTSD scores when depression and combat experience were added to the regression model.
Litz et al., 2018	In a study of the frequency of trauma types reported in a population of service members seeking treatment for PTSD, the "moral injury – self" and "traumatic loss trauma" types reported more severe re-experiencing, guilt, and sadness, compared to the "life threat – self" type. Compared to the "life threat – self" type, the "violence trauma" type reported more peri and post traumatic sadness. The "Moral injury – others" type reported greater betrayal/humiliation and more frequent aggressive behaviours. Participants reported 'non-threat' events to be a more distressing war zone experience than 'threat' events. 'Life threat-self' events were mostly endorsed by participants who were young in age and had low number of service years and exposure to war stressors.
Presseau et al., 2019	Prevalence of trauma types, prevalence of PTSD within these trauma types, and the association between trauma types and mental health and behavioural outcomes in an epidemiological sample of service members was investigated. It was revealed that Life Threat to Self (LTS; 51%) and Life Threat to Others (LTO; 31%) were the most commonly endorsed events. Moral Injury – Self (MIS) and Aftermath of Violence (AV) were the least endorsed events. Nearly 40% of the participants endorsed multiple events. These complex traumas were not associated with more severe symptoms of mental health difficulties. There were also no differences in the PTSD prevalence by the trauma type. Only 10% of those



	<p>exposed to Criterion A events – defined as “exposure to actual or threatened death, serious injury, or sexual violence,” experienced personally, witnessed, or learned about – were diagnosed of PTSD. Although majority of these participants who endorsed Criterion A events reported high use of alcohol, anxiety, depression, suicide ideation, PTSD symptom severity, and combat exposure, 90% were not diagnosed with PTSD. AV was positively correlated with the number of PTSD symptoms. LTO and LTS did not report arousal or anxiety, but LTS was related to depression. Participants who endorsed LTS and AV events in Criterion A trauma types reported severe suicidal ideation and their deployment experience to be more distressing.</p>
Shea et al., 2013	<p>With the recognition of the impact post-deployment risk factors on early intervention and prevention of PTSD, a sample of National Guards and Reserve units were screened for PTSD utilising the Clinician Administered PTSD Scale (CAPS). A few pre-deployment risk factors for the development of PTSD were hypothesised: negative temperament, prior history of stressful life events, and perceived inadequacy of training. Out of a sample of 238 participants, 12.6% met full criteria of PTSD within the first six months post-deployment. Participants with PTSD scored differently in all aspects of the aforementioned risk factors compared to those without PTSD, except on unit support and perceived preparation and training. Demographic factors such as gender, ethnicity, and age were not significant predictors of PTSD. Negative temperament and pre-deployment life events independently predicted PTSD while deployment-related variables such as combat experience, and life and family concerns were found to be significant predictors of PTSD. The final model included post deployment variables such as ‘support’ and ‘life events’. Lower levels of post-deployment support significantly predicted beyond the effects of pre- deployment and deployment variables (i.e., post-deployment life stress was marginally significant). Combat experiences and life and family concerns still remained significant predictors of PTSD in the final model, but pre-deployment variables were not. Perceived adequacy of pre-deployment preparation and training, and life and family concerns during deployment revealed a significant interaction with the severity of combat experience. Full criteria of PTSD were fulfilled by 26.6% participants who endorsed higher levels of life and family concerns, and increased levels of combat exposure, but only 2.4% were diagnosed with PTSD when they endorsed higher levels of combat exposure with lower levels of life and family stress. When a low level of combat experience was endorsed by the participants, the interaction with perceived adequacy of pre-deployment training revealed a higher level of perceived adequacy of training. Participants with high levels of combat exposure reported a similar proportion of PTSD (17%; 9/53 and 9/52), independent of the perceived adequacy of training level. Seven percent of the participants with low exposure and perceived adequacy of training reported PTSD whilst participants with low exposure and high perceived adequacy of training did not report PTSD.</p>
Shea et al., 2017	<p>Exposure to combat is a widely regarded risk factor of PTSD. Amongst a sample of recently deployed and returned National Guards and reserve members, over 50% of the exposure were: being attacked or ambushed, receiving small arms fire, exposure to improvised explosive devices, seeing dead bodies/remains, and/or knowing someone seriously injured or killed. The less commonly reported was: having killed the enemy (11%) and endorse the killing of non-combatants (1.3%) where “having killed” was not significantly associated with guilt and/or other symptom measures. Hyperarousal symptoms, depression, re-experiencing, avoidance, and anxiety were uniquely and significantly associated with “danger to oneself” and “exposure to death or serious injury of others”. Both “danger to oneself” and “exposure to death or serious injury of others” were significantly associated with the total CAPS score.</p>
Sørensen et al., 2016	<p>The impact of pre-deployment cognitive abilities on the risk of developing PTSD symptoms or non-resilient PTSD trajectories. The resilient categories (low stable) exhibited low distress levels across six assessments before, during, and after deployment. However, they did not have higher mean pre-deployment cognitive ability scores compared with the mild distress trajectory which exhibited moderate symptoms that decreased after deployment. PTSD symptoms measured as a PCL score revealed an inverse association with pre-deployment cognitive ability. This association remained significant when adjusted for perceived war zone stress, pre-deployment symptoms levels, pre-deployment traumatic life events, and education. Another association was revealed between lower pre-deployment cognitive ability scores and the relieved-worsening trajectory which comprised of a subgroup with PTSD symptoms. This subgroup reported the highest level of pre-deployment emotional problems, the lowest level of perceived social support, the highest level of danger/injury score during the mission, the highest proportion of combat soldiers, a higher proportion of soldiers being wounded or injured during the mission, and a higher proportion of having killed an enemy.</p>
Worthen et al., 2021	<p>Perceived difficulties in controlling violent behaviour were reported by 3.0% of study participants (male: 3.3% and female: 1.7%). Adjusted prevalence ratios (aPR) reported associations between violent behaviour and deployment traumas (aPR = 1.67, 95% confidence interval [CI] = [1.34, 2.08]), PTSD (aPR = 9.95, 95% CI = [5.09, 19.48]), and PTSD symptom severity (aPR for each additional PTSD symptom = 1.07, 95% CI = [1.06, 1.09]). With each additional deployment, perceived difficulties in controlling violent behaviour were increased by 1.67% times (95% CI = [1.34, 2.08]), 80% higher (aPR = 1.80, 95% CI = [1.33, 2.43]) for additional deployment trauma, and 7% higher (aPR = 1.07, 95% CI = [1.06, 1.09]) for each additional symptom of PTSD. A 38% of lower prevalence (aPR = 0.62, 95% CI = [0.52, 0.76]) in violent behaviour was associated with social support. No significant association was reported for the relationship between violent behaviour and alcohol abuse (aPR = 1.94, 95% CI = [0.92, 4.09]). The findings did not differ by age, gender, and/or marital status either, but reported differences for educational status (high school or less: 6.0%; some college: 2.9%; college or more: 1.8%; p = .008) and race (White: 2.0%; non-White: 4.0%; decline to state: 10.6%; p &lt; .001).</p>

Table 3: Implications for Practice, Policy, and Research

Theme	Practical and Policy Implications	Suggestions for Future Research
Before Deployment	Due to the apparent relationship between mental health difficulties and personal and/or situational characteristics, first respondents should be evaluated for their mental health status, and risk and protective factors, prior to recruitment, preferably at multiple time points. This procedure shall then only recruit individuals who are considered to be the best fit for the task.	More inclusive research with diverse demographic factors across multiple geographical locations for different types of crisis and types of deployment should be conducted to assess the nature, severity, and influence of different personal and/ or situational characteristics that may manifest as and/or enhance mental health difficulties of first respondents, should they be exposed to crisis situations.
During Deployment	It is recommended to assess mental health difficulties, and risk and protective factors, which may diminish or enhance the said difficulties of first respondents, during deployment. In doing so, the authorities will not miss to assess any individual as they are nevertheless highly monitored during active service, and the recovery rate will also improve as symptoms may be treated at inception.	There is an apparent lack of inclusive research (as noted above) to assess mental health status, and risk and protective factors of first respondents during deployment, possibly due to circumstances of the atmosphere. However, research during this period should be highly encouraged as it is when first respondents are exposed to crisis situations at a real time and intensity.
After Deployment	Scholars and practitioners have identified the most common mental health difficulties of first respondents, post exposure to a crisis incident. Personal and situational risk and/or protective factors are listed which may be helpful in recruitment, training, and/or trauma recovery interventions.	The majority of research studies in this domain had been conducted after deployment. However, there is an evident lack of longitudinal or follow-up studies. Therefore, scholars and practitioners are uninformed of policy and/or practical implications required for prolonged periods of intervention.

Synthesis Without Meta-analysis (SWiM) reporting items

The citation for the Synthesis Without Meta-analysis explanation and elaboration article is: Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, Hartmann-Boyce J, Ryan R, Shepperd S, Thomas J, Welch V, Thomson H. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline BMJ 2020;368:l6890 <http://dx.doi.org/10.1136/bmj.l6890>

<b>SWiM is intended to complement and be used as an extension to PRISMA</b>			
<b>SWiM reporting item</b>	<b>Item description</b>	<b>Page in manuscript where item is reported</b>	<b>Other*</b>
<i>Methods</i>			
<b>1</b> Grouping studies for synthesis	1a) Provide a description of, and rationale for, the groups used in the synthesis (e.g., groupings of populations, interventions, outcomes, study design)	Page 5	
	1b) Detail and provide rationale for any changes made subsequent to the protocol in the groups used in the synthesis	N/A	
<b>2</b> Describe the standardised metric and transformation methods used	Describe the standardised metric for each outcome. Explain why the metric(s) was chosen, and describe any methods used to transform the intervention effects, as reported in the study, to the standardised metric, citing any methodological guidance consulted	Page 4 – Page 5	
<b>3</b> Describe the synthesis methods	Describe and justify the methods used to synthesise the effects for each outcome when it was not possible to undertake a meta-analysis of effect estimates	Page 7	
<b>4</b> Criteria used to prioritise results for summary and synthesis	Where applicable, provide the criteria used, with supporting justification, to select the particular studies, or a particular study, for the main synthesis or to draw conclusions from the synthesis (e.g., based on study design, risk of bias assessments, directness in relation to the review question)	Page 5	

## Synthesis Without Meta-analysis (SWiM) reporting items

SWiM reporting item	Item description	Page in manuscript where item is reported	Other*
5 Investigation of heterogeneity in reported effects	State the method(s) used to examine heterogeneity in reported effects when it was not possible to undertake a meta-analysis of effect estimates and its extensions to investigate heterogeneity	Page 6	
6 Certainty of evidence	Describe the methods used to assess certainty of the synthesis findings	Page 6	
7 Data presentation methods	Describe the graphical and tabular methods used to present the effects (e.g., tables, forest plots, harvest plots).	N/A	
	Specify key study characteristics (e.g., study design, risk of bias) used to order the studies, in the text and any tables or graphs, clearly referencing the studies included	Page 7	
<i>Results</i>			
8 Reporting results	For each comparison and outcome, provide a description of the synthesised findings, and the certainty of the findings. Describe the result in language that is consistent with the question the synthesis addresses, and indicate which studies contribute to the synthesis	Page 7 – Page 15	
<i>Discussion</i>			
9 Limitations of the synthesis	Report the limitations of the synthesis methods used and/or the groupings used in the synthesis, and how these affect the conclusions that can be drawn in relation to the original review question	Page 20 – Page 21	

PRISMA=Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

\*If the information is not provided in the systematic review, give details of where this information is available (e.g., protocol, other published papers (provide citation details), or website (provide the URL)).

Comment #	Reviewer 1 Comments	Author's Response	Tracker
1	On page 5, line 58, was the agreement between the two screening reviewers quantified?	<p>This information has been highlighted – there was 78% agreement. We are grateful for this clarifying observation.</p> <p>Please check page 6, line 3 – line 4 because text placement has changed.</p>	Page 6, line 3 - line 4
2	<p>On page 6, was the agreement on quality ratings quantified?</p> <p>Also there did not appear to be any reporting on the quality ratings of the studies reviewed.</p>	<p>This information is indicated.</p> <p>Please check page 6, line 8 because text placement has changed.</p> <p>This information is indicated.</p> <p>Please check page 6, line 4 - line 8 because text placement has changed.</p> <p>To clarify further, quality ratings of the ‘Critical Appraisal Checklist for Systematic Reviews and Research Syntheses’ (Joanna Briggs Institute; 2017) is scored qualitatively as, “Include”, “Exclude” or “Seek further info”. There is no quantified outcome for quality ratings. Only the agreement between the two reviewers for each study can be quantified (which is indicated), depending on the overall qualitative rating (i.e., include/exclude/seek further info) for each study</p>	<p>Page 6, line – 8</p> <p>Page 6, line 4 – line 8</p>
3	On page 7, line 24, how was Grounded Theory utilized to summarize findings from quantitative research studies?	<p>The Grounded Theory was not used. This information was already removed in the previous submission after clarifying as a ‘Thematic Narrative Analysis’. This was in response to an earlier request from a reviewer.</p> <p>Please check Page 7, line 16 – line 17 because text placement has changed.</p>	Page 7, line 16 – line 17

4	<p>If the authors wish to include the models of understanding response to crisis situations, then the discussion section should have indicated how the overall review further supported certain models and which models were not supported by the evidence.</p>	<p>We are very grateful for this observation and can note that the review, helpfully, supported all the models. To assist with location, this information is indicated as follows:</p> <ul style="list-style-type: none"> <li>• Self-Determination Theory (SDT): Page 17, line 8, page 18, line 12, page 19 line 7 – line 8 &amp; page 19, line 21</li> <li>• Pluralistic Trauma Theory (PTT): Page 17, line 8; page 18, line 6, page 18, line 12, page 19 line 7 – line 8, page 19, line 20</li> <li>• Job Demand-Control Support (JDCS): Page 18, line 1</li> <li>• Effort-Reward Imbalance model: Page 18, line 4 &amp; page 19, line 22</li> </ul>	<p>Page 17, line 7 – line 13 &amp; Page 18, line 1 – line 14 &amp; Page 19 line 4 – line 8 &amp; Page 19 line 20 – line 23</p>
---	---	--	--