

Central Lancashire Online Knowledge (CLoK)

Title	Recovery of DNA from bone without demineralization
Type	Article
URL	https://clock.uclan.ac.uk/44238/
DOI	https://doi.org/10.1016/j.fsigss.2022.09.031
Date	2022
Citation	Iyavoo, Sasitaran and Goodwin, William H (2022) Recovery of DNA from bone without demineralization. Forensic Science International: Genetics Supplement Series.
Creators	Iyavoo, Sasitaran and Goodwin, William H


It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1016/j.fsigss.2022.09.031>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Recovery of DNA from bone without
demineralization

Sasitaran Iyavoo, William H. Goodwin



PII: S1875-1768(22)00027-0

DOI: <https://doi.org/10.1016/j.fsigss.2022.09.031>

Reference: FSIGSS1888

To appear in: *Forensic Science International: Genetics Supplement Series*

Received date: 18 September 2022

Accepted date: 28 September 2022

Please cite this article as: Sasitaran Iyavoo and William H. Goodwin, Recovery of DNA from bone without demineralization, *Forensic Science International: Genetics Supplement Series*, (2022)
doi:<https://doi.org/10.1016/j.fsigss.2022.09.031>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier.

Recovery of DNA from bone without demineralization

Sasitaran Iyavoo^{a,b} and William H. Goodwin^a

^a School of Natural Sciences, University of Central Lancashire, Preston, United Kingdom.

^b School of Chemistry, College of Science, University of Lincoln, Lincolnshire, United Kingdom.

Corresponding author: WHGoodwin@uclan.ac.uk

Keywords: DNA extraction; bone; decalcification; demineralization; human identification; forensic

Abstract

This study assessed the performance of five different DNA extraction methods for the recovery of DNA from bone: ChargeSwitch[®] gDNA Plant Kit, DNA IQ[™] System Kit, DNeasy[®] Blood & Tissue Kit, PrepFiler[®] BTA Forensic DNA Extraction Kit and phenol-chloroform-isoamyl alcohol. DNA was extracted from pig rib and femur bones that was fresh, had undergone surface decomposition for three months, and had undergone surface decomposition for one year. Extracted DNA was analyzed using real-time PCR and amplification of an in-house PCR multiplex that assessed the quality and quantity of DNA and for the presence of inhibitors.

The phenol-chloroform-based method consistently yielded the highest amounts of DNA and DNA IQ the lowest; however, all methods produced relatively high yields of DNA from both pig rib and femur samples that could be amplified without any detected inhibition.

The data demonstrate that with reasonable quality bone samples any of the tested methods can isolate DNA that can be successfully analyzed. The effective use of internal PCR controls is also demonstrated.

1. Introduction

Bone samples represent one of the most challenging sample types for forensic genetics. The hard nature of bones necessitates extra steps compared to the cellular material and soft tissues, and additional challenges can present in the form of PCR inhibitors that can co-extract with DNA.

A body of work has demonstrated the advantages of demineralization when extracting DNA from bone samples that have been exposed to environmental insult, and often contain low amounts of highly degraded DNA [1]. However, the extra processes involved in demineralization add to the time and cost of processing samples, and in many cases, when working with samples that are in a relatively good condition, are unnecessary.

2. Materials and Methods

2.1 Samples and Preparation

Pig bone (femur and rib) were used in this study as a model for human bones. They were recovered from animals exposed to the environment for 0, 3 and 12 months at the University of Central

Lancashire's TRACES facility (Taphonomic Research in Anthropology: Centre for Experimental Studies) (UK).

Where relevant soft tissue/marrow and algae/bacteria on the bone was removed using scalpels, dental burrs, and abrasive paper; the bones were then soaked in 5% sodium hypochlorite for 15 min, rinsed with water and air-dried. Portions of each bone weighing between 1 g and 2 g were prepared and pulverized under liquid nitrogen.

2.2 DNA extraction

Samples were extracted using ChargeSwitch® gDNA Plant Kit (Thermo Fisher Scientific), DNA IQ™ System Kit (Promega), DNeasy® Blood & Tissue Kit (Qiagen), PrepFiler® BTA Forensic DNA Extraction Kit (Thermo Fisher Scientific) and a conventional phenol-chloroform method, according to the manufacturer's instructions and published methods [2]. Approximately 250 mg of pulverized bone was used for each sample. The lysis step was carried out on 50 mg and was repeated five times for each sample. Lysed samples were pooled and concentrated using an Amicon 30 KDa filtration unit (Merck Millipore) before DNA extraction. Extractions were carried out in triplicate and the final elution volume was set at 100 µl in each extraction.

2.3 DNA Quantification and Amplification

The extracted DNA samples were quantified with GoTaq® qPCR Master Mix (Promega) amplifying a 70 bp target using Applied Biosystems® 7500 Real-Time System. Primers used were designed to amplify both human and porcine DNA: F-5'-CCTCAAAGTCATGGGCAGC-3' and R: 5'-GACTCTCCAGGTCAGTAGG-3'. The standard DNA dilution series was made using control DNA 9947A (Thermo Fisher Scientific).

Reactions comprised 6.25 µl 2x GoTaq® qPCR Master Mix, 4.75 µl H₂O, 0.50 µl primers and 1.00 µl template DNA. The thermal cycler conditions were: 95 °C for 2 min, followed by 40 cycles of 95 °C for 15 s and 60 °C for 1 min.

The DNA concentration for each sample was estimated in ng/µl. All DNA extracts were amplified using an in-house multiplex (4-plex & Internal Amplification Controls) [3].

3. Results and discussion

A total of 30 extractions were undertaken for each of the three timepoints, with each condition tested in triplicate. DNA could be recovered from all pig bones with all methods, however, clear differences existed between the methods in terms of the quantity of DNA recovered (Figure 1). Phenol-chloroform-isoamyl alcohol extraction method consistently performed better than other extraction methods in terms of quantity of DNA extracted; this was followed by ChargeSwitch® gDNA Plant Kit, DNeasy® Blood & Tissue Kit, PrepFiler® BTA Forensic DNA Extraction Kit and the DNA IQ™ System Kit. The results for time zero were very similar to the 3-month time point shown below.




Figure 1: Boxplots showing the recovery of DNA extracted from rib and femur bones recovered from a) animals that had been exposed to the environment for 3 months, and b) animals that had been exposed to the environment for 1 year. 250 mg of starting material were used. CS.: ChargeSwitch, DNA: DNA IQ, DNe: DNeasy, PC: Phenol-chloroform, PF: PrepFiler. Each sample type and method were repeated three times.

Despite the varying amounts of DNA extracted, amplification was successful for all DNA extracts, with fragments amplified between 70 bp to 384 bp; no inhibition was detected in any of the extracts (Figure 2). Inhibition may present more of a problem when processing buried bone, which, depending on the characteristics of the soil, is more likely to accumulate PCR inhibitors.




Figure 2: Example of an electropherogram generated from extracted DNA of rib bone samples recovered from an animal that had been exposed to the environment for 1 year. The blue peaks correspond to fragments of 70 bp, 194 bp, 305 bp and 384 bp. In addition, two internal amplification controls (IACs) of 90 bp and 410 bp, were included and can be seen as red peaks – both IAC fragments have amplified which indicates that no PCR inhibitors are present.

4. Conclusions

The data presented illustrates that several methods are suitable for recovering DNA from bone that is not highly degraded. With such samples demineralization is not necessary, allowing quicker processing of samples in comparison to methods that incorporate demineralization or total demineralization. The samples used in this study were aged for up to one year and therefore further work is required to evaluate the suitability of not incorporating a demineralization step in samples that show higher levels of degradation.

Acknowledgments

This work was supported by a grant from the Public Service Department of Malaysia.

Conflict of interest statement

The authors declare no conflict of interests.

References

- [1] O.M. Loreille, T.M. Diegoli, J.A. Irwin, M.D. Coble, T.J. Parsons, High efficiency DNA extraction from bone by total demineralization, *Forensic Sci. Int. Genet.* 1(2) (2007) 191-195.
- [2] S. Iyavoo, S. Hadi, W. Goodwin, Evaluation of five DNA extraction systems for recovery of DNA from bone, *Forensic Sci. Int. Genet. Suppl. Ser.* 4(1) (2013) e174-e175.
- [3] M.S. Nazir, S. Iyavoo, S. Alimat, N. Zahra, S.H. Sanqoor, J.A. Smith, C. Moffatt, W. Goodwin, Development of a multiplex system to assess DNA persistence in taphonomic studies, *Electrophoresis* 34(24) (2013) 3352-3360.