



Article

Clinical effectiveness, cost effectiveness and service users' perceptions of early, well-resourced communication therapy following a stroke, a randomised controlled trial (The ACT NoW Study)

Bowen, A, Hesketh, A, Patchick, E, Young, A, Davies, L, Vail, A, Long, A, Watkins, Caroline Leigh, Wilkinson, M, Pearl, G, Tyrrell, P and Lambon, Ralph M

Available at <http://clock.uclan.ac.uk/4768/>

Bowen, A, Hesketh, A, Patchick, E, Young, A, Davies, L, Vail, A, Long, A, Watkins, Caroline Leigh ORCID: 0000-0002-9403-3772, Wilkinson, M et al (2012) Clinical effectiveness, cost effectiveness and service users' perceptions of early, well-resourced communication therapy following a stroke, a randomised controlled trial (The ACT NoW Study). Health Technology Assessment, 16 (26). pp. 1-159. ISSN 1366-5278

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.3310/hta16260>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

Clinical effectiveness, cost-effectiveness and service users' perceptions of early, well-resourced communication therapy following a stroke: a randomised controlled trial (the ACT NoW Study)

A Bowen,^{1*} A Hesketh,¹ E Patchick,¹ A Young,¹ L Davies,¹ A Vail,¹ A Long,² C Watkins,³ M Wilkinson,¹ G Pearl,⁴ M Lambon Ralph¹ and P Tyrrell^{1,5} on behalf of the ACT NoW investigators

¹University of Manchester MAHSC (Manchester Academic Health Science Centre), Manchester, UK

²University of Leeds, Leeds, UK

³University of Central Lancashire, Preston, UK

⁴Speakeasy, Bury, UK

⁵Salford Royal NHS Foundation Trust, Salford, UK


*Corresponding author



Executive summary

Health Technology Assessment 2012; Vol. 16: No. 26
DOI: 10.3310/hta16260

Health Technology Assessment
NIHR HTA programme
www.hta.ac.uk



Executive summary

Background

After a stroke, around one-third of people experience persisting problems with aphasia or dysarthria, limiting their ability to communicate through speech, writing or gesture. This restricts everyday activities and social participation, has adverse psychological effects and negatively impacts on families. Early, regular and intensive therapy may capitalise on brain plasticity, augmenting the natural recovery observed in the first few months.

The impact of speech and language (SL) therapy for this population is unknown owing to an absence of robust research evidence. In 2002, the UK's National Institute for Health Research (NIHR) identified the need to evaluate the clinical effectiveness and cost-effectiveness using therapy that was likely to be feasible for routine NHS delivery and comparing this with an attention control (AC), i.e. patient contact but not specific therapy.

Aim

The ACT NoW (Assessing the effectiveness of Communication Therapy in the North West) study was commissioned by the NIHR Health Technology Assessment programme and began with a feasibility study followed by a definitive study of the clinical effectiveness, cost-effectiveness and service user and carer views of an *early* well-resourced flexible communication intervention delivered by SL therapists, compared with an equivalent amount of contact but not therapy provided by employees badged as 'visitors'. This aimed to examine whether there is an added benefit of early SL therapy over and above a combination of natural recovery and receiving regular empathic attention.

Design

Extensive feasibility work was completed (2004–6), including the development of novel outcome measures. The definitive study (2006–10) consisted of a multicentre, parallel-group randomised controlled trial (RCT), economic evaluation and qualitative study. Randomisation was by an independent trials service. Primary analysis compared groups on 6-month outcomes using an intention-to-treat approach. Cost-effectiveness, cost-effectiveness acceptability analysis and net benefit analyses were used to relate costs and outcomes and explore the value for money of SL therapy, including a survey to determine societal preferences for waiting time and outcomes. The qualitative study used individual interviews with a subset of trial participants to explore service users' and carers' perceptions of process and outcome. Collaborative working partnerships with two groups were central to the design and conduct of this study: a research user group of service users ('the RUG') and a visionary group of NHS SL therapists willing to take on this challenging study.

Setting

The study was based in 12 NHS stroke services in England. Recruitment took place during the inpatient phase. Interventions were delivered across the stroke pathway (hospital and community).

Participants

The NHS SL therapists screened over 2000 people admitted with stroke with possible communication problems and considered one-fifth to be eligible for the trial on the grounds that they were likely to benefit from the research intervention. Exclusion criteria were pragmatic (out of area, unable to communicate in English) or clinical [therapy deemed unsuitable (e.g. pre-existing dementia, learning difficulties) for end-of-life care, serious health (including cognitive) problems, global communication problems or, alternatively, communication problems resolved or likely to without intervention].

One hundred and seventy people were randomly allocated to SL therapy (85) or AC (85), a consent rate of 44% of those invited. RUG-developed, aphasia-friendly consent materials proved useful in maximising trial participation. Participants ranged in age from 32 to 97 years (mean 70 years) and 56% were men. Almost all had aphasia (90%), 39% had dysarthria and 29% had both. Half were classified as having a severe activity restriction (disability) in terms of their baseline communication, suggesting that the sample included the full range from mild to severe. Half had dysphagia (impaired swallowing).

Some people declined follow-up, resulting in 81 and 72 cases, respectively, available for analysis in the SL and AC groups. The economic evaluation examined data from the trial participants in addition to a survey returned by 278 members of the public. Twenty-two trial participants and 10 carers took part in the qualitative study. Carers also participated in the RCT ($n=135$). They were typically female family members in the same household, not in paid employment and were younger than the stroke participants.

Intervention and control treatments

The SL therapy was a consensus-based, best practice, flexible intervention developed by NHS SL therapists for delivery in usual care settings, but better resourced. This allowed commencement as soon as clinically indicated and, if required, up to three contacts per week for up to 16 weeks, following participants across their stroke pathway. Adherence to the therapy manual was ensured through inspection of written records, observation of delivery and regular peer group meetings.

The study found that therapy was delivered as intended, flexibly tailored to individual need and on average started 2 weeks after stroke, involved 22 contacts (18 hours) delivered over 13 weeks, in both hospital and community settings. Half of the activities coded were direct intervention. Forty-two per cent of contacts were delivered by therapists of Band 7/8. Therapists sometimes felt that it was too early for an individual to engage intensively and 43% were referred for continuing NHS SL therapy after the study.

Attention control was offered at the same intensity by employed visitors who did not provide therapy or any communication strategies. Visitors had excellent social skills and general competency and were trained to deliver social attention absent of any intuitive form of communication therapy or strategy. They followed a manual allowing everyday activities (e.g. conversation, TV, music), but visits were mostly led by participants. Visitors were monitored to ensure adherence to the protocol.

Main outcome measures

The primary outcome was functional communicative ability 6 months post randomisation, rated on the activity subscale of the Therapy Outcome Measure activity subscale (TOM) by a blinded independent SL therapist. Secondary outcomes were participants' perceptions of their own functional communication and quality of life on the Communication Outcomes After Stroke scale (COAST); carers' perceptions of participants' functional communication using the first 15 questions on the Carer COAST; carers' own quality of life with the relevant five questions from the Carer COAST; and carer 'well-being' using the Carers of Older People in Europe (COPE) Index. Serious adverse events (SAEs) were recorded (deaths, further stroke, readmission to hospital). For the economic evaluation participants completed the European Quality of Life-5 Dimensions (EQ-5D) health status questionnaire and service use data were collected from medical records at 6 months and from carers. Preferences for different outcomes of communication therapy and willingness to wait for therapy were measured by a discrete choice survey completed by members of the public.

Results

External validity was good. Those who participated were similar in their measured characteristics to those who declined. Internal validity was also good. The control group had more disability at baseline but this was adjusted for in sensitivity analyses.

Speech and language therapy services struggled with staffing the intervention but overall succeeded in providing the intended (early but flexible) intervention at an average amount that was higher than most NHS services. Most importantly, they provided continuity by following participants across the stroke pathway, whereas in usual care they would be placed on waiting lists following transfer to the community. There was high uptake of both therapy and control visits by service users. This was slightly lower for the latter, reflecting patient choice; however, an adequate control was provided. An observational comparison of TOM activity scores at baseline and 6 months suggested a clinically meaningful level of improvement in functional communication of 0.8 points [95% confidence interval (CI) 0.6 to 1.0 points] regardless of group allocation.

Primary analysis estimated a difference of 0.25 (95% CI -0.19 to 0.69) points on the primary outcome (TOM) in favour of SL therapy. Sensitivity analyses suggested that this estimated difference was due to the imbalance in baseline severity and the imputation of values for deaths. Per-protocol analyses rejected a possible dilution of therapy from the 18 control participants who rejected their allocation and received some NHS SL therapy. These findings appear robust and exclude the possibility of a clinically significant difference of 0.5 on the TOM in either direction. There was also no evidence of an added benefit of SL therapy on any of the secondary outcomes, including patient- and carer-reported measures. There was no statistically significant difference in SAEs between the groups [odds ratio of 0.42 (95% CI 0.16 to 1.1)]. Although SAEs were less frequent in the therapy group, they were rare (15 vs 7). Subgroup analyses (by aphasia

vs dysarthria or by level of severity of communication impairment) produced no evidence of a differential subgroup effect.

The likely cost-effectiveness of therapy was at the upper end of the acceptable willingness-to-pay thresholds of the National Institute for Health and Clinical Excellence. However, there were several limitations to the economic evaluation and primary and sensitivity analyses indicated a high level of uncertainty, suggesting that it is not possible to conclude whether therapy is more or less cost-effective than AC.

The qualitative study found that, regardless of whether they saw a visitor or an SL therapist, participants highly regarded this experience, which made a positive impact on their lives. The amount of contact and the interpersonal skills/personal qualities of the person providing it (visitor or SL therapist) were identified as important drivers for recovery that built confidence and developed a positive mood. Users believed that an important mechanism for recovery of communicative ability and growing awareness of residual disability was repeated practice of everyday communicative activities with a professional who showed empathy and interest in their individual needs. Carers expressed strongly positive views about the support that survivors had received, whether from the visitor or SL therapist. They identified the importance not just of regular contact, but of that contact coming from someone outside of the survivor's family/social milieu. The outsider provided interaction and communication opportunities that challenged the person they cared for in ways they could not because it forced an engagement with the unfamiliar. Carers did not identify any primary benefit to themselves, but rather secondary benefits from seeing the survivor make progress.

Conclusions

The ACT NoW study provides robust evidence and a definitive answer to a clinically important question, with good generalisability of its findings. Functional communicative ability at 6 months had improved by a clinically meaningful amount for people in both groups. However, there was no evidence of an added benefit of early communication therapy from SL therapists for people with communication disability or their carers over and above that from AC and natural recovery, when both were provided at a higher level than in typical standard practice. Therefore, there is no evidence to recommend enhancing the provision of early communication therapy by a qualified SL therapist over and above usual care, and the evidence suggests that the latter should be reorganised.

To determine whether the benefits were due to therapy rather than time with a therapist, both groups were offered early and well-resourced time. Service users valued the early and frequent contact from professional visitors/therapists outside of their family and friends who showed interest in their individual needs. These impacted positively on their confidence and mood, providing repeated practice of everyday communication. In terms of clinical implications, the study did not evaluate the early role of SL therapists after stroke. Much of SL therapists' early workload is for the assessment and treatment of dysphagia and in usual practice communication and swallowing are managed together. We evaluated one aspect (communication therapy), and only after a diagnosis had been made and provided to the user, family and multidisciplinary team (MDT), thereby precluding conclusions about the value of diagnosis by an SL therapist.

Less definitive were the results on adverse events and the economic evaluation. There remains the possibility that therapy reduced the rate of deaths, further strokes and rehospitalisation and this may have warranted further research had a mechanism been established (i.e. had therapy improved communication), which it was not. Early enhanced SL therapy for communication is

likely to be cost-effective only if decision-makers are prepared to pay \geq £25,000 to gain one unit of utility. However, uncertainty over cost-effectiveness is of no practical concern given the lack of evidence of clinical effectiveness.

There may be divided opinion over whether AC was the most suitable choice for this study. Uncertainty over the relative contribution of natural recovery versus the early, regular attention provided by therapists or visitors would have been eliminated by a no-contact control. There were two good reasons for not commissioning this. First, if therapy proved more effective (as most people expected) it would remain unknown whether the active mechanism was the therapy or the psychosocial effect of providing attention. Second, it is highly unlikely that a no-contact control trial would have been feasible given ethical concerns and lack of equipoise for clinicians and potential participants.

An alternative control would have been usual care. Assuming an effect along a continuum for therapy per se as opposed to attention, with usual care theoretically in between the two ACT NoW groups, our finding of no difference between the two extremes would be replicated when comparing the extreme and midpoint.

Recommendations for research

Research should assess a reorganised SL therapy communication service that uses a stepped care model of intervention considering skill mix and timing. SL therapists' early role could be around diagnosis, communicating this to the user and MDT and supervising assistants for regular visits similar to those provided for the AC group in ACT NoW. Intervention would later step up to direct SL therapist contact for those with persisting need and include the part of the stroke pathway referred to as Life After Stroke. Usual care by NHS SL therapists would be an appropriate comparator.

Further research should investigate whether the ACT NoW SL therapy was delivered too soon in the stroke pathway, by evaluating its effectiveness with a chronic clinical population, those with persisting communication problems months and years post stroke. There is huge unmet need in this population yet considerable uncertainty about service delivery and an understandable tendency to assign scarce resources to those in the first 6 months of recovery. A future study would challenge the unlikely but sometimes cited suggestion that recovery is only possible in the short term.

Specific promising interventions should be subjected to RCTs, for example conversation partner training. Therapy for people with dysarthria is a neglected area of stroke rehabilitation research, warranting investigation. Aphasia research must be generalisable to the target population, including people without English as their first language.

Future economic evaluation needs to find ways of capturing valid baseline EQ-5D data from acutely ill and communication-impaired participants and of ensuring sufficient resources to chase missing data from incomplete or inaccessible NHS records.

Trial registration

This trial is registered as ISRCTN78617680.

Funding

This project was funded by the NIHR Health Technology Assessment programme and will be published in full in *Health Technology Assessment*; Vol. 16, No. 26. See the HTA programme website for further project information. The Stroke Association funded part of the excess treatment costs.

Publication

Bowen A, Hesketh A, Patchick E, Young A, Davies L, Vail A, *et al.* Clinical effectiveness, cost-effectiveness and service users' perceptions of early, well-resourced communication therapy following a stroke: a randomised controlled trial (the ACT NoW Study). *Health Technol Assess* 2012;**16**(26).

NIHR Health Technology Assessment programme

The Health Technology Assessment (HTA) programme, part of the National Institute for Health Research (NIHR), was set up in 1993. It produces high-quality research information on the effectiveness, costs and broader impact of health technologies for those who use, manage and provide care in the NHS. 'Health technologies' are broadly defined as all interventions used to promote health, prevent and treat disease, and improve rehabilitation and long-term care.

The research findings from the HTA programme directly influence decision-making bodies such as the National Institute for Health and Clinical Excellence (NICE) and the National Screening Committee (NSC). HTA findings also help to improve the quality of clinical practice in the NHS indirectly in that they form a key component of the 'National Knowledge Service'.

The HTA programme is needs led in that it fills gaps in the evidence needed by the NHS. There are three routes to the start of projects.

First is the commissioned route. Suggestions for research are actively sought from people working in the NHS, from the public and consumer groups and from professional bodies such as royal colleges and NHS trusts. These suggestions are carefully prioritised by panels of independent experts (including NHS service users). The HTA programme then commissions the research by competitive tender.

Second, the HTA programme provides grants for clinical trials for researchers who identify research questions. These are assessed for importance to patients and the NHS, and scientific rigour.

Third, through its Technology Assessment Report (TAR) call-off contract, the HTA programme commissions bespoke reports, principally for NICE, but also for other policy-makers. TARs bring together evidence on the value of specific technologies.

Some HTA research projects, including TARs, may take only months, others need several years. They can cost from as little as £40,000 to over £1 million, and may involve synthesising existing evidence, undertaking a trial, or other research collecting new data to answer a research problem.

The final reports from HTA projects are peer reviewed by a number of independent expert referees before publication in the widely read journal series *Health Technology Assessment*.

Criteria for inclusion in the HTA journal series

Reports are published in the HTA journal series if (1) they have resulted from work for the HTA programme, and (2) they are of a sufficiently high scientific quality as assessed by the referees and editors.

Reviews in *Health Technology Assessment* are termed 'systematic' when the account of the search, appraisal and synthesis methods (to minimise biases and random errors) would, in theory, permit the replication of the review by others.

The research reported in this issue of the journal was commissioned by the HTA programme as project number 02/11/04. The contractual start date was in October 2004. The draft report began editorial review in January 2011 and was accepted for publication in May 2011. As the funder, by devising a commissioning brief, the HTA programme specified the research question and study design. The authors have been wholly responsible for all data collection, analysis and interpretation, and for writing up their work. The HTA editors and publisher have tried to ensure the accuracy of the authors' report and would like to thank the referees for their constructive comments on the draft document. However, they do not accept liability for damages or losses arising from material published in this report.

The views expressed in this publication are those of the authors and not necessarily those of the HTA programme or the Department of Health.

Editor-in-Chief: Professor Tom Walley CBE
Series Editors: Dr Martin Ashton-Key, Professor Aileen Clarke, Dr Tom Marshall, Professor John Powell, Dr Rob Riemsma and Professor Ken Stein
Associate Editor: Dr Peter Davidson
Editorial Contact: edit@southampton.ac.uk

ISSN 1366-5278 (Print)

ISSN 2046-4924 (Online)

ISSN 2046-4932 (DVD)

© Queen's Printer and Controller of HMSO 2012. This work was produced by Bowen *et al.* under the terms of a commissioning contract issued by the Secretary of State for Health.

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics (COPE) (<http://www.publicationethics.org/>).

This journal may be freely reproduced for the purposes of private research and study and may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NETSCC, Health Technology Assessment, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.

Published by Prepress Projects Ltd, Perth, Scotland (www.prepress-projects.co.uk), on behalf of NETSCC, HTA.

Printed on acid-free paper in the UK by Charlesworth Press.