

Central Lancashire Online Knowledge (CLoK)

Title	Are male soccer players accumulating sufficient load across varying microcycle structures? Examining the load, wellness and training/match ratios of a professional team
Type	Article
URL	https://clock.uclan.ac.uk/49644/
DOI	##doi##
Date	2023
Citation	Oliveira, Rafael, Canário-Lemos, Rui, Morgans, Ryland, Peixoto, Rafael, Vilaça-Alves, José and Brito, João Paulo (2023) Are male soccer players accumulating sufficient load across varying microcycle structures? Examining the load, wellness and training/match ratios of a professional team. <i>Journal of Men's Health</i> , 19 (9). pp. 26-35. ISSN 1875-6867
Creators	Oliveira, Rafael, Canário-Lemos, Rui, Morgans, Ryland, Peixoto, Rafael, Vilaça-Alves, José and Brito, João Paulo

It is advisable to refer to the publisher's version if you intend to cite from the work. ##doi##

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

ORIGINAL RESEARCH

Are male soccer players accumulating sufficient load across varying microcycle structures? Examining the load, wellness and training/match ratios of a professional team

Rafael Oliveira^{1,2,3,*}, Rui Canário-Lemos^{4,5}, Ryland Morgans⁶, Rafael Peixoto^{4,5}, José Vilaça-Alves^{3,4,5}, João Paulo Brito^{1,2,3}

¹Sports Science School of Rio Maior, Polytechnic Institute of Santarém, 2040-413 Rio Maior, Portugal

²Life Quality Research Centre, 2040-413 Rio Maior, Portugal

³Research Center in Sport Sciences, Health Sciences and Human Development, 5001-801 Vila Real, Portugal

⁴University of Trás-os-Montes e Alto Douro, 5001-801 Vila Real, Portugal

⁵Research Group in Strength Training and Fitness Activities, GEETFAA, 5001-801 Vila Real, Portugal

⁶Football Performance Hub, University of Central Lancashire, PR1 2HE Preston, UK

***Correspondence**

rafaeloliveira@esdrm.ipsantarem.pt
(Rafael Oliveira)

Abstract

Professional soccer involves varying numbers of training sessions and matches each week, which can influence load distribution. Understanding the exact distribution may allow appropriate load periodisation and planning for players. Thus, this study aimed to (i) compare accumulated load and wellness between weeks with different numbers of training sessions and (ii) compare training/match ratio (TMr) of external and internal load between weeks with different numbers of training sessions. Ten players with a minimum of 45 minutes of weekly match-play were analysed over 16 weeks. The microcycle structures consisted of three (3dW), four (4dW), five (5dW) and six (6dW) training sessions plus match-day per week. The following measures were used for analysis: duration, fatigue, quality of sleep, muscle soreness, stress, mood, rating of perceived exertion (RPE), session-RPE (s-RPE), high-speed running distance (HSR), sprint distance (SPD), number of accelerations (ACC) and decelerations (DEC). Accumulated wellness/load were calculated by adding all training and match sessions, while TMr was calculated by dividing accumulated load by match data. The main results showed that accumulated wellness and load were significantly different, with moderate to very large effect sizes, except regarding mood, duration, s-RPE, SPD during 5dW vs. 6dW and s-RPE, HSR, SPD, ACC and DEC during 3dW vs. 4dW (all $p > 0.05$). Moreover, 6dW was significantly higher than 4dW regarding TMr of duration ($p < 0.05$, moderate effect size), RPE, HSR and SPD (all $p < 0.05$ with very large effect sizes) and for 3dW of HSR and ACC ($p < 0.05$ with very large effect sizes). This study showed that 5dW and 6dW had higher training measures than 3dW or 4dW. Additionally, higher wellness was presented in the microcycles with higher training frequencies. These findings suggest that physical load and wellness were not adjusted according to the number of training sessions within a microcycle.

Keywords

External load; Fatigue; Internal load; Mood; Load quantification; Muscle soreness; Sleep; Stress; Training load; Well-being

1. Introduction

Quantifying wellness, training and match load/demands in soccer players is a common practice [1–3]. Specifically, the monitorisation of athletes include quantifying training/match demands (e.g., locomotor/mechanical and psychophysiological) and the wellness and readiness of players [4]. On the one hand, wellness is usually measured by questionnaires, as previously proposed by Hooper and Mackinnon [5] using the Hooper Index or by McLean *et al.* [6]. While the Hooper Index includes fatigue, quality of sleep, muscle soreness and stress, measured *via* a seven-point scale [6], the wellness questionnaire by McLean *et al.* [6] includes the same subjective items

plus mood status, measured on a five-point scale. Regardless of the questionnaire, wellness variables depend on the load applied [4]. Meanwhile, locomotor/mechanical demands or physical demands are associated with external load/intensity monitoring using global positioning system (GPS) variables (e.g., distances covered at various running speeds or accelerations), whereas psychophysiological demands are associated with internal load/intensity monitoring using subjective or objective measures such as rating of perceived exertion (RPE) and heart rate [2, 7].

Internal and external training stimuli may vary for players according to the number of training/match sessions per week/microcycle, the aims/objectives of training sessions and

the periodisation/planning strategies of the coach [8]. For instance, a recent systematic review reporting external and internal load in professional soccer revealed that three to six training sessions were performed per week [9], highlighting that different strategies were applied. Other factors that impact load include the use of specific drills and games during training (e.g., small-sided games, long sprints, repeated sprints, interval training and medium- to large-sized games) [10]. Regarding match-play, load quantification can vary due to the dynamics of matches and their contextual factors [11].

Previous studies have suggested that the number of training sessions affects the training load [12, 13]. Oliveira *et al.* [13] showed similar average values for running distance measures and RPE/session-RPE across five different weeks/microcycles with four training sessions and one, two or three matches. Meanwhile, Anderson *et al.* [12] showed that microcycles with only one match accumulated a total distance of ~14,000 meters during training sessions and ~11,000 meters during matches. Moreover, Oliveira *et al.* [13] reported a range for high-speed running distance (HSR, 19–24 km/h) between 17–398 meters during training sessions, while distances of 465–681 meters were recorded during matches. Furthermore, Anderson *et al.* [12] also reported a range for HSR (19.8–25 km/h) of 8–104 meters during training sessions, while distances of 682–727 meters were covered during match-play.

A method to improve the understanding of the physical demands induced by training and competitive match-play is to calculate the training/match ratio (TMr) [8, 14]. This ratio is calculated by dividing the accumulated weekly load by the match load [8]. If the TMr provides a value lower than one, it suggests that the accumulated load of the week is lower than the match load; a value above one suggests that the training load is greater than the match load [8].

Despite the practical implication of such analyses, only four studies were found to make such an investigation [8, 14–16]. Specifically, one study [8] described the TMr of different external load measures—total distance, running distance, HSR, sprint distance (SPD), player load, number of high accelerations (ACC), and number of high decelerations (DEC)—during a full professional soccer season while analysing the variations between varying types of weeks (three, four and five training sessions/week). The study showed that weeks with five training sessions had higher values for all external load ratios than weeks with three or four training sessions. Additionally, HSR distance and SPD measures presented substantially lower ratios than other variables such as total running distance, ACC, DEC and player load [8].

Another study compared the loads of a professional soccer team among training days and matches and between starters and non-starters using TMr. Specifically, in weeks with four training sessions, the weekly training load represented a load equal to 4.4 matches, high accelerations represented a load of 3.9 matches, high decelerations represented a load of 3.3 matches, and HSR represented 2.1 matches [14]. Furthermore, Modric *et al.* [15] analysed the relationship between TMr and match outcomes (wins, draws and losses). However, whereas previous studies [8, 14] used accumulated weekly data divided by match data, Modric *et al.* [15] divided the data for each training session by match data. Finally, Szigeti *et al.* [16] also

used TMr to analyse external load in under-17 soccer players and their main findings highlighted that ACC represented 2.84 of match load, while HSR represented 0.95 of a match in weeks with three training sessions.

None of the previous studies mentioned above included wellness and internal load. Moreover, those that included external load did not provide full details of participant inclusion for TMr calculations [8] and, contrastingly, included an additional session to replace the official match for non-starters [14]. Therefore, the present study aimed to (i) compare accumulated external and internal load and wellness between weeks with different numbers of training sessions and (ii) compare the TMr of external and internal load between weeks with different numbers of training sessions. According to the previous study of Clemente *et al.* [8], who observed similar load distributions across the week, it is speculated that weeks with more training sessions may contribute to higher TMr values. Thus, it was hypothesised that weeks with fewer training sessions are associated with lower accumulated load and TMr values.

2. Materials and methods

2.1 Design

In this observational study, soccer players were monitored daily for wellness measures (sleep quality, muscle soreness, fatigue, stress and mood), internal and external load. The study lasted 16 weeks from the 2022/23 in-season period (July to November) and comprised 70 training sessions and 15 official matches.

Following the same procedures used in a similar study, all weeks with one official match and three or more training sessions were included in this analysis [8]. This decision was made to reduce the variability among comparisons. Thus, only 15 weeks were included in the analysis, as one microcycle only included two training sessions. The week types were classified based on the number of training sessions: weeks with three training sessions (3dW, $n = 3$), weeks with four training sessions (4dW, $n = 3$), weeks with five training sessions (5dW, $n = 4$), and weeks with six training sessions (6dW, $n = 5$).

2.2 Participants

Ten professional male soccer players (age: 25.30 ± 2.11 years; body mass: 71.90 ± 6.23 kg; body height: 179.4 ± 6.88 cm; fat mass: $8.69 \pm 0.95\%$; professional experience: 7.5 ± 2.07 years) participated in the current study [17–20]. Players belonged to a European soccer team that played in the national first division. Of the players included, four were defenders, three were midfielders, and three were attackers.

The eligibility criteria for participant inclusion were as follows: (i) participating in 80% of all training sessions (for the full session duration) [21], (ii) completing wellness and training reports over the data collection period and (iii) participating in a minimum of 45 minutes of the weekly match [8].

Prior to data collection, the club, coaches and participants were fully informed of the study design and signed an informed consent form.

2.3 Wellness quantification

The wellness questionnaire developed by McLean *et al.* [6] was applied individually 30 minutes before each training/match session through a Google form specifically designed. The questionnaire uses a scale of 1–5 arbitrary units (A.U.) and contains five questions about fatigue, quality of sleep, muscle soreness, stress and mood (5 = very fresh, very restful, very great, very relaxed and very positive mood, respectively; 1 = always tired, insomnia, very sore, highly stressed and highly annoyed/irritable/down, respectively). All players were already familiar with the questionnaire from the previous season.

2.4 Internal load quantification

The CR-10 Borg's scale [22] was used to monitor the players' rating of perceived exertion (RPE). Following the usual training procedures, 20–30 minutes after each session, every player provided a perceived exertion value using a Google form by answering the following question: "How intense was the training session?" The scale varied from 0 to 10 A.U. (0 = nothing to all, 0.5 = extremely weak, 1 = very weak, 2 = weak, 3 = moderate, 4 = somewhat strong, 5 = strong, 7 = very strong, and 10 = extremely strong).

RPE was used to measure internal perception of effort. In addition, the duration of the entire training session or match, in minutes, was multiplied by the RPE to generate the session-RPE (s-RPE), measured in A.U. [23, 24]. All players were already familiar with the questionnaire from the previous season.

2.5 External load quantification

Locomotor demands were measured using a GPS Vector S7 (Catapult Innovations, Melbourne, Australia). The same unit was used for each player throughout the analysis period to avoid inter-unit bias. The unit was placed on the upper back of each player 30 minutes before each session (training and match) and removed immediately after the session.

The GPS Vector S7, sampling at 10 Hz, was used to monitor the locomotor demands of players during all sessions. This device was previously validated for accuracy and reliability regarding various measures, such as distance, velocity and average acceleration [25]. The following measures were used for analysis: (i) high-speed running distance (HSR, 20–25 km/h), sprint distance (SPD, >25 km/h) [26], number of accelerations (ACC, >2 m/s²) and number of decelerations (DEC, <2 m/s²) [27].

2.6 Accumulated wellness/load and training/match ratio

Accumulated wellness/load consisted of the sum of each measure during all training sessions of the microcycle and was calculated per player, thus providing the weekly load for each measure (match included) [28–33].

Moreover, accumulated load was calculated without match data to determine the TMr for all internal and external measures. Ratios were then calculated by dividing accumulated load (without match data) by match data (TMr = weekly

load/match demands) [8, 14]. Consequently, the following measures were obtained: RPE ratio (RPEr); session RPEr (s-RPEr); high-speed running distance ratio (HSRr); sprint distance ratio (SPDr); accelerations ratio (ACCr) and decelerations ratio (DECr). The same ratio was calculated for session duration (Dr) by dividing accumulated duration (without match data) by the match duration. All TMr calculations of load and duration measures provided clear descriptions of the microcycle structures applied.

2.7 Statistical analysis

Descriptive statistics are presented as mean \pm standard deviation. The normality of the different variables was analysed (and not confirmed) using the Shapiro-Wilk test. Thus, Friedman's test was used to compare the different week types, while the Wilcoxon test was used for pairwise comparisons. Significant results were considered at $p < 0.05$.

When a significant result was detected, Hedges' effect size was calculated to determine the effect magnitude based on the difference between two means divided by the standard deviation according to the data. The results were categorised based on the following criteria: <0.2 = trivial effect, 0.2–0.6 = small effect, 0.6–1.2 = moderate effect, 1.2–2.0 = large effect, and >2.0 = very large effect [34].

All statistical procedures were executed in IBM SPSS Statistics for Windows (version 23.0, IBM Corp, Armonk, NY, USA).

3. Results

Following the Friedman test, all variables showed $p < 0.001$, except for SPD ($p = 0.002$). Table 1 presents the pairwise comparisons for accumulated training demands (match data included) and wellness for all variables.

The Friedman test was also applied for TMr; all variables showed $p < 0.001$ except for Dr ($p = 0.004$), s-RPE ($p = 0.013$) and SPDr ($p = 0.001$). Fig. 1 represents the mean match data, accumulated weekly data and TMr considering the different weeks' schedules. Table 2 represents the pairwise comparisons for all TMr.

4. Discussion

This study aimed to (i) compare accumulated load and wellness between weeks with different numbers of training sessions and (ii) compare the TMr values of external and internal load between weeks with a different number of training sessions. The hypothesis that weeks with fewer training sessions would present lower accumulated load and TMr was confirmed. Specifically, the main results of the study show that accumulated wellness and load demands were higher in the weeks with the most training sessions and progressively decreased in weeks with fewer training sessions (6dW > 5dW > 4dW > 3dW), with moderate to very large effect sizes. Although the results were insignificant, TMr showed the same tendency (see Fig. 1 and Table 2), while the main findings reported the highest values for 6dW compared to 4dW for RPEr, Dr, HSRr and SPDr and compared to 3dW for HSRr

TABLE 1. Comparisons of different microcycles for accumulated wellness and load demands.

Variable	3dW	4dW	5dW	6dW	<i>p</i> -value	Effect size
Quality of sleep (A.U.)	16.60 ± 1.77 ^{a,b,c}	18.76 ± 1.50 ^{b,c}	23.05 ± 1.44 ^c	26.20 ± 2.88	3dW vs. 4dW: <0.001	3dW vs. 4dW: 0.83
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 2.47
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 3.67
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 2.79
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 2.85
					5dW vs. 6dW: 0.002	5dW vs. 6dW: 1.21
Fatigue (A.U.)	13.28 ± 2.31 ^{a,b,c}	16.85 ± 2.42 ^{b,c}	20.29 ± 2.80 ^c	23.61 ± 3.14	3dW vs. 4dW: <0.001	3dW vs. 4dW: 1.43
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 2.65
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 3.53
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 1.21
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 2.20
					5dW vs. 6dW: <0.001	5dW vs. 6dW: 1.06
Muscle Soreness (A.U.)	13.13 ± 2.31 ^{a,b,c}	16.87 ± 2.65 ^{b,c}	19.73 ± 3.24 ^c	22.55 ± 3.15	3dW vs. 4dW: <0.001	3dW vs. 4dW: 1.44
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 2.89
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 3.39
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 1.54
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 2.14
					5dW vs. 6dW: 0.018	5dW vs. 6dW: 0.74
Stress (A.U.)	14.17 ± 2.02 ^{a,b,c}	17.72 ± 2.51 ^{b,c}	21.60 ± 3.15 ^c	24.0 ± 3.98	3dW vs. 4dW: <0.001	3dW vs. 4dW: 1.48
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 2.61
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 2.85
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 1.28
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 1.72
					5dW vs. 6dW: 0.046	5dW vs. 6dW: 0.62
Mood (A.U.)	14.00 ± 2.00 ^{a,b,c}	17.35 ± 2.28 ^{b,c}	21.71 ± 2.78	24.29 ± 3.69	3dW vs. 4dW: <0.001	3dW vs. 4dW: 1.49
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 2.99
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 3.19
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 1.62
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 2.06
					5dW vs. 6dW: 0.056	5dW vs. 6dW: 0.73
RPE (A.U.)	18.77 ± 3.07 ^{a,b,c}	23.70 ± 2.79 ^{b,c}	31.71 ± 3.33 ^c	36.98 ± 3.12	3dW vs. 4dW: 0.042	3dW vs. 4dW: 1.66
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 3.87
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 5.70
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 2.45
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 4.28
					5dW vs. 6dW: <0.001	5dW vs. 6dW: 1.63
Duration (min)	154.78 ± 24.55 ^{a,b,c}	208.50 ± 26.42 ^{b,c}	273.94 ± 23.71	294.51 ± 26.05	3dW vs. 4dW: 0.004	3dW vs. 4dW: 2.03
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 4.99
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 5.36
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 2.74
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 3.29
					5dW vs. 6dW: 0.147	5dW vs. 6dW: 0.79
s-RPE (A.U.)	969.42 ± 251.08 ^{b,c}	1256.52 ± 253.94 ^{b,c}	1721.61 ± 318.76	1868.51 ± 266.19	3dW vs. 4dW: 0.144	3dW vs. 4dW: 1.13
					3dW vs. 5dW: 0.001	3dW vs. 5dW: 2.36
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 3.38
					4dW vs. 5dW: 0.004	4dW vs. 5dW: 1.46
					4dW vs. 6dW: 0.002	4dW vs. 6dW: 2.30
					5dW vs. 6dW: 0.419	5dW vs. 6dW: 0.55

TABLE 1. Continued.

Variable	3dW	4dW	5dW	6dW	p-value	Effect size
HSR (m)	535.12 ± 165.88 ^{b,c}	744.26 ± 248.34 ^{b,c}	946.03 ± 190.41 ^c	1142.07 ± 297.13	3dW vs. 4dW: 0.360	3dW vs. 4dW: 0.84
					3dW vs. 5dW: 0.002	3dW vs. 5dW: 2.16
					3dW vs. 6dW: 0.002	3dW vs. 6dW: 2.05
					4dW vs. 5dW: 0.006	4dW vs. 5dW: 1.05
					4dW vs. 6dW: 0.001	4dW vs. 6dW: 1.34
Sprint Distance (m)	124.05 ± 49.75 ^{b,c}	172.02 ± 92.84 ^c	238.41 ± 58.09	274.6 ± 77.26	5dW vs. 6dW: 0.023	5dW vs. 6dW: 0.66
					3dW vs. 4dW: >0.999	3dW vs. 4dW: 0.53
					3dW vs. 5dW: 0.008	3dW vs. 5dW: 1.98
					3dW vs. 6dW: 0.013	3dW vs. 6dW: 1.98
					4dW vs. 5dW: 0.061	4dW vs. 5dW: 1.10
Accelerations (nr)	150.27 ± 32.82 ^{b,c}	195.08 ± 29.24 ^{b,c}	261.25 ± 3.10 ^c	305.12 ± 41.93	4dW vs. 6dW: 0.001	4dW vs. 6dW: 1.31
					3dW vs. 4dW: 0.093	3dW vs. 4dW: 1.51
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 3.17
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 3.72
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 9.61
Decelerator (nr)	141.42 ± 32.85 ^{b,c}	183.57 ± 24.88 ^{b,c}	246.48 ± 29.16 ^c	282.21 ± 33.54	4dW vs. 6dW: <0.001	4dW vs. 6dW: 2.64
					3dW vs. 4dW: 0.092	3dW vs. 4dW: 1.51
					3dW vs. 5dW: <0.001	3dW vs. 5dW: 3.55
					3dW vs. 6dW: <0.001	3dW vs. 6dW: 4.18
					4dW vs. 5dW: <0.001	4dW vs. 5dW: 2.16
					4dW vs. 6dW: <0.001	4dW vs. 6dW: 2.96
					5dW vs. 6dW: 0.007	5dW vs. 6dW: 1.06
					5dW vs. 6dW: 0.025	5dW vs. 6dW: 1.07

RPE: rating of perceived exertion using the CR-10 Borg's scale; s-RPE: multiplication of time of session by the score of RPE; A.U.: arbitrary units; m: meters; min: minutes; HSR: high speed running (20–25 km/h); a denotes significant difference from 4dW; b, denotes significant difference from 5dW, c, denotes significant difference from 6dW (all, $p < 0.05$); bold: significant results.

and ACCr.

Regarding the first aim, wellness was higher in weeks with more training sessions considering that in the questionnaire applied (based on a scale ranging from 0–5), the value of 5 A.U. suggested very fresh, very restful, very great, very relaxed and very positive mood. Similarly, a previous study examining youth soccer players revealed that higher external and internal intensities were associated with improved sleep (quality and quantity) and feeling rested [35]. Contrastingly, another study on youth soccer players showed that high-intensity training did not impact the following night's sleep quality [36]. A further study on professional soccer players reported that sleep quality was not impacted by higher-intensity sessions (match included) [37]. However, these studies did not have identical designs, and more research is needed to confirm the results of the present study.

Regarding the second aim of TMr analysis, Clemente *et al.* [8] observed professional soccer players and showed that ACC and DEC ($>3 \text{ m/s}^2$) presented values of 2.2 ± 1.8 and 1.6 ± 0.9 , respectively, during 3dW which then reached 4.1 ± 1.6 and 3.4 ± 1.9 , respectively, during 5dW. Moreover, the TMr of HSR was 1.1 ± 0.8 during 3dW and 2.3 ± 1.5 during 5dW. However, when applying a different approach in professional soccer players and considering the comparison of starters versus non-starters in microcycles with four training

sessions and a match, Stevens *et al.* [14] reported the following values for starters and non-starters, respectively: HSR, 2.1/1.5; medium ACC, 3.1/2.6; high ACC, 3.9/3.6; medium DEC, 3.4/3.0; high DEC 3.3/2.7.

Recently, Szigeti *et al.* [16] found that 3dW showed 2.84 of ACCr and 0.95 of HSRr, while other measures such as SPDr and DEC presented values between 1 and 2 A.U. in under-17 soccer players. The values reported in these previous studies are much higher than those presented in the current study regardless of the number of training sessions per week, which may be associated with the different periodisation/planning practices of the coaches or the possibility of supplementary sessions. Additionally, to our knowledge, the present work is the first study in which TMr was calculated using both RPE and s-RPE, and thus, no comparison can be made based on the ratios.

Moreover, some TMr values were lower than 1.0 A.U., which suggests that the accumulated training load was lower than the match load. Such values occurred in the weeks with three and four training sessions for the measures of s-RPEr, HSRr, and SPDr. The same scenario occurred for ACCr and DEC (but only during 3dW). Nevertheless, in the present study, higher TMr values for ACC and DEC were found regardless of the number of training sessions, which suggests that more small-sided games were performed during training,




FIGURE 1. Accumulative weekly training load, match load and training/match ratios. c denotes significant difference from 6dW ($p < 0.05$). RPE: rating of perceived exertion; A.U.: arbitrary units; TMR: training/match ratio; HSR: high speed running; SPD: sprint distance; ACC: number of accelerations; DEC: decelerations ratio.

as this type of training increases the number of ACC/DEC and decreases HSR and SPD covered [38, 39]. Although this variable was not controlled in the present study, coaches must manage the number of small-sided games performed during different training weeks since they can induce higher TMR values for ACC and DEC. Moreover, additional training should be performed in weeks with low HSR and SPD covered.

Despite its contributions, the present study had some limitations. The main limitation is the small sample size derived from only one team and a restricted period of 16 weeks. Moreover, playing position differences were not considered even

though some positions, such as wingers and wide defenders, generate higher accumulated values, TMR or perceived different wellness statuses than other positions since these positions require greater effort and more running than other positions [29]. Moreover, playing status (starters versus non-starters) was not considered due to the restricted inclusion criteria of the present study. It would be worthwhile to investigate whether non-starters are completing enough training to participate in matches. Consequently, future studies should analyse larger sample sizes, consider different playing statuses, and analyse regular weeks with one match versus congested periods with

TABLE 2. Comparisons of the different microcycles for TMr.

Variable	3dW	4dW	5dW	6dW	p-value	Effect size
RPEr (A.U.)	1.64 ± 0.73	1.75 ± 0.72 ^c	3.16 ± 1.55	3.22 ± 1.19	3dW vs. 4dW: >0.999 3dW vs. 5dW: 0.064 3dW vs. 6dW: 0.107 4dW vs. 5dW: 0.435 4dW vs. 6dW: 0.003 5dW vs. 6dW: >0.999	3dW vs. 4dW: 0.14 3dW vs. 5dW: 1.71 3dW vs. 6dW: 1.88 4dW vs. 5dW: 1.58 4dW vs. 6dW: 1.75 5dW vs. 6dW: 0.04
Dr (A.U.)	1.36 ± 0.98	1.67 ± 0.85 ^c	2.55 ± 1.52	2.35 ± 1.02	3dW vs. 4dW: >0.999 3dW vs. 5dW: 0.447 3dW vs. 6dW: 0.179 4dW vs. 5dW: >0.999 4dW vs. 6dW: 0.038 5dW vs. 6dW: >0.999	3dW vs. 4dW: 0.29 3dW vs. 5dW: 1.08 3dW vs. 6dW: 0.92 4dW vs. 5dW: 0.88 4dW vs. 6dW: 0.72 5dW vs. 6dW: 0.17
s-RPEr (A.U.)	0.87 ± 0.84	0.92 ± 0.54	2.12 ± 2.90	1.40 ± 0.67	3dW vs. 4dW: >0.999 3dW vs. 5dW: >0.999 3dW vs. 6dW: 0.788 4dW vs. 5dW: >0.999 4dW vs. 6dW: 0.081 5dW vs. 6dW: >0.999	3dW vs. 4dW: 0.05 3dW vs. 5dW: 1.47 3dW vs. 6dW: 0.57 4dW vs. 5dW: 2.32 4dW vs. 6dW: 0.80 5dW vs. 6dW: 0.74
HSRr (A.U.)	0.31 ± 0.25 ^c	0.59 ± 0.28 ^c	1.31 ± 1.04	1.25 ± 0.65	3dW vs. 4dW: 0.220 3dW vs. 5dW: 0.110 3dW vs. 6dW: 0.008 4dW vs. 5dW: 0.558 4dW vs. 6dW: 0.010 5dW vs. 6dW: >0.999	3dW vs. 4dW: 1.02 3dW vs. 5dW: 1.32 3dW vs. 6dW: 4.70 4dW vs. 5dW: 3.72 4dW vs. 6dW: 2.39 5dW vs. 6dW: 0.08
SPDr (A.U.)	0.41 ± 0.60	0.49 ± 0.34 ^c	1.44 ± 1.05	1.28 ± 0.72	3dW vs. 4dW: >0.999 3dW vs. 5dW: 0.126 3dW vs. 6dW: 0.071 4dW vs. 5dW: 0.262 4dW vs. 6dW: 0.024 5dW vs. 6dW: 0.999	3dW vs. 4dW: 0.12 3dW vs. 5dW: 1.68 3dW vs. 6dW: 1.33 4dW vs. 5dW: 3.61 4dW vs. 6dW: 2.36 5dW vs. 6dW: 0.20
ACCr (A.U.)	0.92 ± 0.48 ^c	1.34 ± 0.65	2.06 ± 1.00	2.22 ± 1.23	3dW vs. 4dW: 0.441 3dW vs. 5dW: 0.064 3dW vs. 6dW: 0.036 4dW vs. 5dW: 0.879 4dW vs. 6dW: >0.999 5dW vs. 6dW: >0.999	3dW vs. 4dW: 0.79 3dW vs. 5dW: 2.18 3dW vs. 6dW: 2.51 4dW vs. 5dW: 0.99 4dW vs. 6dW: 1.18 5dW vs. 6dW: 0.14
DECr (A.U.)	0.81 ± 0.47	1.04 ± 0.50	1.87 ± 1.20	1.90 ± 0.99	3dW vs. 4dW: 0.999 3dW vs. 5dW: 0.177 3dW vs. 6dW: 0.053 4dW vs. 5dW: 0.713 4dW vs. 6dW: >0.999 5dW vs. 6dW: >0.999	3dW vs. 4dW: 0.44 3dW vs. 5dW: 2.17 3dW vs. 6dW: 2.18 4dW vs. 5dW: 1.50 4dW vs. 6dW: 1.56 5dW vs. 6dW: 0.02

RPEr: rating of perceived exertion ratio; Dr: duration ratio; s-RPEr: session rating of perceived exertion ratio; HSRr: high-speed running distance ratio; SPDr: sprint distance ratio; ACCr: accelerations ratio; DECr: decelerations ratio; A.U.: arbitrary units. a denotes significant difference from 4dW; b, a denotes significant difference from 5dW, c, a denotes significant difference from 6dW (all $p < 0.05$); Bold: significant results.

more than one match per week as previously suggested [14].

Furthermore, despite players' familiarisation with the wellness questionnaire in the previous season, reliability was not calculated, which can be considered a limitation. Additionally, the generalisation of these results to other teams, countries, competitive standards and ages is not recommended; thus, further replication studies are required. For instance, a recent study failed to find any external load difference between an under-18 and a first team [40], but an analysis of TMr would provide more insights for coaches. Considering that previous research observed training load variations and the following match outcome [41] and correlations between TMr and the match result [15], the inclusion of these contextual variables should be considered in future longitudinal studies. Finally, a detailed description of training drills in future research may improve practical implementation.

Although these results may depend on the analysed team, this study showed that training sessions were not adjusted according to weekly variations in terms of training sessions. This suggests that coaches need to consider modifying the training load to provide a balance across different types of microcycles. For instance, the present study revealed non-logical load application considering the weeks with few sessions, which presented shorter durations and, consequently, lower loads. Finally, TMr analysis facilitates the interpretation and contextualisation of data and, consequently, allows the training prescription to be planned accordingly to achieve the appropriate load. It also allows coaches and their staff to communicate with each other or with players as previously suggested [14].

5. Conclusions

In conclusion, weeks with fewer training sessions presented lower accumulated load and TMr. Specifically, accumulated wellness and load demands were higher in the weeks with more training sessions and progressively decreased in weeks with fewer training sessions ($6dW > 5dW > 4dW > 3dW$). On the one hand, higher wellness values are associated with better sleep quality and mood, as well as lower fatigue, muscle soreness and stress, which was associated with higher accumulated load and vice versa. On the other hand, high load values revealed non-logical load application considering that weeks with fewer sessions presented lower duration and consequently lower load. Coaches can plan weeks with fewer training sessions when the priority is for players to recover, as wellness values increased in such weeks during the present study. Of note, the present study analysed the main players on the team who participated in a minimum of 45 minutes in each match. Thus, for teams that only perform three or four training sessions, coaches should consider additional exercises. This could be particularly important for players who do not accumulate playing time.

Moreover, TMr showed the same tendency of higher values for weeks with more sessions, which followed the previous order ($6dW > 5dW > 4dW > 3dW$). However, some TMr, such as RPEr, HSRr, and SPD_r during 3dW and 4dW, as well as ACCr and DEC_r during 3dW, were lower than 1.0 A.U., which suggests that accumulated training load was lower

than match load. This also suggests that coaches may need to provide tailored individualised stimuli of HSR and SPD in weeks with 3dW and 4dW, as well as stimuli of ACCr and DEC in weeks with 3dW, to appropriately manage TMr.

AVAILABILITY OF DATA AND MATERIALS

Due to issues of participant consent related to the new data protection law from 25 May 2018 from the Portuguese data protection law n°. 58/2019 of 08 August, in accordance with the Council and European Parliament (EU) Regulation 2016/679, 27 April 2016, on the protection of individuals regarding the processing of personal data and on the free movement of such data, data will not be shared publicly. Interested researchers may contact the corresponding author.

AUTHOR CONTRIBUTIONS

RO, RCL and JPB—Conceptualization. RO and RCL—methodology. RCL—data collection. RO—formal analysis. RO and RCL—data curation. RO, RCL, RP, JVA, RM and JPB—Writing—original draft preparation; Writing—review and editing. RO, RM and JPB—supervision. RO and JPB—funding. All authors have read and agreed to the published version of the manuscript.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study followed the ethical guidelines for human study as suggested by the Declaration of Helsinki. Furthermore, the study was approved by the Research Ethics Committee of the Polytechnic Institute of Santarém, Santarém, Portugal (N°24-2022ESDRM). All players were informed about study procedures and oral and written consent were obtained from the participants.

ACKNOWLEDGMENT

The authors would like to thank the team's coaches and players for their cooperation during all data collection procedures.

FUNDING

This project was supported by the Portuguese Foundation for Science and Technology, I.P., Grant/Award Numbers: UIDP/04748/2020. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

CONFLICT OF INTEREST

The authors declare no conflict of interest. Rafael Oliveira is serving as one of the Editorial Board members of this journal. We declare that Rafael Oliveira had no involvement in the peer review of this article and has no access to information regarding its peer review. Full responsibility for the editorial

process for this article was delegated to DM.

REFERENCES

- [1] Halson SL. Monitoring training load to understand fatigue in athletes. *Sports Medicine*. 2014; 44: S139–S147.
- [2] Miguel M, Oliveira R, Loureiro N, García-Rubio J, Ibáñez SJ. Load measures in training/match monitoring in soccer: a systematic review. *International Journal of Environmental Research and Public Health*. 2021; 18: 2721.
- [3] Williams S, Trewartha G, Cross MJ, Kemp SPT, Stokes KA. Monitoring what matters: a systematic process for selecting training-load measures. *International Journal of Sports Physiology and Performance*. 2017; 12: S2101–S2106.
- [4] Gabbett TJ, Nassis GP, Oetter E, Pretorius J, Johnston N, Medina D, *et al.* The athlete monitoring cycle: a practical guide to interpreting and applying training monitoring data. *British Journal of Sports Medicine*. 2017; 51: 1451–1452.
- [5] Hooper SL, Mackinnon LT. Monitoring overtraining in athletes: recommendations. *Sports Medicine*. 1995; 20: 321–327.
- [6] McLean BD, Coutts AJ, Kelly V, McGuigan MR, Cormack SJ. Neuromuscular, endocrine, and perceptual fatigue responses during different length between-match microcycles in professional rugby league players. *International Journal of Sports Physiology and Performance*. 2010; 5: 367–383.
- [7] Bourdon PC, Cardinale M, Murray A, Gastin P, Kellmann M, Varley MC, *et al.* Monitoring athlete training loads: consensus statement. *International Journal of Sports Physiology and Performance*. 2017; 12: S2161–S2170.
- [8] Clemente FM, Rabbani A, Conte D, Castillo D, Afonso J, Clark CCT, *et al.* Training/match external load ratios in professional soccer players: a full-season study. *International Journal of Environmental Research and Public Health*. 2019; 16: 3057.
- [9] Oliveira R, Martins A, Moreno-Villanueva A, Brito JP, Nalha M, Rico-González M, *et al.* Reference values for external and internal load training intensity monitoring in professional male soccer players: a systematic review. *International Journal of Sports Science & Coaching*. 2022; 17: 1506–1530.
- [10] Buchheit M, Simpson BM. Player-tracking technology: half-full or half-empty glass? *International Journal of Sports Physiology and Performance*. 2017; 12: S235–S241.
- [11] Trewin J, Meylan C, Varley MC, Cronin J. The influence of situational and environmental factors on match-running in soccer: a systematic review. *Science and Medicine in Football*. 2017; 1: 183–194.
- [12] Anderson L, Orme P, Di Michele R, Close GL, Morgans R, Drust B, *et al.* Quantification of training load during one-, two- and three-game week schedules in professional soccer players from the English Premier League: implications for carbohydrate periodisation. *Journal of Sports Sciences*. 2016; 34: 1250–1259.
- [13] Oliveira R, Brito J, Martins A, Mendes B, Calvete F, Carriço S, *et al.* In-season training load quantification of one-, two- and three-game week schedules in a top European professional soccer team. *Physiology & Behavior*. 2019; 201: 146–156.
- [14] Stevens TGA, de Ruiter CJ, Twisk JWR, Savelsbergh GJP, Beek PJ. Quantification of in-season training load relative to match load in professional Dutch Eredivisie football players. *Science and Medicine in Football*. 2017; 1: 117–125.
- [15] Modric T, Jelacic M, Sekulic D. Relative training load and match outcome: are professional soccer players actually undertrained during the in-season? *Sports*. 2021; 9: 139.
- [16] Szigeti G, Schuth G, Revisnyei P, Pasic A, Szilas A, Gabbett T, *et al.* Quantification of training load relative to match load of youth national team soccer players. *Sports Health*. 2022; 14: 84–91.
- [17] Nobari H, Oliveira R, Clemente FM, Adsuar JC, Pérez-Gómez J, Carlos-Vivas J, *et al.* Comparisons of accelerometer variables training monotony and strain of starters and non-starters: a full-season study in professional soccer players. *International Journal of Environmental Research and Public Health*. 2020; 17: 6547.
- [18] Nobari H, Praça GM, Clemente FM, Pérez-Gómez J, Carlos Vivas J, Ahmadi M. Comparisons of new body load and metabolic power average workload indices between starters and non-starters: a full-season study in professional soccer players. *Proceedings of the Institution of Mechanical Engineers, Part P: Journal of Sports Engineering and Technology*. 2021; 235: 105–113.
- [19] Fernandes R, Brito JP, Vieira LHP, Martins AD, Clemente FM, Nobari H, *et al.* In-season internal load and wellness variations in professional women soccer players: comparisons between playing positions and status. *International Journal of Environmental Research and Public Health*. 2021; 18: 12817.
- [20] Oliveira R, Ceylan HI, Brito JP, Martins A, Nalha M, Mendes B, *et al.* Within- and between-mesocycle variations of well-being measures in top elite male soccer players: a longitudinal study. *Journal of Men's Health*. 2022; 18: 94.
- [21] Clemente FM, Mendes B, Nikolaidis PT, Calvete F, Carriço S, Owen AL. Internal training load and its longitudinal relationship with seasonal player wellness in elite professional soccer. *Physiology & Behavior*. 2017; 179: 262–267.
- [22] Borg G. Perceived exertion and pain scales. 1st edn. Human Kinetics: Champaign IL, USA. 1998.
- [23] Foster C, Florhaug JA, Franklin J, Gottschall L, Hrovatin LA, Parker S, *et al.* A new approach to monitoring exercise training. *The Journal of Strength and Conditioning Research*. 2001; 15: 109–115.
- [24] Foster C, Hector LL, Welsh R, Schragger M, Green MA, Snyder AC. Effects of specific versus cross-training on running performance. *European Journal of Applied Physiology and Occupational Physiology*. 1995; 70: 367–372.
- [25] Crang ZL, Duthie G, Cole MH, Weakley J, Hewitt A, Johnston RD. The inter-device reliability of global navigation satellite systems during team sport movement across multiple days. *Journal of Science and Medicine in Sport*. 2022; 25: 340–344.
- [26] Houtmeyers KC, Jaspers A, Brink MS, Vanrenterghem J, Varley MC, Helsen WF. External load differences between elite youth and professional football players: ready for take-off? *Science and Medicine in Football*. 2021; 5: 1–5.
- [27] Gonçalves LGC, Kalva-Filho CA, Nakamura FY, Rago V, Afonso J, Bedo BLS, *et al.* Effects of match-related contextual factors on weekly load responses in professional Brazilian soccer players. *International Journal of Environmental Research and Public Health*. 2020; 17: 5163.
- [28] Nobari H, Fani M, Pardos-Mainer E, Pérez-Gómez J. Fluctuations in well-being based on position in elite young soccer players during a full season. *Healthcare*. 2021; 9: 586.
- [29] Clemente FM, Silva R, Castillo D, Arcos AL, Mendes B, Afonso J. Weekly load variations of distance-based variables in professional soccer players: a full-season study. *International Journal of Environmental Research and Public Health*. 2020; 17: 3300.
- [30] Hader K, Rumpf MC, Hertzog M, Kilduff LP, Girard O, Silva JR. Monitoring the athlete match response: can external load variables predict post-match acute and residual fatigue in soccer? A systematic review with meta-analysis. *Sports Medicine-Open*. 2019; 5: 48.
- [31] Clemente FM, Martinho R, Calvete F, Mendes B. Training load and well-being status variations of elite futsal players across a full season: comparisons between normal and congested weeks. *Physiology & Behavior*. 2019; 201: 123–129.
- [32] Fessi MS, Nouira S, Dellal A, Owen A, Elloumi M, Moalla W. Changes of the psychophysical state and feeling of wellness of professional soccer players during pre-season and in-season periods. *Research in Sports Medicine*. 2016; 24: 375–386.
- [33] Nobari H, Fani M, Clemente FM, Carlos-Vivas J, Pérez-Gómez J, Ardigo LP. Intra- and inter-week variations of well-being across a season: a cohort study in elite youth male soccer players. *Frontiers in Psychology*. 2021; 12: 671072.
- [34] Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Medicine & Science in Sports & Exercise*. 2009; 41: 3–13.
- [35] Silva AF, Oliveira R, Cataldi S, Clemente FM, Latino F, Badicu G, *et al.* Weekly variations of well-being and interactions with training and match intensities: a descriptive case study in youth male soccer players. *International Journal of Environmental Research and Public Health*. 2022; 19: 2935.

- [36] Robey E, Dawson B, Halson S, Gregson W, Goodman C, Eastwood P. Sleep quantity and quality in elite youth soccer players: a pilot study. *European Journal of Sport Science*. 2014; 14: 410–417.
- [37] Oliveira R, Brito JP, Martins A, Mendes B, Marinho DA, Ferraz R, *et al.* In-season internal and external training load quantification of an elite European soccer team. *PLOS ONE*. 2019; 14: e0209393.
- [38] Dalen T, Sandmæl S, Stevens TGA, Hjelde GH, Kjøenes TN, Wisløff U. Differences in acceleration and high-intensity activities between small-sided games and peak periods of official matches in elite soccer players. *Journal of Strength and Conditioning Research*. 2021; 35: 2018–2024.
- [39] Clemente FM, Sarmiento H, Rabbani A, Van Der Linden CMIN, Kargarfard M, Costa IT. Variations of external load variables between medium- and large-sided soccer games in professional players. *Research in Sports Medicine*. 2019; 27: 50–59.
- [40] Morgans R, Bezuglov E, Orme P, Burns K, Rhodes D, Babraj J, *et al.* The physical demands of match-play in academy and senior soccer players from the scottish premiership. *Sports*. 2022; 10: 150
- [41] Oliveira R, Brito JP, Loureiro N, Padinha V, Ferreira B, Mendes B. Does the distribution of the weekly training load account for the match results of elite professional soccer players? *Physiology & Behavior*. 2020; 225: 113118.

How to cite this article: Rafael Oliveira, Rui Canário-Lemos, Ryland Morgans, Rafael Peixoto, José Vilaça-Alves, João Paulo Brito. Are male soccer players accumulating sufficient load across varying microcycle structures? Examining the load, wellness and training/match ratios of a professional team. *Journal of Men's Health*. 2023; 19(9): 26-35. doi: 10.22514/jomh.2023.083.