

Central Lancashire Online Knowledge (CLoK)

Title	Psycho-social factors associated with type two diabetes remission through						
	lifestyle intervention: A scoping review						
Туре	Article						
URL	https://clok.uclan.ac.uk/id/eprint/49737/						
DOI	https://doi.org/10.1371/journal.pone.0294344						
Date	2023						
Citation	Hart, Kathryn Jane, Kubilius, Andrew and Clark, Martin (2023) Psycho-social						
	factors associated with type two diabetes remission through lifestyle						
	intervention: A scoping review. PLOS ONE, 18 (11).						
Creators	Hart, Kathryn Jane, Kubilius, Andrew and Clark, Martin						

It is advisable to refer to the publisher's version if you intend to cite from the work. https://doi.org/10.1371/journal.pone.0294344

For information about Research at UCLan please go to http://www.uclan.ac.uk/research/

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <u>http://clok.uclan.ac.uk/policies/</u> Check for updates

GOPEN ACCESS

Citation: Hart KJ, Kubilius A, Clark M (2023) Psycho-social factors associated with type two diabetes remission through lifestyle intervention: A scoping review. PLoS ONE 18(11): e0294344. https://doi.org/10.1371/journal.pone.0294344

Editor: Meisam Akhlaghdoust, Shahid Beheshti University of Medical Sciences, ISLAMIC REPUBLIC OF IRAN

Received: August 8, 2023

Accepted: October 28, 2023

Published: November 16, 2023

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: https://doi.org/10.1371/journal.pone.0294344

Copyright: © 2023 Hart et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its <u>Supporting information</u> files.

RESEARCH ARTICLE

Psycho-social factors associated with type two diabetes remission through lifestyle intervention: A scoping review

Kathy J. Hart[®], Andrew Kubilius[®], Martin Clark[®]*

Department of Psychology and Computer Science, University of Central Lancashire, Preston, Lancashire, United Kingdom

So These authors contributed equally to this work.

* MClark11@uclan.ac.uk

Abstract

Objectives

There is growing evidence that type 2 diabetes (T2D) can be put into remission through lifestyle intervention. Current focus on remission in terms of physiological considerations and biomedical scales, means there is limited understanding of the role psycho-social factors play in moderating the efficacy of lifestyle interventions for T2D remission. In the current review we aimed to synthesise the emerging literature on psycho-social factors associated with T2D remission, specifically from lifestyle interventions.

Methods

Five databases (EMBASE, MEDLINE, CINAHL ultimate, PsychINFO and PsycArticles) were searched to identify studies from 2009 onwards that reported remission outcomes from lifestyle interventions in participants \geq 18years old, with a clinical diagnosis of T2D. Studies included were of an interventional or observational design and restricted to English language. Screening and data extraction was performed independently by two reviewers using prespecified criteria.

Results

In total 6106 studies were screened, 36 studies meeting the inclusion criteria were included. Studies were globally diverse, with 30 (83%) being published \geq 2017. Psycho-social scales were under-utilised with 22 (67%) of studies failing to include any psycho-social measures. Single arm, prospective studies were most frequently utilised, however study quality was perceived to be heterogeneous. Further disparity in the quality, content and delineation of the psycho-social interventions was also identified. Education and self-monitoring interventions were the most frequently incorporated. Self-monitoring was also identified as an important facilitator to remission, in addition to social support.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Conclusions

Our findings indicate that psycho-social factors in T2D remission are under-explored. We have identified a number of methodological issues (comparability, conflicting remission criteria and poorly defined psycho-social interventions) suggesting knowledge gaps which could inform the methodology of future research design. There is significant opportunity for future research to incorporate the social model of disease, conceptualise remission more holistically, and build a more comprehensive evidence base to guide clinical practice.

Introduction

In 2021 an estimated 536 million of the global population were affected by type two diabetes (T2D) [1] Saeedi and colleagues [2] estimate this will rise to 700 million by 2045. Physiologically individuals with T2D are at risk of multiple serious morbidities, whilst psychologically, depression, anxiety, diabetes distress and diabetes stigma are all prevalent and associated with poorer health outcomes [3,4].

There has been a paradigmatic shift in how T2D is conceptualised and treated [5]. Historically, it was considered a chronic condition requiring increased medication as it progressed [6–8]. Recently, evidence from bariatric studies, and from dietary and lifestyle interventions, has now elucidated that with sufficient weight loss, remission from diabetes could be an achievable outcome [9–12]. Diabetes remission has been established as a primary research priority by the James Lind Alliance [13,14], however this remains an emergent field with significant heterogeneity as to how remission and the associated clinical parameters should be delineated [3,15–18].

In addition to a lack of consensus regarding a consistent definition of diabetes remission, there is also a lack of scrutiny of the psycho-social factors which could moderate or mediate remission outcomes. 'Psychosocial' is defined by Martikainen et al. [19] as "the influence of social factors on an individual's mind or behaviour, and to the interrelation of behavioural and social factors". These factors play a key role in mediating the effects of social determinants on an individual's health, impacting physical health directly as protective or risk factors, and also through their influence on health behaviours such as physical activity levels and diet [20]. In diabetes management research e.g. Tharek et al [21] has already established the critical role of multiple psycho-social factors in self-care and glycaemic control. Self-efficacy [21–23], social support [24–28], digital health literacy [13,29,30] and motivation [14,31,32] have all been found to positively impact health outcomes. There are multiple psycho-social scales, both generic and diabetes specific, e.g. Polonsky et al [33,34]; Bijl et al [35], which have been developed to measure the impact of these factors and aid the delivery of person-centred care but it is unclear how widely or frequently these have been applied in remission studies.

Within current consensus statements [18] remission is narrowly focused on "physiological considerations and monitoring of biomedical indices", to the detriment of our understanding of psycho-social factors [36] (p 406). Multiple studies, e.g. Nicolucci et al [37]; Whicher et al [38] and Wylie et al [39] also identify that the psychological and emotional needs of individuals with T2D are consistently an unmet priority. This limits our understanding of the support individuals pursuing remission may require. It also means that remission, in the absence of appropriate tailored support, may negatively impact an individual's physical or mental health [3,36].

The primary research objective is to identify the psycho-social factors associated with T2D remission through lifestyle intervention. The secondary objectives are to identify the psycho-social scales used within remission studies; assess the method and design of these studies; identify the psycho-social interventions incorporated; determine how the reported psycho-social factors may influence remission outcomes and finally to identify gaps within the evidence base. It is hoped this review will inform future research design and aid practice of Health Care Professionals (HCP) working with individuals with T2D in identifying the potential barriers and facilitators to remission.

Methods

This scoping review was conducted following guidance from The JBI Manual for Evidence Synthesis [40], Peters et al. [41] and Arksey and O'Malley [42]. Reporting is based on the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) [43], with the completed PRISMA-ScR being provided in supporting information (see <u>S1 Table</u>). The objectives, methods and inclusion criteria for this review were all determined *a priori* and registered on the Open Science Framework (OSF) prior to literature screening: https://osf.io/pu3k4/?view_only=7661ed50e1fe4f31afe53c71239ce9c7.

Eligibility criteria

In line with the JBI manual (2020), inclusion criteria were formulated according to the Population, Concept and Context (PCC) mnemonic, see Table 1. This establishes the scope of the review in identifying the desired study population (adults with T2D), the phenomenon of interest (remission) and any specific restrictions. Of fundamental importance to comparability was the need for authors to include remission outcomes according to whichever criteria they specified. This is a complexity inherent in this field of study where neither a descriptor nor consensus to remission criteria has been agreed. Studies were restricted to 2009 onwards when the first preliminary discussion on remission descriptors and criteria, was published [15]. No geographical restrictions were imposed. Participants under the age of 18 were excluded as is common within existing diabetes remission reviews (see Captieux et al [44] and Goldenberg et al [45]). The focus on this review is on the psycho-social domain including behavioural change through lifestyle or health behaviour intervention. Studies included were interventions that sought to change health related behaviours, diet or exercise, in the absence of new pharmaceutical agents or surgery.

PCC	
Population	Population: T2DM Include: pre-existing clinical diagnosis of T2D, adult Exclude: patients who 'self-identify' with 2TD, diagnosis through screening within study, < 18 years old, any other type of diabetes (e.g., type one, gestational, MODY), combined study population of pre- diabetes and T2D where results cannot be extracted separately
Concept	Concept: Remission Include: diabetes remission through health behaviour intervention Exclude: remission from diabetic complications, studies where remission outcomes are not reported, remission through surgical or pharmaceutical means Outcome: psycho-social scales, psycho-social interventions, psycho-social factors reported by participants
Criteria	Context: Include: interventional or observational studies, ≥ 2009, English language, any geographical area Exclude: secondary literature (reviews, commentary pieces etc.)

Table 1. Population, Concept and Context (PCC) inclusion criteria [46].

https://doi.org/10.1371/journal.pone.0294344.t001

Search strategy and information sources

Preliminary searches of CINAHL and MEDLINE were conducted with a Senior Librarian at the University of Central Lancashire (UCLan) and a search strategy developed *a priori* in June 2022. Key search terms were extracted from consensus remission statements [16–18] (see S2 Table). The electronic databases EMBASE, MEDLINE, CINAHL ultimate (see S3 Table), PsychINFO and PsycArticles were searched in September- October 2022. Key terms were added verbatim to each database with minor adjustments made to index terms and database specific terminology where necessary. Additional records were identified through web searches and hand searching reference lists. Grey literature was searched using Google Scholar.

Data management and software

MS Excel was used to calculate statistical data and compile frequency charts. References were collated, uploaded and de-duplicated using the reference managing software EndNote X9 [47].

Study selection

Two reviewers independently screened all titles, abstracts and full texts excluding studies that did not meet the inclusion criteria. Any discrepancies in inclusion were resolved firstly through discussion. Where we were unable to achieve a consensus, the text was referred to a third reviewer for independent evaluation.

Data extraction

Data was independently extracted and summarised within two data extraction forms, developed *a priori*. The forms were piloted using a sample of three studies [48-50]. The first form (see S4 Table) focused on data pertaining to study design, psycho-social scales, and the impact of psycho-social factors on remission. The second form, on the psycho-social interventions (See S5 Table).

Data analysis

Studies using the following designs: random crossover trials, open-label cluster randomised trials and open-label, parallel-group and randomised controlled trials (RCTs), were grouped together as RCTs for analysis. For the psycho-social interventions, studies that linked to the same RCTs, and therefore the same interventions, were also grouped together. Results for both interventional and observational studies were synthesised in the same manner. In line with the scoping review methodology, risk of bias has not been formally evaluated. However, indication has been made to potential quality issues, data inconsistency and where further research should be directed to address deficits.

Results

Initial database searches identified 6100 records, with a further six identified through hand searching reference lists. Searches for grey literature did not yield any additional records. After de-duplication 5038 records remained, limited to 141 after title and abstract screening. After full text screening, 36 records were identified which met the inclusion criteria, see Fig 1. Of the studies included 30 (83%) were published \geq 2017, reflecting the emergent nature of the field. The sample was globally diverse, with studies being conducted in UK (10), USA (9), India (3), Italy (2), Thailand (2), Barbados (1), Canada (1), Brazil (1), the Netherlands (1), Switzerland (1), Israel (1), Denmark (1), Quatar (1), South Africa (1) and China (1).



Fig 1. PRISMA flow diagram for the electronic search and screening process [51].

https://doi.org/10.1371/journal.pone.0294344.g001

Psycho-social scales

Analysis of the 33 studies which were quantitative or mixed method (qualitative studies due to their nature were unlikely to include psycho-social scales and were therefore excluded) revealed that 22 (67%) failed to include any psycho-social measures (see Fig 2). Of the studies which did include psycho-social measures, the majority were limited to one quality of life scale, either the 36 item Short Form Health Survey (SF-36) or the EuroQol (EQ-5D) battery. Only three studies utilised diabetes specific scales [52–54].

Study designs

Of the 36 studies included, 13 (36%) were retrospective in design (Fig 3). Where the study involved retrospective analysis of an intervention, with the exception of the qualitative studies, the design of the original study was used to inform findings. Particularly significant Randomised Controlled Trials (RCTs) such as the seminal DiRECT study have generated multiple primary [11,12] and secondary [8] analyses. RCT's (n = 10) comprised 36% of the sample. Single arm prospective studies (n = 13), usually pilot or feasibility with n \leq 45, were the most utilised study design (50%).

Whilst participants in all reviewed papers had an existing clinical diagnosis of T2D, study populations were otherwise heterogeneous [55]. Restrictions to inclusion criteria in the papers reviewed were based on multiple factors: duration of diagnosis, being non-insulin taking and/



Fig 2. The number of psycho-social scales utilised in all studies reviewed.

https://doi.org/10.1371/journal.pone.0294344.g002



Fig 3. The number of studies reviewed that confer to one of the seven designs.

https://doi.org/10.1371/journal.pone.0294344.g003

or naïve to glucose lowering medication, age, existing diabetes complications, existing mental health diagnosis, BMI and HbA1c level.

Psycho-social interventions

Interventional studies were grouped and analysed (n = 27), Evans et al. [56] was used to guide identification of intervention specifics. Insufficient detail within the papers necessitated scrutiny of clinical trial registrations, protocols and other cited papers, however in some cases these were inaccessible as indicated on Fig 4. A lack of precision in terminology or the use of generic expressions such as 'care', 'support' and 'counselling' in the absence of context or further detail, resulted in a lack of clarity indication.

Key: + signifies evidence of the intervention,—no evidence of the intervention, ? lack of clarity around the intervention and * unable to access protocol. There was a clear disparity in the content of the psycho-social interventions, with some offering no evidence of any psycho-social intervention (n = 4), and others being multifaceted incorporating 5 or more features (n = 9) (Fig 4). The most commonly used intervention was education (n = 18); however, it was also the most under-described. Self-monitoring (n = 16), the next most frequently adopted intervention, was more explicit in both the measures used (CGM, food diary, weight checks, accelerometers) and how these impacted other dimensions such as goal setting. There was evidence of theoretical underpinning or supportive studies being cited in (n = 11).

Influence of psycho-social factors on remission outcomes

Whilst there is only limited quantitative data from psycho-social scales presented in the reviewed papers, emergent positive relationships were apparent. With participation in the interventions(in which some achieved remission and the majority demonstrably improved their condition), being associated with increased quality of life [8,11,12,57–59], increased self-efficacy [60], significantly increased self-esteem [52], increased work and social adjustment [52], reduced binge eating [8,52], reduced problem areas in diabetes management [54] and finally increased diabetes knowledge and empowerment [53]. There were inconsistencies in relation to anxiety and depression [52,57,60,61].

From the qualitative and mixed method studies, the most prominent factor was the role of social support highlighting both a facilitatory [62] and inhibitory influence on remission outcomes [63]. Where the support was positive and continuous, participants commented that this was beneficial to motivation, accountability and adherence. However, where support was lacking or inconsistent, participants noted the significant challenges of navigating negative feedback and social saboteurs [63,64]. A further key observation related to perceptions of self-image and identity, and how this shifted in relation to weight loss and the appraisal of others, [63,65]. A final shared observation related specifically to the shifting cognition associated with total diet replacement (TDR) and subsequent food reintroduction and included themes relating to decision making, self-regulation, adaption and autonomy [63,65,66].

Discussion

In this scoping review we analysed 36 studies based on lifestyle or health behaviour intervention which reported T2D remission outcomes. To our knowledge this is the first review to specifically address the psycho-social factors associated with remission and broaden the focus from the dominant bio-medical perspective.

Our findings confirm that psycho-social factors in T2D remission are often overlooked [36] or receive minimal acknowledgement. Our review indicated that only 11 out of 33 interventional studies included psycho-social scales and 23 out of 27 studies employed psycho-social

Intervention	Self- monitoring	Goal setting	Cognitive restructuring	Education	Individual sessions	Group sessions	Peer support online	Involvement family/friends	Cited studies/ theoretical underpinning
Study									
Ades et al (2015)	+	+	+	+	?	+	-	-	+
Athinarayan et al (2019)	+	?	?	+	?	?	+	-	+
Bhatt et al (2017)	+	-	-	?	+	-	-	-	-
Bynoe et al * (2019)	-	-	-	-	-	-	-	-	-
Correa et al (2022)	-	-	-	+	-	-	-	-	-
Cox et al (2019)	-	?	?	+	+	-	-	-	-
Dave et al (2019)	+	?	?	+	+	-	-	+	-
De Hoogh et al* (2021)	+	?	?	+	+	-	-	-	-
Esposito et al (2014)	-	-	-	+	-	-	-	-	-
Gregg et al (2012)	+	+	+	+	+	+	-	-	+
Lagger et al (2018)	?	+	+	+	+	+	-	-	+
Lean et al (2018), Lean et al (2019), Rechakova et al (2021) Thom et al (2021) (DIRECT study)	÷	-	-	+	+	-	-	-	-
Marples et al (2022)	-	+	?	+	+	+	-	-	+
Mottalib et al (2015)	+	+	+	+	?	+	-	-	+
Oser et al (2022)	+	?	+	+	-	-	-	-	+
Rechakova et al (2017) Rechakova et al (2019) Steven et al (2016) (Counterbalance study)	+	÷	?	+	+	-	-		+
Rein et al	+	_	_	-	2	_	_	_	_
(2022) Reid-Larsen et al (2019)	+	-	-	+	?	+	+	-	+
Romana et al (2019)	-	-	-	-	-	-	-	-	-
Sarathi et al (2017)	-	-	-	-	-	-	-	-	-
Taheri et al (2020)	+	+	+	+	+	-	-	?	+
Umphonsathien et al * (2019)	+	-	-	-	-	-	-	-	-
Umphonsathien et al* (2022)	+	-	-	-	?	-	-	-	-
Unwin et al (2020)	-	+	+	+	÷	+	-	+	+
Webster et al (2019)	-	-	-	-	-	-	-	-	-
Yancy et al (2019)	+	?	?	+	-	+	-	+	-
Zou et al (2022)	-	-	-	-	+	-	-	-	-

Fig 4. Evidence of psycho-social interventional features in all the studies reviewed.

https://doi.org/10.1371/journal.pone.0294344.g004

interventions, although in some instances these only just met the inclusion criteria. There was significant study heterogeneity and a number of comparability issues: variable trial quality, conflicting remission criteria, differences in remission rates and variability in study populations. Furthermore, psycho-social interventions tended to be poorly delineated which made analysis more challenging.

Psycho-social scales were underused and indeed absent in the majority of studies included. The average number of biomedical scales utilised per study was 12.15, in comparison psychosocial scales averaged at only 0.76. Where psycho-scales were utilised the most frequent were SF36 scale or the EQ-5D battery which measure quality of life. However, these are non-diabetes specific, and the latter has been subject to criticism for its generic nature and insufficient focus on mental and social aspects of health [67]. Effectively they give little indication to participant's cognitive, emotional or behavioural states, beyond a general concept of 'wellness'. As psycho-social factors are critical both in diabetes management [13,22,25,68] and weight loss [69] (the greatest predictor of remission) their underuse within these studies, means there is little quantitative evidence base to identify barriers or facilitators of remission. The inclusion of a broader number of scales at baseline, for example in motivation, self-efficacy or selfesteem, could potentially identify those who may find behavioural change more challenging and suggest potential for preliminary intervention such as motivational interviewing [70] or self-efficacy training [71] both of which have proven efficacy in promoting behaviour regulation. Clearly though this is still an emerging field, and as time progresses there is evidence of greater scrutiny of psycho-social factors. In the studies we reviewed between 2012 and 2017, psycho-social scales were completely absent. From 2018 onwards there has been a slow progression to increased inclusion.

The presence of psycho-social scales within the retrospective studies (n = 13) was dependent and, in many cases, limited by the design of the initial research. This was particularly apparent in the retrospective cohort studies reliant on routine clinical data, as psycho-social measures are rarely utilised in primary care despite these factors being an essential component of person-centred care [72]. There are also known quality and methodological issues inherent with the use of retrospective cohort studies [73] and case studies [74,75]. The most frequent study design, single arm (n = 13/36), measured the initial efficacy of a lifestyle intervention, however the lack of a control arm means that spontaneous recovery or placebo effects cannot be ruled out in relation to remission or psycho-social factors [76]. Furthermore, as the majority of participants either self-selected or in some instances were subject to a compliance check prior to participation [58] study populations may have had higher motivation levels than the general diabetes population.

Randomised Control Trials (RCTs) are widely considered to be the gold standard in research due to the fact they provide more rigour in examining the relationship between intervention and outcome, in addition to minimising bias [77]. Perhaps the most conclusive of these in terms of the relationship between psycho-social factors and remission is the RCT by Taheri et al. [57] due to the higher number of those achieving remission (61% in the intervention arm compared to 12% in the control group). However, within the intervention arm of the included RCTs remitters and non-remitters were rarely separated. As a result, there could be a number of psychological processes underpinning remission that are not being explored. Existing studies have already indicated that duration of diabetes moderates ability to achieve remission which is usually attributed to beta cell dysfunction and pancreatic capacity [49,78–81], but the psychological processes implicit within this are under investigated. Steven et al. [81] note that remission may be more difficult to achieve for those who have failed at repeated attempts at weight loss. Arguably with longer duration of diabetes, coping mechanisms,

cognitive structures and eating behaviours will all have become more entrenched over time, resistance to change may therefore be psychological as much as it is physiological.

Analysis of the qualitative studies identified numerous psycho-social factors for example social support [82] and psychoeducation [83], which support pre-existing knowledge of facilitators to effective diabetes management. Our findings also suggest additional psycho-social factors uniquely associated with weight loss. Dietary restriction, particularly through TDR can result in dramatic weight loss over a short period of time, significantly impacting the appearance of the individual. It is suggested that this leads to challenges with perception of self and self-image [63,65]. It further invites more opportunity for negative societal comment, or social saboteurs [84]. Targeted psychological support may be required to support this process of physical and psychological change. A further facilitatory factor widely incorporated in the psycho-social interventions was self-monitoring (present in 16 of the 27 studies reviewed), specifically the role of biofeedback [62]. The use of CGM for example was evaluated by participants in Oser et al [53] as being the most useful component of the intervention. Whilst self-monitoring was identified as having an empowering effect [53] and there is literature to support the efficacy of both self-weighing [85] and self-monitoring blood glucose [86,87] there appears to be a degree of trepidation in health policy recommendations. The National Institute for Health and Care Excellence (NICE) [88] state that only those advised by a doctor or nurse should selfmonitor their blood glucose. This may serve to disempower individuals who are pursuing remission in the community and is an important focus for further research.

Historically psycho-social interventions have been poorly delineated [89,90] this field proves no exception. Use of terms such as 'support', 'care' or 'counselling' without context or reference to theoretical underpinning complexifies analysis and threatens future replicability. In the majority of the studies lifestyle interventions were incorporated with some degree of psychosocial intervention, meaning the individual impact of either of these mechanisms cannot be extrapolated. This limits understanding of the optimum conditions required to support an individual to achieve and sustain remission. Consideration was given to the potential impact of Covid 19 on the nature and facilitation of the psycho-social interventions. We reviewed the end of each study's data collection period and whilst the majority were pre-Covid, Marples and colleagues [52] and Oser and colleagues [53] both discuss accessibility issues and digital adaptions due to the Covid restrictions. This is a pertinent point for future researchers to consider.

Furthermore, in a number of studies secondary focus/analysis was not on psychosocial factors, but instead rested on cost efficacy of the interventions and the prospects of savings for the respective healthcare services [78,91–93]. Focus on the financial implications of diabetes care may reinforce stigmatising rhetoric whereby the individual is blamed both for their own 'moral failure' to prevent what is perceived as a lifestyle disease and for the subsequent cost burden this incurs. There is a danger that remission could be seen as a panacea, a means of restoring normoglycaemia without costly pharmaceuticals or the need for sustained primary care and concerningly, Holman and colleagues [94] have recently indicated that once a patient is clinically coded as in remission routine diabetes care checks are reduced.

Strengths of this review include the globally diverse sample which minimises publication bias and the inclusion of a broad range of study types. These may previously have been overlooked by systematic reviews focused on diet with more rigid design exclusion criteria in place [95–97]. Clarity of process and reporting was achieved by adhering to both JBI [40] recommendations and the PRISMA-ScR [43] statement for scoping reviews. We adopted a systematic approach with all data extraction and screening being carried out independently by two reviewers. Limitations to this study include the restriction to English language, inability to access a number of protocols and the fact a more extensive search of grey literature could have been implemented by accessing dissertation and theses repositories. Whilst under 18s were excluded from this review, given the increasing prevalence of T2D in young people and the paucity of existing research, this may represent an important focus for future research.

Conclusion

This research aimed to investigate the psycho-social factors associated with type two diabetes remission through lifestyle intervention. Established as a research priority by the James Lind alliance and with significant interest from the diabetes community, this review has demonstrated that there is a clear need to address the psycho-social domain with renewed vigour. Currently, heterogeneity in study design and quality, variance in the study populations, a lack of consensus to remission criteria and poorly delineated terms are impeding replicability and the establishment of a sound knowledge base. For T2D remission to be implemented effectively within clinical practice, data from psycho-social scales and the evaluation of psychosocial interventions is critical to ensuring that advice and support offered to patients is both evidence based and person-centred. There are significant opportunities to tailor future research to reflect a more social model of disease which recognises the challenges of obesogenic environments, societal, family and cultural demands on individuals choosing to pursue remission. Rather than solely a bio-medical procedure, remission should be recognised, and treated, as a holistic process.

Supporting information

S1 Table. Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) checklist. (DOCX)

S2 Table. Search terms used for the database searches of EMBASE, MEDLINE, CINAHL ultimate, PsychINFO and PsycArticles. (DOCX)

S3 Table. Results from database search of CINAHL Ultimate (conducted 26/9/22). (DOCX)

S4 Table. Data extraction form one- detailing study, country, participants, design, lifestyle intervention, number of psycho-social scales used and the effect of these on remission outcomes.

(DOCX)

S5 Table. Data extraction form two- detailing author and related studies, psycho-social interventions and cited studies which underpin the interventions. (DOCX)

Author Contributions

Conceptualization: Kathy J. Hart, Martin Clark. Data curation: Kathy J. Hart, Andrew Kubilius. Formal analysis: Kathy J. Hart. Investigation: Kathy J. Hart, Andrew Kubilius. Methodology: Kathy J. Hart. Project administration: Kathy J. Hart.

Supervision: Martin Clark.

Visualization: Kathy J. Hart.

Writing - original draft: Kathy J. Hart.

Writing - review & editing: Kathy J. Hart, Martin Clark.

References

- Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, et al. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. Diabetes Res Clin Pract. 2022; 183:109119. Epub 2021/12/10. https://doi.org/10.1016/j.diabres.2021. 109119 PMID: 34879977.
- Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9(th) edition. Diabetes Res Clin Pract. 2019; 157:107843. Epub 2019/09/ 14. https://doi.org/10.1016/j.diabres.2019.107843 PMID: 31518657.
- MacKay D, Chan C, Dasgupta K, Dominy C, Gagner M, Jin S, et al. Remission of Type 2 Diabetes: Diabetes Canada Clinical Practice Guidelines Expert Working Group. Can J Diabetes. 2022; 46(8):753–61 e8. Epub 2022/12/26. https://doi.org/10.1016/j.jcjd.2022.10.004 PMID: 36567079.
- Perrin N, Bodicoat DH, Davies MJ, Robertson N, Snoek FJ, Khunti K. Effectiveness of psychoeducational interventions for the treatment of diabetes-specific emotional distress and glycaemic control in people with type 2 diabetes: A systematic review and meta-analysis. Prim Care Diabetes. 2019; 13 (6):556–67. Epub 2019/05/02. https://doi.org/10.1016/j.pcd.2019.04.001 PMID: 31040069.
- Chang TJ. Mechanisms and the strategy for remission of type 2 diabetes mellitus. J Diabetes Investig. 2023. Epub 2023/01/11. https://doi.org/10.1111/jdi.13948 PMID: 36625268.
- Al-Mrabeh A. Pathogenesis and remission of type 2 diabetes: what has the twin cycle hypothesis taught us? Cardiovasc Endocrinol Metab. 2020; 9(4):132–42. Epub 2020/11/24. https://doi.org/10.1097/XCE. 000000000000201 PMID: 33225228.
- 7. Taylor R, Barnes A. From new understanding of type 2 diabetes to practical management. Diabetologia. 2018; 61:273–83.
- Thom G, Messow CM, Leslie WS, Barnes AC, Brosnahan N, McCombie L, et al. Predictors of type 2 diabetes remission in the Diabetes Remission Clinical Trial (DiRECT). Diabet Med. 2021; 38(8): e14395. Epub 2020/09/02. https://doi.org/10.1111/dme.14395 PMID: 32870520.
- Cummings DE, Arterburn DE, Westbrook EO, Kuzma JN, Stewart SD, Chan CP, et al. Gastric bypass surgery vs intensive lifestyle and medical intervention for type 2 diabetes: the CROSSROADS randomised controlled trial. Diabetologia. 2016; 59(5):945–53. Epub 2016/03/18. https://doi.org/10.1007/ s00125-016-3903-x PMID: 26983924.
- Gregg EW, Chen H, Wagenknecht LE, Clark JM, Delahanty LM, Bantle J, et al. Association of an intensive lifestyle intervention with remission of type 2 diabetes. JAMA: Journal of the American Medical Association. 2012; 308(23):2489–96. https://doi.org/10.1001/jama.2012.67929 PMID: 23288372. Corporate Author: Look AHEAD Research Group. Language: English. Entry Date: 20130201. Revision Date: 20211029. Publication Type: journal article.
- Lean ME, Leslie WS, Barnes AC, Brosnahan N, Thom G, McCombie L, et al. Primary care-led weight management for remission of type 2 diabetes (DiRECT): an open-label, cluster-randomised trial. Lancet (London, England). 2018; 391(10120):541–51. https://doi.org/10.1016/S0140-6736(17)33102-1 PMID: 29221645.
- Lean MEJ, Leslie WS, Barnes AC, Brosnahan N, Thom G, McCombie L, et al. Durability of a primary care-led weight-management intervention for remission of type 2 diabetes: 2-year results of the DiRECT open-label, cluster-randomised trial. The Lancet Diabetes & Endocrinology. 2019; 7(5):344–55. https:// doi.org/10.1016/S2213-8587(19)30068-3 PMID: 30852132
- Foley PJ. Effect of low carbohydrate diets on insulin resistance and the metabolic syndrome. Curr Opin Endocrinol Diabetes Obes. 2021; 28(5):463–8. Epub 2021/09/02. https://doi.org/10.1097/MED. 000000000000659 PMID: 34468401 Media, for which he receives financial compensation. He declares no other conflicts of interest.
- 14. Taylor R, Barnes A. Can type 2 diabetes be reversed and how can this best be achieved? James Lind Alliance research priority number one. Diabetic Medicine. 2019; 36(3):308–15. https://doi.org/10.1111/ dme.13851 PMID: 30378706

- Buse JB, Caprio S, Cefalu WT, Ceriello A, Del Prato S, Inzucchi SE, et al. How do we define cure of diabetes? Diabetes care. 2009; 32(11):2133–5. <u>https://doi.org/10.2337/dc09-9036</u> PMID: 19875608
- 16. DiabetesAustralia. POSITION STATEMENT (OCTOBER 2021) Type 2 diabetes remission. 2021.
- 17. Nagi D, Hambling C, Taylor R. Remission of type 2 diabetes: a position statement from the Association of British Clinical Diabetologists (ABCD) and the Primary Care Diabetes Society (PCDS). British Journal of Diabetes. 2019; 19(1):73–6.
- Riddle MC, Cefalu WT, Evans PH, Gerstein HC, Nauck MA, Oh WK, et al. Consensus Report: Definition and Interpretation of Remission in Type 2 Diabetes. Diabetes Care. 2021; 44(10):2438–44. Epub 2021/ 09/01. https://doi.org/10.2337/dci21-0034 PMID: 34462270.
- 19. Martikainen P, Bartley M, Lahelma E. Psychosocial determinants of health in social epidemiology. Oxford University Press; 2002. p. 1091–3.
- Bell R. Psychosocial pathways and health outcomes: Informing action on health inequalities. The Institute of Health Equity, 2017.
- 21. Tharek Z, Ramli AS, Whitford DL, Ismail Z, Mohd Zulkifli M, Ahmad Sharoni SK, et al. Relationship between self-efficacy, self-care behaviour and glycaemic control among patients with type 2 diabetes mellitus in the Malaysian primary care setting. BMC Fam Pract. 2018; 19(1):39. Epub 2018/03/11. https://doi.org/10.1186/s12875-018-0725-6 PMID: 29523075.
- Kato A, Fujimaki Y, Fujimori S, Izumida Y, Suzuki R, Ueki K, et al. A qualitative study on the impact of internalized stigma on type 2 diabetes self-management. Patient Educ Couns. 2016; 99(7):1233–9. Epub 2016/11/23. https://doi.org/10.1016/j.pec.2016.02.002 PMID: 27873575.
- Saad AMJ, Younes ZMH, Ahmed H, Brown JA, Al Owesie RM, Hassoun AAK. Self-efficacy, self-care and glycemic control in Saudi Arabian patients with type 2 diabetes mellitus: A cross-sectional survey. Diabetes Res Clin Pract. 2018; 137:28–36. Epub 2017/12/27. <u>https://doi.org/10.1016/j.diabres.2017</u>. 12.014 PMID: 29278711.
- 24. Gao J, Wang J, Zheng P, Haardörfer R, Kegler MC, Zhu Y, et al. Effects of self-care, self-efficacy, social support on glycemic control in adults with type 2 diabetes. BMC family practice. 2013; 14(1):1–6. https://doi.org/10.1186/1471-2296-14-66 PMID: 23705978
- Indelicato L, Dauriz M, Santi L, Bonora F, Negri C, Cacciatori V, et al. Psychological distress, self-efficacy and glycemic control in type 2 diabetes. Nutr Metab Cardiovasc Dis. 2017; 27(4):300–6. Epub 2017/03/10. https://doi.org/10.1016/j.numecd.2017.01.006 PMID: 28274728.
- Song Y, Song HJ, Han HR, Park SY, Nam S, Kim MT. Unmet needs for social support and effects on diabetes self-care activities in Korean Americans with type 2 diabetes. Diabetes Educ. 2012; 38(1):77– 85. Epub 2012/01/10. https://doi.org/10.1177/0145721711432456 PMID: 22222514.
- Tang TS, Brown MB, Funnell MM, Anderson RM. Social support, quality of life, and self-care behaviors amongAfrican Americans with type 2 diabetes. Diabetes Educ. 2008; 34(2):266–76. Epub 2008/04/01. https://doi.org/10.1177/0145721708315680 PMID: 18375776.
- Zhang X, Norris SL, Gregg EW, Beckles G. Social support and mortality among older persons with diabetes. Diabetes Educ. 2007; 33(2):273–81. Epub 2007/04/12. https://doi.org/10.1177/0145721707299265 PMID: 17426302.
- Shan R, Sarkar S, Martin SS. Digital health technology and mobile devices for the management of diabetes mellitus: state of the art. Diabetologia. 2019; 62(6):877–87. Epub 2019/04/10. https://doi.org/10.1007/s00125-019-4864-7 PMID: 30963188.
- Tefera YG, Gebresillassie BM, Emiru YK, Yilma R, Hafiz F, Akalu H, et al. Diabetic health literacy and its association with glycemic control among adult patients with type 2 diabetes mellitus attending the outpatient clinic of a university hospital in Ethiopia. PLoS One. 2020; 15(4):e0231291. Epub 2020/04/ 09. https://doi.org/10.1371/journal.pone.0231291 PMID: 32267893.
- Shigaki C, Kruse RL, Mehr D, Sheldon KM, Bin G, Moore C, et al. Motivation and diabetes self-management. Chronic Illn. 2010; 6(3):202–14. Epub 2010/08/03. <u>https://doi.org/10.1177/1742395310375630</u> PMID: 20675362.
- Swanson V, Maltinsky W. Motivational and behaviour change approaches for improving diabetes management. Practical Diabetes. 2019; 36(4):121–5. https://doi.org/10.1002/pdi.2229
- Polonsky WH, Anderson BJ, Lohrer PA, Welch G, Jacobson AM, Aponte JE, et al. Assessment of Diabetes-Related Distress. Diabetes Care. 1995; 18(6):754–60. <u>https://doi.org/10.2337/diacare.18.6.754</u> PMID: 7555499
- Polonsky WH, Fisher L, Earles J, Dudl RJ, Lees J, Mullan J, et al. Assessing Psychosocial Distress in Diabetes: Development of the Diabetes Distress Scale. Diabetes Care. 2005; 28(3):626–31. https://doi. org/10.2337/diacare.28.3.626 PMID: 15735199

- Bijl JV, Poelgeest-Eeltink AV, Shortridge-Baggett L. The psychometric properties of the diabetes management self-efficacy scale for patients with type 2 diabetes mellitus. J Adv Nurs. 1999; 30(2):352–9. Epub 1999/08/24. https://doi.org/10.1046/j.1365-2648.1999.01077.x PMID: 10457237.
- 36. Speight J, Skinner TC, Huber JW, Lake AJ, Messina R, Mocan A, et al. A PSAD Group response to the consensus report on the definition and interpretation of remission in type 2 diabetes: a psychosocial perspective is needed. Diabetologia. 2022; 65(2):406–8. Epub 2021/11/16. <u>https://doi.org/10.1007/s00125-021-05615-z</u> PMID: 34778926.
- Nicolucci A, Kovacs Burns K, Holt RI, Comaschi M, Hermanns N, Ishii H, et al. Diabetes Attitudes, Wishes and Needs second study (DAWN2): cross-national benchmarking of diabetes-related psychosocial outcomes for people with diabetes. Diabet Med. 2013; 30(7):767–77. Epub 2013/05/29. https:// doi.org/10.1111/dme.12245 PMID: 23711019.
- Whicher CA, O'Neill S, Holt RIG. Diabetes in the UK: 2019. Diabet Med. 2020; 37(2):242–7. Epub 2020/ 01/05. https://doi.org/10.1111/dme.14225 PMID: 31901175.
- Wylie TAF, Shah C, Connor R, Farmer AJ, Ismail K, Millar B, et al. Transforming mental well-being for people with diabetes: research recommendations from Diabetes UK's 2019 Diabetes and Mental Well-Being Workshop. Diabet Med. 2019; 36(12):1532–8. Epub 2019/06/10. https://doi.org/10.1111/dme. 14048 PMID: 31177573.
- Peters MDJ GC, McInerney P, Munn Z, Tricco AC, Khalil H. JBI Manual for Evidence Synthesis Chapter 11: Scoping reviews (2020 version). Aromataris E MZ, editor. Adelaide2020.
- Peters MDJ, Marnie C, Tricco AC, Pollock D, Munn Z, Alexander L, et al. Updated methodological guidance for the conduct of scoping reviews. JBI Evid Synth. 2020; 18(10):2119–26. Epub 2020/10/11. https://doi.org/10.11124/JBIES-20-00167 PMID: 33038124.
- Arksey H, O'Malley L. Scoping studies: towards a methodological framework. International Journal of Social Research Methodology. 2005; 8(1):19–32. https://doi.org/10.1080/1364557032000119616
- Tricco AC L E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. Annals of Internal Medicine. 2018; 169(7):467–73. https://doi.org/10.7326/M18-0850 PMID: 30178033
- Captieux M, Prigge R, Wild S, Guthrie B. Defining remission of type 2 diabetes in research studies: A systematic scoping review. PLoS Med. 2020; 17(10):e1003396. Epub 2020/10/29. <u>https://doi.org/10. 1371/journal.pmed.1003396 PMID: 33112845.</u>
- 45. Goldenberg JZ, Day A, Brinkworth GD, Sato J, Yamada S, Jonsson T, et al. Efficacy and safety of low and very low carbohydrate diets for type 2 diabetes remission: systematic review and meta-analysis of published and unpublished randomized trial data. BMJ. 2021; 372:m4743. Epub 2021/01/15. https:// doi.org/10.1136/bmj.m4743 PMID: 33441384.
- 46. Breuing J, Pieper D, Neuhaus AL, Hess S, Lutkemeier L, Haas F, et al. Barriers and facilitating factors in the prevention of diabetes type 2 and gestational diabetes in vulnerable groups: A scoping review. PLoS One. 2020; 15(5):e0232250. Epub 2020/05/14. https://doi.org/10.1371/journal.pone.0232250 PMID: 32401778.
- 47. Team TE. EndNote. EndNote X9 ed. Philadelphia, PA: Clarivate Analytics 2013.
- Ades PA, Savage PD, Marney AM, Harvey J, Evans KA. Remission of recently diagnosed type 2 diabetes mellitus with weight loss and exercise. Journal of Cardiopulmonary Rehabilitation & Prevention. 2015; 35(3):193–7. https://doi.org/10.1097/HCR.00000000000106 PMID: 25636149. Language: English. Entry Date: 20150923. Revision Date: 20150923. Publication Type: Journal Article. Journal Subset: Allied Health.
- 49. Athinarayanan SJ, Adams RN, Hallberg SJ, McKenzie AL, Bhanpuri NH, Campbell WW, et al. Long-Term Effects of a Novel Continuous Remote Care Intervention Including Nutritional Ketosis for the Management of Type 2 Diabetes: A 2-Year Non-randomized Clinical Trial. Frontiers in endocrinology. 2019; 10:348. https://doi.org/10.3389/fendo.2019.00348 PMID: 31231311.
- Bhatt AA, Choudhari PK, Mahajan RR, Sayyad MG, Pratyush DD, Hasan I, et al. Effect of a Low-Calorie Diet on Restoration of Normoglycemia in Obese subjects with Type 2 Diabetes. Indian journal of endocrinology and metabolism. 2017; 21(5):776–80. https://doi.org/10.4103/ijem.IJEM_206_17 PMID: 28989891.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. Bmj. 2021. https://doi.org/10.1136/ bmj.n71 PMID: 33782057
- 52. Marples O, Resca L, Plavska J, Hassan S, Mistry V, Mallik R, et al. Real-World Data of a Group-Based Formula Low Energy Diet Programme in Achieving Type 2 Diabetes Remission and Weight Loss in an Ethnically Diverse Population in the UK: A Service Evaluation. Nutrients. 2022; 14(15):3146. <u>https://doi.org/10.3390/nu14153146 PMID: 35956322</u>
- Oser TK, Cucuzzella M, Stasinopoulos M, Moncrief M, McCall A, Cox DJ. An Innovative, Paradigm-Shifting Lifestyle Intervention to Reduce Glucose Excursions With the Use of Continuous Glucose

Monitoring to Educate, Motivate, and Activate Adults With Newly Diagnosed Type 2 Diabetes: Pilot Feasibility Study. JMIR Diabetes. 2022; 7(1):e34465. https://doi.org/10.2196/34465 PMID: 35050857

- 54. Yancy WS Jr, Crowley MJ, Dar MS, Coffman CJ, Jeffreys AS, Maciejewski ML, et al. Comparison of Group Medical Visits Combined With Intensive Weight Management vs Group Medical Visits Alone for Glycemia in Patients With Type 2 Diabetes: A Noninferiority Randomized Clinical Trial. JAMA Internal Medicine. 2020; 180(1):70–9. https://doi.org/10.1001/jamainternmed.2019.4802 PMID: 31682682
- Zou W, Luo K, Hu Z, Zhang X, Feng C, Ye D, et al. Diabetes remission after a lifestyle-medicine intervention on type 2 diabetes in lean and obese Chinese subjects: a prospective study. Ann Palliat Med. 2022; 11(4):1462–72. Epub 2022/05/07. https://doi.org/10.21037/apm-22-369 PMID: 35523754.
- 56. Evans TS, Hawkes RE, Keyworth C, Newson L, Radley D, Hill AJ, et al. How is the NHS Low-Calorie Diet Programme expected to produce behavioural change to support diabetes remission: An examination of underpinning theory. Br J Diabetes. 2022; 22(1):20–9. Epub 2022/09/02. <u>https://doi.org/10. 15277/bjd.2022.341</u> PMID: 36045887.
- 57. Taheri S, Zaghloul H, Chagoury O, Elhadad S, Ahmed SH, El Khatib N, et al. Effect of intensive lifestyle intervention on bodyweight and glycaemia in early type 2 diabetes (DIADEM-I): an open-label, parallel-group, randomised controlled trial. The lancet Diabetes & endocrinology. 2020; 8(6):477–89. <u>https://doi.org/10.1016/S2213-8587(20)30117-0 PMID: 32445735</u>.
- 58. Umphonsathien M, Prutanopajai P, Aiam ORJ, Thararoop T, Karin A, Kanjanapha C, et al. Immediate and long-term effects of a very-low-calorie diet on diabetes remission and glycemic control in obese Thai patients with type 2 diabetes mellitus. Food Sci Nutr. 2019; 7(3):1113–22. Epub 2019/03/29. https://doi.org/10.1002/fsn3.956 PMID: 30918654.
- 59. Umphonsathien M, Rattanasian P, Lokattachariya S, Suansawang W, Boonyasuppayakorn K, Khovidhunkit W. Effects of intermittent very-low calorie diet on glycemic control and cardiovascular risk factors in obese patients with type 2 diabetes mellitus: A randomized controlled trial. Journal of diabetes investigation. 2022; 13(1):156–66. https://doi.org/10.1111/jdi.13619 PMID: 34176234.
- Cox N, Gibas S, Salisbury M, Gomer J, Gibas K. Ketogenic diets potentially reverse Type II diabetes and ameliorate clinical depression: A case study. Diabetes Metab Syndr. 2019; 13(2):1475–9. Epub 2019/07/25. https://doi.org/10.1016/j.dsx.2019.01.055 PMID: 31336509.
- Lagger G, Correia JC, Fernandes P, Sittarame F, Golay A. A 5-dimension patient education program targeting type 2 diabetes remission. Therapeutic Patient Education / Education Therapeutique du Patient. 2018; 10(2):N.PAG-N.PAG. https://doi.org/10.1051/tpe/2018018 PMID: 133120197. Language: English. Entry Date: 20181123. Revision Date: 20200205. Publication Type: Article.
- Rehackova L, Araújo-Soares V, Adamson AJ, Steven S, Taylor R, Sniehotta FF. Acceptability of a verylow-energy diet in Type 2 diabetes: patient experiences and behaviour regulation. Diabetic medicine: a journal of the British Diabetic Association. 2017; 34(11):1554–67. https://doi.org/10.1111/dme.13426 PMID: 28727247.
- 63. Bynoe K, Unwin N, Taylor C, Murphy MM, Bartholomew L, Greenidge A, et al. Inducing remission of Type 2 diabetes in the Caribbean: findings from a mixed methods feasibility study of a low-calorie liquid diet-based intervention in Barbados. Diabetic medicine: a journal of the British Diabetic Association. 2020; 37(11):1816–24. https://doi.org/10.1111/dme.14096 PMID: 31365159.
- 64. Webster CC, Murphy TE, Larmuth KM, Noakes TD, Smith JA. Diet, Diabetes Status, and Personal Experiences of Individuals with Type 2 diabetes Who Self-Selected and Followed a Low Carbohydrate High Fat diet. Diabetes, metabolic syndrome and obesity: targets and therapy. 2019; 12:2567–82. https://doi.org/10.2147/DMSO.S227090 PMID: 31827331.
- 65. Rehackova L, Araújo-Soares V, Steven S, Adamson AJ, Taylor R, Sniehotta FF. Behaviour change during dietary Type 2 diabetes remission: a longitudinal qualitative evaluation of an intervention using a very low energy diet. Diabetic medicine: a journal of the British Diabetic Association. 2020; 37(6):953–62. https://doi.org/10.1111/dme.14066 PMID: 31269276.
- 66. Rehackova L, Rodrigues AM, Thom G, Brosnahan N, Barnes AC, McCombie L, et al. Participant experiences in the Diabetes REmission Clinical Trial (DiRECT). Diabetic Medicine. 2022; 39(1):1–12. https://doi.org/10.1111/dme.14689 PMID: 154218938. Language: English. Entry Date: 20211230. Revision Date: 20211230. Publication Type: Article.
- Chen G, Olsen JA. Filling the psycho-social gap in the EQ-5D: the empirical support for four bolt-on dimensions. Qual Life Res. 2020; 29(11):3119–29. Epub 2020/07/11. <u>https://doi.org/10.1007/s11136-020-02576-5 PMID: 32648198.</u>
- 68. Taylor R, Barnes AC. Can type 2 diabetes be reversed and how can this best be achieved? James Lind Alliance research priority number one. Diabet Med. 2019; 36(3):308–15. Epub 2018/11/01. https://doi. org/10.1111/dme.13851 PMID: 30378706.
- 69. Bishop J, Irby MB, Isom S, Blackwell CS, Vitolins MZ, Skelton JA. Diabetes prevention, weight loss, and social support: program participants' perceived influence on the health behaviors of their social support

system. Fam Community Health. 2013; 36(2):158–71. Epub 2013/03/05. https://doi.org/10.1097/FCH. 0b013e318282b2d3 PMID: 23455686.

- 70. Frost H, Campbell P, Maxwell M, O'Carroll RE, Dombrowski SU, Williams B, et al. Effectiveness of Motivational Interviewing on adult behaviour change in health and social care settings: A systematic review of reviews. PLoS One. 2018; 13(10):e0204890. Epub 2018/10/20. https://doi.org/10.1371/journal.pone. 0204890 PMID: 30335780.
- Jiang X, Wang J, Lu Y, Jiang H, Li M. Self-efficacy-focused education in persons with diabetes: a systematic review and meta-analysis. Psychol Res Behav Manag. 2019; 12:67–79. Epub 2019/02/19. https://doi.org/10.2147/PRBM.S192571 PMID: 30774486.
- Young-Hyman D, de Groot M, Hill-Briggs F, Gonzalez JS, Hood K, Peyrot M. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. Diabetes Care. 2016; 39(12):2126–40. Epub 2016/11/24. https://doi.org/10.2337/dc16-2053 PMID: 27879358.
- Healy P, Devane D. Methodological considerations in cohort study designs. Nurse Res. 2011; 18 (3):32–6. Epub 2011/05/13. https://doi.org/10.7748/nr2011.04.18.3.32.c8461 PMID: 21560924.
- 74. Crowe S, Cresswell K, Robertson A, Huby G, Avery A, Sheikh A. The case study approach. BMC Med Res Methodol. 2011; 11:100. Epub 2011/06/29. https://doi.org/10.1186/1471-2288-11-100 PMID: 21707982.
- 75. Diefenbach T. Are case studies more than sophisticated storytelling?: Methodological problems of qualitative empirical research mainly based on semi-structured interviews. Quality & Quantity. 2008; 43 (6):875–94. https://doi.org/10.1007/s11135-008-9164-0
- 76. Evans SR. Clinical trial structures. Journal of experimental stroke & translational medicine. 2010; 3 (1):8. https://doi.org/10.6030/1939-067x-3.1.8 PMID: 21423788
- Hariton E, Locascio JJ. Randomised controlled trials—the gold standard for effectiveness research: Study design: randomised controlled trials. BJOG. 2018; 125(13):1716. Epub 2018/06/20. <u>https://doi.org/10.1111/1471-0528.15199</u> PMID: 29916205.
- Dave R, Davis R, Davies JS. The impact of multiple lifestyle interventions on remission of type 2 diabetes mellitus within a clinical setting. Obesity Medicine. 2019; 13:59–64. <u>https://doi.org/10.1016/j.obmed.</u> 2019.01.005
- 79. de Hoogh IM, Oosterman JE, Otten W, Krijger A-M, Berbée-Zadelaar S, Pasman WJ, et al. The Effect of a Lifestyle Intervention on Type 2 Diabetes Pathophysiology and Remission: The Stevenshof Pilot Study. Nutrients. 2021; 13(7):2193-. https://doi.org/10.3390/nu13072193 PMID: 34202194. Language: English. Entry Date: 20210804. Revision Date: 20210805. Publication Type: Article.
- Ried-Larsen M, Johansen MY, MacDonald CS, Hansen KB, Christensen R, Wedell-Neergaard A-S, et al. Type 2 diabetes remission 1 year after an intensive lifestyle intervention: A secondary analysis of a randomized clinical trial. Diabetes, obesity & metabolism. 2019; 21(10):2257–66. <u>https://doi.org/10.1111/dom.13802</u> PMID: 31168922.
- Steven S, Hollingsworth KG, Al-Mrabeh A, Avery L, Aribisala B, Caslake M, et al. Very low-calorie diet and 6 months of weight stability in type 2 diabetes: Pathophysiological changes in responders and nonresponders. Diabetes Care. 2016; 39(5):808–15. https://doi.org/10.2337/dc15-1942 PMID: 27002059
- McCoy MA, Theeke LA. A systematic review of the relationships among psychosocial factors and coping in adults with type 2 diabetes mellitus. Int J Nurs Sci. 2019; 6(4):468–77. Epub 2019/11/16. https:// doi.org/10.1016/j.ijnss.2019.09.003 PMID: 31728402.
- Thorpe CT, Fahey LE, Johnson H, Deshpande M, Thorpe JM, Fisher EB. Facilitating healthy coping in patients with diabetes: a systematic review. Diabetes Educ. 2013; 39(1):33–52. Epub 2012/10/18. https://doi.org/10.1177/0145721712464400 PMID: 23073967.
- Rogerson D, Soltani H, Copeland R. The weight-loss experience: a qualitative exploration. BMC Public Health. 2016; 16:371. Epub 2016/05/05. https://doi.org/10.1186/s12889-016-3045-6 PMID: 27142984.
- Steinberg DM, Tate DF, Bennett GG, Ennett S, Samuel-Hodge C, Ward DS. The efficacy of a daily selfweighing weight loss intervention using smart scales and e-mail. Obesity (Silver Spring). 2013; 21 (9):1789–97. Epub 2013/03/21. https://doi.org/10.1002/oby.20396 PMID: 23512320.
- **86.** Fisher WA, Cornman DH, Kohut T, Schachner H, Stenger P. What primary care providers can do to address barriers to self-monitoring of blood glucose. Clinical Diabetes. 2013; 31(1):34–42.
- Griauzde DH, Ling G, Wray D, DeJonckheere M, Mizokami Stout K, Saslow LR, et al. Continuous Glucose Monitoring With Low-Carbohydrate Nutritional Coaching to Improve Type 2 Diabetes Control: Randomized Quality Improvement Program. J Med Internet Res. 2022; 24(2):e31184. Epub 2022/02/ 03. https://doi.org/10.2196/31184 PMID: 35107429.
- 88. NICE. Type 2 diabetes in adults: management NICE guideline [NG28] 2022 [updated 29 June 2022].
- **89.** Michie S, Ashford S, Sniehotta FF, Dombrowski SU, Bishop A, French DP. A refined taxonomy of behaviour change techniques to help people change their physical activity and healthy eating

behaviours: the CALO-RE taxonomy. Psychol Health. 2011; 26(11):1479–98. Epub 2011/06/17. https://doi.org/10.1080/08870446.2010.540664 PMID: 21678185.

- 90. Rehackova L, Arnott B, Araujo-Soares V, Adamson AA, Taylor R, Sniehotta FF. Efficacy and acceptability of very low energy diets in overweight and obese people with Type 2 diabetes mellitus: a systematic review with meta-analyses. Diabet Med. 2016; 33(5):580–91. Epub 2015/10/23. <u>https://doi.org/10. 1111/dme.13005 PMID: 26490082</u>.
- Unwin D, Khalid AA, Unwin J, Crocombe D, Delon C, Martyn K, et al. Insights from a general practice service evaluation supporting a lower carbohydrate diet in patients with type 2 diabetes mellitus and prediabetes: a secondary analysis of routine clinic data including HbA1c, weight and prescribing over 6 years. BMJ nutrition, prevention & health. 2020; 3(2):285–94. https://doi.org/10.1136/bmjnph-2020-000072 PMID: 33521540.
- Venker BT, Dunn JP, Stephenson K. The Rising Cost of Diabetes: Can DiRECT Tip the Scales? Obesity (Silver Spring). 2018; 26(12):1866–7. Epub 2018/11/13. https://doi.org/10.1002/oby.22349 PMID: 30417975.
- Xin Y, Davies A, McCombie L, Briggs A, Messow CM, Grieve E, et al. Type 2 diabetes remission: economic evaluation of the DiRECT/Counterweight-Plus weight management programme within a primary care randomized controlled trial. Diabet Med. 2019; 36(8):1003–12. Epub 2019/04/27. https://doi.org/ 10.1111/dme.13981 PMID: 31026353.
- 94. Holman N, Khunti K, Wild SH, Sattar N, Knighton P, Young B, et al. Care processes in people in remission from type 2 diabetes: A cohort study using the National Diabetes Audit. Diabet Med. 2023; 40(3): e15016. Epub 2022/11/29. https://doi.org/10.1111/dme.15016 PMID: 36440921.
- **95.** Jacob E, Avery A. Energy-restricted interventions are effective for the remission of newly diagnosed type 2 diabetes: A systematic review of the evidence base. Obes Sci Pract. 2021; 7(5):606–18. Epub 2021/10/12. https://doi.org/10.1002/osp4.504 PMID: 34631138.
- 96. Snorgaard O, Poulsen GM, Andersen HK, Astrup A. Systematic review and meta-analysis of dietary carbohydrate restriction in patients with type 2 diabetes. BMJ Open Diabetes Res Care. 2017; 5(1): e000354. Epub 2017/03/21. https://doi.org/10.1136/bmjdrc-2016-000354 PMID: 28316796.
- 97. van Zuuren EJ, Fedorowicz Z, Kuijpers T, Pijl H. Effects of low-carbohydrate- compared with low-fatdiet interventions on metabolic control in people with type 2 diabetes: a systematic review including GRADE assessments. Am J Clin Nutr. 2018; 108(2):300–31. Epub 2018/07/15. <u>https://doi.org/10.1093/ ajcn/ngy096</u> PMID: 30007275.