

Central Lancashire Online Knowledge (CLOK)

Title	OTHR-17 MANAGEMENT AND OUTCOMES OF SELLAR, SUPRASELLAR, AND PARASELLAR MASSES IN LOW- AND MIDDLE-INCOME COUNTRIES: A SCOPING REVIEW AND PROPORTIONAL META-ANALYSIS OF 6584 MASSES
Type	Article
URL	https://clock.uclan.ac.uk/53199/
DOI	https://doi.org/10.1093/noajnl/vdae147.028
Date	2024
Citation	Takoutsing, Berjo Dongmo, Egiz, Abdullah Mohammed abousaleh ma, Dalle, David Ulrich, Dada, Olaoluwa Ezekiel, Ikwuegbuenyi, Chibuikem A, Ooi, Setthasorn Zhi Yang, Gillespie, Conor S, Tunde, Olobatoke A, Yong, Brian Ming Ou et al (2024) OTHR-17 MANAGEMENT AND OUTCOMES OF SELLAR, SUPRASELLAR, AND PARASELLAR MASSES IN LOW- AND MIDDLE-INCOME COUNTRIES: A SCOPING REVIEW AND PROPORTIONAL META-ANALYSIS OF 6584 MASSES. <i>Neuro-Oncology Advances</i> , 6 (Sup2). ii7-ii7. ISSN 2632-2498
Creators	Takoutsing, Berjo Dongmo, Egiz, Abdullah Mohammed abousaleh ma, Dalle, David Ulrich, Dada, Olaoluwa Ezekiel, Ikwuegbuenyi, Chibuikem A, Ooi, Setthasorn Zhi Yang, Gillespie, Conor S, Tunde, Olobatoke A, Yong, Brian Ming Ou, Korch, Moniba, Kesici, Özgür, de Koning, Rosaline, Khan, Mehdi, Erhabor, Joshua, Kotecha, Jay, Bocco, Alberic Fabrice Sewa, Bandyopadhyay, Soham and Bankole, Nourou Dine Adeniran

It is advisable to refer to the publisher's version if you intend to cite from the work.
<https://doi.org/10.1093/noajnl/vdae147.028>

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

sidered to be the gold standard means of diagnosis for these masses, their use in LMICs is less reported. Despite new minimally invasive treatment options, endoscopic surgery is commonly used in LMICs. These translates to the fact that more effort is required in the effective care of patients with these masses in LMICs.

Abstract citation ID: vdae147.028

OTHR-17

MANAGEMENT AND OUTCOMES OF SELLAR, SUPRASELLAR, AND PARASELLAR MASSES IN LOW- AND MIDDLE-INCOME COUNTRIES: A SCOPING REVIEW AND PROPORTIONAL META-ANALYSIS OF 6584 MASSES

Berjo Dongmo Takoutsing^{1,2}, Abdullah Egiz³, David Ulrich Dalle^{4,1}, Olaoluwa Ezekiel Dada^{5,1}, Chibuikem A. Ikwuegbuenyi¹, Setthasorn Zhi Yang Ooi⁶, Conor S. Gillespie⁷, Olobatoke A. Tunde^{8,1}, Brian Ming Ou Yong⁹, Moniba Korch¹⁰, Özgür Kesici¹¹, Rosaline de Koning¹², Mehdi Khan¹³, Joshua Erhabor¹⁴, Jay Kotecha¹⁵, Alberic Fabrice Sewa Bocco¹⁶, Soham Bandyopadhyay^{17,18,19}, Nourou Dine Adeniran Bankole²⁰; ¹Association of Future African Neurosurgeons, Yaounde, Cameroon. ²Winners Foundation, Yaounde, Cameroon. ³School of Medicine, University of Central Lancashire, Preston, United Kingdom. ⁴North Ossetia State Medical Academy, Vladikavkaz, Russian Federation. ⁵College of Medicine, University of Ibadan, Ibadan, Nigeria. ⁶Cardiff University School of Medicine, University Hospital of Wales Main Building, Cardiff, United Kingdom. ⁷Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool, United Kingdom. ⁸College of Medicine University of Lagos, Lagos, Nigeria. ⁹School of Medicine, International Medical University, Kuala Lumpur, Malaysia. ¹⁰Cadi Ayyad University, Faculty of Medicine and Pharmacy, Marrakech, Morocco. ¹¹Ankara University, School of Medicine, Ankara, Turkey. ¹²University of Oxford Medical Sciences Division, Oxford, United Kingdom. ¹³UCL Medical School, London, United Kingdom. ¹⁴University of Exeter Medical School, Exeter, United Kingdom. ¹⁵Leicester Royal Infirmary, Leicester, United Kingdom. ¹⁶Department of Neurosurgery, Ibn Rochd University Hospital, Casablanca, Morocco. ¹⁷Clinical Neurosciences, Clinical & Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton, United Kingdom. ¹⁸Wessex Neurological Centre, University Hospital Southampton NHS Foundation Trust, Southampton, United Kingdom. ¹⁹Oxford University Global Surgery Group, Nuffield Department of Surgical Sciences, University of Oxford, Oxford, United Kingdom. ²⁰Clinical Investigation Center (CIC), 1415, INSERM, Department of Interventional Neuroradiology, Tours University Hospital, Tours, France

OBJECTIVE: To aggregate data on treatment modalities, management approaches, and clinical outcomes of sellar, suprasellar, and parasellar masses in LMICs. **METHODS:** We conducted a scoping review as per the Arksey and O'Malley framework. MEDLINE, Embase, Global Index Medicus, and African Journals Online were searched and included articles between 2000 and 2021 screened against predefined eligibility criteria. Pooled statistics were calculated using measures of central tendency and spread. A proportional meta-analysis was conducted to pool the mortality rates. **RESULTS:** Of the 3526 articles generated by the search strategy, 173 articles were included. The mean age was 35.5 ± 15 years, and most were females (56.47%). Sellar masses predominated (85%; 95% CI = 93.16–118.58), and the most commonly used neuroimaging modality to diagnose these masses was an MRI (65.31%). Surgical resection was mostly utilised with endoscopic surgery (78%; 95% CI = 17.29–154.96) predominating over microsurgery (22%; 95% CI = 11.51–47.73). The transsphenoidal approach was mostly used for both endoscopic surgery and microsurgery (92.13% vs 93.21%), and the extent of resection was gross total resection in most cases ($n = 3611$). Non-surgical management included hormonal therapies ($n = 2080$), chemotherapy ($n = 96$ patients), and radiosurgery ($n = 357$). New onset diabetes insipidus (34.27%), followed by postoperative infection (27.95%), were the most commonly reported postoperative complications. The pooled overall mortality rate was 8.1% (95% CI: 0.031–0.146). **CONCLUSION:** Although MRIs are con-