

Central Lancashire Online Knowledge (CLOK)

Title	Forelimb muscle activity during level and progressive incline and decline walking in dogs and implications for rehabilitation
Type	Article
URL	https://clock.uclan.ac.uk/id/eprint/56547/
DOI	
Date	2025
Citation	Cain, Rachel Mariël, Stanford, Kate, Drum, Marti, Richards, James, Levine, David, Millis, Darryl and Ursini, Tena (2025) Forelimb muscle activity during level and progressive incline and decline walking in dogs and implications for rehabilitation. <i>Frontiers in Veterinary Science</i> .
Creators	Cain, Rachel Mariël, Stanford, Kate, Drum, Marti, Richards, James, Levine, David, Millis, Darryl and Ursini, Tena

It is advisable to refer to the publisher's version if you intend to cite from the work.

For information about Research at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLOK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

Forelimb muscle activity during level and progressive incline and decline walking in dogs and implications for rehabilitation

1 Cain, Rachel Mariël^{1*}; Stanford, Kate^{1*}; Drum, Marti¹; Richards, Jim³; Levine, David²; Millis,
2 Darryl¹; and Ursini, Tena⁴

3 ¹ Department of Small Animal Clinical Sciences, University of Tennessee College of Veterinary
4 Medicine, Knoxville, TN, United States

5 ²Department of Physical Therapy, University of Tennessee at Chattanooga, Chattanooga, TN, United
6 States

7 ³Allied Health Research Unit, University of Lancashire, Lancashire, United Kingdom

8 ⁴Department of Large Animal Clinical Sciences, College of Veterinary Medicine, Equine
9 Performance and Rehabilitation Center, University of Tennessee, Knoxville, TN, United States

10 * Correspondence:

11 Tena Ursini
12 tursini@utk.edu

13 **Keywords: electromyography¹, rehabilitation², supraspinatus³, biceps brachii⁴, triceps brachii⁵,**
14 **deltoideus⁶, incline walk⁷, decline walk⁸**

15 Abstract

16 Introduction – Shoulder pathologies are a common reason for presentation to veterinary sports
17 medicine and rehabilitation practices. Currently there are no standardized rehabilitation protocols for
18 shoulder injuries but controlled walking, either on flat ground or on inclines/declines, is
19 recommended in nearly every recovery rehabilitation program. The objective of this study was to
20 evaluate the peak and average muscle activity of commonly targeted forelimb muscle groups using
21 fine-wire and surface electromyography (EMG) during treadmill walking at five treadmill positions.
22 Our hypothesis was that the forelimb muscle activity would be significantly higher during decline
23 walking than level walking and significantly lower during incline walking.

24 Methods - Fine-wire and surface EMG of the supraspinatus, deltoideus, biceps brachii and lateral
25 head of the triceps brachii muscles were performed during treadmill walking at 0%, 5% incline, 10%
26 incline, 5% decline, and 10% decline. The average and peak muscle enveloped EMG signals for 10
27 gait cycles were compared between the treadmill positions.

28 Results - Significant main effects were seen during decline walking for average supraspinatus muscle
29 activity ($p < 0.001$), and both average and peak deltoideus muscle activity ($p = 0.021$, $p < 0.001$)
30 respectively. There were no significant differences for peak or average lateral triceps brachii or
31 biceps brachii activity between treadmill positions.

32 Conclusion - Decline walking significantly increases muscle activity in the supraspinatus and
33 deltoideus muscles of dogs. This study provides new insights regarding the muscle activity of the
34 thoracic limbs in dogs during various treadmill positions at the walk. The impact of incremental
35 treadmill positions on the average supraspinatus EMG activity, along with increases in the average

and peak EMG activity of the deltoideus muscle during decline walking should be considered when developing a therapeutic exercise plan in canine patients with shoulder injuries.

1 Introduction

Shoulder pathologies are increasingly recognized as causes of lameness and functional impairment, particularly in sporting and working dogs (1, 2). Owner-based surveys, observational, and retrospective studies have identified the shoulder as a common joint injured in agility, flyball, racing, canicross, and sled dogs (3-8). Consequently, canine sports medicine and rehabilitation practitioners frequently diagnose and treat shoulder conditions in companion, sporting and working dog patients.

Supraspinatus and bicipital tendinopathies are two commonly identified shoulder ailments of dogs, and both are often managed conservatively with rehabilitation (1-3). In almost every rehabilitation program, walking appears to be a valuable therapeutic exercise despite its simplicity (2, 9). Walking exercises on level ground, inclines, and declines are commonly used in canine rehabilitation for a wide variety of conditions such as hip dysplasia (10), stifle injuries (11, 12), biceps tenosynovitis (13), and neurological conditions such as intervertebral disc disease (14, 15). Performing rehabilitation exercises on treadmills may provide unique advantages during exercise including precise control over velocity and the amount of incline or decline. Treadmills also reduce external influences such as variable terrain or environmental factors and can also be used for gait analysis (16-18).

Walking on incline or decline surfaces involves adjustments of the head, trunk, and limbs and has been reported to alter muscle activity in dogs. Surface EMG of dogs walking on an inclined treadmill at 5% (20), and 7% (21) increased activity of the biceps femoris and middle gluteal muscles compared with level walking and decline walking (20, 21). Although there is some data on uphill and downhill walking, no comparisons of the incremental effects of incline and decline walking have been reported.

Most of the canine electromyography (EMG) studies during specific therapeutic exercises have assessed hindlimb muscle activity (17, 22-24). Studies examining EMG of the forelimbs are more limited. Garcia et al (25) evaluated the muscle activity of the elbow flexor and extensor muscle groups in normal walking and trotting dogs using surface EMG to provide a reference for clinical evaluation and study of locomotor abnormalities. Janas et al (26) evaluated the muscle activity of biceps brachii and lateral triceps muscles during walking, trotting, and selected therapeutic exercises and showed the mean and peak EMG amplitude of both muscles increased during walking and trotting compared to stance. In addition, Cullen et al (27, 28) evaluated the electromyographic activity of the left biceps brachii, supraspinatus, infraspinatus, and triceps brachii muscles using fine-wire EMG while performing dynamic, highly specific agility-related tasks. However, there is a gap in the current literature on the forelimb muscle activity when considering early recovery exercises such as incline and decline treadmill walking. This EMG data may help in developing progressive therapeutic exercise programs for forelimb injuries, especially with the current prevalence of shoulder pathologies.

Inertial measurement units (IMUs) have been used as a convenient wearable method over marker based kinematic systems for human gait analysis, and several canine studies have evaluated and verified the use of IMUs for gait analysis in dogs. IMU technology seemingly provides accurate and reliable determination of stride parameters in dogs (29-32).

Objective and Hypothesis: The objective of this study was to evaluate the peak and average muscle activity of commonly targeted forelimb muscle groups using fine-wire and surface EMG during treadmill walking at five treadmill positions: 0%, 5% incline, 10% incline, 5% decline, and 10% decline. We hypothesized that the forelimb muscle activity would be significantly greater during decline walking than level walking and significantly less during incline walking.

2 Materials and Methods

All dogs had complete neurologic and orthopedic examinations performed by a board-certified sports medicine and rehabilitation veterinarian prior to enrollment; dogs were excluded if they showed any signs of visible lameness or pain upon palpation of the joints, spine, or skeletal muscles or if they had any gait abnormality at the walk or trot, posture abnormality, or any other orthopedic or neurologic conditions. To confirm that dogs had no lameness, kinetic data were obtained using a force platform (AMTI, Watertown, MA). Four valid trials for each side of the dog were obtained at a trot. Velocity and acceleration of the dog was maintained between 1.7 and 2.1 m/s and ± 0.40 m/s². Mean peak vertical force values were used to identify weight-bearing left-right asymmetry for each dog; dogs were excluded from the study if there was > 5% asymmetry between the forelimbs or hindlimbs. Eighteen dogs were screened and two were excluded due to lameness on the kinetic analysis and abnormal orthopedic examination and one dog was excluded due to abnormal orthopedic examination but had a normal kinetic analysis.

Twelve dogs were objectively and subjectively clinically sound and qualified for the study. This study was approved by the University of Tennessee Institutional Animal Care and Use Committee (protocol #3075) and was performed in accordance with AAALAC and USDA guidelines.

2.1 EMG and IMU Sensors

We evaluated muscle activity bilaterally in the supraspinatus, biceps brachii, scapular portion of deltoideus, and lateral head of triceps brachii muscles. Muscle bellies were located by manual palpation. Large areas around the sensor placement were clipped and then shaved with a razor, and the skin was cleaned with chlorhexidine scrub and isopropyl alcohol to remove oil, dirt, and debris.

Two dogs with behavioral concerns required sedation with IV low dose Dexmedetomidine for comfort during instrumentation of the EMG sensors. The remaining dogs were able to undergo sensor placement without sedation. Muscle activity was recorded from six surface EMG Avanti sensors on the skin over the middle of the left and right biceps brachii, deltoideus, and triceps brachii, and two fine wire EMG Avanti sensors (Delsys, Natick, MA, USA) which were placed into the left and right supraspinatus muscles. IMU data were also collected from the right biceps brachii for event detection (Figure 1). Supplemental documents contain the description of all sensor placements. The same two examiners were responsible for sensor placement throughout the study to ensure consistent accurate positioning. Double sided adhesive tape (Trigno Systems adhesive interfaces) and topical adhesive (Loctite Gel Control) were used to fix the sensors to the skin. For fine-wire EMG placement, a subcutaneous area over the supraspinatus site of needle insertion was desensitized with 0.25 ml of 2% lidocaine per side. Caution was taken to remain superficial to prevent alterations in muscle function as previously reported (33). Briefly, a pre-sterilized 27-gauge, 30 mm length needle with pre-loaded paired fine wire electrodes (RhythmLink, Columbia, SC) was aseptically inserted into the supraspinatus.

2.2 Exercises

The treadmill positions were selected and performed for each dog in a random order to remove any order bias. Dogs were asked to walk on a land treadmill at 3.0 km/h in the following positions: level, 5% incline, 10% incline, 5% decline and 10% decline in a randomized order, and all dogs had prior exposure to the treadmill to ensure familiarization (16, 34, 35). Familiarization was different for each dog, but most dogs were habituated after a single training session prior to enrollment in the study. The incline and decline levels of 5% and 10% respectively were transitioned with the treadmill moving, and the dog was removed from the treadmill to switch between incline and decline positions. All treadmill positions were recorded during the same test period, and after the dog had acclimatized to each new treadmill position, EMG signals were collected for 30 seconds and a minimum of 10 gait cycles at a steady walk. Left and right front leg initial contact events were identified from the IMU sensors on the right biceps brachii, and a gait cycle was defined to start at the peak vertical deceleration directly after swing phase which was identified from the sagittal plane gyroscope data.

2.3 Data Processing

The EMG signals were exported from Trigno Discover (Delsys, Natick, MA, USA) into Visual3d (HAS Motion, Kingston, CA) where they were high pass filtered with a cut off frequency of 40Hz to reduce movement artifacts. The signals were then full wave rectified, and low pass filtered at 15Hz to produce an enveloped EMG signal. Peak and average enveloped EMG values were then normalized to the maximum observed signal across all treadmill positions for each muscle separately (36).

2.4 Statistical Analysis

Average values from the normalized peak and average enveloped EMG signals for each muscle from the 10 gait cycles at the 5 treadmill positions were tested for normality and found suitable for parametric analysis using Shapiro-Wilk tests. Repeated Measures ANOVA tests were performed to explore the effect of the 5 treadmill positions. When significant main effects were seen within a muscle, post hoc pairwise comparisons were performed to determine differences between 0%, 5% incline, 10% incline, 5% decline, and 10% decline positions using least significant difference tests. All statistical analyses were performed using SPSS version 29 (IBM, USA), and the alpha level was set at ($p < 0.05$).

3 Results

Data were collected from twelve dogs and EMG data from nine dogs were deemed acceptable due to signal dropout or technical difficulties. Our participants included 1 German Short-haired Pointer, 1 Wire Haired Pointer, 1 Doberman, 1 German Shepherd, 1 Dutch Shepherd, 1 Golden Retriever, and 6 Mixed Breed dogs between 21.4 – 35.2 kgs with body condition scores of 4-6/9, and ages 1-10 years old.

The EMG data from the right side versus left side showed no significant difference allowing for left and right measurements to be pooled. To limit the effect of skin motion artifact on the EMG signal, one author (JR) extensively experienced in analysing EMG data reviewed each trial for each dog to ensure the quality of data included for analysis. Trials with excessive motion artifact was excluded from final analysis. Therefore, of the possible 18 data sets obtained bilaterally from each muscle in 9 dogs, the final data set included 8 EMG observations from the triceps, 7 EMG observations from the biceps, 18 EMG observations from the deltoids and 14 EMG observations from the supraspinatus.

In this study, the peak muscle activity represented the highest amount of muscle activation, and the average muscle activity was an indication of total muscle activation during the gait cycle (33).

3.1 Supraspinatus muscle

Significant main effects were seen for average supraspinatus muscle activity ($p < 0.001$) (Table 1). Post hoc pairwise comparisons for average supraspinatus activity showed significant increases between 0% to 5% decline ($p = 0.019$, 14.5%), 0% to 10% decline ($p < 0.001$, 29.4%), 5% decline to 10% decline ($p < 0.001$, 13.0%), and significant decreases between 5% incline to 10% incline ($p = 0.023$, 10.3%), (Tables 2,3), (Figure 2). In addition, significant differences were seen for average supraspinatus activity between various treadmill positions including; 5% incline vs 10% decline ($p = 0.004$), 10% incline vs 5% decline ($p = 0.003$), 10% incline vs 10% decline ($p < 0.001$), (Table 2).

3.2 Scapular portion of deltoideus muscle

Significant main effects were seen for both average and peak deltoideus muscle activity ($p = 0.021$, $p < 0.001$) respectively (Table 1). Post hoc pairwise comparisons for peak deltoideus activity showed significant increases between 0% to 5% decline ($p = 0.045$, 21.0%), and 0% to 10% decline ($p = 0.016$, 27.2%), (Figure 3). For average deltoideus activity significant increases were seen between 0% to 5% decline ($p = 0.003$, 29.4%), and 0% to 10% decline ($p < 0.001$, 39.2%), (Tables 2, 3) (Figure 4). In addition, significant differences were seen for average deltoideus activity between various treadmill positions including 5% incline vs 10% decline ($p = 0.009$), 10% incline vs 5% decline ($p = 0.034$), 10% incline vs 10% decline ($p = 0.005$), (Table 2).

3.3 Lateral head of triceps brachii muscle

No significant differences were seen for peak or average triceps brachii muscle activity between treadmill positions (Table 1). However similar trends were seen in peak triceps brachii activity when compared to average changes in supraspinatus activity, with a 19.8% increase for the 5% to 10% decline walking, and a 17.9% decrease for the 5% to 10% incline walking, and for average triceps brachii activity with a 13.1% decrease for the 5% to 10% incline walking, and 6.4% increase for the 5% to 10% decline walking (Table 3).

3.4 Biceps brachii muscle

No significant differences were seen for peak or average biceps brachii muscle activity between treadmill positions (Table 1).

4 Discussion

Incremental effects of incline and decline treadmill walking have not previously been evaluated, and only a few studies (25-27) have evaluated muscle activity in the canine forelimb. Multiple surveys of sporting dog injuries have reported a higher incidence of injuries to the shoulder, highlighting the need for further studies evaluating the muscular response with common exercises used for rehabilitating shoulder injuries and return to sport conditioning programs (6-8). The supraspinatus and deltoideus were chosen based on their actions of flexion and extension of the shoulder, their role in shoulder joint stabilization, and the frequency of injury to the supraspinatus tendon in dogs (2, 37, 38). The biceps brachii and the triceps brachii were chosen because of their direct involvement with flexion and extension of both the shoulder and elbow, the incidence of biceps tendinopathy, and the triceps brachii being an anti-gravity muscle of the forelimb. The data from this study partially support the hypothesis of increased muscle activity with incremental decline walking, with differences seen in the deltoideus and supraspinatus muscles, but not in the lateral head of the triceps brachii or biceps brachii muscles. We also expected to see decreased muscle activity with incremental incline

walking; however, this was only seen in the average supraspinatus muscle activity between the 5% and 10% incline.

4.1 Supraspinatus Muscle Activity and Clinical Implications

The supraspinatus is active throughout the stance phase, stabilizing the shoulder joint for as long as the limb is load bearing (39). In this study, the supraspinatus showed significant increases in average muscle activity during decline walking. Throughout decline walking, the increased activity of the supraspinatus muscle may be attributed to the braking momentum of the body when going downhill, requiring greater muscle activation with steeper grades. This is consistent with the findings by Cullen et al (27), who found the highest supraspinatus activation when descending the A-frame. Of particular clinical interest were the incremental changes of the average supraspinatus activity. During decline walking, an additional 13.0% increase in muscle activity was observed at the 10% decline versus the 5% decline position. Conversely, during incline walking, the muscle activity of the supraspinatus decreased by an additional 10.3% at the 10% incline versus the 5% incline position. Direct comparisons of decline walking to incline walking, though statistically significant, did not offer any additional clinical relevance over the direct comparisons to 0 %.

These results reveal an ability to increase or decrease supraspinatus muscle activity via incremental changes of decline or incline positions respectively during walking. This information may be beneficial when considering the use of treadmill walking for rehabilitation following supraspinatus injuries. For example, a rehabilitation plan for a supraspinatus injury may start initially with walking on a 10% treadmill incline to decrease the supraspinatus activity, and as healing continues, slowly progress to level walking, followed by 5% decline and then 10% decline on a treadmill to systematically increase the load on the muscle. Adjustments to the treadmill position based on these findings may also be useful when designing patient protocols to address muscular strength training, pain and shoulder function during the recovery period.

4.2 Scapular Portion of Deltoideus Muscle Activity and Clinical Implications

The scapular portion of the deltoideus muscle is active during the second half of the stance phase where it supports retraction of the limb (39). In this study, the deltoideus muscle activity showed significant increases in both peak and average muscle activity during decline walking, with progressively increased activity as the decline becomes steeper, suggesting a role in braking while going down an incline. Peak deltoideus activity was notable with a 21.0% increase at 5% decline and a 27.2% increase at 10% decline. Average values were remarkable with a 29.4% increase at 5% decline and a 39.2% increase at 10% decline. However, unlike the supraspinatus, no differences were seen within the deltoideus during incline walking. Again, direct comparisons of decline walking to incline walking, though statistically significant, did not offer any additional clinical relevance over the direct comparisons to 0 %.

Although the deltoideus does not typically experience direct injury, as the primary flexor of the shoulder and a shared joint stabilizer, it may contribute to shoulder injuries if weak. Decline walking at both the 5% and 10% positions significantly increase the activity of the deltoideus muscle indicating greater peak activation and average muscle activity which could be considered clinically important to increase deltoideus strength. As the forelimb experiences braking forces in a retracted position during decline walking (40), it is plausible the notable increase in both peak and average muscle activity was due to the rise in vertical load and subsequent braking when walking downhill. As dogs redistribute their weight, the moment produced by the forelimb retractors attempts to shift

mass onto the hindlimbs and use the shoulders to control the descent despite the increased load (40). During uphill walking, the deltoideus helps to support the shoulder joint, contributing to overall stability and coordination of the body as the dog moves forward and up the incline. However, our data suggests that the deltoideus was not influenced during incline walking at 5% and 10% inclines. This is likely due to the decreased forelimb load and subsequent reduced propulsive forces influencing additional deltoideus muscle contraction during incline walking.

4.3 Lateral Head of Triceps Brachii Muscle Activity and Clinical Implications

In our study, there were no significant differences for peak or average muscle activity for the lateral head of triceps brachii between treadmill positions. However, the percentage differences between treadmill positions for triceps brachii followed a similar trend as the supraspinatus. The triceps brachii muscle activity increased between the 5% decline to 10% decline. Inversely, the muscle activity decreased between the 5% incline to 10% incline. However, these results were not statistically significant thus these findings should be interpreted with caution when considering clinical implementation.

The lateral head of the triceps brachii muscle is responsible for extension of the elbow joint and is important to counter gravity as well as stabilize the joint (39). It has been assumed that increased forelimb muscle contraction is required during decline walking because the dog braces and brakes when walking downhill, and thus it has been assumed that decline walking may be useful to strengthen the triceps brachii muscles (9). This study did not fully support this hypothesis. Triceps brachii muscle activity while climbing an incline could also be expected to be greater due to the increased elbow extension and subsequent propulsion during incline walking noted in the kinematic study by Carr et al (41). However, the degrees of incline incorporated in this study was closer to that reported in the kinematic study by Holler et al, which did not show a significant change in elbow extension while walking (42). Given the Carr et al study used a 70% grade, the Holler et al study used an 11% grade, and this study used a 10% grade, this might suggest that steeper incline and decline positions are needed to influence triceps brachii activity.

It is plausible that the smaller number of usable observations for the triceps brachii negatively impacted this analysis. Therefore, a greater sample size is required to explore the effect of incline and decline slopes on triceps brachii muscle activity to confirm that a change from the 0% to 10% decline does not significantly change activity levels. Additional studies evaluating EMG activity of the forelimb musculature are needed to definitively support or refute the results presented in this study with a focus on incline or decline walking and trotting. Future studies should aim to recruit a larger sample size to ensure adequate power. In addition, the consideration of EMG evaluation of the lateral and/or long head of the triceps brachii would allow for a more complete analysis of triceps brachii muscle activity.

4.4 Biceps Brachii Muscle Activity and Clinical Implications

In this study, there were no significant differences for peak or average biceps brachii activity between treadmill positions. As a shoulder extensor and stabilizer, it was anticipated the biceps brachii would experience increased load during decline walking resulting in increased muscle activity. Additionally, as a synergist for the triceps brachii with co-contraction of both muscles during the stance phase of gait (43), it would be assumed the biceps brachii would also experience a positive propulsive effect during incline walking. However, in this study, incline and decline positions of the land treadmill at the walk did not significantly influence biceps brachii muscle activity. This is consistent with the

findings from Cullen et al, where biceps brachii muscle activity was lower during descending A-frame (27). Clinically this may indicate that incline and decline walking of 10% or less may not assist in targeting and subsequently strengthening the biceps brachii muscle. This lack of muscle activation may suggest that incline and decline walking need not be excluded from exercise protocols for canine patients with injuries to the biceps brachii as the muscle activity does not appear to be significantly influenced by incline and decline walking.

However, as with the triceps brachii, it is plausible that the smaller number of usable observations for the biceps brachii negatively impacted this analysis. However, post hoc sample size calculations from the biceps brachii results, set to achieve 80% power and 5% level of significance, indicated that 271 and 68 EMG observations would be required to indicate a significant difference between 0% and 10% decline walking in peak and average biceps brachii muscle activity respectively. Again, these large sample sizes likely indicate that biceps brachii muscle activity does not significantly change from the 0% to 10% decline, despite the smaller sample size reported here.

Future research efforts should aim to recruit a larger sample size to ensure an adequate power analysis. Additionally, the consideration of fine-wire EMG over surface EMG for evaluation of the biceps brachii may result in a greater numbers of usable EMG observations.

4.5 Limitations

Despite following previously published methods, the signal obtained from the triceps brachii and the biceps brachii muscles did not yield consistently reliable EMG output in all dogs. This resulted in a decreased number of observations which could have affected the power of the analysis particularly on biceps brachii and triceps brachii. In addition the variability in the responses within each of the dogs may have masked the effects of the treadmill positions. Given the large required sample size calculated during post hoc power calculations for each muscle indicate there likely is no significant biologic change in muscle activity. Suspected reasons for decreased usable signal include movement artifact due to skin displacement. Additionally, muscle activity while walking on a treadmill may not be directly comparable to muscle activity during over ground walking. However, rehabilitation programs are commonly performed on treadmills making the findings of this study clinically relevant.

5 Conclusions

Given the prevalence and significance of canine shoulder injuries, specifically supraspinatus and biceps tendinopathies, this study showed implementing a 5% incline, 10% incline, 5% decline or 10% decline during walking could significantly impact shoulder muscle activity and therefore be used with confidence in customized progressive rehabilitation and conditioning programs. This study supports the use of decline walking to activate the supraspinatus and deltoideus muscles and provides new insights regarding the muscle activity of the thoracic limbs during various treadmill positions at the walk. Average EMG activity of the supraspinatus muscle significantly increased during decline walking and decreased during incline walking. The impact of incremental treadmill positions on the average supraspinatus activity, along with increases in the average and peak EMG activity of the deltoideus muscle during decline walking should be considered when developing a therapeutic exercise plan in canine patients with shoulder injuries.

6 Conflict of Interest

330 No conflict of interest to declare. The research was conducted in the absence of any commercial or
331 financial relationships that could be construed as a potential conflict of interest.

332 **7 Author Contributions**

333 RC, KS, JR, DL, and TU contributed to the conception, design, data acquisition, data analysis, and
334 writing and editing of the manuscript.

335 DM and MD contributed to the conception, design, and writing and editing of the manuscript.

336 **8 Funding**

337 Details of all funding sources should be provided, including grant numbers if applicable. Please
338 ensure to add all necessary funding information, as after publication this is no longer possible.

339 **9 Acknowledgments**

340 The authors would like to acknowledge the contributions of Dawn Rector and Christina Woods.
341 Special thanks to The Veterinary Orthopedic Laboratory (VOL) which supports research using a
342 variety of gait analysis tools for studies in the areas of physical rehabilitation and exercise in dogs as
343 well as the UTCVM equine biomechanics research lab.

344 **10 References**

- 345 1. Henderson AL, Latimer C, Millis DL. Rehabilitation and physical therapy for selected
346 orthopedic conditions in veterinary patients. *The Veterinary clinics of North America Small animal*
347 *practice*. 2015;45(1):91-121.
- 348 2. Marcellin-Little DJ, Levine D, Canapp SO, Jr. The canine shoulder: selected disorders and
349 their management with physical therapy. *Clinical techniques in small animal practice*.
350 2007;22(4):171-82.
- 351 3. Stokes R, Dycus D. The Shoulder Joint and Common Abnormalities. *The Veterinary clinics*
352 *of North America Small animal practice*. 2021;51(2):323-41.
- 353 4. von Pfeil DJ, Liska WD, Nelson S, Jr., Mann S, Wakshlag JJ. A survey on orthopedic injuries
354 during a marathon sled dog race. *Veterinary medicine (Auckland, NZ)*. 2015;6:329-39.
- 355 5. Brunke MW, Levine D, Marcellin-Little DJ, Oliver KE, Barnhard JA, Tringali AA.
356 *Musculoskeletal Problems in Sporting Dogs. Advances in small animal care*. 2023;4(1):53-60.
- 357 6. Pechette Markley A, Shoben AB, Kieves NR. Internet-based survey of the frequency and
358 types of orthopedic conditions and injuries experienced by dogs competing in agility. *Journal of the*
359 *American Veterinary Medical Association*. 2021;259(9):1001-8.
- 360 7. Blake S, Melfi V, Tabor G, Wills AP. Frequency of injuries and orthopaedic conditions
361 sustained by flyball dogs. *Comparative exercise physiology*. 2024;20(1):89-96.
- 362 8. Lafuente P, Whyte C. A Retrospective Survey of Injuries Occurring in Dogs and Handlers
363 Participating in Canicross. *Vet Comp Orthop Traumatol*. 2018;31(5):332-8.

- 364 9. Millis DL, Levine D. Canine rehabilitation and physical therapy. Second edition. ed.
365 Philadelphia, PA: Elsevier; 2014. xvi, 760 pages p.
- 366 10. Dycus DL, Levine D, Ratsch BE, Marcellin-Little DJ. Physical Rehabilitation for the
367 Management of Canine Hip Dysplasia: 2021 Update. The Veterinary clinics of North America Small
368 animal practice. 2022;52(3):719-47.
- 369 11. Akaraphutiporn E, Kwananocha I, Meechai C, Suksomboonwong P, Aramsriprasert S,
370 Meethong O, et al. Comparative analysis of post-operative rehabilitation approaches for medial
371 patellar luxation in small-breed dogs. Vet World. 2024;17(3):550-7.
- 372 12. Brantberg I, Grooten WJA, Essner A. The Effect of Therapeutic Exercise on Body Weight
373 Distribution, Balance, and Stifle Function in Dogs following Stifle Injury. Animals (Basel).
374 2023;14(1).
- 375 13. Lane DM, Pfeil DV, Kowaleski MP. Synthesis of surgeon and rehabilitation therapist
376 treatment methods of bicapital tenosynovitis in dogs allows development of an initial consensus
377 therapeutic protocol. Journal of the American Veterinary Medical Association. 2024;262(2):1-8.
- 378 14. Jeong IS, Piao Z, Rahman MM, Kim S, Kim NS. Canine thoracolumbar intervertebral disk
379 herniation and rehabilitation therapy after surgical decompression: A retrospective study. J Adv Vet
380 Anim Res. 2019;6(3):394-402.
- 381 15. Hodgson MM, Bevan JM, Evans RB, Johnson TI. Influence of in-house rehabilitation on the
382 postoperative outcome of dogs with intervertebral disk herniation. Vet Surg. 2017;46(4):566-73.
- 383 16. Gustås P, Pettersson K, Honkavaara S, Lagerstedt AS, Byström A. Kinematic and
384 temporospatial assessment of habituation of Labrador retrievers to treadmill trotting. Vet J. 2013;198
385 Suppl 1:e114-9.
- 386 17. Bockstahler BA, Prickler B, Lewy E, Holler PJ, Vobornik A, Peham C. Hind limb kinematics
387 during therapeutic exercises in dogs with osteoarthritis of the hip joints. American journal of
388 veterinary research. 2012;73(9):1371-6.
- 389 18. Fischer S, Nolte I, Schilling N. Adaptations in muscle activity to induced, short-term
390 hindlimb lameness in trotting dogs. PloS one. 2013;8(11):e80987.
- 391 19. Vilar JM, Miró F, Rivero MA, Spinella G. Biomechanics. Biomed Res Int.
392 2013;2013:271543.
- 393 20. Lauer SK, Hillman RB, Li L, Hosgood GL. Effects of treadmill inclination on
394 electromyographic activity and hind limb kinematics in healthy hounds at a walk. American journal
395 of veterinary research. 2009;70(5):658-64.
- 396 21. Miró F, Galisteo AM, Garrido-Castro JL, Vivo J. Surface Electromyography of the
397 Longissimus and Gluteus Medius Muscles in Greyhounds Walking and Trotting on Ground Flat, Up,
398 and Downhill. Animals (Basel). 2020;10(6).

- 399 22. Bockstahler BB, Gesky R, Mueller M, Thalhammer JG, Peham C, Podbregar I. Correlation of
400 surface electromyography of the vastus lateralis muscle in dogs at a walk with joint kinematics and
401 ground reaction forces. *Vet Surg.* 2009;38(6):754-61.
- 402 23. Breitfuss K, Franz M, Peham C, Bockstahler B. Surface Electromyography of the Vastus
403 Lateralis, Biceps Femoris, and Gluteus Medius Muscle in Sound Dogs During Walking and Specific
404 Physiotherapeutic Exercises. *Vet Surg.* 2015;44(5):588-95.
- 405 24. McLean H, Millis D, Levine D. Surface Electromyography of the Vastus Lateralis, Biceps
406 Femoris, and Gluteus Medius in Dogs During Stance, Walking, Trotting, and Selected Therapeutic
407 Exercises. *Front Vet Sci.* 2019;6:211.
- 408 25. Garcia TC, Sturges BK, Stover SM, Aoki K, Liang JM, Reinhardt KB, et al. Forelimb
409 brachial muscle activation patterns using surface electromyography and their relationship to
410 kinematics in normal dogs walking and trotting. *Comparative exercise physiology.* 2014;10(1):13-22.
- 411 26. DL JKM, editor *Surface Electromyography of the Biceps Brachii and Lateral Triceps*
412 *Muscles in Dogs during Walking, Trotting, and Selected Therapeutic Exercises.* *Vet Comp Orthop*
413 *Traumatol;* 2020; Stuttgart · New York: Thieme; 2020.
- 414 27. Cullen KL, Dickey JP, Brown SH, Nykamp SG, Bent LR, Thomason JJ, et al. The magnitude
415 of muscular activation of four canine forelimb muscles in dogs performing two agility-specific tasks.
416 *BMC veterinary research.* 2017;13(1):68.
- 417 28. Cullen KL, Dickey JP, Brown SHM, Nykamp SG, Bent LR, Thomason JJ, et al. A validated
418 approach for collecting fine-wire electromyographic recordings in four canine shoulder muscles
419 during highly dynamic tasks. *Comparative exercise physiology.* 2015;11(2):65-74.
- 420 29. Zhang X, Jenkins GJ, Hakim CH, Duan D, Yao G. Four-limb wireless IMU sensor system for
421 automatic gait detection in canines. *Sci Rep.* 2022;12(1):4788.
- 422 30. Duerr FM, Pauls A, Kawcak C, Haussler K, Bertocci G, Moorman V, et al. Evaluation of
423 inertial measurement units as a novel method for kinematic gait evaluation in dogs. *Vet Comp*
424 *Orthop Traumatol.* 2016;29(6):475-83.
- 425 31. Jenkins GJ, Hakim CH, Yang NN, Yao G, Duan D. Automatic characterization of stride
426 parameters in canines with a single wearable inertial sensor. *PloS one.* 2018;13(6):e0198893.
- 427 32. Vitt MA, Rendahl A, Pracht SE, Knotek BM, Lascelles BDX, Gordon-Evans W, et al. Nine-
428 axis inertial measurement unit output discriminates activities of varying intensity in the dog.
429 *American journal of veterinary research.* 2023;84(3).
- 430 33. Ursini T, Shaw K, Levine D, Richards J, Adair HS. Electromyography of the Multifidus
431 Muscle in Horses Trotting During Therapeutic Exercises. *Front Vet Sci.* 2022;9:844776.
- 432 34. Owen MR, Richards J, Clements DN, Drew ST, Bennett D, Carmichael S. Kinematics of the
433 elbow and stifle joints in greyhounds during treadmill trotting -- an investigation of familiarisation.
434 *Veterinary and comparative orthopaedics and traumatology.* 2004;17(3):141-5.

35. Fanchon L, Grandjean D. Habituation of healthy dogs to treadmill trotting: repeatability assessment of vertical ground reaction force. *Research in veterinary science*. 2009;87(1):135-9.
36. St. George L, Roy SH, Richards J, Sinclair J, Hobbs SJ. Surface EMG signal normalisation and filtering improves sensitivity of equine gait analysis. *Comparative exercise physiology*. 2019;15(3):173-86.
37. Piras LA, Olimpo M, Lafuente P, Tomba A, Del Magno S, Lardone E, et al. Surgical Treatment of Nonmineralized Supraspinatus Tendinopathy in Dogs: A Retrospective Long-Term Follow-Up. *Animals (Basel)*. 2023;13(4).
38. Leeman JJ, Shaw KK, Mison MB, Perry JA, Carr A, Shultz R. Extracorporeal shockwave therapy and therapeutic exercise for supraspinatus and biceps tendinopathies in 29 dogs. *Vet Rec*. 2016;179(15):385.
39. Fischer MS, Lilje KE, Lauströer J, Andikfar A. *Dogs in motion*. 2nd edition ed. Dortmund: VDH Service; 2014.
40. Granatosky MC, Bryce CM, Hanna J, Fitzsimons A, Laird MF, Stilson K, et al. Inter-stride variability triggers gait transitions in mammals and birds. *Proc Biol Sci*. 2018;285(1893):20181766.
41. Carr JG, Millis DL, Weng HY. Exercises in canine physical rehabilitation: range of motion of the forelimb during stair and ramp ascent. *Journal of small animal practice*. 2013;54(8):409-13.
42. Holler PJ, Brazda V, Dal-Bianco B, Lewy E, Mueller MC, Peham C, et al. Kinematic motion analysis of the joints of the forelimbs and hind limbs of dogs during walking exercise regimens. *American journal of veterinary research*. 2010;71(7):734-40.
43. Tokuriki M. Electromyographic and joint-mechanical studies in quadrupedal locomotion. I. Walk. *Nihon Juigaku Zasshi*. 1973;35(5):433-6.

11 Supplementary Material

Supplementary Material should be uploaded separately on submission, if there are Supplementary Figures, please include the caption in the same file as the figure. Supplementary Material templates can be found in the Frontiers Word Templates file.

Please see the [Supplementary Material](#) section of the [Author guidelines](#) for details on the different file types accepted.

12 Data Availability Statement

The datasets [GENERATED/ANALYZED] for this study can be found in the [NAME OF REPOSITORY] [LINK]. Please see the “Availability of data” section of [Materials and data policies in the Author guidelines](#) for more details.

Table 1: Mean and standard deviation (sd) for Average and Peak EMG normalised to the maximum observed signal with main effects (p value) and effects sizes (np2) between treadmill positions.

	Level Mean (sd)	5% Incline Mean (sd)	10% Incline Mean (sd)	5% Decline Mean (sd)	10% Decline Mean (sd)	p value (np²)
Muscle Activity						
Peak Triceps	0.348 (0.122)	0.387 (0.159)	0.318 (0.147)	0.315 (0.112)	0.378 (0.054)	0.343 (0.14)
Average Triceps	0.158 (0.043)	0.177 (0.120)	0.154 (0.087)	0.157 (0.092)	0.167 (0.073)	0.761 (0.06)
Peak Biceps	0.337 (0.088)	0.347 (0.080)	0.376 (0.104)	0.314 (0.082)	0.357 (0.078)	0.246 (0.19)
Average Biceps	0.179 (0.042)	0.179 (0.038)	0.178 (0.043)	0.185 (0.042)	0.198 (0.037)	0.228 (0.20)
Peak Deltoid	0.288 (0.099)	0.299 (0.097)	0.297 (0.010)	0.348 (0.094)	0.366 (0.103)	0.021 (0.15)
Average Deltoid	0.126 (0.030)	0.137 (0.040)	0.131 (0.039)	0.163 (0.050)	0.176 (0.049)	<0.001 (0.32)
Peak Supraspinatus	0.497 (0.090)	0.509 (0.109)	0.489 (0.093)	0.504 (0.113)	0.542 (0.131)	0.057 (0.16)
Average Supraspinatus	0.214 (0.047)	0.224 (0.049)	0.201 (0.034)	0.245 (0.048)	0.277 (0.061)	<0.001 (0.51)

471

472 Table 2: Pairwise comparisons for Average and Peak EMG where significant main effects were seen
473 between positions.

	Mean Differenc e	p value	Lower Bound	Upper Bound
Maximum Deltoid Activity				
Level v 5 degrees incline	-0.011	0.681	-0.067	0.045
Level v 10 degrees incline	-0.009	0.727	-0.065	0.047
Level v 5 degrees decline	-0.060	0.045	-0.12	-0.001
Level v 10 degrees decline	-0.079	0.016	-0.14	-0.017
5 degrees incline v 10 degrees incline	0.002	0.916	-0.031	0.035
5 degrees incline v 5 degrees decline	-0.049	0.111	-0.111	0.013

5 degrees incline v 10 degrees decline	-0.068	0.062	-0.139	0.004
10 degrees incline v 5 degrees decline	-0.051	0.114	-0.116	0.014
10 degrees incline v 10 degrees decline	-0.069	0.068	-0.144	0.006
5 degrees decline v 10 degrees decline	-0.018	0.428	-0.065	0.029
Average Deltoid Activity				
Level v 5 degrees incline	-0.010	0.291	-0.030	0.010
Level v 10 degrees incline	-0.005	0.540	-0.022	0.012
Level v 5 degrees decline	-0.037	0.003	-0.06	-0.014
Level v10 degrees decline	-0.050	<0.001	-0.072	-0.027
5 degrees incline v 10 degrees incline	0.005	0.376	-0.007	0.018
5 degrees incline v 5 degrees decline	-0.027	0.055	-0.054	0.001
5 degrees incline v 10 degrees decline	-0.039	0.009	-0.067	-0.011
10 degrees incline v 5 degrees decline	-0.032	0.034	-0.061	-0.003
10 degrees incline v 10 degrees decline	-0.045	0.005	-0.074	-0.016
5 degrees decline v 10 degrees decline	-0.013	0.110	-0.028	0.003
Average Supraspinatus Activity				
Level v 5 degrees incline	-0.010	0.287	-0.03	0.010
Level v 10 degrees incline	0.013	0.143	-0.005	0.031
Level v 5 degrees decline	-0.031	0.019	-0.056	-0.006
Level v10 degrees decline	-0.063	<0.001	-0.090	-0.036
5 degrees incline v 10 degrees incline	0.023	0.023	0.004	0.042
5 degrees incline v 5 degrees decline	-0.021	0.096	-0.046	0.004
5 degrees incline v 10 degrees decline	-0.053	0.004	-0.085	-0.020

10 degrees incline v 5 degrees decline	-0.044	0.003	-0.070	-0.018
10 degrees incline v 10 degrees decline	-0.076	<0.001	-0.106	-0.046
5 degrees decline v 10 degrees decline	-0.032	<0.001	-0.048	-0.016

Table 3: Percentage differences between positions (negative indicates second value greater than first). Bold indicates values of noteworthiness.

	% difference					
	Level - 5 Incline	Level - 10 Incline	Level – 5 Decline	Level – 10 Decline	5 incline – 10 incline	5 decline -10 decline
Muscle Activity						
Peak Triceps	11.23	-8.62	-9.34	8.62	-17.85	19.81
Average Triceps	11.87	-2.77	-0.82	5.49	-13.09	6.36
Peak Biceps	2.96	11.31	-7.05	5.77	8.11	13.80
Average Biceps	0.27	-0.44	3.41	10.57	-0.72	6.92
Peak Deltoid	3.85	3.26	21.02	27.32	-0.56	5.19
Average Deltoid	8.22	4.03	29.35	39.24	-3.87	7.64
Peak Supra	2.45	-1.61	1.40	9.22	-3.96	7.70
Average Supra	4.71	-6.02	14.47	29.37	-10.25	13.01

Figure Captions

Figure 1: Left lateral view of one study subject showing the experimental set up of the surface and fine wire electromyographic sensors.

487 Figure 2: Pairwise comparisons for Average Supraspinatus electromyography where significant main
488 effects were seen between positions.

489 Figure 3: Pairwise comparisons for Peak Deltoid electromyography where significant main effects
490 were seen between positions.

491 Figure 4: Pairwise comparisons for Average Deltoid electromyography where significant main
492 effects were seen between positions.

493